

Telitacicept for systemic lupus erythematosus with post-surgical papillary thyroid carcinoma: A case report

JINHUI TAN¹, HAI HUANG², LINGHUA TAN³ and BO LI¹

¹Department of Rheumatology and Immunology, People's Hospital of Longhua, Shenzhen, Guangdong 518109, P.R. China;

²Department of Health Management, People's Hospital of Longhua, Shenzhen, Guangdong 518109, P.R. China;

³Department of Health Management, Jiangmen Wuyi Hospital of Chinese Medicine, Jiangmen, Guangdong 529099, P.R. China

Received July 24, 2024; Accepted November 5, 2024

DOI: 10.3892/br.2025.1926

Abstract. Systemic lupus erythematosus (SLE) is a chronic autoimmune disease with a complex etiology primarily linked to abnormalities in B lymphocytes within the human body, resulting in the production of numerous pathogenic autoantibodies. Telitacicept is a relatively novel humanized, recombinant transmembrane activator, calcium modulator and cyclophilin ligand interactor fused with the Fc portion (TACI-Fc). It works by competitively inhibiting the TACI site, neutralizing the activity of B-cell lymphocyte stimulator and A proliferation-inducing ligand. This, in turn, inhibits the development and survival of plasma cells and mature B cells. A 28-year-old female was admitted to the Department of Rheumatology and Immunology (People's Hospital of Longhua; Shenzhen, China) in June 2021 due to systemic edema for more than a month and hair loss lasting for a week. After comprehensive examination, the patient was diagnosed with SLE with hematological system involvement, serositis, lupus nephritis and secondary antiphospholipid syndrome. After receiving medications including glucocorticoids, mycophenolate mofetil and cyclosporine, the patient's white blood cells, platelets, hemoglobin, urinary protein and multiple serositis returned to normal. However, the levels of complement 3 (C3) and C4 did not significantly improve. Subsequently, the patient underwent thyroid ultrasound examination, which suggested thyroid nodules. After thyroid puncture biopsy, the patient was diagnosed with papillary thyroid carcinoma (PTC). After surgical resection, the patient was confirmed to have PTC by pathological biopsy, with no lymph node metastasis. At two

months after surgery, the patient was treated with telitacicept, and the complement levels not only returned to normal but also remained stable for a long time. The present case was the first to report the use of telitacicept for the successful treatment of a patient with SLE with post-surgical PTC, providing a potential therapeutic option for SLE with a prior history of carcinoma. The role of telitacicept in this field requires further research and attention.

Introduction

Systemic lupus erythematosus (SLE) is an autoimmune disorder that features the generation of numerous autoantibodies and immune complexes, potentially compromising diverse bodily systems. B cells are purportedly pivotal in the etiology of SLE (1), since the abnormal proliferation, differentiation and maturation of these cells lead to the secretion of autoantibodies that inflict harm on various bodily systems. The cytokines B-lymphocyte stimulator (BLyS) and A proliferation-inducing ligand (APRIL) occupy pivotal positions in the intricate process of B-cell maturation and differentiation (2).

Telitacicept is an important biological agent that can effectively suppress the proliferation and differentiation of B cells and plasma cells, by specifically targeting both BLyS and APRIL. Furthermore, it exhibits the potential to dampen the activity of long-lived plasma cells (3). Telitacicept was approved in March 2021 in China for the treatment of SLE (4), but it has not been widely accepted due to its high cost.

Thyroid carcinoma is the most common malignancy of the endocrine system, exhibiting a gradual increase in its incidence rate in high-income countries to certain middle-income countries. Globally, in 2020, the estimated number of new cases of thyroid cancer was ~449,000 in women and ~137,000 in men, corresponding to age-standardised incidence rates of ~10.1 per 100,000 women and ~3.1 per 100,000 men. In each sex, incidence rates are five times greater in high and very high Human Development Index countries than in low and medium Human Development Index countries (5). Previous meta-analyses and observational studies have shown that the risk of thyroid carcinoma is increased in patients with SLE (6-8). The present case documents a 28-year-old female patient who developed papillary thyroid carcinoma (PTC) 1 year after being diagnosed with SLE. Following surgical removal of the tumor, the

Correspondence to: Dr Bo Li or Mr. JinHui Tan, Department of Rheumatology and Immunology, People's Hospital of Longhua, 38 Jinglong Construction Road, Shenzhen, Guangdong 518109, P.R. China

E-mail: friendlybo@163.com

E-mail: 15015093514@163.com

Key words: systemic lupus erythematosus, telitacicept, a proliferation-inducing ligand, B-cell lymphocyte stimulator, papillary thyroid carcinoma

patient was administered with telitacicept for treatment. To the best of our knowledge, the present case was the first reporting the instance of a patient suffering from both SLE and PTC undergoing postoperative telitacicept treatment.

Case report

The patient, a 28-year-old female, was diagnosed with SLE at Department of Rheumatology and Immunology, People's Hospital of Longhua (Shenzhen, China) in June 2021, having experienced systemic edema for >1 month and hair loss for a week. After undergoing comprehensive examinations, the patient was diagnosed with lupus-related hematological damage, which was evidenced by leukopenia, thrombocytopenia, autoimmune hemolytic anemia and polyserositis accompanied by pericardial and pleural effusion. Additional complications included lupus nephritis and secondary antiphospholipid syndrome. After treatment with high-dose steroids, mycophenolate mofetil (MMF) and cyclosporine A (CsA) (Fig. 1), the patient's white blood cells, platelets, hemoglobin, urinary protein and polyserositis normalized (Fig. 2). Complement 3 (C3) levels improved but could not be maintained within the normal range for a prolonged period, whilst C4 levels remained unchanged (Fig. 1). During the course of treatment (Fig. 1), the patient developed herpes zoster, which improved after reducing the dosage of MMF and CsA and administering antiviral treatment with famciclovir. The dosage of MMF was adjusted from twice a day, 0.75 g each time to once a day, 0.5 g. The dosage of CsA has also been adjusted from twice a day (bid), 0.05 g each time to once a day (qd), 0.05 g. Famciclovir is administered three times a day, with 0.25 grams per administration.

On December, 2022, a thyroid ultrasound (Fig. S1) revealed a solid nodule near the isthmus of the left lobe of the thyroid gland, classified as American College of Radiology Thyroid Imaging Reporting and Data System (ACR TI-RADS) (9) grade 4, along with a mixed cystic-solid nodule in the right lobe, classified as ACR TI-RADS grade 2. Thyroid puncture pathology confirmed the diagnosis of PTC in the left lobe (data not shown). In May 2023, a total left thyroidectomy, isthmectomy, subtotal right thyroidectomy and lymph node dissection in the central region of the left neck were performed at Department of Thyroid and Breast Surgery, People's Hospital of Longhua (Shenzhen, China). The specimens were sectioned into book-leaf-like shapes every 3 mm, and all suspected lesions visible to the naked eye were sampled. All specimens were fixed with 4% neutral formaldehyde, routinely embedded in paraffin, sliced at 4 μ m, stained with H&E according to standard procedures and observed under a light microscope. Fig. 3A shows that the thyroid follicular epithelium exhibits papillary and branching proliferation, featuring a fibrovascular core. The surface is lined with a single layer of tumor cells. The cell nuclei are large, oval, densely packed and exhibit a ground-glass appearance. Nuclear grooves and small clear nucleoli can be seen (red rectangular shape). Left thyroid and isthmus are consistent with PTC. The final diagnosis was small PTC in the left lobe (pT1N0M0, stage I; Fig. 3) (10). Postoperative blood examination revealed a significant decrease in thyroid-stimulating hormone levels, therefore, oral administration of levothyroxine sodium tablets at a dose of

100 μ g qd was required. Owing to the failure of conventional drug therapy to successfully restore the patient's complement levels, the treatment plan has been adjusted accordingly since July 2023. Next, CsA was discontinued, and treatment with prednisone at a dosage of 10 mg qd, MMF at 0.75 g bid and hydroxychloroquine (HCQ) at 0.2 g qd was continued. In addition, the patient also received a subcutaneous injection of 160 mg of telitacicept once a week (4). After 1 month, the C3 and C4 levels returned to normal. Due to the recovery of the patient's C3 and C4 levels and the relatively high price of telitacicept, the dosage of telitacicept for subcutaneous injection was reduced to 80 mg once a week starting from September 2023 (Fig. 1).

During the medication period, regular follow-up visits were made to the Department of Thyroid and Breast Surgery, People's Hospital of Longhua (Shenzhen, China), where thyroid function and ultrasound examinations were conducted at half-year intervals. Thyroid stimulating hormone gradually normalized (data not shown) and no thyroid nodules or enlarged lymph nodes were observed.

In December 2023, there was one episode of cough, nasal congestion and yellow nasal discharge, suggestive of acute upper respiratory tract infection. Cefuroxime was administered twice daily, with 0.5 g each time, and compound pseudoephedrine hydrochloride was administered twice daily, with one tablet each time. After administering both orally for 3 days, the aforementioned symptoms improved. In late December 2023, a urine nitrite test yielded positive results (data not shown), but there were no symptoms of frequent urination, urgency or pain. A complete urine culture test indicated the presence of *Klebsiella pneumoniae* (data not shown). After taking levofloxacin tablets for 14 days, once a day at a dosage of 0.5 g each time, the urine nitrite test became negative.

Discussion

The reasons for the comorbidity of SLE and thyroid carcinoma remain elusive. Antonelli *et al* (11) previously found that the incidence of thyroid carcinoma in patients with SLE is significantly elevated compared with that in age-matched normal controls, particularly in those with evidence of thyroid autoimmunity. In another study, Okosieme *et al* (12) discovered that patients with autoimmune thyroid disease and thyroid carcinoma exhibit similar yet distinct thyroid globulin antigen epitopes compared with those with positive anti-thyroid globulin antibodies, suggesting that autoimmune mechanisms may serve a role in the development of thyroid carcinoma. In addition, SLE and thyroid carcinoma are more prevalent in women. Previous studies indicate a significant increase in estrogen receptor expression within thyroid carcinoma tissues and in the peripheral blood of patients with SLE (13-15). Estrogen may exert a significant role in the pathogenesis of both conditions, though the exact mechanism remains elusive. Therefore, further research is necessary to determine whether there is an association between the comorbidity of these two conditions.

Although inheritance, hormones and the environment have been identified as risk factors for developing SLE (16,17), abnormal activation of the immune system is the core mechanism, characterized by excessive B-cell responses and the




Figure 1. Treatment medication process and laboratory indicators. GCs, glucocorticoids; HCQ, hydroxychloroquine; MMF, mycophenolate mofetil; CsA, cyclosporine A; C3, complement 3; C4, complement 4; bid, twice daily; qd, once a day; qw, once a week.




Figure 2. Trends in levels of PLT (black line), Hb (red line) and WBC (blue line). The reference values for PLT, Hb and WBC are 125 to 350 $\times 10^9/l$, 115 to 150 g/l and 3.5 to 9.5 $\times 10^9/l$, respectively. PLT, platelets; Hb, hemoglobin; WBC, white blood cells.

production of autoantibodies (18). Telitacicept is a TACI-Fc fusion protein that, by competitively inhibiting the TACI site, neutralizes the activity of BLyS and APRIL, thereby inhibiting the development and survival of plasma cells and mature B cells. It not only inhibits the transformation of immature B lymphocytes into mature B lymphocytes, but can also prevent the transition of mature B cells into plasma cells. In addition, it can affect the secretion of autoantibodies from abnormal plasma cells, thereby controlling disease progression (19). BLyS binds to B-cell activating factor receptor 3 and functions through the alternative/noncanonical NF- κ B signaling

pathway, effectively blocking its activity (3). The application of telitacicept in the treatment of SLE has resulted in a higher response rate of the Systemic Lupus Erythematosus Disease Activity Index, significantly reducing immunoglobulin levels and increasing complement levels (20).

To date, only two biological agents have been approved for the treatment of SLE, namely belimumab (21) and telitacicept (4). There have been multiple reports (20,22-28) whereby patients with SLE without a history of carcinoma have significantly elevated complement levels after treatment with telitacicept. The present case showed a significant increase in complement levels after the addition of telitacicept, similar to the treatment effects observed in previously reported cases of patients with SLE without concurrent cancer treated with telitacicept (20,22-28).

To the best of our knowledge, no previous study has documented the use of these agents in patients with a history of carcinoma who also suffer from SLE. However, biological agents used to treat other rheumatic diseases, such as rheumatoid arthritis (RA), psoriasis (PS), inflammatory bowel disease (IBD) and ankylosing spondylitis (AS) have been marketed earlier (29). In addition, there have been reports on the use of biological agents in patients with a history of carcinoma who also suffer from these rheumatic diseases. A Danish population-based cohort study previously showed that among patients with RA with a history of carcinoma, treatment with biological disease-modifying antirheumatic drugs was not associated with an increased risk of a second malignant neoplasm (30). Furthermore, two large prospective studies, one originating from the British Society for Rheumatology Biologics (31) and the other from the German RA Observation of Biologic




Figure 3. Thyroid pathology results. (A) It is evident that the thyroid follicular epithelium exhibits papillary and branching proliferation, featuring a fibrovascular core. The surface is lined with a single layer of tumor cells. The cell nuclei are large, oval, densely packed and exhibit a ground-glass appearance. Nuclear grooves and small clear nucleoli can be seen (red rectangular shape). Left thyroid and isthmus are consistent with papillary thyroid carcinoma. (B) In the left thyroid and isthmus, calcification (yellow ellipses) and punctured lesion tissue (red ellipses) are visible. One lymph node is present in the capsular region, without evidence of carcinoma metastasis (0/1). (C) Examination of lymph nodes in the central region of the left neck revealed five lymph nodes, all without metastasis (0/5) (hematoxylin and eosin staining; magnification, x50).

Therapy registry (32), have been conducted to investigate the utilization of biological drugs in patients diagnosed with RA. However, neither studies reported any increases in carcinoma recurrence in patients with a previous history of malignancy and under biological treatments. Another recent study has suggested that the administration of biologics targeting TNF- α , IL-17, IL-23 and IL-12 for the treatment of PS in patients with a prior history of carcinoma appears to be a safe approach (33). A multicenter real-life experience added evidence to the safety of secukinumab in patients with PS with a personal history of carcinoma (34). Secukinumab, as the world's first approved fully human IL-17A inhibitor, can specifically neutralize IL-17A from multiple sources and inhibit its pro-inflammatory activity (35). Initially, secukinumab was approved by the United States Food and Drug Administration (FDA) for the treatment of patients with moderate to severe plaque psoriasis, AS, and active psoriatic arthritis (PsA) (36). A previous meta-analysis included 34 studies (17 IBD, 14 RA, 2 PS and 1 AS), consisting of 24,328 patients and 85,784 person-years of follow-up. Rates of carcinoma recurrence were found to be similar among individuals not on immunosuppression, receiving anti-TNFs, immunomodulators and combination immunosuppression (37).

Whilst several existing guidelines recommend avoiding immunosuppression for 5 years after index carcinoma (38-42), a number of studies have indicated it may be safe to initiate these agents earlier than 5 years in certain patients (37,43). The majority of thyroid carcinoma mortality cases are from the non-papillary histological subtypes, whereas PTC has a 5-year survival rate of >90% (44). In addition, thyroid carcinoma deaths have fallen by 50% in women and by 33% in men over the past 40 years (45). Considering the present SLE case with hypocomplementemia and PTC who underwent surgery, after the treatment with MMF and CsA failed to achieve satisfactory results, the patient was effectively treated with telitacicept. During the postoperative follow-up, no tumor recurrence or lymph node metastasis was observed. Furthermore, the safety data derived from patients suffering from RA, PS, IBD and AS cannot be universally applied to patients with SLE due to various reasons, primarily because each disease exhibits a unique pathogenic mechanism.

In conclusion, to the best of our knowledge, the present case was the first to report the use of telitacicept for the successful treatment of a patient with SLE with post-surgical PTC,

providing a potential therapeutic option for SLE with a prior history of carcinoma. However, the present case is only based on a single case and therefore the conclusion drawn may not be broadly representative and cannot be directly generalized to a broader patient population. Therefore, further research, particularly prospective randomized controlled trials, are warranted to verify the present findings and ensure patient safety.

Acknowledgements

Not applicable.

Funding

The present report was supported by the Scientific Research Projects of Medical and Health Institutions of Longhua District, Shenzhen (grant no. 2022046).

Availability of data and materials

The data generated in the present study may be requested from the corresponding author.

Authors' contributions

JT and BL designed and supervised the study. JT, HH and LT performed data collection and analysis. HH and LT performed data analysis and interpretation. All authors confirm the authenticity of all the raw data. All authors have read and approved the final version of the manuscript.

Ethics approval and consent to participate

The present study was approved by the Ethics Committee of People's Hospital of Longhua (Shenzhen, China; approval no. 059A[2024]). The patient provided written informed consent for the case study to be published.

Patient consent for publication

The patient who participated in the study provided written informed consent for the publication of any associated data and images.

Competing interests

The authors declare that they have no competing interests.

References

- Ma K, Du W, Wang X, Yuan S, Cai X, Liu D, Li J and Lu L: Multiple functions of B cells in the pathogenesis of systemic lupus erythematosus. *Int J Mol Sci* 20: 6021, 2019.
- Möckel T, Basta F, Weinmann-Menke J and Schwarting A: B cell activating factor (BAFF): Structure, functions, autoimmunity and clinical implications in Systemic Lupus Erythematosus (SLE). *Autoimmun Rev* 20: 102736, 2021.
- Shi F, Xue R, Zhou X, Shen P, Wang S and Yang Y: Telitacept as a BLyS/APRIL dual inhibitor for autoimmune disease. *Immunopharmacol Immunotoxicol* 43: 666-673, 2021.
- Dhillon S: Telitacept: First approval. *Drugs* 81: 1671-1675, 2021.
- Pizzato M, Li M, Vignat J, Laversanne M, Singh D, La Vecchia C and Vaccarella S: The epidemiological landscape of thyroid cancer worldwide: GLOBOCAN estimates for incidence and mortality rates in 2020. *Lancet Diabetes Endocrinol* 10: 264-272, 2022.
- Zhang M, Wang Y, Wang Y, Bai Y and Gu D: Association between systemic lupus erythematosus and cancer morbidity and mortality: Findings from cohort studies. *Front Oncol* 12: 860794, 2022.
- Song L, Wang Y, Zhang J, Song N, Xu X and Lu Y: The risks of cancer development in systemic lupus erythematosus (SLE) patients: A systematic review and meta-analysis. *Arthritis Res Ther* 20: 270, 2018.
- Bernatsky S, Ramsey-Goldman R, Labrecque J, Joseph L, Boivin JF, Petri M, Zoma A, Manzi S, Urowitz MB, Gladman D, *et al*: Cancer risk in systemic lupus: an updated international multi-centre cohort study. *J Autoimmun* 42: 130-135, 2013.
- Tessler FN, Middleton WD and Grant EG: Thyroid imaging reporting and data system (TI-RADS): A user's guide. *Radiology* 287: 29-36, 2018.
- Amin MB, Greene FL, Edge SB, Compton CC, Gershenwald JE, Brookland RK, Meyer L, Gress DM, Byrd DR and Winchester DP: The eighth edition AJCC cancer staging manual: Continuing to build a bridge from a population-based to a more 'personalized' approach to cancer staging. *CA Cancer J Clin* 67: 93-99, 2017.
- Antonelli A, Mosca M, Fallahi P, Neri R, Ferrari SM, D'Ascanio A, Ghiri E, Carli L, Miccoli P and Bombardieri S: Thyroid cancer in systemic lupus erythematosus: a case-control study. *J Clin Endocrinol Metab* 95: 314-318, 2010.
- Okosieme OE, Evans C, Moss L, Parkes AB, Premawardhana LD and Lazarus JH: Thyroglobulin antibodies in serum of patients with differentiated thyroid cancer: Relationship between epitope specificities and thyroglobulin recovery. *Clin Chem* 51: 729-734, 2005.
- Inui A, Ogasawara H, Naito T, Sekigawa I, Takasaki Y, Hayashida Y, Takamori K and Ogawa H: Estrogen receptor expression by peripheral blood mononuclear cells of patients with systemic lupus erythematosus. *Clin Rheumatol* 26: 1675-1678, 2007.
- Zeng Q, Chen G, Vlantis A, Tse G and van Hasselt C: The contributions of oestrogen receptor isoforms to the development of papillary and anaplastic thyroid carcinomas. *J Pathol* 214: 425-433, 2008.
- Kansakar E, Chang YJ, Mehrabi M and Mittal V: Expression of estrogen receptor, progesterone receptor, and vascular endothelial growth factor-A in thyroid cancer. *Am Surg* 75: 785-789; discussion 789, 2009.
- Lahita RG: The role of sex hormones in systemic lupus erythematosus. *Curr Opin Rheumatol* 11: 352-356, 1999.
- Cooper GS, Dooley MA, Treadwell EL, St Clair EW, Parks CG and Gilkeson GS: Hormonal, environmental, and infectious risk factors for developing systemic lupus erythematosus. *Arthritis Rheum* 41: 1714-1724, 1998.
- Kiriakidou M and Ching CL: Systemic lupus erythematosus. *Ann Intern Med* 172: ITC81-ITC96, 2020.
- Fan Y, Gao D and Zhang Z: Telitacept, a novel humanized, recombinant TACI-Fc fusion protein, for the treatment of systemic lupus erythematosus. *Drugs Today (Barc)* 58: 23-32, 2022.
- Wu D, Li J, Xu D, Merrill JT, van Vollenhoven RF, Liu Y, Hu J, Li Y, Li F, Huang C, *et al*: Telitacept in patients with active systemic lupus erythematosus: Results of a phase 2b, randomised, double-blind, placebo-controlled trial. *Ann Rheum Dis* 83: 475-487, 2024.
- Navarra SV, Guzmán RM, Gallacher AE, Hall S, Levy RA, Jimenez RE, Li EK, Thomas M, Kim HY, León MG, *et al*: Efficacy and safety of belimumab in patients with active systemic lupus erythematosus: A randomised, placebo-controlled, phase 3 trial. *Lancet* 377: 721-731, 2011.
- Cheng J, Peng Y, Wu Q, Wu Q, He J and Yuan G: Efficacy and safety of telitacept therapy in systemic lupus erythematosus with hematological involvement. *Clin Rheumatol* 43: 2229-2236, 2024.
- Huang L, Qian G, Zhang H, Li Q, Chen L, Tang X and Zhao H: Efficacy of telitacept in a systemic lupus erythematosus patient with suboptimal response to Belimumab: A case report. *Lupus* 33: 172-175, 2024.
- Fan Q, Ji H, Liu Y, Jia C, Zou L and Yang H: Refractory lupus hepatitis successfully treated with telitacept who failed to belimumab: A case report and literature review. *Lupus* 33: 414-419, 2024.
- Ma X, Fu X, Cui B and Lin H: Telitacept for recalcitrant cutaneous manifestations of systemic lupus erythematosus: A case report and review of the literature. *Tohoku J Exp Med* 258: 219-223, 2022.
- Li S, Deng S, Wen S, Peng S, Jiang N, Li B, Chen B, Yuan Y, Wu Q, Tao Y, *et al*: Telitacept treatment refractory lupus nephritis: A case report. *Case Rep Nephrol Dial* 14: 42-47, 2024.
- Huang X, Lin F and Chen H: Efficacy and safety of telitacept in patients with lupus nephritis: A single-center, real-world retrospective study. *Clin Exp Nephrol* 28: 902-909, 2024.
- Chen R, Fu R, Lin Z, Huang C and Huang W: The efficacy and safety of telitacept for the treatment of systemic lupus erythematosus: A real life observational study. *Lupus* 32: 94-100, 2023.
- Sarzi-Putini P, Ceribelli A, Marotto D, Batticciotto A and Atzeni F: Systemic rheumatic diseases: From biological agents to small molecules. *Autoimmun Rev* 18: 583-592, 2019.
- Dreyer L, Cordtz RL, Hansen IMJ, Kristensen LE, Hetland ML and Mellemkjaer L: Risk of second malignant neoplasm and mortality in patients with rheumatoid arthritis treated with biological DMARDs: A danish population-based cohort study. *Ann Rheum Dis* 77: 510-514, 2018.
- Dixon WG, Watson KD, Lunt M, Mercer LK, Hyrich KL and Symmons DP; British Society For Rheumatology Biologics Register Control Centre Consortium; British Society for Rheumatology Biologics Register: Influence of anti-tumor necrosis factor therapy on cancer incidence in patients with rheumatoid arthritis who have had a prior malignancy: Results from the British society for rheumatology biologics register. *Arthritis Care Res (Hoboken)* 62: 755-763, 2010.
- Strangfeld A, Hierse F, Rau R, Burmester GR, Krummel-Lorenz B, Demary W, Listing J and Zink A: Risk of incident or recurrent malignancies among patients with rheumatoid arthritis exposed to biologic therapy in the German biologics register RABBIT. *Arthritis Res Ther* 12: R5, 2010.
- Mastorino L, Dapavo P, Avallone G, Merli M, Cariti C, Rubatto M, Pala V, Quaglini P and Ribero S: Biologic treatment for psoriasis in cancer patients: Should they still be considered forbidden? *J Dermatolog Treat* 33: 2495-2502, 2022.
- Pellegrini C, Esposito M, Rossi E, Gisondi P, Piaserico S, Dapavo P, Conti A, Gambardella A, Burlando M, Narcisi A, *et al*: Secukinumab in patients with psoriasis and a personal history of malignancy: A multicenter real-life observational study. *Dermatol Ther (Heidelb)* 12: 2613-2626, 2022.
- Blauvelt A: Safety of secukinumab in the treatment of psoriasis. *Expert Opin Drug Saf* 15: 1413-1420, 2016.
- Frieder J, Kivelevitch D and Menter A: Secukinumab: A review of the anti-IL-17A biologic for the treatment of psoriasis. *Ther Adv Chronic Dis* 9: 5-21, 2018.
- Gupta A, Peyrin-Biroulet L and Ananthakrishnan AN: Risk of cancer recurrence in patients with immune-mediated diseases with use of immunosuppressive therapies: An updated systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 22: 499-512.e6, 2024.
- Siegel CA, Finlayson SR, Sands BE and Tosteson AN: Adverse events do not outweigh benefits of combination therapy for Crohn's disease in a decision analytic model. *Clin Gastroenterol Hepatol* 10: 46-51, 2012.

39. Scott FI, Vajravelu RK, Bewtra M, Mamtani R, Lee D, Goldberg DS and Lewis JD: The benefit-to-risk balance of combining infliximab with azathioprine varies with age: A markov model. *Clin Gastroenterol Hepatol* 13: 302-309.e11, 2015.
40. Buchbinder R, Barber M, Heuzenroeder L, Wluka AE, Giles G, Hall S, Harkness A, Lewis D, Littlejohn G, Miller MH, *et al*: Incidence of melanoma and other malignancies among rheumatoid arthritis patients treated with methotrexate. *Arthritis Rheum* 59: 794-799, 2008.
41. Allegretti JR, Barnes EL and Cameron A: Are patients with inflammatory bowel disease on chronic immunosuppressive therapy at increased risk of cervical high-grade dysplasia/cancer? A meta-analysis. *Inflamm Bowel Dis* 21: 1089-1097, 2015.
42. Bernheim O, Colombel JF, Ullman TA, Laharie D, Beaugerie L and Itzkowitz SH: The management of immunosuppression in patients with inflammatory bowel disease and cancer. *Gut* 62: 1523-1528, 2013.
43. Shelton E, Laharie D, Scott FI, Mamtani R, Lewis JD, Colombel JF and Ananthakrishnan AN: Cancer recurrence following immune-suppressive therapies in patients with immune-mediated diseases: A systematic review and meta-analysis. *Gastroenterology* 151: 97-109.e4, 2016.
44. Karadaghy OA, Kallogjeri D and Piccirillo JF: Development of a new clinical severity staging system for patients with nonmetastatic papillary thyroid carcinoma. *JAMA Otolaryngol Head Neck Surg* 143: 1173-1180, 2017.
45. Torregrossa L, Poma AM, Macerola E, Rago T, Vignali P, Romani R, Proietti A, Di Stefano I, Scuotri G, Ugolini C, *et al*: The Italian consensus for the classification and reporting of thyroid cytology: Cytohistologic and molecular correlations on 37,371 nodules from a single institution. *Cancer Cytopathol* 130: 899-912, 2022.



Copyright © 2025 Tan et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.