

Application of protein chip combined with SELDI-TOF-MS detection to investigate serum protein expression in patients with silicosis fibrosis

GUITAO LIU^{1,2}, JUNFENG YU³, CUIDONG LI², XIAOYUN ZHOU², LIPING NIE²,
YANHUA WEI², WENYU WANG¹, YING ZHANG¹, NUZIGULI NUSILAITI¹,
PING HUA¹, XIAOHUA WANG², WENLONG WEI² and XINYAN LI⁴

¹Department of Occupational Disease Prevention and Control, The Fifth Affiliated Hospital of Xinjiang Medical University, Urumqi, Xinjiang 830011; ²Department of Respiratory Diseases, The Fifth Affiliated Clinical Medical College of Xinjiang Medical University, Xinjiang Medical University, Urumqi, Xinjiang 830054; ³Dermatological Department; ⁴Endocrinology Department, The Fifth Affiliated Hospital of Xinjiang Medical University, Urumqi, Xinjiang 830011, P.R. China

Received December 1, 2017; Accepted August 14, 2018

DOI: 10.3892/etm.2019.7166

Abstract. The present study aimed to observe the identification of biomarkers of silicosis based on the differentially expressed serum proteins between normal healthy individuals and patients with silicosis fibrosis. A total number of 20 patients with clinically diagnosed silicosis were screened, which were designated as the foundation treatment group. In addition, 20 age-matched healthy patients attending a check-up at the physical examination department were selected. Serum samples were obtained and a combined protein chip with surface-enhanced laser desorption ionization flight mass spectrometry was applied to perform serum analysis. Data preprocessing, screening differences in peak, hierarchical cluster analysis, Principal Component Analysis, construction of a decision tree model, and prediction based on the differences between peaks corresponding to proteins were performed to analyze the data. The results revealed differences in the proteins in serum between the normal group and the group prior to foundation treatment prediction. The corresponding names of the protein peak, predicted protein, and gene name were as follows: M1948_00, complement c3 frag, C3; M2017_02, amyloid- β 4 protein, APP; and M2879_56, hepcidin, HAMP. Differentially expressed serum proteins in the normal group and the basis treatment group were predicted, including M2017_02, amyloid- β 4 protein, APP; M2879_56, hepcidin, HAMP; and M3224_97, fibrinogen- α chain frags, FGA. The

differentially expressed serum proteins in the group prior to basis treatment and the group following basis treatment were predicted, including M2001_69, amyloid- β 4 protein, APP; M2017_02, amyloid- β 4 protein, APP, M4144_81, plasma protease c1 inhibitor frag, and SERPING1. In conclusion, there were differences in the proteins in serum between the patients with silicosis fibrosis and healthy individuals.

Introduction

Pneumoconiosis follows the long-term inhalation of productive dust, which is the primary contributor to pulmonary fibrosis of systemic disease. It is also a progressive fibrotic lung disease, which is caused by occupational exposure to mineral dust and fibers (1). It is one of the most serious occupational disease hazards in China (2). Currently, occupational hazards exist in >8,000 enterprises in Xinjiang, China; >1,300,00 individuals may be contact with occupational hazards and there are >16,000 patients with various occupational diseases. All types of occupational disease hazard factors are increasing with the process of western development and, although there are certain protective measures, there are increasing numbers of patients with silicosis. According to incomplete current statistics, silicosis not only causes serious damage, due to silicon dust exposure, to the health of workers, but also severely reduces the labor force and quality of life, and causes economic loss to the country (3). According to statistics, the economic loss caused by occupational diseases each year is over 140 million RMB. Further investigations on diagnosis are required owing to new challenges in silicosis (4).

Surface-enhanced laser desorption ionization flight mass spectrometry (SELDI-TOF-MS) technology was used in the present study to detect the difference in serum protein between patients with silicosis fibrosis and normal healthy individuals, which is a novel effective method for the clinical diagnosis of silicosis by identifying markers for silicosis.

Correspondence to: Dr Xinyan Li, Endocrinology Department, The Fifth Affiliated Hospital of Xinjiang Medical University, 118 Henan Road, Urumqi, Xinjiang 830011, P.R. China
E-mail: xinyanlii@126.com

Key words: silicosis fibrosis, surface-enhanced laser desorption ionization flight mass spectrometry, different protein in serum

Materials and methods

Serum sample information. A total of 60 male patients with silicosis were selected and, according to inclusion criteria and exclusion criteria, 20 patients with clinical cases of silicosis diagnosed within one period were screened and determined as the foundation treatment group, designated as group A. Therefore, of the 60 individuals in total recruited for the study in The Fifth Affiliated Hospital of Xinjiang Medical University (Urumqi, China), 20 patients with silicosis were included. In addition, 20 age-matched healthy patients attending check-ups at the physical examination department in the Fifth Affiliated Clinical Medical College of Xinjiang Medical University were selected as a normal healthy group, designated as group C. The mean age of the three groups were 75.7±4.32 years (group A), 76.1±5.62 years (group B) and 74.8±4.31 years (group C). All samples were collected between June 2014 and July 2015. Groups A and B were patients with silicosis, but the treatment options were different. The treatment of patients in Group A (basic treatment) was as follows: Oxygen therapy, coughing, and expectorant, anti-asthmatic and anti-inflammatory therapy. The treatment of patients in Group B was as follows: basic treatment + 2 ml Deworming Vernonia Injection (repellent spotted chrysanthemum injection; Anhui Golden Sun Biochemical Pharmaceutical Co., Ltd., Fuyang, China) in 10 ml of 0.9% physiological saline for nebulization. Venous blood (5 ml) was centrifuged at 900 x g for 10 min at 4°C; the serum was removed and stored in a freezer at -80°C. Fasting morning venous blood samples were obtained from the normal group (group C), the pre-foundation treatment group (A-Q) and the post-foundation treatment group (A-H). There were a total of 60 samples from the three groups, as each group had 20 biological repeats. SELDI-TOF-MS detection analysis was performed on the samples. The foundation treatment for silicosis comprised chemotherapy using poly 2-vinyl pyridine nitroxide (5). Patients with silicosis can be rapidly detected by detecting the plasma levels of tumor necrosis factor- α and matrix metalloproteinase 9 (6). The changes in the expression levels of numerous proteins following foundation treatment occur in the same patient and correlate with disease status. The inclusion criteria were as follows: History of dust exposure; early symptoms or signs; X-ray examination as the main basis for the diagnosis of silicosis; laboratory tests for the examination of early diagnostic indicators; pulmonary function tests showing pulmonary ventilation disorders. The exclusion criteria comprised patients who did not meet the inclusion criteria, did not attend follow-up or did not sign the informed consent forms.

Equipment and instrument. The Ciphergen® SELDI-TOF-MS mass spectrometer (protein fingerprint device) was used in the present study (Ciphergen Biosystems, Inc., Fremont, CA, USA). In order to identify the biomarkers, the Protein Chip SELDI system was used, which can rapidly produce a protein molecular weight map from a large number of complex biological samples. Using surface enhanced laser desorption ion technology, it is able to capture, detect and measure the molecular weight of peptides and proteins in complex biological samples (4).

SELDI protein chip. The covalent coupling of selected proteins or other target molecules can be implemented on the surface of

a chip. A particular subgroup of complex protein samples can be captured by the chip through simple chemistry or protein interaction.

Following incubation, protein that was not combined with other ingredients from the chip surface was cleared. Only those specifically binding proteins were retained for further analysis. Following the elution step, the organic solution of energy absorption molecules was added. The chip in the SELDI reading machine was analyzed, which involved a type of time of flight mass spectrometry. Once in a gaseous state, charged protein molecules under the effect of a separation voltage show rapid movement, termed 'flight'. The separation voltage for all the molecules in the sample has the same effect, with differences in time of flight according to the different molecular weight. The SELDI reading machine recorded the time of flight, and converted this to a molecular weight.

Contrast strategy and analysis of the content. A total of three comparative analyses with the following comparisons, respectively, were performed, and data was obtained for further bioinformatics analysis: C, vs. A-Q; C, vs. A-H; and A-Q, vs. A-H. The data analysis process included data preprocessing, differences in peak screening, hierarchical cluster analysis, Principal Component Analysis (PCA), construction of a decision tree model, and protein prediction based on differences between the peak corresponding to proteins.

Statistical analysis. SPSS 17.0 software (SPSS, Inc., Chicago, IL, USA) was used for the quantitative data, with two sets of equal variances, using a Student's t-test. Two sets of heterogeneity of variance, with rank and inspection, disordered classification data were analyzed using the χ^2 test. The Wilcoxon sum rank test was used with a capacity of <10. The specific steps of the test were as follows: Firstly, data of two samples were mixed and ranked from the smallest to the largest (the smallest data order is 1 and the largest data order is $n_1 + n_2$). Secondly, the ranks of data in the smaller capacity sample are add, that is, the rank sum, denoted by \check{T} . Thirdly, the T value was compared with the critical value at a certain level of significance in the rank sum test table. If $T_1 < T < T_2$, the difference between the two samples is insignificant; $T < T_1$ or $T \geq T_2$ indicates that the two samples are significantly different. The Wilcoxon rank sum test was used in the present study to obtain P-values. The difference in protein peaks was determined by calculating the P-value to determine the significance of differences of each protein peak. The permutation tests were performed with statistical tools in SIMCA 14.0 software (Umetrics AB, Umea, Sweden). $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Data preprocessing. Raw data analyses were performed using Ciphergen Protein Chip software correction processing to obtain the peak data. A mass-to-charge ratio <1,000 of the peak was considered the substrate peak, and the subsequent data was filtered based on the peak.

Difference in peak filtering. The peak between two groups was compared using the Wilcoxon sum rank test, using the peak

Table I. C, vs. A-Q group differences in peak filtering.

SAMP_GRP	P-value	q-value	VIP	C average	A-Q average
M1004_63	2.9018x10 ⁻¹¹	4.38x10 ⁻⁹	2.8325	9.337262319	20.59812918
M1009_81	2.884x10 ⁻⁶	1.4043x10 ⁻⁵	1.18109	31.51986384	40.84034414
M1019_45	4.3527x10 ⁻¹⁰	1.5438x10 ⁻⁸	2.63561	4.718782081	12.56350599
M1031_99	6.529x10 ⁻¹⁰	1.6425x10 ⁻⁸	3.48656	7.811703062	21.54302217
M1083_37	4.9091x10 ⁻⁶	2.0583x10 ⁻⁵	1.12792	0.294565816	2.199163708
M1098_33	1.3412x10 ⁻⁷	1.1292x10 ⁻⁶	1.67063	0.38657576	3.615140978
M1101_43	9.9241x10 ⁻⁹	1.07x10 ⁻⁷	2.14677	0.927844358	5.631289799
M1103_90	2.0167x10 ⁻⁹	3.3824x10 ⁻⁸	1.64852	0.479273296	3.302777091
M1114_98	4.0713x10 ⁻⁷	2.6719x10 ⁻⁶	1.5397	0.418107229	3.207923021
M1123_40	3.028x10 ⁻⁸	2.8566x10 ⁻⁷	1.64771	0.54015768	3.571119791
M1127_55	1.3276x10 ⁻⁸	1.3359x10 ⁻⁷	1.91652	0.784175155	4.616498053
M1138_12	9.6475x10 ⁻⁶	3.6341x10 ⁻⁵	1.25759	0.461889022	2.446496536
M1144_16	2.0167x10 ⁻⁹	3.3824x10 ⁻⁸	2.7147	1.740541284	9.570840078
M1146_35	6.529x10 ⁻¹⁰	1.6425x10 ⁻⁸	2.77186	2.612216217	10.34906681
M1150_36	3.9464x10 ⁻⁹	5.4153x10 ⁻⁸	1.87836	1.903402107	6.508579836
M1163_78	0.00010426	0.00029951	1.28628	0.268622126	2.446015217
M1190_81	2.8978x10 ⁻⁵	9.678x10 ⁻⁵	1.21026	0.31970599	2.341884927
M1207_98	6.529x10 ⁻¹⁰	1.6425x10 ⁻⁸	2.49502	0.550691748	6.396113855
M1216_31	2.7567x10 ⁻¹⁰	1.3978x10 ⁻⁸	2.92377	2.903470782	11.45476404
M1230_37	4.9091x10 ⁻⁶	2.0583x10 ⁻⁵	1.5803	0.390190586	3.294667162
M1234_44	1.5646x10 ⁻⁵	5.4923x10 ⁻⁵	1.72882	1.132329262	5.002732013
M1259_38	4.0713x10 ⁻⁷	2.6719x10 ⁻⁶	1.72134	0.668984452	3.864822477
M1324_75	4.1219x10 ⁻⁶	1.8269x10 ⁻⁵	1.32713	0.326495903	2.353783547
M1351_32	0.00025797	0.0006279	1.07649	0.222331714	1.765712861
M1368_84	2.0167x10 ⁻⁹	3.3824x10 ⁻⁸	2.00209	0.614290076	4.832761142
M1435_91	2.4028x10 ⁻⁶	1.2236x10 ⁻⁵	1.64372	0.304051349	3.099785952
M1454_17	9.9241x10 ⁻⁹	1.07x10 ⁻⁷	2.09005	0.237729927	4.569096508
M1475_93	0.00015512	0.0004181	1.1731	0.172207775	2.014840182
M1505_23	4.5102x10 ⁻⁵	0.00014227	1.27917	0.903329951	3.306640883
M1625_88	1.6891x10 ⁻⁷	1.3726x10 ⁻⁶	1.48823	0.535477091	3.408169271
M1700_90	2.1171x10 ⁻⁷	1.6498x10 ⁻⁶	1.37375	0.23234938	2.32507649
M1948_00	0.00015512	0.0004181	1.18882	0.126504245	1.85957764
M2017_02	2.8292x10 ⁻⁹	4.2705x10 ⁻⁸	2.28118	3.299510237	10.03986804
M2879_56	0.00022766	0.00057042	1.78224	6.897042594	14.34943243
M2959_08	7.5712x10 ⁻⁷	4.5713x10 ⁻⁶	2.03316	0.916531467	5.060081501
M3224_97	0.00093342	0.00185385	1.06247	2.49928245	1.090001717
M3246_10	2.4028x10 ⁻⁶	1.2236x10 ⁻⁵	1.87892	0.396642163	4.2861341
M3322_38	0.04595423	0.04441369	1.54968	7.748435198	4.898142987
M3825_21	0.00074741	0.00154007	1.08572	2.514007357	0.827869797
M3900_31	0.00022766	0.00057042	1.24374	2.663706342	0.890604754
M4421_61	5.2082x10 ⁻⁵	0.00016044	1.69686	4.732555028	1.951991338
M4478_27	3.2864x10 ⁻⁷	2.3028x10 ⁻⁶	1.54532	2.864591283	0.276282046
M4715_56	3.8989x10 ⁻⁵	0.00012563	1.51004	3.657925544	1.239881596

C, normal group; A-Q, pre-foundation treatment group.

P-value to judge whether there were significant differences between the two groups. By selecting different peaks, differences between peaks were compared and the screening results for C, vs. A-Q (Table I), C, vs. A-H (Table II), and A-Q, vs. S A-H (Table III) were obtained. As shown in Table I,

the normal group and pre-foundation treatment group had statistically significant differences between peaks. As shown in Table II, the normal group and post-foundation treatment group had statistically significant differences between peaks. As shown in Table III, between the pre- and post-foundation

Table II. C, vs. A-H group differences in peak filtering.

SAMP_GRP	P-value	q-value	VIP	C average	A-H average
M1004_63	1.06x10 ⁻⁷	8.18 x10 ⁻⁶	2.75206	9.337262319	18.36057094
M1009_81	6.91x10 ⁻⁵	0.000491	1.4842	31.51986384	40.48819307
M1019_45	3.45x10 ⁻⁶	6.65x10 ⁻⁵	1.97222	4.718782081	9.320910566
M1031_99	6.53x10 ⁻⁸	8.18x10 ⁻⁶	3.3601	7.811703062	18.49940089
M1083_37	0.00033	0.001496	1.01317	0.294565816	1.699066783
M1098_33	0.000595	0.002364	1.56859	0.38657576	2.544233659
M1101_43	6x10 ⁻⁵	0.000455	2.06365	0.927844358	4.239463466
M1103_90	1.14x10 ⁻⁵	0.000135	1.32065	0.479273296	2.349881329
M1114_98	6x10 ⁻⁵	0.000455	1.58215	0.418107229	2.796677459
M1123_40	0.010314	0.020685	1.14424	0.54015768	2.276438971
M1127_55	0.001767	0.005778	1.68324	0.784175155	3.442394353
M1138_12	0.010314	0.020685	1.29039	0.461889022	2.551331219
M1144_16	9.1x10 ⁻⁵	0.000562	2.53058	1.740541284	6.707315793
M1146_35	0.000258	0.001265	2.35356	2.612216217	7.147118481
M1150_36	0.000228	0.001156	1.31248	1.903402107	4.223835486
M1190_81	0.006145	0.015189	1.18846	0.31970599	1.670230569
M1207_98	6.91x10 ⁻⁶	9.69x10 ⁻⁵	1.95582	0.550691748	3.644110346
M1216_31	1.65x10 ⁻⁶	4.25x10 ⁻⁵	2.35906	2.903470782	8.011714226
M1225_21	0.008712	0.018838	1.15032	0.055092064	1.468164449
M1230_37	9.1x10 ⁻⁵	0.000562	1.47151	0.390190586	2.300611105
M1234_44	0.000136	0.000774	1.93723	1.132329262	5.024199001
M1351_32	0.014297	0.025343	1.11582	0.222331714	1.538636715
M1368_84	3.36x10 ⁻⁵	0.000317	1.76946	0.614290076	3.075023606
M1454_17	6.91x10 ⁻⁵	0.000491	1.6592	0.237729927	2.389504296
M1625_88	2.88x10 ⁻⁶	6.04x10 ⁻⁵	1.7615	0.535477091	4.070970461
M2017_02	0.001291	0.004424	1.25703	3.299510237	6.380801455
M2879_56	0.002643	0.008118	1.43552	6.897042594	11.9477084
M2959_08	0.001435	0.004811	1.52917	0.916531467	2.90088487
M3224_97	0.035011	0.049694	1.06234	2.49928245	1.509255098
M3246_10	1.13x10 ⁻⁶	3.47x10 ⁻⁵	1.83767	0.396642163	3.275699253
M3322_38	0.004681	0.0126	1.88645	7.748435198	4.837010977
M3825_21	0.011207	0.021603	1.16951	2.514007357	1.159904936
M3900_31	0.011207	0.021603	1.09896	2.663706342	1.345515395
M4412_23	0.012166	0.022765	1.03269	2.483203325	1.25835564
M4421_61	0.003211	0.00949	1.89073	4.732555028	2.263517789
M4715_56	9.65x10 ⁻⁶	0.000122	1.86603	3.657925544	0.858452439
M4770_22	0.009484	0.019762	1.05903	1.036151502	0.130670044
M4797_31	0.022719	0.03575	1.37116	4.054278147	1.971892733
M4825_09	0.014297	0.025343	1.30691	3.330875754	2.065185447
M4832_49	0.000372	0.00164	1.29409	2.046751735	0.412735635
M5254_17	0.000201	0.001052	1.5746	1.256545457	4.629496349
M5341_27	2.64x10 ⁻⁷	1.36x10 ⁻⁵	4.89611	9.675443865	26.47358094
M5775_96	0.037519	0.052238	1.1959	2.1058316	0.742538154

C, normal group; A-H, post-foundation treatment group.

treatment group, there were statistically significant differences in peaks.

Hierarchical clustering analysis. The hierarchical cluster analysis was performed on the differences in peaks, with a

clustering diagram showing the association between the samples. Each line represents a peak in the diagram (Fig. 1), and each column represents a sample. The red and green colors represent sample testing content is higher and lower, respectively. The comparison results for each group are shown

Table III. A-Q, vs. A-H group differences in peak filtering.

SAMP_GRP	P-value	q-value	VIP	A-Q average	A-H average
M1004_63	0.039989	0.415819	1.16467	20.59813	18.36057
M1019_45	0.002325	0.220852	2.23625	12.56351	9.320911
M1123_40	0.044054	0.427081	1.27539	3.57112	2.276439
M1146_35	0.036234	0.403855	2.52543	10.34907	7.147118
M1150_36	0.000708	0.220852	1.73778	6.50858	4.223835
M1207_98	0.013617	0.299041	2.71017	6.396114	3.64411
M1216_31	0.00639	0.247157	2.00268	11.45476	8.011714
M1245_23	0.036234	0.403855	2.37039	-0.30271	0.870859
M1259_38	0.048441	0.437643	1.91349	3.864822	2.087152
M1324_75	0.021484	0.333673	1.45305	2.353784	1.244105
M1368_84	0.044054	0.427081	2.00074	4.832761	3.075024
M1435_91	0.013617	0.299041	1.76205	3.099786	1.304288
M1475_93	0.048441	0.437643	1.73211	2.01484	0.937137
M1505_23	0.026642	0.363733	1.35806	3.306641	2.081841
M1661_24	0.019234	0.317797	2.17012	1.212314	0.238766
M1694_85	0.004221	0.220852	1.66347	1.245148	0.288131
M1700_90	0.019234	0.317797	2.19981	2.325076	1.021293
M2001_69	0.013617	0.299041	1.81131	2.097512	0.993857
M2017_02	0.002712	0.220852	2.50106	10.03987	6.380801
M4144_81	0.003654	0.220852	1.30244	-0.13272	0.656736
M4478_27	0.003654	0.220852	2.15012	0.276282	2.172395
M4861_39	0.048441	0.437643	1.32359	0.092989	0.802765
M5642_48	0.007296	0.254484	1.30207	0.955681	0.245088
M7564_68	0.012079	0.291543	1.54656	0.551073	1.201719

A-Q, pre-foundation treatment group; A-H, post-foundation treatment group.

in heat maps: C, vs. A-Q (Fig. 1A), C, vs. A-H (Fig. 1B), and A-Q, vs. A-H (Fig. 1C). Shown in Fig. 1A is the association between the normal group and pre-foundation treatment group of samples, with red indicating sample testing content was higher and green indicating the sample testing content was lower. In Fig. 1B, the association between the normal group and post-foundation treatment group of samples is shown. In Fig. 1C, the association between pre- and post-foundation treatment group samples is shown.

PCA analysis. Using SIMCA software (version 14), the PCA analysis results were obtained. The scoring results for the total and the comparisons in each group are shown. As shown in Fig. 2A, the normal group and pre-foundation treatment group were distinguished. As shown in Fig. 2B, the normal group and post-foundation treatment group were distinguished.

Orthogonal Projections to Latent Structures Discriminant Analysis (OPLS-DA) model construction. The C, vs. A-Q group OPLS-DA scoring chart is shown in Fig. 3A, the displacement test (permutation test) diagram is shown in Fig. 3B, and the OPLS-DA load diagram is shown in Fig. 3C. It was shown that there was a significant difference in the protein level between the normal group and pre-foundation treatment group. The

C, vs. A-H group OPLS-DA scoring diagram is shown in Fig. 4A, the displacement test (permutation test) diagram is shown in Fig. 4B, and the OPLS-DA load diagram is shown in Fig. 4C. The protein level was significantly different between the normal group and post-foundation treatment group. The A-Q, vs. A-H group OPLS-DA scoring chart is shown in Fig. 5A, the displacement test (permutation test) diagram is shown in Fig. 5B and the OPLS-DA load diagram is shown in Fig. 5C. A-Q and A-H were markedly different (Fig. 5A). The displacement test of the OPLS-DA model demonstrated that the intercept of Q2 was less than 0.05, therefore the models were robust (Fig. 5B). Each point represents a protein that was expressed higher in the sample compared with another (Fig. 5C). Proteins on the left side of the axis were highly expressed in the A-H group and proteins on the right side of the axis were highly expressed in the A-Q group. These results demonstrate that proteins level were markedly different between the normal group and the post-foundation treatment group.

Differential protein prediction. Using Swiss-Prot (2017-10 release; https://www.uniprot.org/uniprot/?query=*&fil=organism%3A%22Homo+sapiens+%28Human%29+%5B9606%5D%22+AND+reviewed%3Ayes) as the standard protein data in the database, self-owned coding texted software was used to compare this with the amino acid molecular weight of protein,




Figure 2. Comparison of score plots of PCA model obtained from C, A-Q and A-H groups. (A) C, vs. A-Q PCA scoring chart. (B) C, vs. A-H PCA scoring chart. (C) A-Q, vs. A-H PCA scoring chart. PCA, Principal Component Analysis; C, normal group; A-Q, pre-foundation treatment group; A-H, post-foundation treatment group.




Figure 3. C, vs. A-Q group OPLS-DA model. (A) C, vs. A-Q group OPLS-DA scoring diagram shows the score plot of the OPLS-DA model obtained from the C and A-Q groups. (B) C, vs. A-Q group permutation test chart. A total of 200 permutations were performed, and the resulting R2 and Q2 values are plotted. Green circle, R2; blue square, Q2. The green line represents the regression line for R2 and the blue line for Q2. (C) C, vs. A-Q group OPLS-DA load diagram of the loading plot of the OPLS-DA model obtained from the C and A-Q groups. OPLS-DA, Orthogonal Projections to Latent Structures Discriminant Analysis; C, normal group; A-Q, pre-foundation treatment group; A-H, post-foundation treatment group.




Figure 4. C, vs. A-H group OPLS-DA model. (A) C, vs. A-H group OPLS-DA scoring chart. (B) C, vs. A-H group displacement test chart. (C) C, vs. A-H group OPLS-DA load diagram. OPLS-DA, Orthogonal Projections to Latent Structures Discriminant Analysis; C, normal group; A-Q, pre-foundation treatment group; A-H, post-foundation treatment group.




Figure 5. A-Q, vs. A-H group model. (A) A-Q, vs. A-H group OPLS-DA scoring chart. (B) A-Q, vs. A-H group permutation test chart. (C) A-Q, vs. A-H group OPLS-DA load diagram. OPLS-DA, Orthogonal Projections to Latent Structures Discriminant Analysis; C, normal group; A-Q, pre-foundation treatment group; A-H, post-foundation treatment group.

Table IV. C, vs. A-Q group differences protein prediction results.

SAMP_GRP	P-value	q-value	VIP	ID	MZ_SELDI	Predicted protein	Theoretical MZ	Gene name
M1948_00	0.000155115	0.000418099	1.18882	M1948_00	1,948	Complement c3 frag	1,865.12	C3
M2017_02	2.83x10 ⁻⁹	4.27x10 ⁻⁸	2.28E+00	M2017_02	2,017.02	Amyloid β a4 protein	1,953.10	APP
M2879_56	0.000227661	0.000570423	1.78224	M2879_56	2,879.56	Hepcidin	2,797.41	HAMP
M3224_97	0.000933418	0.001853851	1.06247	M3224_97	3,224.97	Fibrinogen α -chain frag	3,262.47	FGA
M3246_10	2.40x10 ⁻⁶	1.22x10 ⁻⁵	1.88E+00	M3246_10	3,246.1	Fibrinogen α -chain frag	3,262.47	FGA
M3322_38	0.04595423	0.044413687	1.54968	M3322_38	3,322.38	Fibrinogen α -chain frag	3,262.47	FGA
M3825_21	0.000747408	0.001540074	1.08572	M3825_21	3,825.21	Neurosecretory protein vgf frag	3,688.03	VPF
M3900_31	0.000227661	0.000570423	1.24374	M3900_31	3,900.31	Amyloid β a4 protein	4,074.54	APP
M4421_61	5.2x10 ⁻⁵	0.000160437	1.69686	M4421_61	4,421.61	α -1-antichymotrypsin, α -1-antichymotrypsin frag	4,343.65	SERPINA3
M4478_27	3.29x10 ⁻⁷	2.30x10 ⁻⁶	1.55E+00	M4478_27	4,478.27	α -1-antichymotrypsin, α -1-antichymotrypsin frag	4,343.65	SERPINA3
M4715_56	3.90x10 ⁻⁵	0.000125632	1.51004	M4715_56	4,715.56	Neurosecretory protein vgf frag	4,823.50	VPF
M4797_31	0.005617767	0.008299276	1.0441	M4797_31	4,797.31	Neurosecretory protein vgf frag	4,823.50	VPF
M5895_90	0.003210931	0.005373126	1.36332	M5895_90	5,895.9	Fibrinogen a-chain frag	5,904.22	FGA
M5908_34	5.83x10 ⁻⁶	2.32x10 ⁻⁵	3.48E+00	M5908_34	5,908.34	Fibrinogen achain frag	5,904.22	FGA
M6114_30	0.007330817	0.010059399	1.62262	M6114_30	6,114.3	Apolipoprotein c-i	6,432.35	APOC1
M6435_51	5.83x10 ⁻⁶	2.32x10 ⁻⁵	4.57E+00	M6435_51	6,435.51	Apolipoprotein c-i	6,630.58	APOC1
M6599_07	4.91x10 ⁻⁶	2.06x10 ⁻⁵	1.55E+00	M6599_07	6,599.07	Transthyretin	6,880.00	TTR
M6633_61	0.000835766	0.001682037	4.08017	M6633_61	6,633.61	Transthyretin	6,880.00	TTR
M6814_07	5.21x10 ⁻⁵	0.000160437	1.25676	M6814_07	6,814.07	Transthyretin	6,880.00	TTR
M6842_25	4.91x10 ⁻⁶	2.06x10 ⁻⁵	3.20E+00	M6842_25	6,842.25	Transthyretin	6,880.00	TTR
M6854_85	3.29x10 ⁻⁷	2.30x10 ⁻⁶	2.41E+00	M6854_85	6,854.85	Transthyretin	6,880.00	TTR
M7152_30	0.000530525	0.001161127	1.07687	M7152_30	7,152.3	Transthyretin	6,880.00	TTR
M8600_82	0.018093579	0.02061882	1.40868	M8600_82	8,600.82	Apolipoprotein c-iii	8,764.67	APOC3
M8696_24	0.006144956	0.008869015	2.30513	M8696_24	8,696.24	Apolipoprotein c-iii	8,764.67	APOC3
M8762_75	1.65x10 ⁻⁶	9.07x10 ⁻⁶	2.50E+00	M8762_75	8,762.75	Apolipoprotein c-iii	8,764.67	APOC3
M8823_58	0.00011921	0.000335954	3.04231	M8823_58	8,823.58	Apolipoprotein c-iii	8,764.67	APOC3
M8868_31	0.000667562	0.001405836	1.71114	M8868_31	8,868.31	Apolipoprotein a-ii	9,303.65	APOA2
M8923_78	0.000530525	0.001161127	2.0589	M8923_78	8,923.78	Apolipoprotein a-ii	9,303.65	APOA2
M9138_93	0.00011921	0.000335954	1.16089	M9138_93	9,138.93	Apolipoprotein a-ii	9,303.65	APOA2
M9414_68	7.37x10 ⁻⁹	8.75x10 ⁻⁸	2.70E+00	M9414_68	9,414.68	Apolipoprotein a-ii	9,303.65	APOA2
M9440_33	1.56x10 ⁻⁵	5.49x10 ⁻⁵	1.62E+00	M9440_33	9,440.33	Apolipoprotein a-ii	9,303.65	APOA2
M9633_47	2.88x10 ⁻⁶	1.40x10 ⁻⁵	1.10E+00	M9633_47	9,633.47	Apolipoprotein a-ii	9,303.65	APOA2
M9708_99	0.000291908	0.000688459	1.01733	M9708_99	9,708.99	Apolipoprotein a-ii	9,303.65	APOA2
M13886_7	0.000329848	0.000765972	1.03751	M13886_7	13,886.7	Cystatin-c	13,347.14	CST3
M14042_5	2.40x10 ⁻⁶	1.22x10 ⁻⁵	1.19E+00	M14042_5	14,042.5	Transthyretin	13,761.41	TTR

C, vs. A-Q, P<0.05 (t-test used for data analysis). C, normal group; A-Q, pre-foundation treatment group.

Table V. C, vs. A-H group differences protein prediction results.

SAMP_GRP	P-value	q-value	VIP	ID	MZ_SELDI	Predicted protein	Theoretical MZ	Gene name
M2017_02	0.001290971	0.004424107	1.25703	M2017_02	2,017.02	Amyloid β a4 protein	1,953.10	APP
M2879_56	0.002642631	0.008117783	1.43552	M2879_56	2,879.56	Hepcidin	2,797.41	HAMP
M2959_08	0.001434955	0.004810632	1.52917	M2959_08	2,959.08	No protein matched.		
M3224_97	0.03501066	0.049693616	1.06234	M3224_97	3,224.97	Fibrinogen α -chain frag	3,262.47	FGA
M3246_10	1.13x10 ⁻⁶	3.47x10 ⁻⁵	1.84E+00	M3246_10	3,246.1	Fibrinogen α -chain frag	3,262.47	FGA
M3322_38	0.004681199	0.012599765	1.88645	M3322_38	3,322.38	Fibrinogen α -chain frag	3,262.47	FGA
M3825_21	0.011206767	0.021602908	1.16951	M3825_21	3,825.21	Neurosecretory protein vgf frag	3,688.03	VPF
M3900_31	0.011206767	0.021602908	1.09896	M3900_31	3,900.31	Amyloid β a4 protein	4,074.54	APP
M4412_23	0.012165581	0.022764829	1.03269	M4412_23	4,412.23	α -1-antichymotrypsin, alpha-1-antichymotrypsin frag	4,343.65	SERPINA3
M4421_61	0.003210931	0.00948974	1.89073	M4421_61	4,421.61	α -1-antichymotrypsin, α -1-antichymotrypsin frag	4,343.65	SERPINA3
M4715_56	9.65 x10 ⁻⁶	0.00012165	1.86603	M4715_56	4,715.56	Neurosecretory protein vgf frag	4,823.50	VPF
M4770_22	0.009483607	0.019762305	1.05903	M4770_22	4,770.22	Neurosecretory protein vgf frag	4,823.50	VPF
M4797_31	0.022718728	0.035750307	1.37116	M4797_31	4,797.31	Neurosecretory protein vgf frag	4,823.50	VPF
M4825_09	0.014297273	0.02534287	1.30691	M4825_09	4,825.09	Neurosecretory protein vgf frag	4,823.50	VPF
M4832_49	0.000372206	0.001639974	1.29409	M4832_49	4,832.49	Neurosecretory protein vgf frag	4,823.50	VPF
M5775_96	0.037519175	0.052237795	1.1959	M5775_96	5,775.96	Neurosecretory protein vgf frag	5,904.22	FGA
M5883_70	0.049090325	0.06313916	1.07377	M5883_70	5,883.7	Fibrinogen α -chain frag	5,904.22	FGA
M5895_90	0.04595423	0.060326782	1.2068	M5895_90	5,895.9	Fibrinogen α -chain frag	5,904.22	FGA
M5908_34	0.000530525	0.002164918	3.30565	M5908_34	5,908.34	Fibrinogen α -chain frag	5,904.22	FGA
M5954_71	0.018093579	0.030305345	1.20553	M5954_71	5,954.71	Fibrinogen α -chain frag	5,904.22	FGA
M6114_30	0.012165581	0.022764829	1.64383	M6114_30	6,114.3	Fibrinogen α -chain frag	5,904.22	FGA
M6435_51	0.000667562	0.002573673	4.89498	M6435_51	6,435.51	Apolipoprotein c-i	6,432.35	APOC1
M6484_93	0.019536762	0.032051368	1.05576	M6484_93	6,484.93	Apolipoprotein c-i	6,630.58	APOC1
M6599_07	0.001041218	0.003734178	1.48867	M6599_07	6,599.07	Apolipoprotein c-i	6,630.58	APOC1
M6633_61	0.006144956	0.015189373	4.32524	M6633_61	6,633.61	Transthyretin	6,880.00	TTR
M6814_07	0.000104264	0.000618417	1.4876	M6814_07	6,814.07	Transthyretin	6,880.00	TTR
M6842_25	0.000419438	0.001796749	3.26991	M6842_25	6,842.25	Transthyretin	6,880.00	TTR
M6854_85	0.000667562	0.002573673	2.37706	M6854_85	6,854.85	Transthyretin	6,880.00	TTR
M7065_82	0.011206767	0.021602908	1.08096	M7065_82	7,065.82	Transthyretin	6,880.00	TTR
M7652_53	0.006144956	0.015189373	1.08712	M7652_53	7,652.53	Osteopontin frag	7,658.19	SPPI
M8600_82	0.013194383	0.023938228	2.21464	M8600_82	8,600.82	Apolipoprotein c-iii	8,764.67	APOC3
M8622_98	0.005617767	0.014306888	1.24828	M8622_98	8,622.98	Apolipoprotein c-iii	8,764.67	APOC3
M8696_24	0.006715009	0.016087017	2.86633	M8696_24	8,696.24	Apolipoprotein c-iii	8,764.67	APOC3
M8762_75	5.83x10 ⁻⁶	8.94x10 ⁻⁵	2.75E+00	M8762_75	8,762.75	Apolipoprotein c-iii	8,764.67	APOC3
M8823_58	0.0038885006	0.01098804	3.27878	M8823_58	8,823.58	Apolipoprotein c-iii	8,764.67	APOC3

Table V. Continued.

SAMP_GRP	P-value	q-value	VIP	ID	MZ_SELDI	Predicted protein	Theoretical MZ	Gene name
M8868_31	0.008711881	0.018837648	1.78778	M8868_31	8,868.31	Apolipoprotein a-ii	9,303.65	APOA2
M9138_93	0.011206767	0.021602908	1.00803	M9138_93	9,138.93	Apolipoprotein a-ii	9,303.65	APOA2
M9414_68	6.18x10 ⁻⁷	2.38x10 ⁻⁵	3.03E+00	M9414_68	9,414.68	Apolipoprotein a-ii	9,303.65	APOA2
M9440_33	0.000329848	0.001496086	1.70483	M9440_33	9,440.33	Apolipoprotein a-ii	9,303.65	APOA2
M9633_47	7.94 x10 ⁻⁵	0.000526668	1.14282	M9633_47	9,633.47	Apolipoprotein a-ii	9,303.65	APOA2
M13886_7	0.003533771	0.010223786	1.13858	M13886_7	13,886.7	Cystatin-c	13,347.14	CST3
M14042_5	9.10 x10 ⁻⁵	0.000561632	1.26247	M14042_5	14,042.5	Transferrin	13,761.41	TTR

C, vs. A-H, P<0.05 (t-test used for data analysis). C, normal group; A-H, post-foundation treatment group.

Peaks were mapped to the amino acid sequences based on the molecular weight; a 5% deviation in molecular weight was allowed. Eventually, a list of mapped proteins was compiled into an Excel table. The differences in predicted proteins between C, vs. A-Q (Table IV), C, vs. A-H (Table V), and A-Q, vs. A-H (Table VI) were determined. From Table IV, it was possible to estimate the differences in serum protein between the normal group and pre-foundation treatment group of patients with silicosis. From Table V, differences in serum protein were estimated between the normal group and post-foundation treatment group of patients with silicosis. From Table VI, it was possible to estimate the differences in serum protein between the patients with silicosis pre-foundation treatment and post-foundation treatment, respectively. These results demonstrate that complement c3 frag, amyloid β a4 protein, hepcidin, fibrinogen α -chain frags, plasma protease c1 inhibitor frag and SERPING1 are markers for silicosis.

Discussion

According to previous studies, the pathological changes of silicosis are the main contributor to pulmonary fibrosis (7-11). The exact pathogenesis of silicosis remains to be fully elucidated, however, substantial evidence shows the involvement of alveolar macrophages, cytokines, Clara cells, oxidative stress, the immune system of the body, and SiO₂ can lead to the occurrence and development of silicosis and be important in silicosis (12-17). The proteins that are differentially expressed in the serum of patients with silicosis are associated with silicosis (18,19). There have been no studies reporting on a correlation between protein expression differences and silicosis-associated mortality.

SELDI-TOF-MS, through chemical or biological treatment on the surface of the protein chip, laser desorption ionization mass spectrometry equipment and artificial intelligence data analysis processing software comprised of three parts, is a type of ultramicro, high flux, fully automatic protein screening technology for blood serum, urine, cell extracts or other samples directly added for identification (20). The present study used SELDI-TOF-MS to obtain original data and perform detailed analysis; the main analytical steps included the following: Data pre-processing, differences in peak screening, hierarchical cluster analysis, PCA analysis, construction of a decision tree model, and prediction based on differences in peaks corresponding to proteins. The advantages of this method over current diagnostic methods are the accuracy, speed and low cost of the application clinically (21).

The small number of patients is a limitation of the present study, and further investigations are required to corroborate the preliminary results. In particular, for patients with similar infectious diseases, it is possible that similar diseases have a similar pattern of proteins, which should be included in future investigations. A study demonstrated that there is no correlation between protein expression differences and silicosis-associated mortality (22). From the above results, the group of patients with silicosis pre- and post-foundation treatment and the normal group showed differences in proteins in serum. Therefore, patients with silicosis fibrosis and normal healthy individuals have differences in serum protein expression; in order to further understand these proteins and improve a simple and feasible method for the diagnosis of patients with

Table VI. A-Q, vs. A-H group differences protein prediction results.

SAMP_GRP	P-value	q-value	VIP	ID	MZ_SELDI	Predicted protein	Theoretical MZ	Gene name
M2001_69	0.013616562	0.299041303	1.81131	M2001_69	2001.69	Amyloid β a4 protein	1,953.10	APP
M2017_02	0.00271225	0.220851981	2.50106	M2017_02	2017.02	Amyloid β a4 protein	1,953.10	APP
M4144_81	0.00365448	0.220851981	1.30244	M4144_81	4144.81	Plasma protease c1 inhibitor frag	4,152.87	SERPING1
M4478_27	0.00365448	0.220851981	2.15012	M4478_27	4478.27	α -1-antichymotrypsin, α -1-antichymotrypsin frag	4,343.65	SERPINA3
M4861_39	0.048440933	0.437642642	1.32359	M4861_39	4861.39	Neurosecretory protein vgf frag	4,823.50	VGF
M5642_48	0.007295609	0.254483755	1.30207	M5642_48	5642.48	Fibrinogen α -chain frag	5,904.22	FGA
M7564_68	0.012079239	0.29154291	1.54656	M7564_68	7564.68	Osteopontin frag	7,658.19	SPPI

A-Q, vs. A-H, P<0.05 (t-test used for data analysis). A-Q, pre-foundation treatment group; A-H, post-foundation treatment group.

silicosis fibrosis, further investigations are to be performed for the identification of these differences in proteins. The aim of the present study was to offer occupational doctors a novel and alternative diagnostic test. The novel method, compared with present diagnostic strategies, of the proposed protein chip requires proper equipment, staff training and money. In conclusion, the present study identified complement c3 frag, amyloid β a4 protein, hepcidin, fibrinogen α -chain frags, plasma protease c1 inhibitor frag and SERPING1 are markers for silicosis.

Acknowledgements

Not applicable.

Funding

The present study was supported by the National Natural Science Funds (grant no. 81360416).

Availability of data and materials

The datasets used and/or analysis in the present study are available from the corresponding author on reasonable request.

Authors' contributions

GL contributed to the conception and design of the study, literature research and analysis, and manuscript preparation and review. JU contributed to the design of the study. CL contributed to the experimental and clinical studies, data acquisition and analysis, statistical analysis, manuscript preparation and manuscript review. XZ contributed to the experimental studies and clinical studies, data acquisition and analysis, statistical analysis and manuscript preparation. LN contributed to the experimental and clinical studies and data acquisition. YW, WWa, XW and WWe contributed to the experimental studies. YZ, NN and PH contributed to the clinical studies. XL contributed to the conception of the study, literature research and manuscript editing and review, and is a guarantor of integrity of the entire study.

Ethics approval and consent to participate

This study is approved by the Ethics Committee in the Fifth Affiliated Clinical Medical College of Xinjiang Medical University. Signed informed consent was obtained from all patients.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Jp NA, Imanaka M and Suganuma N: Japanese workplace health management in pneumoconiosis prevention. *J Occup Health* 59: 91-103, 2017.

2. Liu H, Tang Z, Yang Y, Weng D, Sun G, Duan Z and Chen J: Identification and classification of high risk groups for coal workers' Pneumoconiosis using an artificial neural network based on occupational histories: A retrospective cohort study. *BMC Public Health* 9: 366, 2009.
3. Fichtner-Feigl S, Fuss IJ, Young CA, Watanabe T, Geissler EK, Schlitt HJ, Kitani A and Strober W: Induction of IL-13 triggers TGF-beta1-dependent tissue fibrosis in chronic 2,4,6-trinitrobenzene sulfonic acid colitis. *J Immunol* 178: 5859-5870, 2007.
4. Mandal AK, Zhang Z, Ray R, Choi MS, Chowdhury B, Pattabiraman N and Mukherjee AB: Uteroglobin represses allergen-induced inflammatory response by blocking PGD2 receptor-mediated functions. *J Exp Med* 199: 1317-1330, 2004.
5. Ernst H, Rittinghausen S, Bartsch W, Creutzenberg O, Dasenbrock C, Görlitz BD, Hecht M, Kairies U, Muhle H, Müller M, *et al*: Pulmonary inflammation in rats after intratracheal instillation of quartz, amorphous SiO₂, carbon black, and coal dust and the influence of poly-2-vinylpyridine-N-oxide (PVNO). *Exp Toxicol Pathol* 54: 109-126, 2002.
6. Jiang PR, Cao Z, Qiu ZL, Pan JW, Zhang N and Wu YF: Plasma levels of TNF- α and MMP-9 in patients with silicosis. *Eur Rev Med Pharmacol Sci* 19: 1716-1720, 2015.
7. Lustgarten JL, Kimmel C, Ryberg H and Hogan W: EPO-KB: A searchable knowledge base of biomarker to protein links. *Bioinformatics* 24: 1418-1419, 2008.
8. Mahavadi P, Korfei M, Henneke I, Liebisch G, Schmitz G, Gochuico BR, Markart P, Bellusci S, Seeger W, Ruppert C and Guenther A: Epithelial stress and apoptosis underlie Hermansky-Pudlak syndrome-associated interstitial pneumonia. *Am J Respir Crit Care Med* 182: 207-219, 2010.
9. Carneiro PJ, Clevelario AL, Padilha GA, Silva JD, Kitoko JZ, Olsen PC, Capelozzi VL, Rocco PR and Cruz FF: Cruz bosutinib therapy ameliorates lung inflammation and fibrosis in experimental silicosis. *Front Physiol* 8: 159, 2017.
10. Liu H, Cheng Y, Yang J, Wang W, Fang S, Zhang W, Han B, Zhou Z, Yao H, Chao J and Liao H: BBC3 in macrophages promoted pulmonary fibrosis development through inducing autophagy during silicosis. *Cell Death Dis* 8: e2657, 2017.
11. Chen Y, Li C, Lu Y, Zhuang H, Gu W, Liu B, Liu F, Sun J, Yan B, Weng D and Chen J: IL-10-Producing CD1d hi CD5+ regulatory B cells may play a critical role in modulating immune homeostasis in silicosis patients. *Front Immunol* 8: 110, 2017.
12. Bhattacharya S, Dey A, Pal A, Kar S and Saha S: Silicosis in the form of progressive massive fibrosis: A diagnostic challenge. *Indian J Occup Environ Med* 20: 114-117, 2016.
13. Rosengarten D, Fox BD, Fireman E, Blanc PD, Rusanov V, Frucht O, Raviv Y, Shtraichman O, Saute M and Kramer MR: Survival following lung transplantation for artificial stone silicosis relative to idiopathic pulmonary fibrosis. *Am J Ind Med* 60: 248-254, 2017.
14. Mouratis MA and Aidinis V: Modeling pulmonary fibrosis with bleomycin. *Curr Opin Pulm Med* 17: 355-361, 2011.
15. Chiba Y, Kurotani R, Kusakabe T, Miura T, Link BW, Misawa M and Kimura S: Uteroglobin-related protein 1 expression suppresses allergic airway inflammation in mice. *Am J Respir Crit Care Med* 173: 958-964, 2006.
16. Liu P, Wang SX, Chen L, Wei MT, Liang XC, Wang YF and Tu ZG: Changes of Clara cell protein and surfactant protein-D in serum of patients with silicosis. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 25: 18-21, 2007 (In Chinese).
17. Anlar HG, Bacanlı M, İritiş S, Bal C, Kurt T, Tutkun E, Hinc Yılmaz O and Basaran N: Effects of occupational silica exposure on OXIDATIVE stress and immune system parameters in ceramic workers in TURKEY. *J Toxicol Environ Health A* 80: 688-696, 2017.
18. Zhu Z, Yang G, Wang Y, Yang J, Gao A, Niu P and Tian L: Suppression of thioredoxin system contributes to silica-induced oxidative stress and pulmonary fibrogenesis in rats. *Toxicol Lett* 222: 289-294, 2013.
19. Liu H, Fang S, Wang W, Cheng Y, Zhang Y, Liao H, Yao H and Chao J: Macrophage-derived MCP1 mediates silica-induced pulmonary fibrosis via autophagy. *Part Fibre Toxicol* 13: 55, 2016.
20. Fang SC, Zhang HT, Wang CY and Zhang YM: Serum CA125 and NSE: Biomarkers of disease severity in patients with silicosis. *Clin Chim Acta* 433: 123-127, 2014.
21. Lü GC, Yao JM and Zhang JW: Significance of detection of serum oxidant function in patients with silicosis. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 28: 52-53, 2010 (In Chinese).
22. Idec-Sadkowska I, Andrzejak R, Antonowicz-Juchniewicz J and Kaczmarek-Wdowiak B: Trials of casual treatment of silicosis. *Med Pr* 57: 271-280, 2006 (In Polish).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.