Repeated intravenous injection of recombinant human hepatocyte growth factor ameliorates liver cirrhosis but causes albuminuria in rats

KAZUNORI KUSUMOTO¹, AKIO IDO², AKIHIRO MORIUCHI², TOSHIYA KATSURA³, ILDEOK KIM², YUKA TAKAHAMA¹, MASATSUGU NUMATA¹, MAYUMI KODAMA¹, SATORU HASUIKE¹, KENJI NAGATA¹, HIROFUMI UTO¹, KEN-ICHI INUI³ and HIROHITO TSUBOUCHI^{1,2}

¹Department of Internal Medicine II, Faculty of Medicine, University of Miyazaki, 5200 Kihara, Kiyotake, Miyazaki 889-1692; ²Department of Experimental Therapeutics, Translational Research Center, ³Department of Pharmacy, Kyoto University Hospital, 54 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan

Received October 6, 2005; Accepted November 17, 2005

Abstract. Hepatocyte growth factor (HGF) is a promising agent for the treatment of liver cirrhosis because of its mitogenic and anti-fibrotic effects. We investigated the effect of recombinant human HGF (rh-HGF) on cirrhosis development; its pharmacokinetics and nephrotoxicity in rats with liver cirrhosis induced by 4-week treatment with dimethylnitrosamine (DMN). rh-HGF (0.3 mg/kg) was intravenously administered to rats once a day for 4 weeks in parallel with DMN treatment or twice a day for the last 2 weeks of DMN treatment. Repeated doses of rh-HGF increased the liver weight and serum albumin, and reduced serum ALT. The development of hepatic fibrosis was inhibited more efficiently by extended low-dose treatment with rh-HGF. In cirrhotic rats, serum levels of rh-HGF increased and clearance was decreased, leading to an increase in the area under the plasmaconcentration time curve and a decrease in the steady-state volume of distribution. Repeated doses of rh-HGF led to increased urinary albumin excretion, but no rh-HGF-treated animals developed increased serum creatinine levels. Urinary albumin excretion returned to baseline after the cessation of rh-HGF. These results suggest that extended treatment with rh-HGF is required for the attenuation of cirrhosis, and repeated doses of rh-HGF cause adverse effects in extra-hepatic organs. These issues must be resolved before the widespread application of rh-HGF in the treatment of liver cirrhosis.

E-mail: akioido@kuhp.kyoto-u.ac.jp

Introduction

Hepatocyte growth factor (HGF) was originally identified as a potent hepatocyte mitogen from the plasma of patients with fulminant hepatic failure (1,2). We have established an enzymelinked immunosorbent assay (ELISA) to measure human HGF in the serum, and we identified increased serum HGF levels in patients with various liver diseases (3). Additionally, serum HGF levels are a valuable prognostic tool in fulminant hepatic failure (4).

HGF is a multifunctional growth factor that acts as a mitogen, motogen, and morphogen for a variety of cells, including epithelial and endothelial cells (5-8). It is a major agent promoting hepatocyte proliferation (9-11) and acts in concert with transforming growth factor (TGF)- α and heparinbinding epidermal growth factor during liver regeneration (12,13). Additionally, HGF ameliorates hepatic injury via anti-apoptotic effects in animal models of fulminant hepatic failure (14-20) and attenuates hepatic fibrosis in animals with liver cirrhosis (21-25). Consequently, HGF is considered to not only induce liver regeneration but also to inhibit disease progression and ameliorate hepatic fibrosis in patients suffering from intractable liver diseases. Therefore, the clinical uses of recombinant human HGF (rh-HGF) in the treatment of fatal liver diseases, including fulminant hepatic failure, small-forsize grafts in living donor liver transplantation, and liver cirrhosis, are currently under investigation in Japan.

We recently showed that bolus intravenous injection of rh-HGF led to increased serum levels of human HGF, and intravenously administered HGF was primarily distributed to the liver (26). Additionally, despite its short half-life, a single intravenous injection of rh-HGF induced tyrosine phosphorylation of c-Met, a specific receptor for HGF, in liver tissue. However, intravenously administered rh-HGF is thought to be primarily metabolized in the liver, but the effect of liver injury on the pharmacokinetics of rh-HGF is not well understood.

In the present study, we investigated the effect of intravenously administered rh-HGF on hepatic fibrosis and its

Correspondence to: Dr Akio Ido, Department of Experimental Therapeutics, Translational Research Center, Kyoto University Hospital, 54 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto 606-8507, Japan

Key words: hepatocyte growth factor, liver cirrhosis, albuminuria, pharmacokinetics

pharmacokinetics in a rat model of liver cirrhosis. We further examined changes in urinary albumin excretion, a known adverse effect in extra-hepatic organs caused by the repeated administration of rh-HGF.

Materials and methods

Animals. Seven-week-old male SD rats were obtained from Japan SLC, Inc. (Shizuoka, Japan). The animals were maintained under a constant room temperature of 25°C and given free access to water and the indicated diet throughout the study. The protocol for animal studies was approved by the Ethics Committee of University of Miyazaki, Faculty of Medicine (Miyazaki, Japan). All animal experiments were performed after a 1-week acclimation period on a standard diet.

To induce liver cirrhosis, 1% dimethylnitrosamine (DMN) (Wako Pure Chemical Industries Ltd., Tokyo, Japan) dissolved in saline was given intraperitoneally at 1 ml per kg body weight for three consecutive days per week for 4 weeks. Rats were divided into four groups as follows: group C1 (n=7), rats treated with intravenous injection of phosphate-buffered saline (PBS) once a day for 4 weeks in parallel with DMN treatment; group H1 (n=8), rats treated with intravenous injection of rh-HGF (0.3 mg/kg) once a day for 4 weeks in parallel with DMN treatment; group C2 (n=9), rats treated with intravenous injection of PBS twice a day for the latter 2 weeks of DMN treatment; group H2 (n=10), rats treated with intravenous injection of rh-HGF (0.3 mg/kg) twice a day for the latter 2 weeks of DMN treatment.

After 4-week treatment with DMN, rats were sacrificed and serum albumin and alanine aminotransferase (ALT) were examined. The liver, spleen, and kidneys were immediately excised, and the wet weight of these organs was determined. Samples were subjected to histological analysis or frozen in liquid nitrogen and stored at -80°C until analysis. TGF-B levels in liver tissue were measured by a commercially available ELISA kit (R&D Systems Co., MN, USA).

Seventy-percent partial hepatectomy was performed according to a modification of the method of Higgins and Anderson (27). The rats were anesthetized with ether and a two-thirds partial hepatectomy was performed.

Measurement of serum human HGF. A silicone-rubber catheter (0.5x1.0 mm o.d.) was inserted into the jugular vein of normal, partially hepatectomized, or DMN-induced cirrhotic rats, and saline was administered continuously via the catheter using an infusion pump (0.1 ml/h) to prevent obstruction. rh-HGF (0.1 mg/kg) was injected into the femoral vein in <10 sec, and sequential blood samples were obtained via the catheter 5, 10, 20, 30, 60, 90 and 120 min after the injection.

Serum levels of human HGF were determined using a commercially available ELISA kit (Otsuka Pharmaceutical Co., Tokushima, Japan), in which only human HGF and not rat HGF is detected (3,28). Pharmacokinetic parameters were calculated using WinNonlin (Pharsight Co., CA, USA).

Histological examination for liver cirrhosis. Two 5-mm thick slices from the two major liver lobes (left lateral and median lateral lobes) were fixed in 10% formalin. After routine

processing of paraffin sections, $3-\mu$ m thick liver sections were made. To evaluate the degree of fibrosis, sections were stained with 0.1% (w/v) Sirius red F3BA (Sigma, St. Louis, MO, USA) for 1 h in a saturated aqueous solution (1.2% w/v) of picric acid (Wako); the final pH of the solution was 2.0. After staining, slides were rinsed for 2 min in 0.01 N HCl solution to remove unbound dye; following dehydration in alcohol, slides were mounted for observation. The Sirius redpositive areas were measured using an Olympus video micrometer, VM-30 (Olympus, Tokyo, Japan). The results were expressed as a fibrotic rate (%), which was calculated as the ratio of the Sirius red-positive area to the total area examined.

Measurement of urinary albumin excretion. Rats with DMNinduced liver cirrhosis were treated with PBS or rh-HGF as described above. Urine was periodically collected during PBS or DMN treatment, and urine albumin and creatinine were determined.

Repeated doses of rh-HGF (1.0 mg/kg/day) were intravenously administered to normal rats for 5 consecutive days per week for 2 weeks (days 1-5 and 8-12). Urinary levels of albumin and creatinine were examined during HGF treatment and for the following 2 weeks.

Statistical analysis. Unless otherwise specified, data are expressed as mean \pm SD. Statistical parameters were ascertained using StatView J-4.5 software (Abacus Concepts Ind., Berkeley, CA). Differences between means were assessed by using the unpaired Student's t-test. The significance level was set at p<0.05.

Results

Repeated intravenous injection of rh-HGF stimulated liver regeneration and attenuated liver injury in DMN-induced cirrhotic rats. To examine the effects of repeated intravenous injection of rh-HGF on liver regeneration, we measured the liver weights and serum albumin levels in rats with DMNinduced cirrhosis treated with either saline or rh-HGF (Table I). Treatment with rh-HGF for both 2 and 4 weeks concurrent with DMN administration significantly increased both liver weights and serum albumin levels compared with PBS-treated animals. Additionally, compared to untreated rats, serum ALT levels were elevated in DMN-induced cirrhotic rats treated with PBS (groups C1 and C2) (Table I), and treatment with rh-HGF for 2 weeks significantly decreased ALT levels (C2 vs. H2). While ALT values for animals treated with rh-HGF for 4 weeks (group H1) were reduced compared to group C1, this difference was not significant. The reason for the differences in 2- and 4-week rh-HGF treatment is unclear. Nevertheless, repeated intravenous injection of rh-HGF not only stimulated liver regeneration but also attenuated liver injury in DMNinduced cirrhotic rats.

Prolonged treatment with rh-HGF efficiently attenuated hepatic fibrosis in DMN-induced cirrhotic rats. Liver weight and serum ALT are good markers for liver regeneration and injury, but the pathological hallmark of cirrhosis is extensive liver fibrosis. Thus, we wished to examine the effect of rh-HGF on hepatic fibrosis. When rh-HGF was intravenously

	Group			Group		
	C1 (n=7)	H1 (n=8)	p-value	C2 (n=9)	H2 (n=10)	p-value
Body weight (BW g)	270.1±64.6	319.0±32.5	0.105	305.3±38.2	311.0±22.9	0.705
Liver (g/100 g BW)	2.77±1.10	3.91±0.40	0.034	3.32±0.93	4.65±0.52	0.003
Kidney, right (g/100 g BW)	0.42±0.09	0.39±0.05	0.405	0.41 ± 0.07	0.43±0.06	0.584
Albumin (g/dl)	1.81±0.50	2.34±0.19	0.031	2.06±0.43	2.92±0.18	0.0002
ALT (IU/l)	136.7±99.4	78.4±18.6	0.175	94.8±34.2	67.6±12.5	0.048

Table I. Effect of rh-HGF administration on body, liver and right kidney weight, and serum levels of albumin and ALT in rats with DMN-induced cirrhosis.

Group C1, rats treated with PBS once a day for 4 weeks in parallel with DMN administration; group H1, rats treated with 0.3 mg/kg of rh-HGF once a day for 4 weeks in parallel with DMN administration; group C2, rats administered PBS twice a day for the last 2 weeks of DMN treatment; group H2, rats administered 0.3 mg/kg of rh-HGF twice a day for the last 2 weeks of DMN treatment. Results are represented as the mean ± SD.



Figure 1. Representative microscopic appearance of liver tissue in DMNinduced cirrhotic rats treated with or without rh-HGF. Rats with DMNinduced cirrhosis were treated with rh-HGF for 4 or 2 weeks, and liver tissue was stained with Sirius red as described in Materials and methods. (A) Group C1; (B) Group H1; (C) Group C2; (D) Group H2 (magnification x40).

injected for 4 or 2 weeks in conjunction with DMN administration, the development of hepatic fibrosis was substantially inhibited (Fig. 1). Sirius red, a dye specific for fibrotic areas, was used to quantitate the extent of liver fibrosis in PBSor rh-HGF-treated rats (Table II). Repeated injection of rh-HGF for 4 weeks concurrent with DMN treatment (group H1) significantly reduced the areas of hepatic fibrosis compared with PBS-treated rats (group C1), but there were no significant differences between the fibrotic areas of rats treated with PBS and those of rats treated with rh-HGF for 2 weeks (groups C2 and H2, respectively). We also examined the hepatic tissue levels of TGF-B, a key mediator of cirrhosis development (Table II). Administration of rh-HGF reduced TGF-B levels in liver tissue, regardless of the treatment period, but these differences did not achieve statistical significance. Pharmacokinetics of rh-HGF administered intravenously in a bolus in normal, partially hepatectomized, and cirrhotic rats. HGF is primarily metabolized by the liver, but patients most likely to receive rh-HGF have severe liver disease with possible impaired metabolism of a variety of compounds, including rh-HGF. Therefore, we examined changes in serum human HGF concentration sequentially after the single intravenous injection of rh-HGF in rats with normal or abnormal liver function, and we calculated pharmacokinetic parameters (Table III). When compared with normal rats, the serum levels of human HGF were elevated 5 min after injection and the terminal elimination half-life (T_{1/2terminal}) of rh-HGF was prolonged in rats with DMN-induced cirrhosis. Consequently, an increase in the area under the plasma-concentration time curve $(AUC_{0-\infty})$ and a decrease in clearance (CL) were observed in cirrhotic rats. Also, the steady-state volume of distribution (Vdss) was reduced to approximately 50%, and the mean resident time $(MRT_{0-\infty})$ was prolonged. The degree of pharmacokinetic change in rats with DMN-induced cirrhosis was virtually the same as that seen in rats with 70% partial hepatectomy.

Repeated administration of rh-HGF increases albuminuria. While most of the physiological effects of rh-HGF are confined to the liver, some changes in kidney function are seen in animals treated with rh-HGF. As a measure for kidney dysfunction, we examined urinary albumin excretion during the treatment with rh-HGF (Fig. 2). When rats were treated with rh-HGF (0.3 mg/kg, once a day) in parallel with DMN administration, the urinary albumin excretion gradually increased for 7 days from the beginning of HGF treatment, and moderate to high levels of albuminuria were observed during the following 14 days (Fig. 2A). Repeated injection of rh-HGF (0.3 mg/kg, twice a day) for days 15-28 of DMN administration rapidly increased the urinary excretion of albumin, and marked albuminuria continued throughout the experimental period. Conversely, kidney weight was not affected by either 2- or 4-week treatment with rh-HGF (Table I).

		Group			Group		
	C1 (n=7)	H1 (n=8)	p-value	C2 (n=9)	H2 (n=10)	p-value	
Fibrosis (%)	10.30±5.02	4.53±1.64	0.022	7.31±3.84	5.71±2.77	0.319	
TGF-β (ng/g tissue)	30.77±10.26	22.65±6.00	0.098	34.69±12.06	28.03±11.91	0.244	

Table II. Effect of rh-HGF administration on hepatic fibrosis and hepatic levels of TGF-ß in DMN-induced cirrhotic rats.

Results are represented as the mean \pm SD.

Table III. Pharmacokinetics of rh-HGF in normal, partially hepatectomized, and cirrhotic rats.

	Normal (n=4)	DMN-LC (n=5)	70% PH (n=4)
$\overline{C_5^a (ng/ml)}$	89.71±20.57	255.4±181.4	341.2±55.1
T _{1/2terminal} (min)	27.88±2.00	35.41±18.74	39.49±13.79
$AUC_{0-\infty}$ ($\mu g \cdot min/ml$)	1.685±0.342	13.78±16.48	10.84±2.24
CL (ml/min/kg)	62.09±13.82	20.76±16.73	9.511±1.864
Vdss (ml/kg)	903.7±253.5	453.4±174.6	419.6±144.6
$MRT_{0-\infty}$ (min)	14.44±1.07	41.71±33.20	43.79±9.87

Results are represented as the mean \pm SD. ^aConcentration of serum human HGF 5 min after injection. DMN-LC, rats with DMN-induced liver cirrhosis; 70% PH, rats with 70% partial hepatectomy; T_{1/2terminal}, terminal elimination half-life; AUC_{0-∞}, the area under the plasma-concentration time curve; Vdss, steady-state volume of distribution; MRT_{0-∞}, mean resident time.

To clarify whether HGF-induced albuminuria was reversible, we injected rh-HGF (1.0 mg/kg/day) via tail veins into normal rats for different periods of time, and monitored the urinary albumin excretion (Fig. 2B). Administration of rh-HGF for 5 days gradually increased albuminuria and, when rh-HGF injection was stopped on days 6 and 7, the albumin excretion reduced. When rh-HGF was subsequently administered on days 8-12, albuminuria once again worsened but, after complete cessation of rh-HGF injection, the albumin excretion gradually returned to baseline levels by days 21-28. On day 28, serum creatinine levels were unchanged compared to baseline indicating minimal permanent kidney damage (data not shown).

Discussion

HGF is produced by mesenchymal cells as an inactive precursor, pro-HGF (29-31), and the mature protein plays important roles in the regeneration and repair of various organs, including the liver, gastrointestinal tract, and kidney. Following tissue injury, pro-HGF is converted to an active heterodimer consisting of light and heavy chains by the specific serine protease HGF activator (HGFA) (32-35). Additionally, HGFA inhibitor (HAI)-1, which was first purified as an inhibitor of HGFA (36), regulates HGFA activity at sites of tissue injury (37). Recently, Yanagida *et al* reported that the administration of recombinant human HGFA stimulated the regeneration of cirrhotic liver after partial hepatectomy through the activation of endogenous

pro-HGF (38). In the present study, we administered rh-HGF (0.3 mg/kg) in an already active heterodimeric form into cirrhotic rats once a day for 4 weeks (group H1) or twice a day for the last 2 weeks (group H2) of a 4-week DMN treatment. The total amount of rh-HGF administered throughout the experimental period was equivalent between both groups. Administration of rh-HGF stimulated liver regeneration and ameliorated hepatic fibrosis, and treatment with a lower dose for a longer period of time (group H1) was more effective at inhibiting fibrosis than a shorter therapeutic course with a higher dose (group H2). HGF decreases both extracellular matrix (ECM) deposition and mortality in various models of liver cirrhosis in rats (21,22,24,25), and hepatic expression of TGF-ß, a key mediator of the development of liver cirrhosis, is suppressed by HGF treatment (21,23). We examined hepatic TGF-ß levels but, although a trend toward decreased TGF-ß levels was seen in animals treated with rh-HGF, our results were not statistically significant. In contrast, HGF apparently enhances TGF-ß production and DNA synthesis in activated c-Met expressing hepatic stellate cells (HSCs) isolated from rats treated with carbon tetrachloride (39). These seemingly contradictory effects of HGF on TGF-ß production may be explained by in vitro or in vivo experiments and/or differences in animal models of liver cirrhosis.

Hepatic fibrosis is a long-term consequence of chronic or repeated liver injury. Two different cell populations play a key role in the development of liver cirrhosis. Once hepatic injury occurs, quiescent HSCs are activated and undergo phenotypic changes, developing into myofibroblast-like cells



Figure 2. Sequential changes in urinary excretion of albumin in cirrhotic or normal rats treated with repeated intravenous injection of rh-HGF. (A) Administration of rh-HGF induced an increase in albuminuria in rats with DMN-induced liver cirrhosis.•, group H1 (n=4); \triangle , group H2. (B) One mg/kg of rh-HGF was intravenously injected into normal rats (n=4) for days 1-5 and 8-12. Administration of rh-HGF increased urinary excretion of albumin. However, rh-HGF-induced albuminuria was a reversible change, which recovered after cessation of rh-HGF.

able to produce ECM proteins (40,41). In addition to activated HSCs, portal myofibroblasts also express ECM proteins involved in fibrotic changes of the liver (42,43). HGF stimulates the expression of matrix metalloproteinase-1 in human HSCs (44), and HGF inhibits cell growth and induces apoptosis in liver myofibroblasts, leading to the amelioration of hepatic fibrosis in cirrhotic rats (45). In the present study, we observed a more pronounced amelioration of increases in serum ALT levels in 2-week-treated rats (group H2) compared to 4-weektreated rats (group H1). Because the limited administration of HGF efficiently ameliorated experimental hepatic injury and rescued animals from lethal hepatic failure through antiapoptotic effects (15-18), short-term administration and/or a high dose of rh-HGF appears sufficient to prevent tissue injury and resulting ALT release in DMN-induced cirrhotic rats. The ability of rh-HGF to prevent liver injury, maintain hepatic function and suppress the development of cirrhosis is probably mediated, at least in part, by the inhibition of HSC activation. Collectively, these data indicate that HGF prevents the development of cirrhosis through a variety of different mechanisms.

The liver is a primary target of intravenously administered rh-HGF in normal rats, and serum levels of rh-HGF were

elevated in rats with partial hepatectomy or a choline-deficient L-amino acid-defined diet compared to normal rats (26). These findings suggest that rh-HGF administered intravenously is primarily metabolized by the liver. In this study, we further investigated the pharmacokinetics of intravenously injected rh-HGF in rats with DMN-induced cirrhosis. Serum levels of rh-HGF 5 min after injection were elevated, and the CL was reduced, resulting in an increase in the $AUC_{\scriptscriptstyle 0\mathchar`omega}$ and $MRT_{\scriptscriptstyle 0\mathchar`omega}$ and a decrease in the Vdss in cirrhotic rats. HGF in the circulating plasma is efficiently extracted by the liver compared with other HGF target organs, and the liver is responsible for approximately 70% of the overall elimination of rh-HGF (46,47). Furthermore, HGF uptake is three times higher by hepatocytes than other liver non-parenchymal cells (48). Additionally, HGF receptor expression is saturable (48), and elimination of HGF from the systemic circulation is affected by changes in HGF receptor density on the liver cell surface (49). Therefore, a decrease in the number of hepatocytes and a change in c-Met expression in cirrhotic rat livers possibly affected the pharmacokinetics of rh-HGF. Also, the administered dose of rh-HGF (0.1 mg/kg) may be in excess leading to the saturation of receptor-mediated endocytosis and significant increases in serum concentration of rh-HGF.

When rh-HGF was injected intravenously as a bolus, the kidneys, spleen and adrenal glands as well as the liver contained large amounts of rh-HGF (26), and higher concentrations of rh-HGF in the systemic circulation increase the potential of adverse effects. In the present study, an increase in urinary albumin excretion was observed in cirrhotic rats treated with rh-HGF (0.3 mg/kg, once or twice a day). Although repeated doses of rh-HGF (1.0 mg/kg/day) increased urinary albumin excretion in normal rats, the effect was reversible, and treatment with rh-HGF did not affect serum creatinine levels, indicating the absence of permanent kidney injury. Further studies are needed to clarify the renal involvement following repeated doses of rh-HGF. However, since an increase in urinary albumin excretion is more sensitive to changes in kidney function than proteinuria (data not shown), the measurement of urinary albumin may be a useful marker for the detection of HGF-induced nephrotoxicity in clinical applications of rh-HGF.

In the present study, we showed that, although short-term administration of rh-HGF stimulated liver regeneration and ameliorated hepatic injury, prolonged treatment with rh-HGF was more effective in preventing cirrhosis development. However, elevated serum concentrations of rh-HGF in cirrhotic rats have the potential to increase the incidence of adverse events. Therefore, the development of a tissue-specific and/or sustained release delivery system is required for clinical applications of rh-HGF in the treatment of liver cirrhosis. The development of these technologies may also allow the application of rh-HGF in the treatment of diseases of extrahepatic organs, such as inflammatory bowel disease, pulmonary fibrosis, or neurodegenerative disease.

Acknowledgements

We thank Ms. Yuko Nakamura for her technical assistance. This work was supported in part by Grants-in-Aid from the Ministry of Health, Labor and Welfare of Japan.

References

- 1. Gohda E, Tsubouchi H, Nakayama H, *et al*: Human hepatocyte growth factor in plasma from patients with fulminant hepatic failure. Exp Cell Res 166: 139-150, 1986.
- 2. Gohda E, Tsubouchi H, Nakayama H, *et al*: Purification and partial characterization of hepatocyte growth factor from plasma of a patient with fulminant hepatic failure. J Clin Invest 81: 414-419, 1988.
- Tsubouchi H, Niitani Y, Hirono S, *et al*: Levels of the human hepatocyte growth factor in serum of patients with various liver diseases determined by an enzyme-linked immunosorbent assay. Hepatology 13: 1-5, 1991.
- 4. Tsubouchi H, Kawakami S, Hirono S, *et al*: Prediction of outcome in fulminant hepatic failure by serum human hepatocyte growth factor. Lancet 340: 307, 1992.
- Michalopoulos GK: Liver regeneration: molecular mechanisms of growth control. FASEB J 4: 176-185, 1990.
- Dignass AU, Lynch-Devaney K and Podolsky DK: Hepatocyte growth factor/scatter factor modulates intestinal epithelial cell proliferation and migration. Biochem Biophys Res Commun 202: 701-709, 1994.
- 7. Bussolino F, Di Renzo MF, Ziche M, *et al*: Hepatocyte growth factor is a potent angiogenic factor which stimulates endothelial cell motility and growth. J Cell Biol 119: 629-641, 1992.
- Zarneger R and Michalopoulos GK: The many faces of hepatocyte growth factor: from hepatopoiesis to hematopoiesis. J Cell Biol 129: 1177-1180, 1995.
- Fujiwara K, Nagoshi S, Ohno A, *et al*: Stimulation of liver growth by exogenous human hepatocyte growth factor in normal and partially hepatectomized rats. Hepatology 18: 1443-1449, 1993.
- Ishiki Y, Ohnishi H, Muto Y, Matsumoto K and Nakamura T: Direct evidence that hepatocyte growth factor is a hepatotrophic factor for liver regeneration and has a potent antihepatitis effect *in vivo*. Hepatology 16: 1227-1235, 1992.
 Ishii T, Sato M, Sudo K, *et al*: Hepatocyte growth factor
- Ishii T, Sato M, Sudo K, *et al*: Hepatocyte growth factor stimulates liver regeneration and elevates blood protein level in normal and partially hepatectomized rats. J Biochem 117: 1105-1112, 1995.
- 12. Webber EM, Fitzgerald MJ, Brown PI, Barlett MH and Fausto N: Transforming growth factor- α expression during liver regeneration after partial hepatectomy and toxic injury, and potential interactions between transforming growth factor- α and hepatocyte growth factor. Hepatology 18: 1422-1431, 1993.
- 13. Moriuchi A, Hirono S, Ido A, *et al*: Additive and inhibitory effects of simultaneous treatment with growth factors on DNA synthesis through MAPK pathway and G1 cyclins in rat hepatocytes. Biochem Biophys Res Commun 280: 368-273, 2001.
- Okano J, Shiota G and Kawasaki H: Protective action of hepatocyte growth factor for acute liver injury caused by D-galactosamine in transgenic mice. Hepatology 26: 1241-1249, 1997.
- Kosai K, Matsumoto K, Nagata S, Tsujimoto Y and Nakamura T: Abrogation of Fas-induced fulminant hepatic failure in mice by hepatocyte growth factor. Biochem Biophys Res Commun 244: 683-690, 1998.
- Masunaga H, Fujise N, Shiota A, *et al*: Preventive effects of the deleted form of hepatocyte growth factor against various liver injuries. Eur J Pharmacol 342: 267-279, 1998.
- Kosai K, Matsumoto K, Funakoshi H and Nakamura T: Hepatocyte growth factor prevents endotoxin-induced lethal hepatic failure. Hepatology 30: 151-159, 1999.
 Mori I, Tsuchida A, Taiji M and Noguchi H: Hepatocyte growth
- Mori I, Tsuchida A, Taiji M and Noguchi H: Hepatocyte growth factor protects mice against anti-Fas antibody-induced death. Med Sci Res 27: 355-359, 1999.
- Nomi T, Shiota G, Isono M, Sato K and Kawasaki H: Adenovirusmediated hepatocyte growth factor gene transfer prevents lethal liver failure in rats. Biochem Biophys Res Commun 278: 338-343, 2000.
- 20. Xue F, Takahara T, Yata Y, *et al*: Attenuated acute liver injury in mice by naked hepatocyte growth factor gene transfer into skeletal muscle with electroporation. Gut 50: 558-562, 2002.
- 21. Yasuda H, Imai E, Shiota A, Fujise N, Morinaga T and Higashio K: Antifibrogenic effect of a deletion variant of hepatocyte growth factor on liver fibrosis in rats. Hepatology 24: 636-642, 1996.
- Matsuda Y, Matsumoto K, Yamada A, *et al*: Prevention and therapeutic effects in rats of hepatocyte growth factor infusion on liver fibrosis/cirrhosis. Hepatology 26: 81-89, 1997.

- Ueki T, Kaneda Y, Tsutsui H, et al: Hepatocyte growth factor gene therapy of liver cirrhosis in rats. Nat Med 5: 226-230, 1999.
- 24. Sato M, Kakubari M, Kawamura M, Sugimoto J, Matsumoto K and Ishii T: The decrease in total collagen fibers in the liver by hepatocyte growth factor after formation of cirrhosis induced thioacetamide. Biochem Pharmacol 59: 681-690, 2000.
- 25. Oe S, Fukunaka Y, Hirose T, Yamaoka Y and Tabata Y: A trial on regeneration therapy of rat liver cirrhosis by controlled release of hepatocyte growth factor. J Control Release 307: 146-151, 2003.
- Ido A, Moriuchi A, Kim ID, *et al*: Pharmacokinetic study of recombinant human hepatocyte growth factor administered in a bolus intravenously or via portal vein. Hepatol Res 30: 175-181, 2004.
- Higgins GM and Anderson RM: Experimental pathology of the liver: restoration of the liver of the white rat following partial surgical removal. Arch Pathol 12: 186-202, 1931.
- Kawakami S, Tsubouchi H, Nakagawa S, *et al*: Expression of hepatocyte growth factor in normal and carbon tetrachloridetreated monkeys. Hepatology 20: 1255-1260, 1994.
 Gohda E, Kataoka H, Tsubouchi H, Daikuhara Y and
- 29. Gohda E, Kataoka H, Tsubouchi H, Daikuhara Y and Yamamoto I: Phorbol ester-induced secretion of human hepatocyte growth factor by human skin fibroblasts and its inhibition by dexamethasone. FEBS Lett 301: 107-110, 1992.
- 30. Tamura M, Arakaki N, Tsubouchi H, Takada H and Daikuhara Y: Enhancement of human hepatocyte growth factor production by interleukin-1α ar-α by fibroblasts in culture. J Biol Chem 268: 8140-8145, 1993.
- 31. Gohda E, Takebe T, Sotani T, Nakamura S, Minowada J and Yamamoto I: Induction of hepatocyte growth factor by interleukin-γ in human leukemia cells. J Cell Physiol 174: 107-114, 1998.
- 32. Miyazawa K, Shimomura T, Kitamura A, Kondo J, Morimoto Y and Kitamura N: Molecular cloning and sequence analysis of the cDNA for a human serine protease responsible for activation of hepatocyte growth factor. Structural similarity of the protease precursor to blood coagulation factor XII. J Biol Chem 268: 10024-10028, 1993.
- Miyazawa K, Shimomura T, Naka D and Kitamura N: Proteolytic activation of hepatocyte growth factor in response to tissue injury. J Biol Chem 269: 8966-8970, 1994.
- 34. Miyazawa K, Shimomura T and Kitamura N: Activation of hepatocyte growth factor in the injured tissues is mediated by hepatocyte growth factor activator. J Biol Chem 271: 3615-3618, 1996.
- Shimomura T, Miyazawa K, Komiyama Y, Hiraoka H, Naka D, Morimoto Y and Kitamura N: Activation of hepatocyte growth factor by two homologous proteases, blood-coagulation factor XIIa and hepatocyte growth factor activator. Eur J Biochem 229: 257-261, 1995.
 Shimomura T, Denda K, Kitamura A, *et al*: Hepatocyte growth
- 36. Shimomura T, Denda K, Kitamura A, *et al*: Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor. J Biol Chem 272: 6370-6376, 1997.
- 37. Kataoka H, Shimomura T, Kawaguchi T, *et al*: Hepatocyte growth factor activator inhibitor type-1 is a specific cell surface binding protein of hepatocyte growth factor activator (HGFA) and regulates HGFA activity in the pericellular microenvironment. J Biol Chem 275: 40453-40462, 2000.
- 38. Yanagida H, Kaibori M, Hijikawa T, Kwon A-H, Kamiyama Y and Okumura T: Administration of rhHGF-activator via portal vein stimulates the regeneration of cirrhotic liver after partial hepatectomy in rats. J Surg Res doi:10.1016/j.jss.2005.08.002.
- 39. Ikeda H, Nagoshi S, Ohno A, Yanase M, Maekawa H and Fujiwara K: Activated stellate cells express c-met and respond to hepatocyte growth factor to enhance transforming growth factor β1 expression and DNA synthesis. Biochem Biophys Res Commun 250: 769-775, 1998.
- Alcolado R, Arther MJP and Iredale JP: Pathogenesis of liver fibrosis. Clin Sci 92: 103-112, 1997.
- 41. Wu J and Zern MA: Hepatic stellate cells: a target for the treatment of liver fibrosis. J Gastroenterol 35: 665-672, 2000.
- Tuchweber B, Desmouliere A, Bochaton-Piallat ML, Rubbia Brandt L and Gabiani G: Proliferation and phenotypic modulation of portal fibroblasts in the early stages of cholestatic fibrosis in the rats. Lab Invest 74: 265-278, 1996.
 Knittel T, Kobold D, Saile B, Grundmann A, Neubauer K,
- 43. Knittel T, Kobold D, Saile B, Grundmann A, Neubauer K, Piscaglia F and Ramadori G: Rat liver myofibroblasts and hepatic stellate cells: different cell populations of the fibroblast lineage with fibronogenic potential. Gastroenterology 117: 1205-1221, 1999.

- 44. Ozaki I, Zhao G, Mizuta T, *et al*: Hepatocyte growth factor induces collagenase (matrix metalloproteinase-1) via the transcription factor Ets-1 in human hepatic stellate cell line. J Hepatol 36: 169-178, 2002.
- 45. Kim W-H, Matsumoto K, Bessho K and Nakamura T: Growth inhibition and apoptosis in liver myofibroblasts promoted by hepatocyte growth factor leads to resolution from liver cirrhosis. Am J Pathol 166: 1017-1028, 2005.
 46. Liu KX, Kato Y, Narukawa M, *et al*: Importance of the liver in
- 46. Liu KX, Kato Y, Narukawa M, *et al*: Importance of the liver in plasma clearance of hepatocyte growth factor in rats. Am J Physiol 263: G645-G649, 1992.
- 47. Kato M, Kato Y, Nakamura T and Sugiyama Y: Efficient extraction by the liver governs overall elimination of hepatocyte growth factor in rats. J Pharmacol Exp Ther 290: 373-379, 1999.
- 48. Liu KX, Kato Y, Terasaki T, Aoki S, Okumura K, Nakamura T and Sugiyama Y: Contribution of parenchymal and non-parenchymal liver cells to the clearance of hepatocyte growth factor from the circulation in rats. Pharm Res 12: 1737-1740, 1995.
- Liu KX, Kato Y, Kino I, Nakamura T and Sugiyama Y: Ligandinduced downregulation of receptor-mediated clearance of hepatocyte growth factor in rats. Am J Physiol 275: E835-E842, 1998.