

# Cellular Na<sup>+</sup> handling mechanisms involved in airway smooth muscle contraction (Review)

BETTINA SOMMER<sup>1</sup>, EDGAR FLORES-SOTO<sup>2</sup> and GEORGINA GONZALEZ-AVILA<sup>3</sup>

<sup>1</sup>Department of Bronchial Hyperreactivity, National Institute of Respiratory Diseases 'Ismael Cosío Villegas', CP 14080 Mexico City; <sup>2</sup>Department of Pharmacology, Faculty of Medicine, National Autonomous University of Mexico, CP 04510 Mexico City; <sup>3</sup>Biomedical Oncology Laboratory, Department of Chronic-Degenerative Diseases, National Institute of Respiratory Diseases 'Ismael Cosío Villegas', CP 14080 Mexico City, Mexico

Received December 22, 2016; Accepted April 20, 2017

DOI: 10.3892/ijmm.2017.2993

**Abstract.** A decrease in bronchial diameter is designated as bronchoconstriction (BC) and impedes the flow of air through the airway. Asthma is characterized by inflammation of the airways, reversible BC and nonspecific hyperreactivity. These last two symptoms are dependent on airway smooth muscle. Stimuli that trigger contraction can be characterized as chemical (neurotransmitters, cytokines and terpenoids) and physical (volume inspired, air pressure). Both stimuli activate signaling pathways by acting on membrane proteins and facilitating the passage of ions through the membrane, generating a voltage change and a subsequent depolarization. Na<sup>+</sup> plays an important role in preserving the resting membrane potential; this ion is extracted from the cells by the Na<sup>+</sup>/K<sup>+</sup> ATPase (NKA) or introduced into the cytoplasm by the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger (NCX). During depolarization, Na<sup>+</sup> appears to accumulate in specific regions beneath the plasma membrane, generating local concentration gradients which determine the handling of Ca<sup>2+</sup>. At rest, the smooth muscle has a basal tone that is preserved by the continuous adjustment of intracytoplasmic concentrations of Ca<sup>2+</sup> and Na<sup>+</sup>. At homeostasis, the Na<sup>+</sup> concentration is primarily dependent on three structures: the NKA, the NCX and non-specific cation channels (NSCC). These three structures, their functions and the available evidence of the probable role of Na<sup>+</sup> in asthma are described in the present review.

## Contents

1. Introduction
2. Na<sup>+</sup>/K<sup>+</sup> ATPase

3. Na<sup>+</sup>/Ca<sup>2+</sup> exchanger
4. Non-specific cation channels
5. Na<sup>+</sup> voltage dependent currents in airway smooth muscle
6. Evidence of Na<sup>+</sup> participation in asthma
7. The paradox of exercise-induced bronchoconstriction
8. Conclusion

## 1. Introduction

Airway bronchoconstriction (BC) and hyperresponsiveness are hallmarks of asthma, a chronic, irreversible inflammatory disease (1). Asthma-related BC can be induced by exercise, infection or allergen exposure according to the phenotypic characteristics of each patient (2) and is frequently relieved by inhaled corticosteroids and β<sub>2</sub> adrenergic agonists (3). Basically, it can be described as the consequence of the airway smooth muscle (ASM) contraction developed by increases in intracellular Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>i</sub>) in response to agonists or membrane depolarization (4). On the other hand, asthma patients also show airway hyperresponsiveness commonly evaluated by inhalation of methacholine or histamine and this exacerbated ASM response also involves cellular Ca<sup>2+</sup> handling mechanisms (5-8). In addition to Ca<sup>2+</sup>, Na<sup>+</sup> is paramount in ASM contraction. The literature available, although not yet conclusive, indicates a possible link between altered ASM cell Na<sup>+</sup> handling mechanisms and BC and this evidence is discussed in the present review.

It is well known that excitable cells possess a characteristic resting membrane potential, a state of equilibrium between inner and outer ionic concentrations. In airway myocytes, resting membrane potential fluctuates between -40 and -50 mV (9) and reflects many membrane mechanisms that keep the balance between internal and external ionic concentrations. In this tissue, stimuli that trigger contraction can be characterized as chemical (neurotransmitters, cytokines and terpenoids) and physical (inspired air volume, air pressure and temperature). Although different in nature, both types of stimuli act on membrane proteins that facilitate the passage of ions in and out, causing a current and a voltage change. Na<sup>+</sup> transit through the cell membrane has been known to play the leading role

---

*Correspondence to:* Dr Bettina Sommer, Department of Bronchial Hyperreactivity, National Institute of Respiratory Diseases 'Ismael Cosío Villegas', Calzada de Tlalpan 4502, Col. Sección XVI, CP 14080 Mexico City, Mexico  
E-mail: bsommerc@hotmail.com

*Key words:* sodium, calcium, smooth muscle, asthma




Figure 1. Cellular  $\text{Na}^+$  homeostatic mechanisms in airway smooth muscle. The  $\text{Na}^+/\text{K}^+$  ATPase (NKA) introduces 2  $\text{K}^+$  and extracts 3  $\text{Na}^+$  ions spending 1 ATP. The  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX), by following an electrochemical gradient, introduces 3  $\text{Na}^+$  and extrudes 1  $\text{Ca}^{2+}$ . Non-specific cation channels (NSCC) allow  $\text{Na}^+$  and  $\text{Ca}^{2+}$  entrance and the resting membrane potential in healthy airway smooth muscle is maintained by NSCC recognized as TRPC3 (56). SR, sarcoplasmic reticulum; SERCA, sarcoplasmic reticulum  $\text{Ca}^{2+}$  ATPase.

in preserving membrane potential at rest. It is extracted from the cells by the  $\text{Na}^+/\text{K}^+$  ATPase (NKA) pump or introduced into the cytoplasm by the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX) (Fig. 1). At rest, ASM has a basal tension (tone) which results from the continuous adjustment of  $\text{Ca}^{2+}$  and  $\text{Na}^+$  intracytoplasmic concentrations.

During membrane depolarization,  $\text{Na}^+$  seems to accumulate in regions beneath the plasma membrane, generating local concentration gradients that influence cell  $\text{Ca}^{2+}$  handling mechanisms. The local  $\text{Na}^+$  concentration increases rely on three proteins: the NKA, the NCX and non-specific cation channels (NSCC). These proteins are located in precise membrane regions and their function reflects as an augmentation in  $\text{Na}^+$  concentration and a consequent change in the amount of  $\text{Ca}^{2+}$  (10). Increases in  $[\text{Ca}^{2+}]_i$  regulate diverse processes depending on magnitude, duration and cytoplasmic region; in ASM, such cellular processes range from contraction, including protein synthesis, to apoptosis (programmed cell death) (11). Therefore, complex and refined methods to handle  $[\text{Ca}^{2+}]_i$  have evolved. In this regard, the sarcoplasmic reticulum (SR) stores most of the ASM cellular  $\text{Ca}^{2+}$ . This ion is released during contraction, but the SR is never completely depleted from its content. After its emptying, the SR  $\text{Ca}^{2+}$  refilling seems to involve increases in  $\text{Na}^+$  concentrations in sites nearby. This refilling is also related to smooth muscle sustained contraction (12-18). Recent findings in our laboratory, demonstrate that  $\text{Na}^+$  permeates through the L-type voltage dependent  $\text{Ca}^{2+}$  channels when ASM is stimulated with carbachol (19). It is feasible that this  $\text{Na}^+$  entrance contributes to depolarization, favoring  $\text{Ca}^{2+}$  entry for SR refilling and sustained contraction maintenance (Fig. 2).

## 2. $\text{Na}^+/\text{K}^+$ ATPase

The NKA on the plasma membrane extracts 3  $\text{Na}^+$  ions from the cytoplasm and simultaneously introduces 2  $\text{K}^+$  against both ions' concentration gradient. Spending ATP is required, hence the name of ATPase. This pump is very sensitive to inhibition by cardioactive steroids, particularly the group of digitalis and ouabain used to treat congestive heart failure

and arrhythmias (20). Abundant sources report the presence of a 'serum factor' (most probably ouabain) that inhibits the function of NKA (21-26). Endogenous ouabain has been characterized as a steroidal glycoside found in blood plasma and is synthesized by the adrenal glands, hypothalamus and pituitary (27-33). *In vitro*, the release of ouabain in primary cultures of bovine adrenocortical cells is favored by the adrenocorticotrophic hormone (ACTH), angiotensin II, vasopressin and phenylephrine (34,35). In ASM, exogenous ouabain induces contraction by blocking the NKA as it promotes a  $\text{Na}^+$  concentration increase that causes the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX) to reverse ( $\text{NCX}_{\text{REV}}$ ). When in this reverse state,  $\text{NCX}_{\text{REV}}$  introduces  $\text{Ca}^{2+}$  into the cytoplasm, favoring contraction. This mechanism is identical to that exerted by inotropic glycosides in cardiac muscle (36), indicating that this steroid could induce BC *in vivo* (37). The NKA consists of  $\alpha\beta$  heterodimers; the  $\alpha$  catalytic subunit contains the binding sites for  $\text{Na}^+$ ,  $\text{K}^+$ , ATP and ouabain and is phosphorylated during each pump cycle. The  $\beta$  subunit is indispensable for the function of NKA; it stabilizes the conformation of the  $\alpha$  subunit and is the chaperone of the  $\alpha\beta$  complex in its transit between the SR and the plasma membrane. Each one of the four mammalian isoforms of the  $\alpha$  subunit ( $\alpha 1$ - $\alpha 4$ ) is the product of a different gene and they show ~90% sequence identity. Functionally, they are different in their correspondent kinetics, expression pattern and regulatory mechanisms. In most cells, NKA isoform  $\alpha 1$  is expressed with another isoform; for instance, muscles express  $\alpha 1$  and  $\alpha 2$  subunits. It has recently been found that the affinity of ouabain differs for each NKA  $\alpha$  subunit isoform. Thus, its affinity for the  $\alpha 2$  subunit is ~2.5 times less than for the  $\alpha 1$  subunit (38). Furthermore, each NKA subtype is located in specific membrane domains depending on its isotype. While NKAs formed by  $\alpha 1$  subunits are distributed ubiquitously throughout the membrane, the  $\alpha 2/\alpha 3$  isoforms are confined to membrane nanodomains adjacent to the SR. The NCX and many canonical transient receptor potential (TRPC) channels that belong to the NSCC are also located there, as well; both NCX and NSCC are also described herein. The low affinity for ouabain and the membrane location of the NKAs with  $\alpha 2$  subunit compared to those formed by  $\alpha 1$ , may indicate




Figure 2. Airway smooth muscle cell  $\text{Na}^+$  management during stimulation with a cholinergic agonist. Non-selective cationic channels (NSCC) activated by the lowering of the sarcoplasmic reticulum (SR)- $\text{Ca}^{2+}$  content can be denominated store operated channels (SOC), while NSCC corresponding to receptor operated channels (ROCs) are activated by IP3 cascade secondary messengers (i.e., DAG, diacylglycerol). Localized  $\text{Na}^+$  gradients originated by this cation's flux through either NSCC induce the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger's reversal ( $\text{NCX}_{\text{REV}}$ ).  $\text{NCX}_{\text{REV}}$  introduces 1  $\text{Ca}^{2+}$  and extrudes 3  $\text{Na}^+$ , contributing to regulate intracellular  $\text{Na}^+$  concentrations and SR  $\text{Ca}^{2+}$  refilling. Ionic currents flowing through NSCC depolarize the membrane, promoting the opening of the voltage-dependent L-type  $\text{Ca}^{2+}$  channel (L-VDCC). This channel permeates both  $\text{Ca}^{2+}$  and  $\text{Na}^+$ . This  $\text{Na}^+$  entrance probably contributes to membrane depolarization ( $\Delta\text{Em}$ ) to maintain  $\text{Ca}^{2+}$  entrance and SR refilling. Inset: representative fluorometric recording of a bovine tracheal myocyte stimulated with carbachol (CCh, 10  $\mu\text{M}$ ) in normal  $\text{Na}^+$  containing Krebs solution (2 mM  $\text{Ca}^{2+}$ ).  $\text{Na}^+$ -specific fluorophore (SBFI) signal was augmented when the cell received CCh, indicating the augmentation of this cation in the cytoplasm (19). SERCA, SR  $\text{Ca}^{2+}$  ATPase; NKA,  $\text{Na}^+/\text{K}^+$ ATPase; ACh, acetylcholine; GPCR, G protein-coupled receptor; IP3, inositol trisphosphate; IP3R, inositol trisphosphate receptor.

that they are involved in different cellular mechanisms, besides the regulation of the membrane potential at rest. At nanomolar concentrations, ouabain partially inhibits arterial smooth muscle contraction by augmenting the  $\text{Na}^+$  concentration in the microdomains containing NKAs with  $\alpha 2$  subunit well above that in the rest of the cytoplasm. This is due to the low affinity of  $\alpha 2$  pumps for ouabain (38,39) and could have implications in cell physiology. The wide distribution of NKAs with subunit  $\alpha 1$  indicates that they are responsible of controlling  $\text{Na}^+$  concentration in the whole cell, while the  $\alpha 2/\alpha 3$  subunits regulate only the concentrations of the nanodomains where the electrochemical gradient of  $\text{Na}^+$  reverses the NCX function to increase regional  $[\text{Ca}^{2+}]_i$ . In contrast, when the ASM is stimulated by an agonist, the NCX is reversed by the local increase in  $\text{Na}^+$  entry through the NSCC (Fig. 2).

### 3. $\text{Na}^+/\text{Ca}^{2+}$ exchanger

When the  $[\text{Ca}^{2+}]_i$  increases, it is promptly lowered by various calcium handling proteins. One of them is the NCX (40). Although its activity is determined by the surrounding  $\text{Na}^+$  concentration, it contributes significantly to the removal of intracellular  $\text{Ca}^{2+}$ . In a smooth muscle at rest, its function is to extract one  $\text{Ca}^{2+}$  ion from the cytoplasm and introduce 3  $\text{Na}^+$ . In mammals, the NCX is encoded by three genes: NCX1, 2 and 3 (41) and in various tissues, its cDNA is subjected to alternative splicing. This splicing implies the elimination of certain regions of the cDNA and as a result, at least 5 exchanger isoforms are generated (42). It has been reported that the NCX1 isoform has at least 12 splicing variants that define the exchanger's ionic sensitivity and regulation (43). Their tissue

distribution could be related to this characteristic; for instance the NCX1.1 has been found in the heart, while the NCX1.3 is located in the kidney and tracheal smooth muscle (44-46). As illustrated on Fig. 1, the NCX1.3 is located in certain cytoplasmic membrane regions. Local increases in  $\text{Na}^+$  in the areas next to the NCX1.3 induce its function to reverse, favoring the entry of  $\text{Ca}^{2+}$ ; under these circumstances, the NCX is said to be in reverse mode ( $\text{NCX}_{\text{REV}}$ ).  $\text{NCX}_{\text{REV}}$  extracts  $\text{Na}^+$  and introduces  $\text{Ca}^{2+}$  to the cytoplasm (41). In ASM, the  $\text{NCX}_{\text{REV}}$  provides  $\text{Ca}^{2+}$  for SR refilling, a physiological process indispensable for ASM sustained contraction induced by agonists (47,48) (Fig. 2). In this sense, using Patch Clamp techniques, an outwardly rectifying  $\text{Na}^+$  current was characterized in cultured human bronchial smooth cells loaded with  $\text{Na}^+$  and in the presence of extracellular  $\text{Ca}^{2+}$ . siRNA against NCX1 or knockdown of the stromal cell interaction molecule, (STIM1, an SR  $\text{Ca}^{2+}$  sensor) significantly diminished this response, suggesting a functional link between these calcium handling proteins. Visibly, STIM seems to promote SR  $\text{Ca}^{2+}$  refilling through  $\text{NCX}_{\text{REV}}$  activity (49).

### 4. Non-specific cation channels

This group is constituted by ion channels of varied nature that permeate mainly ions such as  $\text{Ca}^{2+}$  and  $\text{Na}^+$  and are designated as NSCC. They can be classified in two groups based on the cellular mechanisms that activate them: store operated (SOC) channels permeate cations in response to a decrease in the SR  $\text{Ca}^{2+}$  content and receptor operated (ROC) channels which, once the agonist (e.g., acetylcholine) occupies its receptor activating a signaling pathway, open in response to secondary

messengers (e.g., diacylglycerol) (50). The components of NSCCs have not been fully defined, but the TRPC channels are considered as an essential part of most of them. In general, all TRPs have six transmembranal regions (TM1-6) and oligomerize to form homotetramers or heterotetramers (51). The TRPC family consists of seven members (1-7) and TRPCs are present in excitable and non-excitable cells allowing nonspecific cation entry through the membrane. The current hence generated depolarizes the membrane, triggering the opening of voltage-dependent channels involved in diverse cellular functions. Thus, the TRPCs may cause action potentials in excitable cells (51). These channels also contribute to an increase in intracellular  $\text{Na}^+$  concentration to induce  $\text{NCX}_{\text{REV}}$  (Fig. 2). TRPCs were initially characterized in *Drosophila*; in this species, certain mutation prevents the passage of  $\text{Ca}^{2+}$  through the phospholipase C coupled channels. Normally, these photoreceptors remain depolarized when exposed to a continuous light source but those cells from specimens with the mutation show a transient response, therefore 'TRPC'. In mammals, the equivalent of the TRPC studied in *Drosophila* is classified into four subfamilies: TRPC1, TRPC2, TRPC3/6/7 and TRPC4/5 (sometimes TRPC1 is included in the subfamily TRPC4/5) (52-54). TRPC1 is considered paradigmatic for the study of SOCs as it has been shown to participate in SR  $\text{Ca}^{2+}$  refilling (capacitative entry) and in the sustained contraction. TRPC1 forms part of many types of NSCCs, from relatively selective to strictly non-selective channels (regarding  $\text{Ca}^{2+}$  permeability) and this diversity may be related to its heteromeric association with other TRPCs. Some studies even suggest that the presence of TRPC1 depends entirely on its interaction with another TRPC (TRPC4) (55). It has also been reported that the association of TRPC1 with two other proteins (STIM1 on the SR membrane and Orai on the plasma membrane) is highly selective for  $\text{Ca}^{2+}$  and that they only permeate a capacitative  $\text{Ca}^{2+}$  current known as calcium release-activated calcium current (ICRAC) (50,56-58).

With regard to those channels operated by the receptor, it has been found that TRPC3 (and most likely TRPC6 and TRPC7) are activated by inositol triphosphate (IP3) and/or the IP3 receptor (IP3R) (59-61). Therefore, TRPC3 is used as a model channel to study what is considered a conformational coupling between the channel and the IP3R; the latter requires at least its N-terminal portion to activate TRPC3 (62). Nevertheless, IP3 presence is not necessary to induce  $\text{Ca}^{2+}$  entry through these channels, as pharmacological methods other than agonist stimulation (thapsigargin or ionomycin) induce the same response. It has also been reported that the presence of TRPC3 is crucial to maintain the resting membrane potential in cells of healthy ASM, while in myocytes from asthmatic airways, TRPC1 seems to play an important role (63). Many of the functions of the different TRPC isoforms remain unclear; it has even been proposed that TRPCs are not indispensable for SOCs, but rather to ROCs due to their sensitivity to diacylglycerol (64).

## 5. $\text{Na}^+$ voltage dependent currents in airway smooth muscle

The recent evidence presented by Bradley *et al* (65) regarding the expression of a voltage-dependent  $\text{Na}^+$  current in freshly dispersed rabbit bronchial myocytes seemingly mediated by

the Nav1.5  $\alpha$  subunit, cannot be overlooked. When this current in bronchial myocytes was pharmacologically altered (blocked with veratridine), their action potential was prolonged. This finding opens new perspectives, as it is possible that airway inflammatory conditions like those prevailing in asthma could modify the voltage-dependent inactivation of the current promoting a bronchospasm.

Former studies (66,67) have already shown that cultured human bronchial smooth muscle cells express  $\text{Na}^+$  currents. Nakajima *et al* (68) were able to characterize and diminish Nav1.7 channel expression with dexamethasone, suggesting that these channels may be important in airway remodeling in asthma.

## 6. Evidence of $\text{Na}^+$ participation in asthma

Multiple studies of patients with atopic asthma and/or allergic rhinitis report the presence of plasmatic, endogenous ouabain, indicating a probable intracellular  $\text{Na}^+$  accumulation (21-26). In fact, the study by Tribe *et al* (25) shows an increased intracellular  $\text{Na}^+$  concentration in mononuclear cells from healthy individuals incubated with plasma of asthmatic blood donors. Apparently, endogenous ouabain acts not only on ASM NKAs, but also on inflammatory cells perhaps contributing to consolidate airway hyperresponsiveness. Knox *et al* (69) postulate that the difference lies on the tissue, since ASM contracts *in vitro* in response to exogenous ouabain but does not predispose tissue to hyperresponsiveness. Moreover, Knudsen *et al* (70) demonstrated that ouabain addition to rat peritoneal mast cells induced histamine release while simultaneously blocking intracellular  $\text{Rb}^+$  increase (used as surrogate to evaluate  $\text{K}^+$  cell entry through  $\text{Na}^+/\text{K}^+$  ATPase activity).

The origin of endogenous ouabain found in atopic asthmatic plasma has not been clarified completely, but documentary evidence indicates that mammals produce an endogenous ouabain analog. It is a hormone found in blood plasma, synthesized mainly by the adrenal, hypothalamus and pituitary glands (27-33). *In vitro*, the release of ouabain in primary cultures of bovine adrenocortical cells is favored by the addition of ACTH, angiotensin II, vasopressin and phenylephrine (34,35).

On the other hand, ouabain has been claimed to have anti-inflammatory capabilities by suppressing  $\text{TNF}\alpha$  production (71). In this regard, the high ouabain generation seen in atopic asthma may be associated with a compensatory process intended to regulate circulating TNF concentrations (72).

## 7. The paradox of exercise-induced bronchoconstriction

Exercise-induced bronchoconstriction (EIB) is currently defined as a syndrome where exercise or an increase in ventilation triggers airflow obstruction that may last 30-90 min if not attended. Although it occurs most frequently in asthmatic patients, cross sectional studies show that only a portion of patients with asthma have EIB when tested with a specific challenge test (73). The indirect challenges commonly used to define EIB are exercise itself, eucapnic voluntary hyperpnea, hypertonic (4.5%) saline, mannitol and adenosine (74). These 'specific challenge tests' differ from conventional methacholine or histamine challenges. Theoretically, exercise

increases the amount of inhaled air but augments airway lining fluid osmolarity by dehydration, favoring lung mast cell activation and pro-inflammatory mediator release from the epithelium involving a soluble phospholipase A (75,76). Most of these pro-inflammatory mediators can induce ASM contraction directly or favor a primed response to other agonists (hyperractivity). It is conceivable that airway lining fluid  $\text{Na}^+$  content could augment osmolarity in the airways during exercise. In this regard, Schmitt *et al* (77) established that, in airway epithelia from asthmatic patients with EIB, transepithelial nasal potential carried out by  $\text{Cl}^-$  and  $\text{Na}^+$  is not modified during exercise, indicating that a  $\text{Na}^+$  handling mechanism is at stake. Also associated with impaired cellular  $\text{Na}^+$  homeostasis, is the inhibition of the NKA in atopy and EIB. Mast cells express NKAs and NCXs (78); the  $[\text{Ca}^{2+}]_i$  increase induced by  $\text{NCX}_{\text{REV}}$  in these cells favors the release of inflammatory mediators, pointing out this exchanger's role in inflammation. In fact, the use of an  $\text{NCX}_{\text{REV}}$  inhibitor (KB-R7943) significantly reduced the amount of histamine released by stimulated peritoneal mast cells which correlated with a decrease in  $[\text{Ca}^{2+}]_i$  (79).

Plenty of evidence suggests that  $\text{Na}^+$  intake worsens EIB; a low-sodium diet improved spirometric results after physical activity in exercise-induced asthma patients, while subjects in each of the control groups (healthy individuals under high- or low-sodium diet) had no changes in their spirometric values (80,81). Furthermore, high serum concentrations of antioxidant vitamins, selenium, calcium, chloride and iron, were associated with better  $\text{FEV}_1$  values, as compared with higher concentrations of potassium and sodium, that were correlated with lower  $\text{FEV}_1$  (82). Asthmatic patients following a low-sodium diet showed reduced inflammation associated with EIB measured by eosinophilic cationic protein, IL-1 $\beta$  and IL-8 concentration in their sputum. This decrease correlated with an improvement in respiratory function assessed by spirometry. In contrast, the group of patients on a high-sodium diet showed a significant worsening of spirometric parameters that correlated with the amount of inflammatory cells and cytokine concentrations in sputum (83). Thus it appears that increasing dietary salt intake worsened inflammation of the airways in asthma patients. However, two separate Cochrane reviews established that there were no significant benefits of salt restriction on the control of asthma. In one of them, Pogson and McKeever (84) none the less claim that 'there was some evidence from the exercise-induced asthma studies that a low-sodium diet may improve lung function after exercise and possibly baseline lung function, but this is based on findings from a very small number of participants'. In the other, Ram and Arden (85) analyzed six randomized controlled trials and did not find any significant effect of dietary salt restriction on asthma treatment or management.

## 8. Conclusion

In conclusion, dietary sodium intake seems not to play a pivotal role in asthma nor in EIB, rather it is more plausible that  $\text{Na}^+$  handling mechanisms are partially responsible for EIB and augmented inflammatory mediator release in both ailments. This argument warrants further and deeper research in ASM sodium handling mechanisms.

## Acknowledgements

Authors are grateful to National Institute of Respiratory Diseases for the financial support in the publication of this manuscript.

## References

1. Woloski JR, Heston S and Escobedo Calderon SP: Respiratory Allergic Disorders. *Prim Care* 43: 401-415, 2016.
2. Bostantzoglou C, Delimpoura V, Samitas K, Zervas E, Kanniss F and Gaga M: Clinical asthma phenotypes in the real world: Opportunities and challenges. *Breathe Sheff* 11: 186-193, 2015.
3. Lommatzsch M and Stoll P: Novel strategies for the treatment of asthma. *Allergo J Int* 25: 11-17, 2016.
4. Koopmans T, Anaparti V, Castro-Piedras I, Yarova P, Irechukwu N, Nelson C, Perez-Zoghbi J, Tan X, Ward JP and Wright DB:  $\text{Ca}^{2+}$  handling and sensitivity in airway smooth muscle: Emerging concepts for mechanistic understanding and therapeutic targeting. *Pulm Pharmacol Ther* 29: 108-120, 2014.
5. Carbajal V, Vargas MH, Flores-Soto E, Martínez-Cordero E, Bazán-Perkins B and Montaña LM: LTD4 induces hyperresponsiveness to histamine in bovine airway smooth muscle: Role of SR-ATPase  $\text{Ca}^{2+}$  pump and tyrosine kinase. *Am J Physiol Lung Cell Mol Physiol* 288: L84-L92, 2005.
6. Liu C, Tazzeo T and Janssen LJ: Isoprostane-induced airway hyperresponsiveness is dependent on internal  $\text{Ca}^{2+}$  handling and Rho/ROCK signaling. *Am J Physiol Lung Cell Mol Physiol* 291: L1177-L1184, 2006.
7. Morin C and Rousseau E: Enhanced  $\text{Ca}^{2+}$  sensitivity in hyper-responsive cultured bronchi is mediated by TNF $\alpha$  and NF-kappaB. *Can J Physiol Pharmacol* 84: 1029-1041, 2006.
8. Sweeney D, Hollins F, Gomez E, Saunders R, Challiss RA and Brightling CE:  $[\text{Ca}^{2+}]_i$  oscillations in ASM: Relationship with persistent airflow obstruction in asthma. *Respirology* 19: 763-766, 2014.
9. Fleischmann BK, Washabau RJ and Kotlikoff MI: Control of resting membrane potential by delayed rectifier potassium currents in ferret airway smooth muscle cells. *J Physiol* 469: 625-638, 1993.
10. Blaustein MP: Sodium ions, calcium ions, blood pressure regulation, and hypertension: A reassessment and a hypothesis. *Am J Physiol* 232: C165-C173, 1977.
11. Orrenius S, Zhivotovsky B and Nicotera P: Regulation of cell death: The calcium-apoptosis link. *Nat Rev Mol Cell Biol* 4: 552-565, 2003.
12. Eder P, Poteser M, Romanin C and Groschner K:  $\text{Na}^+$  entry and modulation of  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchange as a key mechanism of TRPC signaling. *Pflugers Arch* 451: 99-104, 2005.
13. Blaustein MP and Lederer WJ: Sodium/calcium exchange: Its physiological implications. *Physiol Rev* 79: 763-854, 1999.
14. Blaustein MP and Wier WG: Local sodium, global reach: Filling the gap between salt and hypertension. *Circ Res* 101: 959-961, 2007.
15. Poburko D, Fameli N, Kuo KH and van Breemen C:  $\text{Ca}^{2+}$  signaling in smooth muscle: TRPC6, NCX and LNats in nanodomains. *Channels (Austin)* 2: 10-12, 2008.
16. Dai JM, Kuo KH, Leo JM, Paré PD, van Breemen C and Lee CH: Acetylcholine-induced asynchronous calcium waves in intact human bronchial muscle bundle. *Am J Respir Cell Mol Biol* 36: 600-608, 2007.
17. Flores-Soto E, Reyes-García J, Sommer B and Montaña LM: Sarcoplasmic reticulum  $\text{Ca}^{2+}$  refilling is determined by L-type  $\text{Ca}^{2+}$  and store operated  $\text{Ca}^{2+}$  channels in guinea pig airway smooth muscle. *Eur J Pharmacol* 721: 21-28, 2013.
18. Perusquía M, Flores-Soto E, Sommer B, Campuzano-González E, Martínez-Villa I, Martínez-Banderas AI and Montaña LM: Testosterone-induced relaxation involves L-type and store-operated  $\text{Ca}^{2+}$  channels blockade, and PGE $_2$  in guinea pig airway smooth muscle. *Pflugers Arch* 467: 767-777, 2015.
19. Sommer B, Flores-Soto E, Reyes-García J, Díaz-Hernández V, Carbajal V and Montaña LM:  $\text{Na}^+$  permeates through L-type  $\text{Ca}^{2+}$  channel in bovine airway smooth muscle. *Eur J Pharmacol* 782: 77-88, 2016.
20. Lingrel JB: The physiological significance of the cardiotonic steroid/ouabain-binding site of the  $\text{Na}^+$ ,K-ATPase. *Annu Rev Physiol* 72: 395-412, 2010.

21. Agrawal A, Agrawal KP, Ram A, Sondhi A, Chhabra SK, Gangal SV and Mehta D: Basis of rise in intracellular sodium in airway hyperresponsiveness and asthma. *Lung* 183: 375-387, 2005.
22. Chhabra SK, Khanduja A and Jain D: Increased intracellular calcium and decreased activities of leucocyte Na<sup>+</sup>,K<sup>+</sup>-ATPase and Ca<sup>2+</sup>-ATPase in asthma. *Clin Sci (Lond)* 97: 595-601, 1999.
23. Gentile DA and Skoner DP: The relationship between airway hyperreactivity (AHR) and sodium, potassium adenosine triphosphatase (Na<sup>+</sup>,K<sup>+</sup> ATPase) enzyme inhibition. *J Allergy Clin Immunol* 99: 367-373, 1997.
24. Skoner DP, Gentile D and Evans RW: A circulating inhibitor of the platelet Na<sup>+</sup>,K<sup>+</sup> adenosine triphosphatase (ATPase) enzyme in allergy. *J Allergy Clin Immunol* 87: 476-482, 1991.
25. Tribe RM, Barton JR, Poston L and Burney PG: Dietary sodium intake, airway responsiveness, and cellular sodium transport. *Am J Respir Crit Care Med* 149: 1426-1433, 1994.
26. Van Deusen MA, Gentile DA and Skoner DP: Inhibition of the sodium, potassium adenosine triphosphatase enzyme in peripheral blood mononuclear cells of subjects with allergic rhinitis. *Ann Allergy Asthma Immunol* 78: 259-264, 1997.
27. Hamlyn JM, Blaustein MP, Bova S, DuCharme DW, Harris DW, Mandel F, Mathews WR and Ludens JH: Identification and characterization of a ouabain-like compound from human plasma. *Proc Natl Acad Sci USA* 88: 6259-6263, 1991.
28. Ferrandi M, Manunta P, Balzan S, Hamlyn JM, Bianchi G and Ferrari P: Ouabain-like factor quantification in mammalian tissues and plasma: Comparison of two independent assays. *Hypertension* 30: 886-896, 1997.
29. Laredo J, Hamilton BP and Hamlyn JM: Ouabain is secreted by bovine adrenocortical cells. *Endocrinology* 135: 794-797, 1994.
30. Schoner W: Ouabain, a new steroid hormone of adrenal gland and hypothalamus. *Exp Clin Endocrinol Diabetes* 108: 449-454, 2000.
31. Komiyama Y, Nishimura N, Munakata M, Mori T, Okuda K, Nishino N, Hirose S, Kosaka C, Masuda M and Takahashi H: Identification of endogenous ouabain in culture supernatant of PC12 cells. *J Hypertens* 19: 229-236, 2001.
32. el-Masri MA, Clark BJ, Qazzaz HM and Valdes R Jr: Human adrenal cells in culture produce both ouabain-like and dihydro-ouabain-like factors. *Clin Chem* 48: 1720-1730, 2002.
33. Murrell JR, Randall JD, Rosoff J, Zhao JL, Jensen RV, Gullans SR and Hauptert GT Jr: Endogenous ouabain: Upregulation of steroidogenic genes in hypertensive hypothalamus but not adrenal. *Circulation* 112: 1301-1308, 2005.
34. Laredo J, Shah JR, Lu ZR, Hamilton BP and Hamlyn JM: Angiotensin II stimulates secretion of endogenous ouabain from bovine adrenocortical cells via angiotensin type 2 receptors. *Hypertension* 29: 401-407, 1997.
35. Shah JR, Laredo J, Hamilton BP and Hamlyn JM: Effects of angiotensin II on sodium potassium pumps, endogenous ouabain, and aldosterone in bovine zona glomerulosa cells. *Hypertension* 33: 373-377, 1999.
36. Saklani P and Skanes A: Novel anti-arrhythmic medications in the treatment of atrial fibrillation. *Curr Cardiol Rev* 8: 302-309, 2012.
37. Cortijo J, Sarria B, Mata M, Naline E, Advenier C and Morcillo EJ: Effects of ouabain on human bronchial muscle in vitro. *Naunyn Schmiedeberg Arch Pharmacol* 368: 393-403, 2003.
38. Blaustein MP and Hamlyn JM: Signaling mechanisms that link salt retention to hypertension: Endogenous ouabain, the Na(+)-pump, the Na(+)/Ca(2+) exchanger and TRPC proteins. *Biochim Biophys Acta* 1802: 1219-1229, 2010.
39. Katz A, Lifshitz Y, Bab-Dinitz E, Kapri-Pardes E, Goldshleger R, Tal DM and Karlish SJ: Selectivity of digitalis glycosides for isoforms of human Na,K-ATPase. *J Biol Chem* 285: 19582-19592, 2010.
40. Floyd R and Wray S: Calcium transporters and signalling in smooth muscles. *Cell Calcium* 42: 467-476, 2007.
41. DiPolo R and Beaugé L: Sodium/calcium exchanger: Influence of metabolic regulation on ion carrier interactions. *Physiol Rev* 86: 155-203, 2006.
42. Kofuji P, Lederer WJ and Schulze DH: Mutually exclusive and cassette exons underlie alternatively spliced isoforms of the Na/Ca exchanger. *J Biol Chem* 269: 5145-5149, 1994.
43. Quednau BD, Nicoll DA and Philipson KD: Tissue specificity and alternative splicing of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger isoforms NCX1, NCX2, and NCX3 in rat. *Am J Physiol* 272: C1250-C1261, 1997.
44. Mejía-Elizondo R, Espinosa-Tanguma R and Saavedra-Alanis VM: Molecular identification of the NCX isoform expressed in tracheal smooth muscle of guinea pig. *Ann NY Acad Sci* 976: 73-76, 2002.
45. Philipson KD, Nicoll DA, Ottolia M, Quednau BD, Reuter H, John S and Qiu Z: The Na<sup>+</sup>/Ca<sup>2+</sup> exchange molecule: An overview. *Ann NY Acad Sci* 976: 1-10, 2002.
46. Pitt A and Knox AJ: Molecular characterization of the human airway smooth muscle Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *Am J Respir Cell Mol Biol* 15: 726-730, 1996.
47. Janssen LJ, Walters DK and Wattie J: Regulation of [Ca<sup>2+</sup>]<sub>i</sub> in canine airway smooth muscle by Ca(2+)-ATPase and Na<sup>+</sup>/Ca<sup>2+</sup> exchange mechanisms. *Am J Physiol* 273: L322-L330, 1997.
48. Flores-Soto E, Carbajal V, Reyes-García J, García-Hernández LM, Figueroa A, Checa M, Barajas-López C and Montañón LM: In airways ATP refills sarcoplasmic reticulum via P2X smooth muscle receptors and induces contraction through P2Y epithelial receptors. *Pflügers Arch* 461: 261-275, 2011.
49. Liu B, Peel SE, Fox J and Hall IP: Reverse mode Na<sup>+</sup>/Ca<sup>2+</sup> exchange mediated by STIM1 contributes to Ca<sup>2+</sup> influx in airway smooth muscle following agonist stimulation. *Respir Res* 11: 168, 2010.
50. Putney JW and Tomita T: Phospholipase C signaling and calcium influx. *Adv Biol Regul* 52: 152-164, 2012.
51. Pedersen SF, Owsianik G and Nilius B: TRP channels: An overview. *Cell Calcium* 38: 233-252, 2005.
52. Trebak M, Lemonnier L, Smyth JT, Vazquez G and Putney JW Jr: Phospholipase C-coupled receptors and activation of TRPC channels. *Handb Exp Pharmacol* 179: 593-614, 2007.
53. Trebak M, Vazquez G, Bird GS and Putney JW Jr: The TRPC3/6/7 subfamily of cation channels. *Cell Calcium* 33: 451-461, 2003.
54. Vazquez G, Wedel BJ, Aziz O, Trebak M and Putney JW Jr: The mammalian TRPC cation channels. *Biochim Biophys Acta* 1742: 21-36, 2004.
55. Ong HL and Ambudkar IS: The dynamic complexity of the TRPC1 channelosome. *Channels (Austin)* 5: 424-431, 2011.
56. Yuan JP, Lee KP, Hong JH and Muallem S: The closing and opening of TRPC channels by Homer1 and STIM1. *Acta Physiol (Oxf)* 204: 238-247, 2012.
57. Cheng KT, Liu X, Ong HL, Swaim W and Ambudkar IS: Local Ca<sup>2+</sup> entry via Orail1 regulates plasma membrane recruitment of TRPC1 and controls cytosolic Ca<sup>2+</sup> signals required for specific cell functions. *PLoS Biol* 9: e1001025, 2011.
58. Soboloff J, Madesh M and Gill DL: Sensing cellular stress through STIM proteins. *Nat Chem Biol* 7: 488-492, 2011.
59. Zhu X, Jiang M, Peyton M, Boulay G, Hurst R, Stefani E and Birnbaumer L: Trp, a novel mammalian gene family essential for agonist-activated capacitative Ca<sup>2+</sup> entry. *Cell* 85: 661-671, 1996.
60. Kiselyov K, Xu X, Mozhayeva G, Kuo T, Pessah I, Mignery G, Zhu X, Birnbaumer L and Muallem S: Functional interaction between InsP3 receptors and store-operated Htrp3 channels. *Nature* 396: 478-482, 1998.
61. McKay RR, Szymczek-Seay CL, Lievreumont JP, Bird GS, Zitt C, Jüngling E, Lückhoff A and Putney JW Jr: Cloning and expression of the human transient receptor potential 4 (TRP4) gene: Localization and functional expression of human TRP4 and TRP3. *Biochem J* 351: 735-746, 2000.
62. Kiselyov K, Mignery GA, Zhu MX and Muallem S: The N-terminal domain of the IP3 receptor gates store-operated hTrp3 channels. *Mol Cell* 4: 423-429, 1999.
63. Xiao JH, Zheng YM, Liao B and Wang YX: Functional role of canonical transient receptor potential 1 and canonical transient receptor potential 3 in normal and asthmatic airway smooth muscle cells. *Am J Respir Cell Mol Biol* 43: 17-25, 2010.
64. Putney JW: The physiological function of store-operated calcium entry. *Neurochem Res* 36: 1157-1165, 2011.
65. Bradley E, Webb TI, Hollywood MA, Sergeant GP, McHale NG and Thornbury KD: The cardiac sodium current Na(v)1.5 is functionally expressed in rabbit bronchial smooth muscle cells. *Am J Physiol Cell Physiol* 305: C427-C435, 2013.
66. Snetkov VA, Hirst SJ and Ward JP: Ion channels in freshly isolated and cultured human bronchial smooth muscle cells. *Exp Physiol* 81: 791-804, 1996.
67. Jo T, Nagata T, Iida H, Imuta H, Iwasawa K, Ma J, Hara K, Omata M, Nagai R, Takizawa H, *et al.*: Voltage-gated sodium channel expressed in cultured human smooth muscle cells: Involvement of SCN9A. *FEBS Lett* 567: 339-343, 2004.
68. Nakajima T, Jo T, Meguro K, Oonuma H, Ma J, Kubota N, Imuta H, Takano H, Iida H, Nagase T, *et al.*: Effect of dexamethasone on voltage-gated Na<sup>+</sup> channel in cultured human bronchial smooth muscle cells. *Life Sci* 82: 1210-1215, 2008.
69. Knox AJ, Ajao P, Britton JR and Tattersfield AE: Effect of sodium-transport inhibitors on airway smooth muscle contractility in vitro. *Clin Sci (Lond)* 79: 315-323, 1990.

70. Knudsen T, Bertelsen H and Johansen T: Ouabain enhancement of compound 48/80 induced histamine secretion from rat peritoneal mast cells: Dependence on extracellular sodium. *Pharmacol Toxicol* 70: 412-418, 1992.
71. de Vasconcelos DI, Leite JA, Carneiro LT, Piuvezam MR, de Lima MR, de Moraes LC, Rumjanek VM and Rodrigues-Mascarenhas S: Anti-inflammatory and antinociceptive activity of ouabain in mice. *Mediators Inflamm* 2011: 912925, 2011.
72. Choi JP, Kim YS, Kim OY, Kim YM, Jeon SG, Roh TY, Park JS, Gho YS and Kim YK: TNF-alpha is a key mediator in the development of Th2 cell response to inhaled allergens induced by a viral PAMP double-stranded RNA. *Allergy* 67: 1138-1148, 2012.
73. Hallstrand TS: New insights into pathogenesis of exercise-induced bronchoconstriction. *Curr Opin Allergy Clin Immunol* 12: 42-48, 2012.
74. Anderson SD: Indirect challenge tests: Airway hyperresponsiveness in asthma: its measurement and clinical significance. *Chest* 138 (Suppl): 25S-30S, 2010.
75. Hallstrand TS, Moody MW, Wurfel MM, Schwartz LB, Henderson WR Jr and Aitken ML: Inflammatory basis of exercise-induced bronchoconstriction. *Am J Respir Crit Care Med* 172: 679-686, 2005.
76. Hallstrand TS, Lai Y, Ni Z, Oslund RC, Henderson WR Jr, Gelb MH and Wenzel SE: Relationship between levels of secreted phospholipase A<sub>2</sub> groups IIA and X in the airways and asthma severity. *Clin Exp Allergy* 41: 801-810, 2011.
77. Schmitt L, Wiebel M, Frese F, Dehnert C, Zugck C, Bärtsch P and Mairbäurl H: Exercise reduces airway sodium ion reabsorption in cystic fibrosis but not in exercise asthma. *Eur Respir J* 37: 342-348, 2011.
78. Aneiros E, Philipp S, Lis A, Freichel M and Cavalié A: Modulation of Ca<sup>2+</sup> signaling by Na<sup>+</sup>/Ca<sup>2+</sup> exchangers in mast cells. *J Immunol* 174: 119-130, 2005.
79. Praetorius HA, Friis UG, Praetorius J and Johansen T: Evidence for a Na<sup>+</sup>/Ca<sup>2+</sup> exchange mechanism in rat peritoneal mast cells. *Pflugers Arch* 437: 86-93, 1998.
80. Rundell KW and Jenkinson DM: Exercise-induced bronchospasm in the elite athlete. *Sports Med* 32: 583-600, 2002.
81. Gotshall RW, Mickleborough TD and Cordain L: Dietary salt restriction improves pulmonary function in exercise-induced asthma. *Med Sci Sports Exerc* 32: 1815-1819, 2000.
82. McKeever TM, Lewis SA, Smit HA, Burney P, Cassano PA and Britton J: A multivariate analysis of serum nutrient levels and lung function. *Respir Res* 9: 67-89, 2008.
83. Mickleborough TD, Lindley MR and Ray S: Dietary salt, airway inflammation, and diffusion capacity in exercise-induced asthma. *Med Sci Sports Exerc* 37: 904-914, 2005.
84. Pogson Z and McKeever T: Dietary sodium manipulation and asthma. *Cochrane Database Syst Rev* 3: CD000436, 2011.
85. Arden KD and Arden KD: Dietary salt reduction or exclusion for allergic asthma. *Cochrane Database Syst Rev* 2: CD000436, 2004.