

# Shared mechanisms of musculoskeletal dysfunction in type 2 diabetes mellitus: Insights into future therapeutic directions (Review)

ZHENHONG HE<sup>1,2\*</sup>, HANYU LI<sup>1,2\*</sup>, YUHAN LUO<sup>2\*</sup>, YINGYING YU<sup>2</sup>,  
SIYU LI<sup>2</sup>, TAO ZHANG<sup>1</sup> and DINGSU BAO<sup>1</sup>

<sup>1</sup>Department of Orthopedics, The Affiliated Traditional Chinese Medicine Hospital, Southwest Medical University, Luzhou, Sichuan 646000, P.R. China; <sup>2</sup>College of Integrated Traditional Chinese and Western Medicine, Southwest Medical University, Luzhou, Sichuan 646000, P.R. China

Received June 12, 2026; Accepted July 30, 2026

DOI: 10.3892/ijmm.2026.5955

**Abstract.** Type 2 diabetes mellitus (T2DM) profoundly affects the musculoskeletal system through multifactorial mechanisms. The present review systematically summarizes the impact of T2DM on skeletal muscle, tendon, tendon-bone interface (TBI), and bone, highlighting their shared and tissue-specific pathophysiological pathways. In skeletal muscle, impaired insulin signaling and mitochondrial dysfunction lead to reduced protein synthesis, enhanced proteolysis, and muscle atrophy. In tendons, advanced glycation end products accumulation and microangiopathy disrupt collagen metabolism and biomechanics. At the TBI, persistent hyperglycemia induces fibrocartilage fibrosis, impaired angiogenesis and disorganized stress transmission, ultimately weakening load-bearing capacity. In bone, the imbalance between osteoblast and osteoclast activity, coupled with microvascular dysfunction and metabolic acidosis, results in reduced bone formation and increased fragility. Although the manifestations of T2DM differ among skeletal muscle, tendon, TBI and bone, these tissues share common pathological drivers, including metabolic dysregulation, chronic inflammation, oxidative stress, extracellular matrix remodeling, and impaired regenerative capacity. A deeper understanding of these shared mechanisms may facilitate the identification of novel therapeutic targets.

Emerging technologies, including single-cell RNA sequencing, regenerative medicine, and precision medicine, further provide new opportunities for early diagnosis and personalized management of diabetes-related musculoskeletal disorders.

## Contents

1. Introduction
2. Systemic pathophysiology of T2DM in musculoskeletal tissues
3. Tissue-specific pathological mechanisms underlying T2DM musculoskeletal disorders
4. Conclusion and perspective

## 1. Introduction

According to the International Diabetes Federation (IDF), an estimated 537 million adults aged 20-79 years worldwide had diabetes in 2021. Projections indicate this number will rise to 783 million by 2045 (1). Notably, over 90% of these cases are type 2 diabetes mellitus (T2DM) (2). Inadequately managed T2DM can lead to numerous severe complications. Within the musculoskeletal system, T2DM induces multiple pathological alterations affecting bone, joints, muscles and the nervous system. These impairments not only directly compromise mobility but are also intrinsically linked to metabolic dysregulation, chronic inflammation and microvascular dysfunction. Physical exercise is an important intervention for diabetes management. Exercise activates adenosine monophosphate-activated protein kinase (AMPK) signaling, promotes glucose transport and lipid metabolism, and enhances mitochondrial function in skeletal muscle. It also improves insulin sensitivity and blood sugar control (3-8). However, a scientific exercise plan fundamentally relies on a stable and functional musculoskeletal system. Because of the detrimental impact of T2DM on the musculoskeletal system, early intervention is critical. Identifying effective treatment targets and preventive strategies may reduce the need for surgery and significantly

---

*Correspondence to:* Professor Tao Zhang or Professor Dingsu Bao, Department of Orthopedics, The Affiliated Traditional Chinese Medicine Hospital, Southwest Medical University, 182 Chunhui Road, Longmatan, Luzhou, Sichuan 646000, P.R. China  
E-mail: 130225@swmu.edu.cn  
E-mail: 18383089769@163.com

\*Contributed equally

**Key words:** type 2 diabetes mellitus, musculoskeletal system, tendon-bone interface, advanced glycation end products, oxidative stress, inflammation, metabolic dysregulation

improve the quality of life for patients. This approach aligns with current priorities in the prevention and management of major diseases, emphasizing the integration of traditional Chinese and Western medicine, as well as multidisciplinary collaboration. The present review synthesizes current research to explain how T2DM affects the musculoskeletal system and identifies potential intervention targets, thereby laying a solid foundation for future investigations.

## 2. Systemic pathophysiology of T2DM in musculoskeletal tissues

*Hyperglycemia and insulin resistance as the primary instigators.* Hyperglycemia and insulin resistance represent the earliest pathological events that start musculoskeletal degeneration in T2DM. Persistent metabolic dysregulation disrupts glucose utilization, energy metabolism and cellular anabolic signaling. This creates a common pathological basis for abnormalities in skeletal muscle, tendon, tendon-bone interface (TBI) and bone. Although these tissues perform distinct physiological functions, they all exhibit impaired cellular metabolism and tissue homeostasis under chronic hyperglycemic and insulin-resistant conditions (Table I).

Skeletal muscle is the primary site for post-prandial glucose uptake and disposal, playing a central role in maintaining glucose homeostasis (9). Peripheral insulin resistance originating in skeletal muscle is a key driver in the development and progression of T2DM (10). Under normal physiological conditions, insulin promotes glucose uptake by activating glucose transporter type 4 (GLUT4). However, in T2DM, defects in the insulin signaling pathways lead to impaired translocation of GLUT4 to the cell membrane, which represents a core mechanism of insulin resistance (11,12). The resistance results in decreased activity of glycogen synthase in skeletal muscle and reduced glycogen storage. Reduced glycogen synthesis forces cells to rely on lipolysis for energy production. At the same time, impaired mitochondrial function decreases fatty acid oxidation, which exacerbates the accumulation of lipid intermediates (13,14). Among these metabolites, diacylglycerol (DAG) activates protein kinase C (PKC), attenuating insulin-stimulated insulin receptor substrate- phosphorylation and phosphoinositide 3-kinase (PI3K) activity. As a result, GLUT4 movement is further reduced, worsening insulin resistance (15,16). Moreover, prolonged exposure to free fatty acids or a high-fat diet increases DAG accumulation inside muscle cells. A high blood sugar environment further promotes fat synthesis and buildup, forming a vicious cycle of metabolic imbalance that progressively worsens energy metabolism (Fig. 1A) (16).

The metabolic disturbances initiated by hyperglycemia and insulin resistance also extend to tendon tissue. Tendons are highly organized connective tissues composed primarily of type I collagen, together with proteoglycans and glycoproteins (17). Chronic hyperglycemia suppresses the function of tendon stem/progenitor cell (TSPC). It also disrupts the balance between matrix metalloproteinases (MMPs) and tissue inhibitors of metalloproteinases (TIMPs), promoting collagen degradation and preventing effective matrix remodeling (18-20). Consequently, collagen fibers become disorganized, losing elasticity and changing their mechanical

properties. Experimental studies have demonstrated reduced elasticity and impaired mechanical performance in diabetes rats' tendons, highlighting the harmful effects of chronic hyperglycemia on tendon function (21,22). Similarly, the TBI is continuously exposed to the diabetic metabolic environment. As a specialized tissue responsible for transferring loads between tendon and bone, the TBI depends on coordinated cellular differentiation and extracellular matrix (ECM) homeostasis. Persistent hyperglycemia promotes abnormal collagen cross-linking and chronic low-grade inflammation, leading to a progressive worsening of the interface structure and function (23-25). Following an injury, the diabetic TBI heals mostly by forming fibrotic scar tissue rather than regenerating the normal fibrocartilage transition zone (26). The newly formed scar tissue is characterized by disorganized collagen fibers and poor biomechanical properties, leading to reduced load-bearing capacity and compromised interface function.

Bone tissue is likewise profoundly affected by chronic hyperglycemia and insulin resistance. Under diabetic conditions, hyperglycemia directly suppresses the expression of bone-forming markers, including osteocalcin, osteopontin, runt-related transcription factor 2 (Runx2) and alkaline phosphatase. Hyperglycemia also promotes the death of bone-forming cells, resulting in reduced bone formation and poor bone quality (27-30). In addition, insulin deficiency or resistance diminishes the anabolic effects of insulin and insulin-like growth factor-1 on osteoblasts, further impairing matrix synthesis and mineralization (31-33). Elevated glucose levels also induce aberrant differentiation of mesenchymal stem cells, resulting in increased bone marrow adiposity and impaired osteoblast formation (34-37). Furthermore, chronic hyperglycemia enhances the sensitivity of osteoclast precursors to macrophage colony-stimulating factor and receptor activator of nuclear factor  $\kappa$ B ligand (RANKL), thereby facilitating osteoclast differentiation and fusion, and consequently increasing bone resorptive activity (38). These findings indicate that hyperglycemia and insulin resistance constitute the primary early drivers of diabetic musculoskeletal disorders. By disrupting glucose utilization, lipid metabolism, anabolic signaling and lineage commitment, they establish the pathological foundation for subsequent oxidative stress (OS), chronic inflammation, ECM remodeling, vascular dysfunction, and impaired tissue regeneration across skeletal muscle, tendon, the TBI and bone.

*AGEs-RAGE signaling, OS, and chronic inflammation as interconnected pathogenic mediators.* Following chronic hyperglycemia and insulin resistance, the excessive accumulation of advanced glycation end products (AGEs), persistent OS, and chronic inflammation constitute a connected pathogenic network that drives musculoskeletal degeneration in T2DM. Despite the tissue-specific signs of diabetes-related damage, these pathological mechanisms are remarkably similar across skeletal muscle, tendon, the TBI and bone (Table I). Under normal physiological conditions, the formation of AGEs is minimal and efficiently cleared from the body (39). However, chronic hyperglycemia greatly accelerates the non-enzymatic glycation of proteins and lipids, resulting in excessive AGEs generation and progressive accumulation within multiple tissues (40,41). Excessive AGEs alter the structural and

Table I. Shared pathogenic mechanisms of musculoskeletal dysfunction in T2DM.

| Shared pathogenic mechanisms         | Skeletal muscle                                                                                                                                                                                                                                | Tendon                                                                                                                                                                                                      | Tendon-bone interface                                                                                                                                                                              | Bone                                                                                                                                                                                                                                   | (Refs.)                                |
|--------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------|
| Hyperglycemia and insulin resistance | Impaired GLUT4 translocation → reduced glucose uptake; Reduced glycogen synthesis → reliance on lipolysis; DAG accumulation → PKC activation → IRS-1/PI3K Suppression; Impaired mitochondrial oxidative phosphorylation; metabolic dysfunction | TSPC function suppression; Disrupted MMP/TIMP balance → collagen degradation; Reduced collagen synthesis; Diminished tendon elasticity; Altered biomechanical                                               | Abnormal collagen cross-linking; Chronic low-grade inflammation; Fibrotic scar formation; Reduced load-bearing capacity                                                                            | Osteoblast apoptosis; Impaired matrix synthesis and mineralization; Aberrant MSC differentiation → marrow adiposity; Enhanced osteoclast precursor sensitivity to M-CSF and RANKL                                                      | (9,11-17,19,21, 22,25-27,31-34, 36,37) |
| AGEs-RAGE signaling                  | AGEs accumulation in ECM and collagen networks → fibrosis; RAGE activation → ROS and inflammation; Functional impairment and reduced muscle quality                                                                                            | AGEs cross-link collagen → reduced fiber sliding; Increased passive tension; Inhibition of collagen-related gene expression; Glycosylated collagen accumulation                                             | Chronic inflammatory; TGF-β1/Smad3 activation → fibroblast hyperproliferation and fibrosis; Abnormal collagen; Increased ECM stiffness; Impaired mechanical buffering and load transmission        | Reduced matrix elasticity and mechanical strength; Impaired osteoblast differentiation; Osteocyte senescence and apoptosis                                                                                                             | (23,24,42,48-53, 56,58-63)             |
| Oxidative stress                     | ROS; glucose autooxidation; Inflammation; ATP production; Mitochondrial DNA and respiratory chain damage PI3K/Akt/mTOR suppression                                                                                                             | ROS damage tenocytes; Impaired tenocyte proliferation, differentiation, and metabolic function; Collagen degradation; Impaired tendon self-repair capacity                                                  | ROS damage endothelial cells → impaired angiogenic responses; Inflammation and hypoxia vicious cycle; Impaired angiogenesis; Disrupted stem cell differentiation                                   | Osteoblast differentiation↓; Promote osteoclast activity; Hypoxia → premature Increased osteocyte RANKL → osteoclastogenesis                                                                                                           | (70,73-77,97)                          |
| Chronic low-grade inflammation       | IKKβ/NF-κB activation → TNF-α, IL-6, CRP increased; Cytokines impair insulin; UPS activation → accelerated proteolysis; JAK/STAT pathway activation → muscle loss; Impaired muscle regeneration                                                | Immune cell activation; Pro-inflammatory cytokines release; Disrupted collagen fiber organization; M2 polarization → TGF-β /VEGF secretion → fibrosis and ECM deposition; Impaired biomechanical properties | Chronic inflammatory immune responses; Fibroblast hyperproliferation → fibrotic remodeling; Fibrocartilage replaced by scar tissue; Loss of stress-buffering capacity; Impaired force transmission | Inflammatory cytokines promote osteoclastogenesis; Bone marrow adiposity and inflammation; Imbalanced RANKL/OPG ratio → bone resorption; SASP from senescent osteocytes → further inflammation; Trabecular deterioration and bone loss | (38,79,81-84,95, 109,229,230, 261-263) |

Table I. Continued.

| Shared pathogenic mechanisms | Skeletal muscle                                                                                                                                                                    | Tendon                                                                                                                                            | Tendon-bone interface                                                                                                                                                                                          | Bone                                                                                                                                            | (Refs.)                                       |
|------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------|
| Microvascular Dysfunction    | Capillary rarefaction, basement membrane thickening; Endothelial dysfunction; mitochondrial dysfunction and energy deficiency; Impaired glucose utilization; peripheral neuropathy | Aberrant vascularization; Local hypoxia; Tenocyte damage and collagen degradation; Impaired stem cell migration; Delayed post-injury angiogenesis | Reduced expression of CD31 and endomucin; EPC and APC dysfunction; HIF-2 $\alpha$ $\uparrow$ $\rightarrow$ Scx suppression $\rightarrow$ impaired TSPC differentiation; Ectopic calcification; VEGF resistance | Reduced bone marrow capillary density; Reduced blood perfusion; Hypoxia; Trabecular structure disruption; Impaired bone remodeling and turnover | (89-91,93-95,99,115,118-121, 123-129,133,134) |
| Abnormal ECM                 | AGEs cross-linking increased ECM stiffness and fibrosis                                                                                                                            | AMPK/EGR1 inhibition $\rightarrow$ collagen synthesis decreased                                                                                   | MMP/TIMP balance disruption $\rightarrow$ excessive matrix degradation                                                                                                                                         | Osteoclasts create acidic microenvironment $\rightarrow$ matrix                                                                                 | (59,137,138,141,144-148, 150-154,156)         |
| Impaired cell regeneration   | PI3K/Akt/mTOR $\downarrow$ , NF- $\kappa$ B/FoxO $\uparrow$ ; Reduced muscle fiber CSA and atrophy; MuSC dysfunction; Impaired muscle repair                                       | TSPC proliferation $\downarrow$ ; apoptosis $\uparrow$ ; Aberrant osteogenic; Excessive vascularization; Persistent inflammation and fibrosis     | TSPC apoptosis, proliferation $\downarrow$ ; Fibrocartilage thinning and degeneration; Fibrotic scar replaces native transition zone                                                                           | Bone marrow adiposity, osteoblast numbers; Bone formation, Trabecular deterioration, cortical thinning, skeletal fragility $\uparrow$           | (18,19,34-37,111,163-165, 167-172,175)        |

AGE, advanced glycation end product; Akt, protein kinase B; ALP, alkaline phosphatase; AMPK, AMP-activated protein kinase; APC, angiogenic progenitor cell; aP2, adipocyte protein 2; BH4, tetrahydrobiopterin; CKMT2, mitochondrial creatine kinase 2; CRP, C-reactive protein; CSA, cross-sectional area; DAG, diacylglycerol; ECM, extracellular matrix; eNOS, endothelial nitric oxide synthase; EPC, endothelial progenitor cell; FoxO, forkhead box O; GLUT4, glucose transporter type 4; HIF-1 $\alpha$ , hypoxia-inducible factor-1 $\alpha$ ; HIF-2 $\alpha$ , hypoxia-inducible factor-2 $\alpha$ ; IGF-1, insulin-like growth factor-1; IKK $\beta$ , I $\kappa$ B kinase  $\beta$ ; IL-6, interleukin-6; iNOS, inducible nitric oxide synthase; IRS-1, insulin receptor substrate-1; JAK, janus kinase; MAFbx, muscle atrophy F-box; M-CSF, macrophage colony-stimulating factor; Mfn1, mitofusin-1; Mfn2, mitofusin-2; MMP, matrix metalloproteinase; MSC, mesenchymal stem cell; mTOR, mechanistic target of rapamycin; MuRF1, muscle RING-finger protein-1; MuSC, muscle satellite cell; NADPH, nicotinamide adenine dinucleotide phosphate; NF- $\kappa$ B, nuclear factor kappa B; NO, nitric oxide; OCN, osteocalcin; OPG, osteoprotegerin; OPN, osteopontin; PGC-1 $\alpha$ , peroxisome proliferator-activated receptor gamma coactivator-1 alpha; PI3K, phosphoinositide 3-kinase; PKC, protein kinase C; PPAR $\gamma$ , peroxisome proliferator-activated receptor gamma; RAGE, receptor for advanced glycation end products; RANKL, receptor activator of nuclear factor- $\kappa$ B ligand; ROS, reactive oxygen species; Runx2, runt-related transcription factor 2; SASP, senescence-associated secretory phenotype; Scx, scleraxis; Smad3, mothers against decapentaplegic homolog 3; Sox9, SRY-box transcription factor 9; STAT, signal transducer and activator of transcription; TDSC, tendon-derived stem cell; TGF- $\beta$ , transforming growth factor- $\beta$ ; TGase, transglutaminase; TIMP, tissue inhibitor of metalloproteinase; TNF- $\alpha$ , tumor necrosis factor  $\alpha$ ; TSPC, tendon stem/progenitor cell; UPS, ubiquitin-proteasome system; VEGF, vascular endothelial growth factor.

biochemical properties of the ECM, disrupt collagen integrity through non-enzymatic cross-linking, and trigger persistent signaling through the receptor for AGEs (RAGE) (42-45). The sustained activation of this pathway amplifies OS and

inflammatory cascades, leading to mitochondrial dysfunction, impaired cellular differentiation, and cell death in multiple tissue types (46,47). These connected events ultimately compromise the balance between tissue formation and




Figure 1. Mechanisms of skeletal muscle dysfunction in T2DM. (A) It highlights the role of glucose uptake, insulin signaling, and lipid oxidation leading to oxidative stress. This results in protein cross-linking with tissues, affecting the mechanical properties of muscle and triggering ROS generation, inflammation and protein degradation, which collectively contribute to muscle quality decline in T2DM. (B) It focuses on the role of CBMT and NO signaling in endothelial function, which leads to impaired blood vessel integrity. This is linked to the formation of ROS and AGE accumulation, which together contribute to diabetic angiopathy. (C) It involves axonal cytoskeletal glycosylation, AGE accumulation and ROS aggravation, leading to disruption in neurocyte function and contributing to diabetic neuropathy. T2DM, type 2 diabetes mellitus; ROS, reactive oxygen species; NO, nitric oxide; AGEs, advanced glycation end products; RAGE, receptor for AGEs; DAG, diacylglycerol; IRS, insulin receptor substrate.

degradation, weaken the biomechanical properties of muscles, tendons and bone, and destroy the coordinated function of the TBI.

AGE accumulation profoundly alters the mechanical properties of collagen-rich tissues throughout the musculoskeletal system. In skeletal muscle, AGEs accumulate within the muscle ECM and collagen networks, increasing collagen cross-linking, tissue stiffness, fibrosis and functional

impairment (42,48-50). In tendon, AGEs impair biomechanics in multiple species by reducing tendon fibers sliding, resulting in increased passive muscle tension, reduced elasticity and decreased flexibility (51-53). In addition, AGEs directly inhibit the expression of collagen-related genes. Glycosylated collagen accumulates in connective tissues due to its insolubility and increased resistance to collagenase, leading to further AGE deposition (54-56). At the TBI, the persistent accumulation

of AGEs, along with hyperglycemia-induced OS, continuously activates AGEs-RAGE signaling and promotes chronic inflammatory immune responses (23-25). This pathological cascade activates the TGF- $\beta$ 1/Smad3 pathway, causing excessive fibroblast growth and fibrotic tissue remodeling. This is accompanied by abnormal cross-linking of collagen and proteoglycans, increased ECM stiffness, and loss of tissue elasticity, which impairs mechanical buffering and load transmission across the enthesis (24,53,57,58). In bone, excessive accumulation of AGEs within the bone matrix harms collagen structure through non-enzymatic cross-linking, reduces matrix elasticity and mechanical strength, and directly impairs osteoblast differentiation through activating RAGE-dependent signaling pathways (59-63). Major axonal cytoskeletal proteins, such as tubulin, neurofilaments and actin, are vulnerable to glycation under diabetic conditions. As essential components for maintaining axonal structure and function, their modification via glycation may alter structural integrity and functional properties of axons, leading to axonal atrophy, degeneration and slowed axonal transport (64-67). Studies indicate that AGE-modified myelin sheaths in peripheral nerves are more vulnerable to being consumed by macrophages. This process stimulates the release of enzymes and likely contributes to demyelination in diabetic neuropathy (68). Hyperglycemia-induced OS may accelerate the accumulation of AGEs, thereby exacerbating oxidative damage in peripheral nerves associated with neuronal disturbance and impaired Schwann cell signaling (67). In this manner, diabetic neuropathy promotes progressive nerve fiber loss and poor regeneration. Additionally, increased activity in the polyol pathway promotes the formation of 3-deoxyglucosone and methylglyoxal, which are precursors that contribute to the accumulation of AGEs. These AGEs alter neural cell components and, through engagement with the RAGE expressed on neural cells, trigger the production of various cytokines implicated in nerve injury (38,67). Thus, both hyperglycemia and polyol pathway-mediated AGE accumulation, along with AGE-RAGE interactions, are likely to occur within diabetic nervous and vascular systems. This highlights a cooperative relationship between the polyol pathway and AGE/RAGE-dependent signaling in promoting OS and nuclear factor kappa B (NF- $\kappa$ B) activation, which aggravates neurovascular dysfunction in experimental T2DM neuropathy (67).

The binding of AGEs to RAGE represents the central molecular event linking metabolic disturbance with OS and inflammation. By activating nicotinamide adenine dinucleotide phosphate (NADPH) oxidase and the PI3K/protein kinase B (Akt) pathway, AGEs stimulate excessive reactive oxygen species (ROS) generation and activate NF- $\kappa$ B signaling, thereby establishing a self-amplifying cycle between OS and inflammation (41,43-45,69). Simultaneously, AGEs formation, glucose autooxidation, and inflammatory activation collectively induce excessive ROS production. This disrupts intracellular metabolic homeostasis, impairs insulin signaling, and promotes catabolic processes (70). ROS further attack mitochondrial DNA, respiratory chain complexes, and membrane phospholipids, leading to mitochondrial dysfunction (71). As the primary source of cellular energy, dysfunctional mitochondria fail to meet tissue energy demands, resulting in

muscle atrophy, fatigue, impaired cellular differentiation and reduced exercise tolerance (72). Meanwhile, OS suppresses anabolic pathways, including PI3K/Akt/mammalian target of rapamycin (mTOR) signaling, and compromises tissue structural integrity. It also impairs angiogenesis and tissue repair, aggravates hypoxia, and ultimately contributes to delayed or failed healing across musculoskeletal tissues (73-77).

Chronic inflammation is another essential part of this pathogenic network. While T2DM has traditionally been defined by insulin resistance and progressive  $\beta$ -cell dysfunction, increasing evidence highlights chronic low-grade inflammation and immune dysregulation as equally central to disease progression (78). Persistent hyperglycemia, lipid overload, and the metabolic stress caused by compensatory insulin hypersecretion activate innate immune pathways. This environment releases signaling molecules that recruit monocytes to eliminate apoptotic  $\beta$ -cells. Under the regulation of NF- $\kappa$ B signaling, these recruited monocytes primarily differentiate into pro-inflammatory M1 and anti-inflammatory M2 macrophages. This process contributes to  $\beta$ -cell dysfunction and apoptosis. In particular, tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) impairs insulin signaling and diminishes insulin sensitivity (79,80). A vicious inflammatory cycle subsequently emerges. The IkappaB kinase  $\beta$  (IKK $\beta$ )/NF- $\kappa$ B signaling axis serves as a critical mediator in this process. It drives the systemic accumulation of pro-inflammatory mediators such as TNF- $\alpha$ , interleukin-6 (IL-6) and C-reactive protein (81-83). In turn, these cytokines further activate the IKK/NF- $\kappa$ B cascade, impair insulin signaling, and exacerbate  $\beta$ -cell dysfunction and systemic insulin resistance. They also promote the excessive breakdown of structural proteins and the ECM (79,84). Together with persistent ROS production, this inflammatory microenvironment progressively shifts tissue homeostasis toward catabolism, inhibits tissue regeneration, and amplifies structural degeneration throughout skeletal muscle, tendon, the TBI and bone.

Collectively, AGEs-RAGE activation, OS and chronic inflammation form a tightly connected pathological axis linking metabolic abnormalities to musculoskeletal degeneration. Rather than acting independently, these mechanisms reinforce one another through positive feedback, continuously exacerbating mitochondrial dysfunction, ECM abnormalities, impaired cellular differentiation, cell death and biomechanical deterioration. This sustained inflammatory and oxidative microenvironment provides the foundation for the subsequent development of microvascular dysfunction and defective tissue regeneration.

#### *Microvascular dysfunction as a tissue-level amplifier.*

Microvascular dysfunction is a critical pathological consequence of persistent hyperglycemia and chronic metabolic stress. It acts as a tissue-level amplifier that accelerates musculoskeletal degeneration in T2DM (Table I). Under physiological conditions, an intact microvascular network ensures adequate oxygen and nutrient delivery while facilitating metabolite removal to maintain tissue homeostasis. However, prolonged exposure to hyperglycemia, insulin resistance, OS and chronic inflammation progressively impairs endothelial function, resulting in reduced nitric oxide (NO) bioavailability, excessive ROS production, endothelial cell

dysfunction, basement membrane thickening and poor blood vessel formation (76,85-87). Consequently, tissue blood flow declines, chronic hypoxia develops, and regenerative capacity is progressively weakened across skeletal muscle, tendon, the TBI and bone. Rather than representing an isolated pathological process, microvascular dysfunction reinforces metabolic stress, inflammation and ECM abnormalities, thereby establishing a vicious cycle that accelerates tissue degeneration.

Skeletal muscle is highly dependent on an extensive capillary network to support its substantial metabolic demand (88). In T2DM, diabetic microangiopathy features capillary rarefaction, basement membrane thickening, endothelial dysfunction, and reduced skeletal muscle blood flow. The consequent reduction in oxygen and nutrient delivery aggravates mitochondrial dysfunction and energy deficiency, which further impairs glucose utilization and insulin sensitivity (85,89-91). In addition, diabetic microvascular injury extends to the blood vessels of nerves, producing neural ischemia, axonal degeneration and peripheral neuropathy, which disrupt nerve-to-muscle communication (Fig. 1B and C) (92,93).

The detrimental effects of microvascular dysfunction extend to tendon tissue, which naturally has limited blood supply and therefore possesses a relatively poor intrinsic healing capacity (94). Diabetic tendons exhibit a significant increase in blood vessel density, with vessels extending into the central tendon region, leading to an enlarged vascular cross-sectional area (95). This aberrant vascularization disrupts the normal blood supply pattern and microcirculatory balance in tendons, leading to local tissue hypoxia, accumulation of metabolic waste and ROS generation. Ultimately, this compromises collagen fiber stability, increases tissue brittleness, and contributes to tenocyte damage and collagen degradation (21,74,95-97). Concurrently, this pathological environment triggers an inflammatory response characterized by immune cell activation and the release of pro-inflammatory cytokines. These cytokines disrupt the organized architecture of collagen fibers, leading to a decrease in fiber diameter, a reduction in density, fiber separation and impaired biomechanical properties (74). Furthermore, the impaired microenvironment hinders stem cell migration and differentiation and delays post-injury angiogenesis, collectively weakening the tendon's self-repair capacity (74,94,98,99). Besides, the hypoxic environment also directly impairs the proliferation, differentiation and metabolic function of tenocytes (100-102). Although local hypoxia upregulates vascular endothelial growth factor (VEGF) expression to stimulate angiogenesis, endothelial progenitor cell (EPC) dysfunction, reduced VEGF signaling, and the accumulation of ROS and AGEs result in the formation of aberrant, immature blood vessels that fail to meet the mechanical demands of the tendon (103-108). Moreover, hypoxia promotes M2 macrophage polarization, which accelerates fibrosis and scar formation, disrupting the balance of ECM turnover. This imbalance is reflected in aberrant protease activity, such as elevated MMP-13, leading to tendon structural degeneration and functional impairment (95,109). In summary, microangiopathy-induced local hypoxia and inadequate nutrient supply suppress tenocytes metabolic activity and impair the tendon's self-repair capacity and biomechanical properties. Hypoxia further exacerbates vascular endothelial damage through AGE accumulation and OS, increasing

vascular permeability and promoting the release of inflammatory factors, thereby establishing a vicious cycle (Fig. 2).

Successful tendon-bone healing (TBH) proceeds through sequential inflammatory, proliferative and remodeling phases, all of which depend on adequate vascularization (110-112). Newly formed vessels deliver oxygen, nutrients, growth factors, and reparative cells to the healing interface, while insufficient blood supply delays tissue regeneration and compromises biomechanical recovery (77,113,114). Hyperglycemia disrupts vascular homeostasis through persistent OS and endothelial dysfunction. Excessive ROS damage endothelial cells and impair new blood vessel responses, creating a vicious cycle of inflammation and hypoxia (76,115). Demirag *et al* (116) reported that the degree of vascularization positively correlates with histological maturity of the TBI, particularly with the formation of Sharpey's fibers. Likewise, inhibition of VEGF expression at the TBI suppresses blood vessel formation, thereby reducing tendon maturation and biomechanical strength (117). Although VEGF expression is often elevated in diabetes, endothelial responsiveness to VEGF is diminished because of the reduced expression of VEGF receptors, particularly VEGFR2 (115,118,119). Consequently, angiogenic signaling becomes ineffective, resulting in defective neovascularization and delayed healing. Additionally, both CD31 and endomucin have been identified as critical markers involved in vascular development and TBH (120,121). Exosomes derived from bone marrow mesenchymal stem cells (BMSC-Exos) have been shown to markedly enhance the expression of CD31 and endomucin within the interface microenvironment (26,77). By contrast, diabetic conditions significantly impair the function of BMSCs and suppress CD31 and endomucin expression (122). Accumulating evidence also suggests that angiogenic progenitor cells (APCs) are dysfunctional in diabetes. Insulin resistance impairs the PI3K/Akt pathway and increases intracellular ROS levels, reducing APC mobilization, migration and homing capacity (123-129). Hypoxia further alters cell fate at the TBI. The fibrocartilage region is naturally hypovascular and maintains physiological expression of hypoxia-inducible factors (HIFs) (130). At this interface, progenitor cells expressing SRY-box transcription factor 9 (Sox9) and Scleraxis (Scx) give rise to TSPCs and fibrocartilage cells during early musculoskeletal development (131,132). Under diabetic conditions, chronic inflammation and elevated IL-1 $\beta$  drive excessive HIF-2 $\alpha$  activation (79,131). This aberrant upregulation directly suppresses the tendon-specific transcription factor Scx, impairing the tenogenic differentiation of TSPCs and shifting their fate toward osteogenic and chondrogenic lineages, thereby promoting ectopic calcification or ossification at the TBI (131).

Bone remodeling is likewise highly dependent on an intact vascular network, as osteoblasts, osteoclasts, osteocytes, and skeletal stem cells require continuous vascular support to maintain normal bone turnover. Histological and morphological studies demonstrate that diabetes decreases hematopoietic tissue, increases marrow adiposity, and reduces microvascular density (133). Reduced bone marrow capillary density and microangiopathy further limit blood perfusion, leading to local hypoxia that suppresses osteoblast differentiation while stimulating osteoclast activity, ultimately resulting in direct disruption of the bone structure (133,134). Not only can hypoxia




Figure 2. Dysregulation of the AMPK-eNOS/iNOS-NO pathway, coupled with hypoxia-induced excessive VEGF activation, leads to extracellular matrix metabolic disorder, aberrant and immature angiogenesis, activation of inflammatory signaling, and abnormal macrophage polarization. Collectively, these changes suppress cellular activity, promote apoptosis, inhibit proliferation, and impair the differentiation of TSPCs into tenocytes. NO, nitric oxide; eNOS, endothelial NO synthase; iNOS, inducible NO synthase; TSPCs, tendon stem/progenitor cell; AGEs, advanced glycation end products; VEGF, vascular endothelial growth factor.

affect osteoblast development at early stages, failing to provide proper signals for matrix maturation, but under hypoxic conditions, osteocytes can also increase RANKL expression via the HIF-1 $\alpha$  signaling pathway (135,136). This promotes the differentiation of osteoclast precursors into mature osteoclasts and enhances bone resorption activity (136). The dual effects exacerbate the imbalance of bone remodeling, contributing to trabecular thinning and sparsity.

*ECM dysregulation and impaired regeneration as convergent terminal pathways.* ECM dysregulation and impaired regenerative capacity represent the converging pathways through which chronic metabolic disturbances ultimately compromise musculoskeletal integrity in T2DM (Table I). Following persistent hyperglycemia, AGEs-RAGE activation, OS, chronic inflammation and microvascular dysfunction, the coordinated balance between matrix synthesis and degradation is progressively disrupted. These pathological alterations not only impair the structural organization and biomechanical properties of the ECM but also compromise the regenerative potential of resident stem/progenitor cells, resulting in defective tissue remodeling and incomplete functional recovery.

Hyperglycemia suppresses collagen synthesis through multiple pathways. Inhibition of the AMPK and early growth response factor 1 (EGR1) pathway reduces the expression

of collagen-related genes, including collagen type 1 alpha 1 (Colla1), Colla2, and biglycan (137,138). AGE deposition not only compromises cell viability but also disrupts the structure and function of the ECM, interferes with normal collagen cross-linking, and additionally suppresses collagen synthesis. Collectively, insulin resistance, suppression of AMPK/EGR1 signaling, and AGE accumulation reduce collagen synthesis and contribute to tendon degeneration. Collagen degradation is primarily mediated by MMPs, particularly MMP-9 and MMP-13 (139,140). Hyperglycemia increases their expression and activity, leading to excessive collagen breakdown and disruption of collagen fiber organization (141-143). Under physiological conditions, TIMPs maintain a balance with MMPs to preserve ECM integrity (144). However, T2DM reduces TIMP activity, resulting in excessive MMP-mediated collagen degradation (20). This systemic imbalance between synthetic and degradation drives tissue-specific structural disorder. In tendon, T2DM disrupts collagen homeostasis by reducing collagen synthesis while enhancing degradation, resulting in collagen disorganization, altered fiber architecture, and impaired biomechanical properties that predispose tendons to degeneration and injury (15). At the TBI, the ECM of the fibrocartilage region is primarily composed of collagens and proteoglycans, whose balance is tightly regulated by MMPs and TIMPs (131,145,146). Under diabetic conditions,

hyperglycemia, hypoxia and chronic inflammation disrupt the MMP/TIMP balance, resulting in excessive matrix degradation (146-153). Simultaneously, AGEs accumulate within the TBI and form abnormal cross-links with collagen and proteoglycans, reducing tissue elasticity and impairing cell-matrix interactions (50,58,70,154). In bone, osteoclasts degrade the bone matrix by creating an acidic microenvironment and secreting collagenases and other proteolytic enzymes (155). Additionally, the structural integrity of collagen fibers within the bone matrix is disrupted, with compromised fiber architecture and reduced matrix ductility and strength (59,156). These ECM-level abnormalities collectively undermine bone matrix quality and mechanical performance.

Simultaneously, the regenerative capacity of tissue-resident stem cells is severely compromised across all four tissues. Skeletal muscle satellite cells (MuSCs) are essential for muscle repair and regeneration, but increased inflammation and excessive OS impair MuSC function and suppress myogenic differentiation (157-160). Inflammatory mediators such as IL-6 and TNF- $\alpha$  inhibit PI3K/AKT/mTOR signaling while activating NF- $\kappa$ B and forkhead box O (FoxO) pathways, thereby impairing muscle regeneration (159,161,162). The reduction of myogenic regulatory factors, including MyoD1 and myogenin, further impairs muscle regenerative capacity, leading to reduced cross-sectional area of muscle fibers and atrophy (163). In tendon, TSPCs play a central role in maintaining tendon homeostasis and promoting tissue repair (164-166). Under T2DM conditions, TSPC proliferation is reduced while cell death is increased. Hyperglycemia and AGE accumulation alter TSPC fate, promoting abnormal osteogenic and chondrogenic differentiation at the expense of tenogenic differentiation, which contributes to tendon calcification, structural abnormalities and functional impairment (18,19,167,168). Impaired self-renewal, proliferation and differentiation of TSPCs directly compromise tendon regeneration, contributing to collagen disorganization, excessive vascularization, persistent inflammation, fibrosis and reduced biomechanical performance (19). Reduced collagen synthesis and enhanced degradation lead to impaired structural and mechanical properties, while TSPC dysfunction compromises the tendon's self-repair capacity and increases the risk of rupture. At the TBI, tendon-derived stem cells (TDSCs) are critical for fibrocartilage regeneration during tendon-bone healing (111,169-173). However, hyperglycemic conditions directly induce TDSC apoptosis, inhibit their proliferation, and deplete the available TDSC pool for repair (18,174). Consequently, fibrocartilage exposed to repetitive mechanical loading becomes increasingly vulnerable to thinning, structural disorganization and degeneration. Fibrotic scar formation replaces the native fibrocartilage transition zone, resulting in disorganized collagen fibers, poor biomechanical properties, and compromised load-bearing capacity (26). In bone, the osteogenic potential of MSCs is disrupted. Elevated glucose levels promote adipogenic differentiation through activation of adipogenic transcription factors, including peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) and adipocyte protein 2 (aP2), while suppressing osteogenic regulators such as Runx2 and Osterix (34,35). Consequently, MSCs preferentially differentiate into adipocytes rather than osteoblasts, resulting in increased bone marrow adiposity and reduced osteoblast numbers (36,37). The imbalance between

osteoblast-mediated formation and osteoclast-mediated resorption, coupled with preferential adipogenic differentiation of MSCs, leads to reduced bone formation, trabecular deterioration, cortical thinning and increased skeletal fragility (175). Together, these tissue-specific stem/progenitor cell dysfunctions, coupled with systemic ECM remodeling abnormalities, establish a self-perpetuating cycle of structural degradation and failed repair across the entire musculoskeletal system. These findings highlight the critical need for therapeutic strategies that simultaneously target ECM homeostasis and endogenous regenerative mechanisms.

### 3. Tissue-specific pathological mechanisms underlying T2DM musculoskeletal disorders

Although hyperglycemia, AGEs-RAGE signaling, OS, chronic inflammation, microvascular dysfunction and ECM dysregulation collectively drive musculoskeletal degeneration in T2DM, each tissue exhibits distinct biological characteristics. These characteristics determine their unique pathological response to chronic metabolic stress. These tissue-specific mechanisms further contribute to the heterogeneity of diabetic musculoskeletal complications and represent potential targets for precision treatment interventions (Table II).

*Skeletal muscle.* Unlike other musculoskeletal tissues, skeletal muscle serves as the principal organ responsible for post-prandial glucose disposal and whole-body energy metabolism (9). A characteristic feature of diabetic skeletal muscle is the progressive disruption of intracellular energy metabolism. Defective insulin signaling impairs GLUT4 translocation and glucose uptake, forcing muscle fibers to rely increasingly on lipid oxidation (15,16). However, incomplete fatty acid oxidation results in the accumulation of toxic lipid intermediates, particularly DAG, which activates PKC and further suppresses insulin signaling, forming a vicious cycle of metabolic dysfunction (16). In parallel, mitochondrial dysfunction serves as a fundamental mechanism underlying insulin resistance and muscle atrophy (72,176). In patients with T2DM, skeletal muscle exhibits reduced mitochondrial density, impaired oxidative phosphorylation, and abnormal mitochondrial morphology, resulting in diminished adenosine triphosphate (ATP) production and compromised energy metabolism (14). These alterations contribute to the loss of muscle mass, elevated fatigue and reduced exercise tolerance (72). Mitochondrial dynamics are regulated by fusion proteins, including mitofusin-1 (Mfn1), Mfn2 and optic atrophy 1 (Opa1) (177). Reduced expression of Mfn2 and Opa1 in T2DM is associated with impaired mitochondrial fusion and decreased mitochondrial function (178-180). Hyperglycemia promotes excessive mitochondrial fission and ROS generation, further exacerbating mitochondrial damage and OS (97,178,181-184). In addition, mitochondrial creatine kinase 2 (CKMT2), an important regulator of oxidative phosphorylation and mitochondrial respiration (185,186). Decreased CKMT2 impairs phosphocreatine formation and ATP generation, leading to reduced glucose oxidation and worsening energy deficits (186). Together, mitochondrial dysfunction, OS, and impaired substrate utilization contribute to progressive muscle weakness and atrophy.

Table II. Tissue-specific mechanisms, signaling pathways, pathological alterations, and therapeutic strategies in T2DM-associated musculoskeletal disorders.

| Category                                | Skeletal muscle                                                                                                                                                                                                                                                                                                                                                                                                  | Tendon                                                                                                                                                                                                                                                                                                                                               | Tendon-bone interface                                                                                                                                                                                                                                                                                            | Bone                                                                                                                                                                                                                                                                                                   | (Refs.)                                                                                 |
|-----------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| Tissue-Specific Mechanism               | GLUT4 translocation↓<br>→ Impaired insulin-stimulated glucose uptake; Mfn1/Mfn2/Opa1↓, excessive fission<br>→ Mitochondrial dynamics imbalance; CKMT2 dysfunction → oxidative phosphorylation ↓, ATP↓; Type I oxidative fibers → type II glycolytic fibers; UPS-mediated proteolysis → MuRF1/MAFbx ↑ via FoxO; Suppressed mTOR-dependent protein synthesis → PI3K/Akt/mTOR↓; Neuromuscular junction degeneration | Excessive collagen cross-linking through AGEs and TGase-mediated; TSPC dysfunction → proliferation↓, apoptosis↑; dysfunction; Aberrant osteo/chondrogenic differentiation; eNOS/iNOS-NO axis dysregulation → excessive NO, endothelial M2 macrophage polarization → TGF-β/VEGF secretion, fibrosis; Impaired mechanotransduction ECM disorganization | Impaired fibrocartilage regeneration; Defective osteochondral transition; Integrin-mediated mechanotransduction dysfunction → AGEs-modified RGD domain; Integrin α/β; Macrophage imbalance → pro-inflammatory and pro-fibrotic; Disrupted tendon-bone integration; TDSC dysfunction → apoptosis↑, proliferation↓ | Osteoblast dysfunction → Runx2/Osterix↓, OCN/OPN/ALP↓; Enhanced osteoclastogenesis → RANKL sensitivity↑, M-CSF responsiveness↑; Osteocyte senescence and apoptosis → sclerostin↑, SASP↑, Wnt/β-catenin↓; BMSC adipogenic differentiation → PPARγ↑, Runx2↓; Cortical porosity and impaired bone quality | (28,34,36-38, 179-186, 190-192,198, 211,214,215, 217,221,223, 225,255,256, 259,260,263) |
| Major signaling pathways                | mTOR; AMPK; FOXO; NF-κB; CKMT2; IGF-1/PGC-1α; MuRF1/MAFbx; PI3K/Akt/ GLUT4; Mfn1/Mfn2/Opa1                                                                                                                                                                                                                                                                                                                       | NF-κB; MAPK; MMP/TIMP; TGF-β/Smad; Wnt/β-catenin; M2 polarization; eNOS/iNOS-NO; TGase-mediated crosslinking                                                                                                                                                                                                                                         | Integrin α/β; FGF2; VEGF; PI3K/Akt; Wnt/β-catenin; BMP/TGF-β; RANKL/OPG                                                                                                                                                                                                                                          | Notch; RANKL/OPG; PI3K/Akt; GF-β/Smad; Sclerostin/Wnt/β-catenin                                                                                                                                                                                                                                        | (38,72,109, 115,118,119, 137,138,145, 146,178,186, 198,221,229, 230,252,253)            |
| Representative pathological alterations | Sarcopenia; Impaired regenerative capacity; Mitochondrial dysfunction; Insulin resistance; Neuromuscular junction degeneration; Reduced muscle strength and endurance                                                                                                                                                                                                                                            | Tendinopathy; Delayed healing; Increased stiffness, reduced elasticity; Impaired biomechanical properties; Increased rupture risk; Collagen disorganization and fiber separation                                                                                                                                                                     | Delayed TBI healing; Poor fibrocartilage formation; Impaired osseointegration; Mechanotransduction dysfunction; Reduced biomechanical strength; Fibrotic scar replacement                                                                                                                                        | Trabecular microarchitectural deterioration; Cortical porosity; Reduced bone quality; Delayed fracture healing; Increased skeletal fragility                                                                                                                                                           | (14,22,26-30, 37,72,74,84, 133,134,154, 164,175,178, 186,207,208, 244,252-259)          |
| Potential Therapeutic Strategies        | Exercise: AMPK activation, improved insulin sensitivity; Metformin: AMPK activation, anti-inflammation, improved insulin signaling; Insulin sensitizers: thiazolidinediones;                                                                                                                                                                                                                                     | AGEs-RAGE inhibition: reducing cross-linking and inflammation; Antioxidant therapy: reducing OS; TSPC-based therapy: promoting tenogenic differentiation;                                                                                                                                                                                            | MSC/exosome therapy: enhancing CD31/endomucin, angiogenesis; Macrophage immunomodulation: balancing M1/M2; Angiogenic factors: promoting neovascularization;                                                                                                                                                     | Anti-resorptive agents: RANKL inhibition; Anabolic therapies: sclerostininhibition, Wnt activation; Metformin: promotes osteogenesis, inhibits osteoclastogenesis; Wnt activation:                                                                                                                     | (3-8,26,40, 77, 113,210, 266-269, 273-277)                                              |

Table II. Continued.

| Category | Skeletal muscle                                                                        | Tendon                                                                                       | Tendon-bone interface                                                                                   | Bone                                                                                                    | (Refs.) |
|----------|----------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|---------|
|          | Mitochondrial-targeted antioxidants: reducing OS; Senolytics: clearing senescent cells | Anti-fibrotic therapy: targeting TGF- $\beta$ /Smad; Exosome therapy: promoting regeneration | Bioactive scaffolds: providing structural support; Magnesium-based biomaterials: promoting osteogenesis | activation: bone formation; Senolytics: clearing senescent osteocytes; Antioxidant therapy: reducing OS |         |

AGE, advanced glycation end product; Akt, protein kinase B; ALP, alkaline phosphatase; AMPK, AMP-activated protein kinase; BMP, bone morphogenetic protein; CKMT2, mitochondrial creatine kinase 2; ECM, extracellular matrix; eNOS, endothelial nitric oxide synthase; FGF2, fibroblast growth factor 2; FoxO, forkhead box O; GLUT4, glucose transporter type 4; IGF-1, insulin-like growth factor-1; iNOS, inducible nitric oxide synthase; MAFbx, muscle atrophy F-box; M-CSF, macrophage colony-stimulating factor; Mfn1, mitofusin-1; Mfn2, mitofusin-2; MMP, matrix metalloproteinase; MSC, mesenchymal stem cell; mTOR, mechanistic target of rapamycin; MuRF1, muscle RING-finger protein-1; NF- $\kappa$ B, nuclear factor kappa B; NO, nitric oxide; OCN, osteocalcin; OPG, osteoprotegerin; OPN, osteopontin; PGC-1 $\alpha$ , peroxisome proliferator-activated receptor gamma coactivator-1 alpha; PI3K, phosphoinositide 3-kinase; PPAR $\gamma$ , peroxisome proliferator-activated receptor gamma; RAGE, receptor for advanced glycation end products; RANKL, receptor activator of nuclear factor- $\kappa$ B ligand; ROS, reactive oxygen species; Runx2, runt-related transcription factor 2; SASP, senescence-associated secretory phenotype; Smad, mothers against decapentaplegic homolog; STAT, signal transducer and activator of transcription; TDSC, tendon-derived stem cell; TGF- $\beta$ , transforming growth factor- $\beta$ ; TGase, transglutaminase; TIMP, tissue inhibitor of metalloproteinase; TSPC, tendon stem/progenitor cell; UPS, ubiquitin-proteasome system; VEGF, vascular endothelial growth factor; Wnt, wingless-related integration site.

Skeletal muscle loss in T2DM results from both enhanced protein degradation and impaired protein synthesis (187). T2DM is associated with a shift from oxidative type I fibers toward glycolytic type II fibers, reducing oxidative capacity and insulin sensitivity while increasing susceptibility to muscle wasting (84,188-193). Insulin resistance suppresses the anabolic IRS/PI3K/Akt/mTOR pathway while concurrently activating the ubiquitin-proteasome system (UPS), which is the principal pathway responsible for protein degradation in skeletal muscle (72,194,195). The muscle-specific enzymes muscle RING-finger protein-1 (MuRF1) and muscle atrophy F-box (MAFbx/atrogen-1) are markedly upregulated during muscle atrophy (196). The FoxO transcription factors regulate the expression of MuRF1 and MAFbx (196,197). Reduced insulin signaling decreases FoxO phosphorylation, thereby activating the UPS and further amplifying proteolysis (84,198). Furthermore, inflammatory cytokines further enhance UPS activity through NF- $\kappa$ B and janus kinase/signal transducers and activators of transcription (JAK/STAT) signaling pathways, accelerating proteolysis and muscle loss (38,72,84,199-202). Under healthy conditions, insulin activates the PI3K/Akt signaling to promote glucose uptake and stimulate mTORC1-dependent protein synthesis (203,204). In T2DM, impaired signaling suppresses these pathways, resulting in diminished protein synthesis (73,205,206). Combined with diabetic peripheral neuropathy (DPN), neuromuscular junction degeneration and denervation-induced muscle atrophy, these metabolic abnormalities contribute to the progressive decline in muscle strength, endurance, and physical performance characteristic of diabetic sarcopenia.

**Tendon.** Compared with skeletal muscle, tendon pathology in T2DM is characterized primarily by disruption of collagen homeostasis and deterioration of biomechanical properties. Diabetes profoundly alters tendon mechanical behavior,

reducing its ability to withstand and transmit mechanical loads. Structural abnormalities such as collagen disorganization, fibrosis and calcification compromise tendon integrity and increase susceptibility to injury (22,207,208). As mentioned previously, collagen cross-linking is essential for maintaining tendon mechanical properties. Enzymatic cross-linking enhances tendon stiffness and fatigue resistance by stabilizing collagen fibril arrangement. By contrast, non-enzymatic cross-linking, particularly AGEs, are minimal under physiological conditions but accumulate markedly in pathological states such as diabetes (209). Hyperglycemia accelerates non-enzymatic glycosylation, leading to AGE deposition between collagen fibers. This promotes abnormal intermolecular cross-linking that impede fibrillar sliding, increasing tissue stiffness, reducing overall elasticity and ductility (40,210-212). Transglutaminase (TGase) is an enzyme that catalyzes acyl-transfer reactions to form covalent cross-links between proteins, normally enhancing the mechanical strength and fatigue resistance of tissues (213). However, hyperglycemia elevates TGase activity within tenocytes, which exacerbates non-enzymatic cross-linking and impairs dynamic collagen sliding (74,214,215). These aberrant cross-links disrupt normal fibrillar organization and helical structure, ultimately reducing the elastic modulus of the tendon (216-219).

Hyperglycemia also activates endothelial NO synthase and inducible NO synthase, resulting in a short-term increase in the absolute production of NO (220,221). Elevated NO levels inhibit fibroblast proliferation and collagen synthesis, thereby reducing tendon elasticity (222). In the short term, excessive NO impairs vascular endothelial barrier function. The endothelial dysfunction further induces OS via activation of NADPH oxidase and glycolytic pathways, leading to a marked rise in ROS. The overproduction of ROS reduces NO bioavailability through both the degradation of tetrahydrobiopterin and direct oxidation of NO, thereby exacerbating endothelial dysfunction (223-227). Such




Figure 3. Schematic representation of the TBI illustrating its four gradient zones and corresponding cellular compositions. The native TBI exhibits coordinated gradients in cell phenotype, mineralization, and collagen fiber alignment across these distinct regions. TBI, tendon-bone interface.

impairment diminishes the regenerative potential of MSCs, compromising the tendon's repair capacity. Tendon elasticity relies on the organized arrangement of collagen and the homeostasis of elastin. Impaired repair results in abnormal collagen cross-linking and elastin degradation, collectively reducing elasticity and increasing brittleness (18,228). Concurrently, aberrant M2 macrophage polarization promotes the secretion of pro-fibrotic factors (for example, TGF- $\beta$  and VEGF), which stimulate pathological scarring (109,229,230). These changes lead to stiffened tendon fibers, diminished tensile strength, and an elevated risk of rupture (Fig. 2).

**TBI.** The TBI is a specialized transitional tissue that comprises two distinct insertion types, which are direct and indirect insertions, enabling efficient load transfer between tendon and bone (231,232). It consists of tendon, unmineralized fibrocartilage, mineralized fibrocartilage, and bone, forming a graded structure characterized by progressive changes in cellular phenotype, collagen composition, and mineral content (Fig. 3) (112,233-238). The indirect insertion is characterized by Sharpey's fibers, which anchor tendon directly into bone (239). The two insertion types differ primarily in collagen maturity and extent of new bone formation. Biomechanically and biologically, indirect insertions exhibit inferior properties compared with direct insertions (26).

By providing a gradual transition in stiffness, this native fibrocartilage interface minimizes stress concentration between tissues with vastly different mechanical properties, ensuring effective force transmission (240-244). However, this transmission relies heavily on both the structural integrity of the interface and the capacity of resident cells to sense and respond to mechanical stimuli. Biomechanical studies have shown that diabetes significantly reduces the

Young's modulus, stiffness and ultimate failure load of the TBI (22,154). Diabetes-induced fibrosis disrupts the hierarchical architecture by replacing fibrocartilage with scar tissue. This loss of fibrocartilage, alongside reduced vascularization and collagen disorganization, severely compromises the interface's stress-buffering capacity. Consequently, these alterations impair force transmission and contribute to a higher incidence of rotator cuff tears, Achilles tendinopathy, postoperative retears, and delayed healing in diabetic patients (244). Experimental studies further demonstrate that diabetic animals exhibit reduced fibrocartilage formation, poorer collagen organization, and lower biomechanical strength following rotator cuff repair compared with healthy controls (154).

Beyond structural degradation, cells within the TBI depend on mechanotransduction pathways to continuously sense mechanical cues from the extracellular environment (245-246). Integrins, which are heterodimeric transmembrane receptors composed of  $\alpha$  and  $\beta$  subunits, are key mediators of this process. By linking the ECM to the actin cytoskeleton, integrins convert mechanical forces into intracellular biochemical signals that regulate tissue remodeling and repair (247,248). Multiple studies indicate that hyperglycemia impairs this integrin-mediated signaling. AGEs can modify the Arg-Gly-Asp (RGD) domain of fibronectin, disrupting its interaction with integrins, thereby altering ECM stiffness and attenuating mechanotransduction efficiency (249). Furthermore, in human proximal tubular epithelial (HK-2) cells cultured under high-glucose conditions, the expression of specific integrin subunits is significantly reduced (250). Chronic hyperglycemia has also been reported to delay fibronectin and integrin expression, impairing the integration of implants into bone (251). These findings suggest that diabetes downregulates integrin expression, disrupts interactions between integrins and ECM




**Figure 4.** Mechanisms of tendon-bone and bone quality decline in type 2 diabetes mellitus. (A) Hyperglycemia and oxidative stress drive persistent accumulation of AGEs at the TBI, leading to sustained activation of AGEs-RAGE signaling and chronic inflammatory immune responses. This pathological cascade activates the TGF-β1/Smad3 pathway, which promotes fibroblast proliferation and fibrotic remodeling. Meanwhile, abnormal cross-linking of collagen and proteoglycans reduces tissue elasticity and compromises interface integrity. In addition, AGEs modify the RGD domain of fibronectin, disrupting integrin binding, altering extracellular matrix stiffness, and impairing mechanotransduction. These structural abnormalities weaken the mechanical buffering capacity and force transmission of the enthesis. Concurrent endothelial dysfunction and aberrant hypoxia signaling further disrupt stem cell differentiation and tenocyte transcriptional programs. Collectively, these changes promote fibrocartilage calcification and heterotopic ossification, ultimately leading to structural deterioration and functional failure of the TBI. (B) The accumulation of AGEs, lipid metabolic disturbances, and chronic inflammatory microenvironment suppress osteoblast proliferation and differentiation while promoting apoptosis, leading to markedly reduced bone formation. In parallel, AGEs-induced osteocyte senescence and apoptosis upregulate sclerostin expression, thereby inhibiting osteogenic signaling pathways such as Wnt/β-catenin and further impairing bone anabolism. Meanwhile, remodeling of the bone microenvironment, characterized by an imbalanced RANKL/OPG ratio, acidic conditions, and disrupted insulin signaling, promotes osteoclast differentiation, maturation and resorptive activity. Driven by the combined effects of suppressed osteogenesis and enhanced osteoclastogenesis, bone remodeling shifts from homeostatic balance toward negative balance, ultimately resulting in trabecular bone sparsity, cortical thinning and progressive bone loss, which define diabetic bone degeneration and skeletal fragility. AGEs, advanced glycation end products; TBI, tendon-bone interface; RAGE, receptor for AGEs; RGD, Arg-Gly-Asp; ROS, reactive oxygen species; TDSC, tendon-derived stem cell; RANKL, receptor activator of nuclear factor-κB ligand; OPG, osteoprotegerin.

components, and consequently impairs mechanosensation, leading to abnormal mechanical stress transmission at the TBI (Fig. 4A).

**Bone.** Bone exhibits a distinct pathological response to diabetes because its homeostasis depends on the dynamic coupling of osteoblast-mediated bone formation,

osteoclast-mediated bone resorption, and osteocyte-mediated mechanosensation (252-255). A hallmark of diabetic bone disease is uncoupled bone remodeling. As noted previously, hyperglycemia suppresses osteoblast differentiation while simultaneously promoting excessive osteoclast development and bone resorption. Another distinctive mechanism involves osteocyte dysfunction. Osteocytes are the most abundant and essential cellular component of the skeletal system, acting as the primary sensors of mechanical and biochemical stimuli and playing an irreplaceable role in bone remodeling (256). They are derived from osteoblasts, and a subset of osteoblasts becomes embedded within the mineralized bone matrix to differentiate into osteocytes. These cells are interconnected through a complex network and produce a variety of signaling molecules that regulate osteoblast and osteoclast activity (255). Hyperglycemia impairs the mechanosensory function of osteocytes, accelerates cellular aging, and compromises bone strength (37,257). It can also directly induce osteocyte death, disrupting the integrity of the osteocyte network and impairing skeletal mechanical adaptability (258,259). Cellular aging is a stress-responsive cell fate characterized by durable growth arrest, extensive chromatin remodeling, and acquisition of a proinflammatory secretory program known as the senescence-associated secretory phenotype (SASP) (260). The accumulation of AGEs and senescent cells compromises osteocyte viability and function through the activation of RAGE signaling and the secretion of proinflammatory SASP factors, ultimately enhancing the production of sclerostin by osteocytes. When sclerostin binds to Wnt co-receptors, it inhibits bone formation and reduces osteocyte mechanosensitivity, thereby contributing to poor bone quality in diabetic patients (261-263). Meanwhile, hyperglycemia drives BMSCs toward a metabolically stressed adipogenic lineage rather than osteoblasts. This shift not only causes marrow fat accumulation and impaired trabecular formation, but it also triggers inflammation mediated by hyaluronan, ultimately exacerbating trabecular demineralization (37). These alterations collectively impair fracture healing, reduce bone quality, and increase skeletal fragility despite relatively preserved bone mineral density in numerous patients (Fig. 4B).

#### 4. Conclusion and perspective

*Critical appraisal of the evidence, knowledge gaps and future directions.* Although the preceding sections summarize the proposed mechanisms underlying diabetes-related musculoskeletal disorders, the strength of evidence supporting each pathway warrants critical evaluation. Much of the available evidence, particularly for AGEs-RAGE signaling and chronic inflammation, comes from *in vitro* and animal studies. Therefore, its direct translation to human pathophysiology should be interpreted with caution. For instance, AGE accumulation is consistently associated with increased tendon stiffness and reduced elasticity in rodent models. However, evidence from human studies is largely correlative and is often confounded by disease duration and comorbidities (22,52,216). Likewise, mitochondrial dysfunction is considered a central feature of numerous proposed mechanisms. However, current evidence mainly comes from associative studies showing reduced oxidative capacity and abnormal mitochondrial

morphology in diabetic muscle (14,178). Whether these mitochondrial alterations initiate insulin resistance or arise as a consequence of the diabetic metabolic milieu remains unresolved (264).

Notable contradictions exist within literature. For example, although chronic hyperglycemia is widely accepted to impair angiogenesis, some studies report elevated VEGF levels in diabetic tissues such as the retina. This finding suggests a paradoxical upregulation in response to hypoxia under certain pathological conditions (103,104). By contrast, peripheral tissues such as skeletal muscle and tendon exhibit reduced VEGFR2 signaling and poor neovascularization (115,118,119). These observations suggest that the overall angiogenic response depends on the complex interplay of multiple dysregulated pathways and may be tissue dependent. A significant knowledge gap remains in the understanding of the TBI. While the role of TGF- $\beta$ 1 in fibrosis is well-established, the precise molecular triggers that shift tissue repair from regenerative fibrocartilage formation to pathological scarring in diabetes remain largely unknown (244,265). In addition, the specific contributions of different immune cell subpopulations, particularly the dynamic phenotypic switching of macrophages, remain incompletely understood. Most available evidence comes from a limited number of studies. Finally, most current evidence is derived from single-tissue or single-mechanism studies, making it difficult to understand the integrated, system-level response to diabetes. Future research should prioritize longitudinal human studies, apply multi-omics approaches to capture the complexity of the disease, and develop experimental models that simultaneously investigate interactions among skeletal muscle, tendon, bone and their interfaces. Such integrative approaches will facilitate the identification of robust therapeutic targets and their translation into effective interventions for diabetes-related musculoskeletal complications.

*Conventional antidiabetic and anti-osteoporotic therapies bridging metabolism and musculoskeletal health.* Conventional antidiabetic medications and anti-osteoporotic therapies not only improve systemic metabolic homeostasis but also exert pleiotropic effects on musculoskeletal tissues by targeting shared pathological mechanisms underlying T2DM-associated musculoskeletal disorders. Rather than acting solely on glucose metabolism or bone remodeling alone, these therapies serve as an important bridge between metabolic regulation and musculoskeletal health through coordinated modulation of inflammation, OS, AGEs, and stem cell function.

Among conventional antidiabetic agents, metformin has been the cornerstone of first-line pharmacological therapy for T2DM for over six decades because of its well-established efficacy, excellent safety profile, minimal risk of hypoglycemia, favorable effects on body weight, cost-effectiveness, robust long-term clinical evidence, and pleiotropic metabolic benefits (266). Beyond its glucose-lowering effect, metformin activates AMPK, thereby suppressing NF- $\kappa$ B-mediated inflammatory signaling and reducing the production of pro-inflammatory cytokines such as TNF- $\alpha$  and IL-6 (267). It also inhibits the formation and accumulation of AGEs, consequently attenuating AGEs-RAGE signaling, OS, apoptosis, and tissue fibrosis (267-269). Since excessive AGEs

accumulation disrupts the stem cell microenvironment and impairs proliferation and lineage commitment, metformin may indirectly preserve stem cell function by reducing glycation stress (270-272). Moreover, accumulating evidence suggests that metformin promotes osteogenic differentiation while suppressing osteoclast activity through multiple signaling pathways, including AMPK, Nrf2/HO-1, SIRT1 and mTOR (273,274). Combination therapy with metformin and anti-osteoporotic agents has also been shown to improve both glycemic control and diabetes-induced bone loss in experimental models (275). Anti-osteoporotic therapies primarily restore skeletal integrity by re-establishing the balance between bone resorption and bone formation. Antiresorptive agents, including RANKL-neutralizing antibody denosumab, suppress osteoclast differentiation and induce osteoclast apoptosis, thereby reducing excessive bone resorption (276). By contrast, anabolic agents such as sclerostin-neutralizing antibody romosozumab directly stimulate osteoblast differentiation and bone formation (277). Beyond increasing bone mineral density and reducing fracture risk, these therapies improve the musculoskeletal microenvironment by preserving bone remodeling homeostasis, thereby providing a favorable niche for stem cell-mediated tissue regeneration.

Conventional antidiabetic and anti-osteoporotic therapies share complementary mechanisms that extend beyond their primary therapeutic indications. By simultaneously improving metabolic control, suppressing chronic inflammation and OS, limiting AGEs accumulation, and preserving stem cell activity, these agents help maintain musculoskeletal tissue homeostasis and strengthen the link between systemic metabolism and the musculoskeletal system.

*Application of emerging technologies in diabetes.* In recent years, rapid advances in emerging technologies have transformed the prevention, diagnosis, monitoring and management of diabetes. Continuous glucose monitoring, artificial intelligence-assisted decision support systems, digital health tools, biosensors, and advanced drug delivery approaches are reshaping current models of diabetes care. Meanwhile, progress in omics research, big data analytics, and regenerative medicine has provided new insights into disease mechanisms and facilitated the development of personalized therapeutic strategies. The integration of biotechnology, data science and clinical practice marks the beginning of a new stage in precision diabetology, with an emphasis on improving glycemic control, preventing complications, and enhancing the overall quality of life for individuals with diabetes.

*Single-cell RNA sequencing (scRNA-seq) reveals the heterogeneity of the diabetic microenvironment.* With the rapid development of high-throughput sequencing technologies, the genetic and transcriptomic mechanisms underlying diabetes have been progressively uncovered. The precise regulation of glucose homeostasis depends on the coordinated activity of multiple endocrine cell types within the pancreatic islets (278). Human pancreatic islets are composed mainly of  $\beta$  cells (~54%),  $\alpha$  cells (35%) and  $\delta$  cells (11%), along with smaller populations of  $\gamma$ /PP and  $\epsilon$  cells (279,280). Increasing evidence suggests that alterations in the proportion or function of these endocrine cells are closely associated with the pathophysiological processes of monogenic diabetes, T1DM

and T2DM (278). However, because pancreatic islets account for only ~1% of pancreatic tissue and contain rare cell populations such as  $\delta$  and  $\epsilon$  cells, whole-genome sequencing at the single-islet level has long been technically challenging (281). The emergence of scRNA-seq has overcome numerous of these limitations. It enables detailed investigation of the molecular mechanisms underlying diabetes and provides new opportunities for clinical diagnosis and treatment (278,282).

Yang *et al* (283) applied scRNA-seq to pancreatic islets from T1DM mouse models and generated a comprehensive single-cell atlas, identifying 11 major cellular subpopulations. Their results revealed marked heterogeneity among  $\alpha$  and  $\beta$  cell subsets and demonstrated a significant reduction in the number of mature  $\alpha$  and  $\beta$  cells, which was further confirmed by flow cytometry. These observations highlight the potential roles of specific  $\alpha$ - and  $\beta$ -cell subpopulations in T1DM progression (283). Honaradoost *et al* (284) conducted scRNA-seq analyses of peripheral blood mononuclear cells (PBMCs) from patients with T1DM. Their study demonstrated widespread immune dysregulation across 13 immune cell types, consistent with abnormalities observed in pancreatic islets. Based on these findings, the authors developed a T1DM Meta-Gene Z-score (TMZ score) model, providing a framework for personalized therapeutic strategies in diabetes (284). Analysis of pancreatic islets from patients with T2DM further showed that even the rare  $\delta$  and  $\epsilon$  cells specifically expressed important molecules such as the GHRL receptor and GHRL, suggesting that these rare cell subpopulations may play unique roles in the regulation of islet function (285). Diabetic foot ulcers (DFUs) represent another major complication of diabetes, affecting ~19-34% of patients during the course of the disease (286). Human skin is composed primarily of keratinocytes and fibroblasts, along with immune cells, melanocytes, adipocytes and endothelial cells (287,288). Among these, keratinocyte dysfunction is considered a major contributor to delayed wound repair (282,289). Luo *et al* (290) combined bulk RNA sequencing with scRNA-seq to investigate the mechanisms of mitochondrial programmed cell death in DFUs. They identified key genes, including BCL2 and LIPT1, that were downregulated in DFU tissues, with keratinocytes identified as the most affected cell type (290). Similarly, Wang *et al* (291) applied scRNA-seq to explore wound-healing mechanisms in DFUs and identified 1,948 differentially expressed genes between healing and non-healing wounds, including 1,198 upregulated and 685 downregulated genes. Functional enrichment analysis further demonstrated that the CCL2-ACKR1 signaling axis plays a crucial role in regulating endothelial cell activity and promoting wound healing in diabetic ulcers (291). Collectively, these studies suggest that scRNA-seq serves as a powerful tool for decoding the molecular basis of keratinocyte dysfunction and the complexity of wound repair in T2DM. By revealing disease-specific signaling pathways and potential therapeutic targets, this technology provides opportunities for precision medicine in chronic wound management. Beyond mechanistic studies, scRNA-seq is gradually evolving from a powerful tool for elucidating disease mechanisms into an important technological platform for translational diabetes research. By resolving cellular heterogeneity across pancreatic islets, immune cells and diabetic wounds, and identifying disease-associated molecular signatures, scRNA-seq may

provide clinically relevant insights for biomarker discovery, patient stratification and therapeutic target identification. For example, scRNA-seq-based quality assessment of donor islets before transplantation may facilitate donor selection and potentially improve transplantation outcomes. Moreover, the PBMC-derived TMZ score has shown significant associations with established prognostic immune biomarkers and clinical trial drug responses in T1DM, suggesting its potential utility in identifying patient populations that are more likely to benefit from immunomodulatory therapies. Overall, integrating scRNA-seq with advanced computational and multi-omics techniques holds great promise for developing personalized therapeutic interventions based on the molecular profiles of individual cells.

Although the widespread application of scRNA-seq in both basic research and clinical studies continues to expand, several technical challenges remain. The single-cell capture procedure often compromises cellular integrity and viability, highlighting the need for further optimization to improve throughput and precision. In addition, scRNA-seq produces high-dimensional datasets that are inherently noisy, and reducing this noise remains a major challenge for reliable data analysis. Moreover, cell differentiation and proliferation are inherently dynamic processes, making it difficult for current scRNA-seq methods to accurately distinguish between cell types (292). Beyond these technical limitations, the relatively high cost of scRNA-seq also limits its broader adoption as a routine research tool. Clinical translation of scRNA-seq is further challenged by the limited accessibility of disease-relevant tissues, the lack of standardized analytical workflows, and the need for large-scale prospective validation to ensure the reproducibility and clinical interpretability of single-cell findings across diverse patient populations. Continued improvements in sequencing technologies, bioinformatics algorithms, analytical standardization and cost efficiency will be essential for advancing the application of scRNA-seq in diabetes research and clinical practice.

*Targeted delivery systems enhance local therapeutic efficacy.* Precision medicine has become a key strategy for managing a wide range of diseases, and targeted drug delivery systems (TDDS) are now emerging as one of its central technologies in diabetes treatment. Insulin remains the fundamental therapy for maintaining glucose homeostasis in patients with T1DM and advanced T2DM. However, conventional insulin therapy relies primarily on frequent subcutaneous injections, such as rapid-acting insulin before meals and long-acting insulin at bedtime, to mimic physiological secretion patterns (293). Repeated injections often lead to pain, local tissue injury, infection and neuropathy, resulting in poor patient compliance (294). In addition, factors such as injection site, depth and dosage can significantly affect insulin absorption kinetics, further increasing the uncertainty of glycemic control (295). To overcome the limitations of injection-based therapy, research has increasingly focused on non-invasive and intelligent TDDS strategies. These systems enable targeted drug delivery to diseased sites and achieve rhythm-synchronized release in response to physiological demands. By improving local therapeutic efficacy while minimizing systemic side effects and reducing dosage requirements, TDDS provide new opportunities for personalized

diabetes management. They also optimize pharmacokinetics, regulate drug biodistribution and enhance patient adherence, makes them highly promising for the treatment of complex chronic diseases (296).

A typical TDDS consists of three essential components: The therapeutic target, the active pharmaceutical ingredient (API), and the carrier or delivery vehicle. The therapeutic target refers to the specific organ, tissue, or cell that requires intervention, whereas the API provides the therapeutic effect. The carrier transports the API to the target site with spatial and temporal precision. An ideal carrier should exhibit favorable biocompatibility, biodegradability, low toxicity and specific recognition of the target site (297). Various carrier systems, including liposomes, aptamers and tetrahedral framework nucleic acids, have been extensively investigated. These carriers can improve the bioavailability and tissue penetration of therapeutic molecules, thereby enhancing efficacy. According to their delivery mechanisms, TDDS can be classified as active or passive targeting systems. Active targeting relies on ligand-receptor interactions to achieve precise drug localization, whereas passive take advantage of pathological features such as increased permeability to enhance drug accumulation in diseased tissues (297). The combination of micro-nanotechnology with TDDS represents a major advancement in diabetes therapy. Encapsulating drugs within micro- or nanoscale carriers improves drug stability, bioavailability and local concentration, enhancing therapeutic efficacy and reducing adverse reactions (293,298). For example, polymer-based nanoparticle systems for insulin delivery, constructed from biocompatible and biodegradable materials, such as chitosan, poly(lactic-co-glycolic acid) (PLGA) and polycaprolacton, can effectively overcome the limitations of traditional injection therapy. These nanocarriers protect insulin from enzymatic degradation and acidic conditions during oral administration, significantly increasing its bioavailability and enabling controlled and targeted release (299). Similar benefits have been observed with other oral hypoglycemic agents. A PLGA-based metformin delivery system demonstrated remarkable changes in pharmacokinetic properties. Compared with conventional formulations, PLGA-loaded metformin achieved more than a 10-fold increase in mean residence time and a substantially larger apparent distribution volume, even at 1/10 of the standard dose (300). These findings support PLGA as an efficient sustained-release delivery platform with great potential in diabetes treatment.

As  $\beta$  cells play a central role in the onset and progression of diabetes, designing delivery systems capable of specifically targeting the islet microenvironment is crucial for effective therapy. Ghosh *et al* (301) developed functionalized nanoparticles by conjugating the targeting peptide CHVLWSTRKC to a PLGA-b-PEG-COOH polymer and loading it with genistein. These nanoparticles bound to islet endothelial cells with 3-fold greater affinity through specific interaction with the EphA4 receptor, resulting in a 200-fold enhancement in the immunosuppressive activity of genistein (301). These results demonstrate the feasibility and potential of islet-specific drug delivery. In addition to targeting peptides, antibodies and aptamers have been widely used to construct nanoscale delivery systems for gene therapy, demonstrating considerable potential for precise genetic modulation in diabetes. These systems use

antibody- or aptamer-mediated targeting to deliver therapeutic genes into specific cells and employ disease-responsive promoters to achieve controlled gene expression. This approach helps restore disrupted gene networks and recover cellular function (302). Among these ligands, aptamers have received growing attention for their unique advantages as single-stranded DNA or RNA oligonucleotides with high affinity and selectivity (303). Compared with antibodies, aptamers are smaller, enabling deeper tissue penetration, and they exhibit low immunogenicity and high stability *in vivo*, making them suitable for long-term treatment. Their easy synthesis and chemical flexibility also facilitate the development of personalized therapeutic platforms for diabetes (304,305). Yamada *et al* (306) provided strong experimental evidence for the concept by designing a  $\beta$ -cell-targeted nanocarrier system known as  $\beta$ -MEND. This carrier efficiently delivered nucleic acids into pancreatic  $\beta$  cells.  $\beta$ -MEND successfully internalized into MIN6 cells and delivered antisense RNA targeting microRNA (miR)-375, leading to a marked reduction in miR-375 expression. As a result, Pdk1 and Mtpn were upregulated, significantly enhancing insulin secretion (306). These findings demonstrated that aptamer-based nanocarriers can effectively regulate  $\beta$ -cell function and provide a viable route toward gene therapy in diabetes. Collectively, these advances indicate that TDDS are gradually evolving from experimental delivery platforms toward clinically applicable therapeutic strategies. Recent advances in oral nanoplateforms have achieved efficient insulin delivery, glucose-responsive drug release and sustained glycemic control in preclinical models, highlighting the promising translational potential of precision drug delivery. For example, a milk-derived nanovesicle-based oral insulin delivery system achieved ~20% oral bioavailability with sustained glucose-lowering efficacy in diabetic pigs, highlighting the feasibility of non-invasive insulin replacement therapy (307). Similarly, the recently developed the MOP@T@D (MON-Proteins@Transferrin@Deoxycholic acid) oral nanoplateform in a T2DM rat model achieved an oral bioavailability of 10.6%, increased islet area by 26.7% after treatment, restored insulin secretion to 74.6% of the physiological level, and maintained normal blood glucose levels for up to two weeks after treatment cessation (308). By enabling sustained drug release and selective delivery to diseased tissues or cells, these strategies may facilitate individualized therapeutic interventions while reducing systemic adverse effects, thereby providing a feasible route for future clinical translation.

Despite these encouraging advances, TDDS still face significant challenges in clinical translation. Each type of carrier has inherent limitations. Nanoparticles often exhibit poor physical stability and tend to aggregate or undergo oxidative degradation. In addition, their manufacturing processes are complex (309). Polymer-based also face challenges such as complicated synthesis routes, high production costs and limited structural stability. These limitations may lead to aggregation and, in some cases, inflammatory reactions *in vivo* (310).

**Application of artificial intelligence in early diagnosis and prognostic assessment.** As a systemic metabolic disorder, diabetes is not only inherently complex but also closely associated with numerous severe complications, which substantially increase patient morbidity and mortality (311). Therefore,

treatment alone is insufficient to address the full spectrum of health challenges associated with diabetes. Early intervention through prevention and timely diagnosis is equally important. Consequently, the integration of artificial intelligence (AI) into diabetes management, through machine learning, deep learning and other AI-based analytical approaches, has the potential to transform diabetes care. By leveraging advanced data analysis, feature extraction and predictive modeling, AI provides a data-driven framework for screening, diagnosis, treatment, prognostic assessment and preventive interventions, thereby opening new avenues for disease management (312).

AI applications in diabetes primarily focus on four key domains: Screening for diabetes and its complications, clinical decision support systems (CDSS), population risk prediction and patient self-management tools (313,314). In the field of diabetes screening, AI offers the potential for early, non-invasive, rapid, and cost-effective detection of pathological changes (314). Shu *et al* (315) utilized facial texture analysis combined with a support vector machine model, achieving an accuracy of 99.02%, sensitivity of 99.64% and specificity of 98.26% in diabetes detection. Similarly, Li *et al* (316) integrated tongue color and texture features with machine learning models to predict diabetes risk, reporting an average accuracy of 0.821. However, neither study included external validation, limiting the generalizability of the findings. AI-based early detection of diabetic retinopathy (DR) has emerged as a cost-effective alternative, potentially reducing diabetes-related ocular complications and preventable blindness (317). The American Diabetes Association formally recognizes the clinical value of autonomous AI systems in detecting DR and macular edema (318). Keel *et al* (319) demonstrated that an AI-based DR screening model is highly feasible in endocrinology outpatient settings, showing a sensitivity of 92.3% and specificity of 93.7%, and a patient satisfaction rate of 96%, providing strong evidence for integrating AI screening into routine clinical practice. DPN is another common yet often overlooked complication, largely because of limited screening tools and insufficient diagnostic sensitivity (320). Deep learning algorithms trained on corneal confocal microscopy images have shown promise in detecting DPN. Williams *et al* (321) developed a deep learning algorithm that demonstrated rapid and accurate localization for quantifying corneal nerve biomarkers. Preston *et al* (322) developed a ResNet-50-based model that achieved precisions value of 0.83 in healthy volunteers, with 0.92 and 1.0 for non-neuropathic and neuropathic individuals, respectively.

AI can also assist in optimizing treatment strategies for patients with complex diabetes by identifying optimal therapeutic targets and supporting clinical decision-making. Murphree *et al* (323) conducted a retrospective study of 12,147 patients initiating metformin therapy, using a machine learning model to predict 1-year glycemic control outcomes. Baseline HbA1c, initial metformin dose and complications emerged as key predictors of treatment failure. The best-performing model achieved an area under the curve of 0.75, providing evidence for early identification of high-risk patients and personalized treatment planning (323). Additionally, AI-driven CDSS based on treatment pathway modeling can simulate HbA1c responses under different therapeutic strategies in T2DM, offering critical guidance

for individualized care (324). By analyzing patients' lifestyle behaviors, physical and mental health parameters, and social network activities, machine learning-based health recommendation systems can predict disease risks (324). Decision tree models have also been applied to predict the risk of T2DM development in women with gestational diabetes, achieving discrimination performance of 83.0% in training datasets and 76.9% in independent test sets, outperforming traditional fasting glucose monitoring (325). AI models integrating genetic and medical imaging data have demonstrated strong predictive performance for T2DM risk assessment. Huang *et al* (326) employed XGBoost to combine polygenic risk scores, multi-modality imaging risk scores and demographic variables (for example, age, sex and family history of diabetes) in 68,911 participants, achieving an Area Under the Receiver Operating Curve (AUC) of 0.94. This model effectively identified high-risk populations with specific genetic and imaging features, enabling precision prevention and early intervention for T2DM (326). These advances suggest that AI is gradually transitioning from a research-oriented analytical tool to an integrated component of precision diabetes care. In clinical practice, AI-based systems may facilitate earlier identification of individuals at risk, improve screening efficiency for diabetes complications, and support individualized therapeutic decision-making through integration with continuous glucose monitoring, electronic health records, and digital health platforms. Among current AI-enabled therapeutic strategies, personalized insulin adjustment based on reinforcement learning, prediction of treatment responses to glucose-lowering medications, and AI-guided lifestyle interventions have shown promising potential for improving glycemic control and optimizing long-term disease management.

Self-management, particularly maintaining blood glucose within recommended ranges, remains a cornerstone of diabetes care. This involves active glucose monitoring, regulation of physical activity and diet, and appropriate insulin administration (327). Dietary and exercise interventions are primary strategies for preventing T2DM in high-risk populations (328). Personalized applications provide tailored dietary plans, while wearable devices track daily activity and use behavioral incentives to promote healthy habits, thereby helping prevent chronic diseases including T2DM (314). In addition, accurate prediction of blood glucose fluctuations in patients with T2DM facilitates improved glycemic control, reduces hypoglycemic events, and lowers diabetes-related morbidity and mortality, ultimately enhancing quality of life. Deng *et al* (329) developed a predictive model using deep transfer learning and data augmentation on continuous glucose monitoring data recorded every 30 min, forecasting blood glucose 5 min to 1 h ahead with over 95% accuracy and 90% sensitivity. Wang *et al* (330) proposed a reinforcement learning-based dynamic insulin titration regimen, improving glycemic control without increasing hypoglycemia risk, achieving personalized insulin dose optimization for patients with T2DM. These technological advances enable both healthcare providers and patients to proactively manage diabetes, adjusting insulin doses and lifestyle behaviors to prevent hyperglycemia or hypoglycemia and associated complications. Despite the promise of AI in research, its real-world application remains challenging, particularly across diverse populations (331).

Training datasets lacking balanced representation in terms of age, sex, ethnicity, or risk factors may exacerbate biases, leading to misdiagnosis or inappropriate treatment for under-represented groups (332). Therefore, inclusive data collection strategies are essential. Given the potential for AI errors, ethical challenges arise, underscoring the need for robust supervision and clear accountability guidelines in AI-driven diabetes care. Furthermore, ensuring data privacy is critical. Securing data during development and deployment remains challenging, and global-scale implementation intensifies these risks, necessitating stringent protection measures.

### Acknowledgements

Not applicable.

### Funding

The present study was supported by Sichuan Provincial Administration of Traditional Chinese Medicine (grant no. 25ZDZX028) and the Science and Technology Strategic Cooperation Programs of Luzhou Municipal People's Government and Southwest Medical University (grant no. 2025LZXNYDJC06).

### Availability of data and materials

Not applicable.

### Authors' contributions

ZH conceptualized the study. ZH, HL, YL and YY conducted investigation and data visualization, wrote the original draft, and wrote, reviewed and edited the manuscript. SL conducted investigation, and wrote, reviewed and edited the manuscript. TZ supervised the study, conducted project administration, and wrote, reviewed and edited the manuscript. DB conceptualized and supervised the study, conducted project administration, acquired funding, and wrote, reviewed and edited the manuscript. All authors read and approved the final version of the manuscript. Data authentication is not applicable.

### Ethics approval and consent to participate

Not applicable.

### Patient consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### References

1. Liu H, Wu Y, Zhu H, Wang P, Chen T, Xia A, Zhao Z, He D, Chen X, Xu J and Ji L: Association between napping and type 2 diabetes mellitus. *Front Endocrinol (Lausanne)* 15: 1294638, 2024.
2. Chatterjee S, Khunti K and Davies MJ: Type 2 diabetes. *Lancet* 389: 2239-2251, 2017.

3. Hawley JA, Hargreaves M and Zierath JR: Signalling mechanisms in skeletal muscle: Role in substrate selection and muscle adaptation. *Essays Biochem* 42: 1-12, 2006.
4. Ruderman NB, Carling D, Prentki M and Cacicedo JM: AMPK, insulin resistance, and the metabolic syndrome. *J Clin Invest* 123: 2764-2772, 2013.
5. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC and Swain DP: American College of Sports Medicine: Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults. *Med Sci Sports Exerc* 43: 1334-1359, 2011.
6. Mulya A, Haus JM, Solomon TP, Kelly KR, Malin SK, Rocco M, Barkoukis H and Kirwan JP: Exercise training-induced improvement in skeletal muscle PGC-1 $\alpha$ -mediated fat metabolism is independent of dietary glycemic index. *Obesity (Silver Spring)* 25: 721-729, 2017.
7. Kirwan JP, Sacks J and Nieuwoudt S: The essential role of exercise in the management of type 2 diabetes. *Cleve Clin J Med* 84 (7 Suppl 1): S15-S21, 2017.
8. Boul NG, Kenny GP, Haddad E, Wells GA and Sigal RJ: Meta-analysis of the effect of structured exercise training on cardiorespiratory fitness in type 2 diabetes mellitus. *Diabetologia* 46: 1071-1081, 2003.
9. Evans PL, McMillin SL, Weyrauch LA and Witzak CA: Regulation of skeletal muscle glucose transport and glucose metabolism by exercise training. *Nutrients* 11: 2432, 2019.
10. DeFronzo RA and Tripathy D: Skeletal muscle insulin resistance is the primary defect in type 2 diabetes. *Diabetes Care* 32 (Suppl 2): S157-S163, 2009.
11. Abdul-Ghani MA and DeFronzo RA: Pathogenesis of insulin resistance in skeletal muscle. *J Biomed Biotechnol* 2010: 476279, 2010.
12. Shepherd PR and Kahn BB: Glucose transporters and insulin action-implications for insulin resistance and diabetes mellitus. *N Engl J Med* 341: 248-257, 1999.
13. Razak F and Anand SS: Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *Petersen KF, Dufour S, Befroy D, Garcia R and Shulman GI N Engl J Med* 350: 664-671, 2004. *Vasc Med* 9: 223-224, 2004.
14. Phielix E and Mensink M: Type 2 diabetes mellitus and skeletal muscle metabolic function. *Physiol Behav* 94: 252-258, 2008.
15. Timmers S, Schrauwen P and de Vogel J: Muscular diacylglycerol metabolism and insulin resistance. *Physiol Behav* 94: 242-251, 2008.
16. Chow L, From A and Seaquist E: Skeletal muscle insulin resistance: The interplay of local lipid excess and mitochondrial dysfunction. *Metabolism* 59: 70-85, 2010.
17. Ahmed AS, Schizas N, Li J, Ahmed M, Östenson CG, Salo P, Hewitt C, Hart DA and Ackermann PW: Type 2 diabetes impairs tendon repair after injury in a rat model. *J Appl Physiol* (1985) 113: 1784-1791, 2012.
18. Lin YC, Li YJ, Rui YF, Dai GC, Shi L, Xu HL, Ni M, Zhao S, Chen H, Wang C, *et al*: The effects of high glucose on tendon-derived stem cells: Implications of the pathogenesis of diabetic tendon disorders. *Oncotarget* 8: 17518-17528, 2017.
19. Lu P, Dai G, Shi L, Li Y, Zhang M, Wang H and Rui Y: HMGB1 modulates high glucose-induced erroneous differentiation of tendon stem/progenitor cells through RAGE/ $\beta$ -catenin pathway. *Stem Cells Int* 2024: 2335270, 2024.
20. Huo JL, Feng Q, Pan S, Fu WJ, Liu Z and Liu Z: Diabetic cardiomyopathy: Early diagnostic biomarkers, pathogenetic mechanisms, and therapeutic interventions. *Cell Death Discov* 9: 256, 2023.
21. Connizzo BK, Bhatt PR, Liechty KW and Soslowsky LJ: Diabetes alters mechanical properties and collagen fiber re-alignment in multiple mouse tendons. *Ann Biomed Eng* 42: 1880-1888, 2014.
22. Fox AJ, Bedi A, Deng XH, Ying L, Harris PE, Warren RF and Rodeo SA: Diabetes mellitus alters the mechanical properties of the native tendon in an experimental rat model. *J Orthop Res* 29: 880-885, 2011.
23. Lee JM and Veres SP: Advanced glycation end-product cross-linking inhibits biomechanical plasticity and characteristic failure morphology of native tendon. *J Appl Physiol* (1985) 126: 832-841, 2019.
24. Kamml J, Acevedo C and Kammer DS: Advanced-Glycation Endproducts: How cross-linking properties affect the collagen fibril behavior. *J Mech Behav Biomed Mater* 148: 106198, 2023.
25. Kayal RA, Siqueira M, Alblowi J, McLean J, Krothapalli N, Faibish D, Einhorn TA, Gerstenfeld LC and Graves DT: TNF- $\alpha$  mediates diabetes-enhanced chondrocyte apoptosis during fracture healing and stimulates chondrocyte apoptosis through FOXO1. *J Bone Miner Res* 25: 1604-1615, 2010.
26. Qin B, Bao D, Liu Y, Zeng S, Deng K, Liu H and Fu S: Engineered exosomes: A promising strategy for tendon-bone healing. *J Adv Res* 64: 155-169, 2024.
27. Wongdee K and Charoenphandhu N: Osteoporosis in diabetes mellitus: Possible cellular and molecular mechanisms. *World J Diabetes* 2: 41-48, 2011.
28. Medeiros C and Wallace JM: High glucose-induced inhibition of osteoblast like MC3T3-E1 differentiation promotes mitochondrial perturbations. *PLoS One* 17: e0270001, 2022.
29. Sheng N, Xing F, Wang J, Zhang QY, Nie R, Li-Ling J, Duan X and Xie HQ: Recent progress in bone-repair strategies in diabetic conditions. *Mater Today Bio* 23: 100835, 2023.
30. Zhen D, Chen Y and Tang X: Metformin reverses the deleterious effects of high glucose on osteoblast function. *J Diabetes Complications* 24: 334-344, 2010.
31. Sanches CP, Vianna AGD and Barreto FC: The impact of type 2 diabetes on bone metabolism. *Diabetol Metab Syndr* 9: 85, 2017.
32. Jiao H, Xiao E and Graves DT: Diabetes and its effect on bone and fracture healing. *Curr Osteoporos Rep* 13: 327-335, 2015.
33. Wu B, Fu Z, Wang X, Zhou P, Yang Q, Jiang Y and Zhu D: A narrative review of diabetic bone disease: Characteristics, pathogenesis, and treatment. *Front Endocrinol (Lausanne)* 13: 1052592, 2022.
34. Lecka-Czernik B, Gubrij I, Moerman EJ, Kajkenova O, Lipschitz DA, Manolagas SC and Jilka RL: Inhibition of *Osf2/Cbfa1* expression and terminal osteoblast differentiation by *PPARgamma2*. *J Cell Biochem* 74: 357-371, 1999.
35. Keats E and Khan ZA: Unique responses of stem cell-derived vascular endothelial and mesenchymal cells to high levels of glucose. *PLoS One* 7: e38752, 2012.
36. Piccinin MA and Khan ZA: Pathophysiological role of enhanced bone marrow adipogenesis in diabetic complications. *Adipocyte* 3: 263-272, 2014.
37. Wang A, Midura RJ, Vasanthi A, Wang AJ and Hascall VC: Hyperglycemia diverts dividing osteoblastic precursor cells to an adipogenic pathway and induces synthesis of a hyaluronan matrix that is adhesive for monocytes. *J Biol Chem* 289: 11410-11420, 2014.
38. Roy B: Biomolecular basis of the role of diabetes mellitus in osteoporosis and bone fractures. *World J Diabetes* 4: 101-113, 2013.
39. Bansal S, Burman A and Tripathi AK: Advanced glycation end products: Key mediator and therapeutic target of cardiovascular complications in diabetes. *World J Diabetes* 14: 1146-1162, 2023.
40. Shi L, Lu PP, Dai GC, Li YJ and Rui YF: Advanced glycation end products and tendon stem/progenitor cells in pathogenesis of diabetic tendinopathy. *World J Stem Cells* 13: 1338-1348, 2021.
41. Baig MH, Jan AT, Rabbani G, Ahmad K, Ashraf JM, Kim T, Min HS, Lee YH, Cho WK, Ma JY, *et al*: Methylglyoxal and Advanced Glycation End products: Insight of the regulatory machinery affecting the myogenic program and of its modulation by natural compounds. *Sci Rep* 7: 5916, 2017.
42. Mori H, Kuroda A, Ishizu M, Ohishi M, Takashi Y, Otsuka Y, Taniguchi S, Tamaki M, Kurahashi K, Yoshida S, *et al*: Association of accumulated advanced glycation end-products with a high prevalence of sarcopenia and dynapenia in patients with type 2 diabetes. *J Diabetes Investig* 10: 1332-1340, 2019.
43. Yan SD, Schmidt AM, Anderson GM, Zhang J, Brett J, Zou YS, Pinsky D and Stern D: Enhanced cellular oxidant stress by the interaction of advanced glycation end products with their receptors/binding proteins. *J Biol Chem* 269: 9889-9897, 1994.
44. Lander HM, Tauras JM, Ogiste JS, Hori O, Moss RA and Schmidt AM: Activation of the receptor for advanced glycation end products triggers a p21-dependent mitogen-activated protein kinase pathway regulated by oxidant stress. *J Biol Chem* 272: 17810-17814, 1997.
45. Wautier MP, Chappey O, Corda S, Stern DM, Schmidt AM and Wautier JL: Activation of NADPH oxidase by AGE links oxidant stress to altered gene expression via RAGE. *Am J Physiol Endocrinol Metab* 280: E685-E694, 2001.
46. Chen J, Song M, Yu S, Gao P, Yu Y, Wang H and Huang L: Advanced glycation endproducts alter functions and promote apoptosis in endothelial progenitor cells through receptor for advanced glycation endproducts mediate overexpression of cell oxidant stress. *Mol Cell Biochem* 335: 137-146, 2009.

47. Nowotny K, Jung T, Höhn A, Weber D and Grune T: Advanced glycation end products and oxidative stress in type 2 diabetes mellitus. *Biomolecules* 5: 194-222, 2015.
48. Mori H, Kuroda A, Araki M, Suzuki R, Taniguchi S, Tamaki M, Akehi Y and Matsuhisa M: Advanced glycation end-products are a risk for muscle weakness in Japanese patients with type 1 diabetes. *J Diabetes Investig* 8: 377-382, 2017.
49. Ramamurthy B, Höök P, Jones AD and Larsson L: Changes in myosin structure and function in response to glycation. *FASEB J* 15: 2415-2422, 2001.
50. Avery NC and Bailey AJ: The effects of the Maillard reaction on the physical properties and cell interactions of collagen. *Pathol Biol (Paris)* 54: 387-395, 2006.
51. Fessel G, Li Y, Diederich V, Guizar-Sicairos M, Schneider P, Sell DR, Monnier VM and Snedeker JG: Advanced glycation end-products reduce collagen molecular sliding to affect collagen fibril damage mechanisms but not stiffness. *PLoS One* 9: e110948, 2014.
52. Li Y, Fessel G, Georgiadis M and Snedeker JG: Advanced glycation end-products diminish tendon collagen fiber sliding. *Matrix Biol* 32: 169-177, 2013.
53. Gautieri A, Passini FS, Silván U, Guizar-Sicairos M, Carimati G, Volpi P, Moretti M, Schoenhuber H, Redaelli A, Berli M and Snedeker JG: Advanced glycation end-products: Mechanics of aged collagen from molecule to tissue. *Matrix Biol* 59: 95-108, 2017.
54. Kudsi M and Labban L: The prevalence of musculoskeletal complications in type 2 diabetes mellitus. *Open Access Library J* 7: 1-10, 2020.
55. Cagliero E: Rheumatic manifestations of diabetes mellitus. *Curr Rheumatol Rep* 5: 189-194, 2003.
56. DeGroot J, Verzijl N, Budde M, Bijlsma JWJ, Lafeber FP and TeKoppele JM: Accumulation of advanced glycation end products decreases collagen turnover by bovine chondrocytes. *Exp Cell Res* 266: 303-310, 2001.
57. Chang J, Thunder R, Most D, Longaker MT and Lineaweaver WC: Studies in flexor tendon wound healing: Neutralizing antibody to TGF- $\beta$ 1 increases postoperative range of motion. *Plast Reconstr Surg* 105: 148-155, 2000.
58. Khalid M, Petroianu G and Adem A: Advanced glycation end products and diabetes mellitus: Mechanisms and perspectives. *Biomolecules* 12: 542, 2022.
59. Valcourt U, Merle B, Gineys E, Viguet-Carrin S, Delmas PD and Garnero P: Non-enzymatic glycation of bone collagen modifies osteoclastic activity and differentiation. *J Biol Chem* 282: 5691-5703, 2007.
60. Aoki C, Uto K, Honda K, Kato Y and Oda H: Advanced glycation end products suppress lysyl oxidase and induce bone collagen degradation in a rat model of renal osteodystrophy. *Lab Invest* 93: 1170-1183, 2013.
61. Cavati G, Pirrotta F, Merlotti D, Ceccarelli E, Calabrese M, Gennari L and Mingiano C: Role of advanced glycation end-products and oxidative stress in type-2-diabetes-induced bone fragility and implications on fracture risk stratification. *Antioxidants (Basel)* 12: 928, 2023.
62. Miranda C, Giner M, Montoya MJ, Vázquez MA, Miranda MJ and Pérez-Cano R: Influence of high glucose and advanced glycation end-products (ages) levels in human osteoblast-like cells gene expression. *BMC Musculoskelet Disord* 17: 377, 2016.
63. Li Y, Wang L, Zhang M, Huang K, Yao Z, Rao P, Cai X and Xiao J: Advanced glycation end products inhibit the osteogenic differentiation potential of adipose-derived stem cells by modulating Wnt/ $\beta$ -catenin signalling pathway via DNA methylation. *Cell Prolif* 53: e12834, 2020.
64. Ryle C, Leow CK and Donaghy M: Nonenzymatic glycation of peripheral and central nervous system proteins in experimental diabetes mellitus. *Muscle Nerve* 20: 577-584, 1997.
65. Cullum NA, Mahon J, Stringer K and McLean WG: Glycation of rat sciatic nerve tubulin in experimental diabetes mellitus. *Diabetologia* 34: 387-389, 1991.
66. Williams SK, Howarth NL, Devenny JJ and Bitensky MW: Structural and functional consequences of increased tubulin glycosylation in diabetes mellitus. *Proc Natl Acad Sci USA* 79: 6546-6550, 1982.
67. Sugimoto K, Yasujima M and Yagihashi S: Role of advanced glycation end products in diabetic neuropathy. *Curr Pharm Des* 14: 953-961, 2008.
68. Vlassara H, Brownlee M and Cerami A: Recognition and uptake of human diabetic peripheral nerve myelin by macrophages. *Diabetes* 34: 553-557, 1985.
69. Bierhaus A, Chevion S, Chevion M, Hofmann M, Quehenberger P, Illmer T, Luther T, Berentshtein E, Tritschler H, Müller M, *et al*: Advanced glycation end product-induced activation of NF-kappaB is suppressed by alpha-lipoic acid in cultured endothelial cells. *Diabetes* 46: 1481-1490, 1997.
70. Li P, Alenazi KKK, Dally J, Woods EL, Waddington RJ and Moseley R: Role of oxidative stress in impaired type II diabetic bone repair: Scope for antioxidant therapy intervention? *Front Dent Med* 5: 1464009, 2025.
71. Belosludtsev KN, Belosludtseva NV and Dubinin MV: Diabetes mellitus, mitochondrial dysfunction and Ca<sup>2+</sup>-Dependent permeability transition pore. *Int J Mol Sci* 21: 6559, 2020.
72. Lopez-Pedrosa JM, Camprubi-Robles M, Guzman-Rolo G, Lopez-Gonzalez A, Garcia-Almeida JM, Sanz-Paris A and Rueda R: The vicious cycle of type 2 diabetes mellitus and skeletal muscle atrophy: Clinical, biochemical, and nutritional bases. *Nutrients* 16: 172, 2024.
73. Bitar MS and Al-Mulla F: ROS constitute a convergence nexus in the development of IGF1 resistance and impaired wound healing in a rat model of type 2 diabetes. *Dis Model Mech* 5: 375-388, 2012.
74. Lu PP, Chen MH, Dai GC, Li YJ, Shi L and Rui YF: Understanding cellular and molecular mechanisms of pathogenesis of diabetic tendinopathy. *World J Stem Cells* 12: 1255-1275, 2020.
75. Kwan CK, Fu SC and Yung PS: A high glucose level stimulate inflammation and weaken pro-resolving response in tendon cells-A possible factor contributing to tendinopathy in diabetic patients. *Asia Pac J Sports Med Arthrosc Rehabil Technol* 19: 1-6, 2019.
76. Bakker W, Eringa EC, Sipkema P and van Hinsbergh VW: Endothelial dysfunction and diabetes: roles of hyperglycemia, impaired insulin signaling and obesity. *Cell Tissue Res* 335: 165-189, 2009.
77. Huang Y, He B, Wang L, Yuan B, Shu H, Zhang F and Sun L: Bone marrow mesenchymal stem cell-derived exosomes promote rotator cuff tendon-bone healing by promoting angiogenesis and regulating M1 macrophages in rats. *Stem Cell Res Ther* 11: 496, 2020.
78. Pinheiro-Machado E, Gurgul-Convey E and Marzec M: Immunometabolism in type 2 diabetes mellitus: Tissue-specific interactions. *Arch Med Sci* 19: 985-991, 2020.
79. Baker RG, Hayden MS and Ghosh S: NF- $\kappa$ B, inflammation, and metabolic disease. *Cell Metab* 13: 11-22, 2011.
80. Hotamisligil GS, Shargill NS and Spiegelman BM: Adipose expression of tumor necrosis factor- $\alpha$ : Direct role in obesity-linked insulin resistance. *Science* 259: 87-91, 1993.
81. Yin MJ, Yamamoto Y and Gaynor RB: The anti-inflammatory agents aspirin and salicylate inhibit the activity of I(kappa)B kinase-beta. *Nature* 396: 77-80, 1998.
82. Yuan M, Konstantopoulos N, Lee J, Hansen L, Li ZW, Karin M and Shoelson SE: Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikkbeta. *Science* 293: 1673-1677, 2001.
83. Mulder AH, de Langen CDJ, de Regt V and Hogenboom F: Alpha-receptor-mediated modulation of 3H-noradrenaline release from rat brain cortex synaptosomes. *Naunyn Schmiedeberg Arch Pharmacol* 303: 193-196, 1978.
84. Liu Z, Guo Y and Zheng C: Type 2 diabetes mellitus related sarcopenia: A type of muscle loss distinct from sarcopenia and disuse muscle atrophy. *Front Endocrinol (Lausanne)* 15: 1375610, 2024.
85. Hulett NA, Scalzo RL and Reusch JEB: Glucose uptake by skeletal muscle within the contexts of type 2 diabetes and exercise: An integrated approach. *Nutrients* 14: 647, 2022.
86. Mason RM and Wahab NA: Extracellular matrix metabolism in diabetic nephropathy. *J Am Soc Nephrol* 14: 1358-1373, 2003.
87. Brownlee M: Biochemistry and molecular cell biology of diabetic complications. *Nature* 414: 813-820, 2001.
88. Hepple RT: Skeletal muscle: Microcirculatory adaptation to metabolic demand. *Med Sci Sports Exerc* 32: 117-123, 2000.
89. Baum O and Bigler M: Pericapillary basement membrane thickening in human skeletal muscles. *Am J Physiol Heart Circ Physiol* 311: H654-H666, 2016.
90. Siperstein MD, Unger RH and Madison LL: Studies of muscle capillary basement membranes in normal subjects, diabetic, and prediabetic patients. *J Clin Invest* 47: 1973-1999, 1968.
91. D'Souza DM, Al-Sajee D and Hawke TJ: Diabetic myopathy: Impact of diabetes mellitus on skeletal muscle progenitor cells. *Front Physiol* 4: 379, 2013.
92. Mesinovic J, Zengin A, De Courten B, Ebeling PR and Scott D: Sarcopenia and type 2 diabetes mellitus: A bidirectional relationship. *Diabetes Metab Syndr Obes* 12: 1057-1072, 2019.

93. Sharma KR, Cross J, Farronay O, Ayyar DR, Shebert RT and Bradley WG: Demyelinating neuropathy in diabetes mellitus. *Arch Neurol* 59: 758-765, 2002.
94. Tempfer H and Traweger A: Tendon vasculature in health and disease. *Front Physiol* 6: 330, 2015.
95. de Oliveira RR, Martins CS, Rocha YR, Braga AB, Mattos RM, Hecht F, Brito GA and Nasciutti LE: Experimental diabetes induces structural, inflammatory and vascular changes of achilles tendons. *PLoS One* 8: e74942, 2013.
96. Ziobro A, Duchnowicz P, Mulik A, Koter-Michalak M and Broncel M: Oxidative damages in erythrocytes of patients with metabolic syndrome. *Mol Cell Biochem* 378: 267-273, 2013.
97. Rehman K and Akash MSH: Mechanism of generation of oxidative stress and pathophysiology of type 2 diabetes mellitus: How are they interlinked? *J Cell Biochem* 118: 3577-3585, 2017.
98. Kim DH, Noh SU, Chae SW, Kim SJ and Lee YT: Altered differentiation of tendon-derived stem cells in diabetic conditions mediated by macrophage migration inhibitory factor. *Int J Mol Sci* 22: 8983, 2021.
99. Knobloch K: The role of tendon microcirculation in Achilles and patellar tendinopathy. *J Orthop Surg Res* 3: 18, 2008.
100. Yu Y, Lin L, Zhou Y, Lu X, Shao X, Lin C, Yu K, Zhang X, Hong J and Chen Y: Effect of hypoxia on self-renewal capacity and differentiation in human tendon-derived stem cells. *Med Sci Monit* 23: 1334-1339, 2017.
101. Lavagnino M, Oslapas AN, Gardner KL and Arnoczky SP: Hypoxia inhibits primary cilia formation and reduces cell-mediated contraction in stress-deprived rat tail tendon fascicles. *Muscles Ligaments Tendons J* 6: 193-197, 2016.
102. Mahjoor M, Fakouri A, Farokhi S, Nazari H, Afkhami H and Heidari F: Regenerative potential of mesenchymal stromal cells in wound healing: Unveiling the influence of normoxic and hypoxic environments. *Front Cell Dev Biol* 11: 1245872, 2023.
103. Gupta N, Mansoor S, Sharma A, Sapkal A, Sheth J, Falatoonzadeh P, Kuppermann B and Kenney M: Diabetic retinopathy and VEGF. *Open Ophthalmol J* 7: 4-10, 2013.
104. Murata T, Nagai R, Ishibashi T, Inomata H, Ikeda K and Horiuchi S: The relationship between accumulation of advanced glycation end products and expression of vascular endothelial growth factor in human diabetic retinas. *Diabetologia* 40: 764-769, 1997.
105. Van Dyken P and Lacoste B: Impact of metabolic syndrome on neuroinflammation and the blood-brain barrier. *Front Neurosci* 12: 930, 2018.
106. Spinetti G, Kraenkel N, Emanuelli C and Madeddu P: Diabetes and vessel wall remodelling: From mechanistic insights to regenerative therapies. *Cardiovasc Res* 78: 265-273, 2008.
107. Hazarika S, Dokun AO, Li Y, Popel AS, Kontos CD and Annex BH: Impaired angiogenesis after hindlimb ischemia in type 2 diabetes mellitus: Differential regulation of vascular endothelial growth factor receptor 1 and soluble vascular endothelial growth factor receptor 1. *Circ Res* 101: 948-956, 2007.
108. Kolluru GK, Bir SC and Kevil CG: Endothelial dysfunction and diabetes: Effects on angiogenesis, vascular remodeling, and wound healing. *Int J Vasc Med* 2012: 918267, 2012.
109. Zhao C, Ackerman JE, Geary MB, Orner CA, Bawany F and Loiselle AE: Obesity/Type II diabetes alters macrophage polarization resulting in a fibrotic tendon healing response. *PLoS One* 12: e0181127, 2017.
110. Bunker DL, Ilie V, Ilie V and Nicklin S: Tendon to bone healing and its implications for surgery. *Muscles Ligaments Tendons J* 4: 343-350, 2014.
111. Chen J, Jiang C, Yin L, Liu Y, He Y, Li S and Shen H: A review of the role of tendon stem cells in tendon-bone regeneration. *Med Sci Monit* 29: e940805, 2023.
112. Lei T, Zhang T, Ju W, Chen X, Heng BC, Shen W and Yin Z: Biomimetic strategies for tendon/ligament-to-bone interface regeneration. *Bioact Mater* 6: 2491-2510, 2021.
113. Cheng P, Han P, Zhao C, Zhang S, Wu H, Ni J, Hou P, Zhang Y, Liu J, Xu H, *et al*: High-purity magnesium interference screws promote fibrocartilaginous entheses regeneration in the anterior cruciate ligament reconstruction rabbit model via accumulation of BMP-2 and VEGF. *Biomaterials* 81: 14-26, 2016.
114. Yoshikawa T, Tohyama H, Enomoto H, Matsumoto H, Toyama Y and Yasuda K: Expression of vascular endothelial growth factor and angiogenesis in patellar tendon grafts in the early phase after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 14: 804-810, 2006.
115. Fadini GP, Albiero M, Bonora BM and Avogaro A: Angiogenic abnormalities in diabetes mellitus: Mechanistic and clinical aspects. *J Clin Endocrinol Metab* 104: 5431-5444, 2019.
116. Demirag B, Sarisozen B, Ozer O, Kaplan T and Ozturk C: Enhancement of tendon-bone healing of anterior cruciate ligament grafts by blockage of matrix metalloproteinases. *J Bone Joint Surg Am* 87: 2401-2410, 2005.
117. Takayama K, Kawakami Y, Mifune Y, Matsumoto T, Tang Y, Cummins JH, Greco N, Kuroda R, Kurosaka M, Wang B, *et al*: The effect of blocking angiogenesis on anterior cruciate ligament healing following stem cell transplantation. *Biomaterials* 60: 9-19, 2015.
118. Zafar N, Krishnasamy SS, Shah J, Rai SN, Riggs DW, Bhatnagar A and O'Toole TE: Circulating angiogenic stem cells in type 2 diabetes are associated with glycemic control and endothelial dysfunction. *PLoS One* 13: e0205851, 2018.
119. Paine SK, Mondal LK, Borah PK, Bhattacharya CK and Mahanta J: Pro- and antiangiogenic VEGF and its receptor status for the severity of diabetic retinopathy. *Mol Vis* 23: 356-363, 2017.
120. Kusumbe AP, Ramasamy SK and Adams RH: Coupling of angiogenesis and osteogenesis by a specific vessel subtype in bone. *Nature* 507: 323-328, 2014.
121. Ji G, Xu R, Niu Y, Li N, Ivashkiv L, Boström MPG, Greenblatt MB and Yang X: Vascular endothelial growth factor pathway promotes osseointegration and CD31hiEMCNhi endothelium expansion in a mouse tibial implant model. *Bone Joint J* 101-B (7\_Suppl\_C): 108-114, 2019.
122. Fijany A, Sayadi LR, Khoshab N, Banyard DA, Shaterian A, Alexander M, Lakey JRT, Paydar KZ, Evans GRD and Widgerow AD: Mesenchymal stem cell dysfunction in diabetes. *Mol Biol Rep* 46: 1459-1475, 2018.
123. Kahn MB, Yuldasheva NY, Cubbon RM, Smith J, Rashid ST, Viswambharan H, Imrie H, Abbas A, Rajwani A, Aziz A, *et al*: Insulin resistance impairs circulating angiogenic progenitor cell function and delays endothelial regeneration. *Diabetes* 60: 1295-1303, 2011.
124. Sorrentino SA, Bahlmann FH, Besler C, Müller M, Schulz S, Kirchhoff N, Doerries C, Horváth T, Limbourg A, Limbourg F, *et al*: Oxidant stress impairs in vivo reendothelialization capacity of endothelial progenitor cells from patients with type 2 diabetes mellitus. *Circulation* 116: 163-173, 2007.
125. Ceradini DJ, Yao D, Grogan RH, Callaghan MJ, Edelstein D, Brownlee M and Gurtner GC: Decreasing intracellular superoxide corrects defective ischemia-induced new vessel formation in diabetic mice. *J Biol Chem* 283: 10930-10938, 2008.
126. Wheatcroft SB, Shah AM, Li JM, Duncan E, Noronha BT, Crossey PA and Kearney MT: Preserved glucoregulation but attenuation of the vascular actions of insulin in mice heterozygous for knockout of the insulin receptor. *Diabetes* 53: 2645-2652, 2004.
127. Duncan ER, Walker SJ, Ezzat VA, Wheatcroft SB, Li JM, Shah AM and Kearney MT: Accelerated endothelial dysfunction in mild prediabetic insulin resistance: The early role of reactive oxygen species. *Am J Physiol Endocrinol Metab* 293: E1311-E1319, 2007.
128. Duncan ER, Crossey PA, Walker S, Anilkumar N, Poston L, Douglas G, Ezzat VA, Wheatcroft SB, Shah AM and Kearney MT: Effect of endothelium-specific insulin resistance on endothelial function in vivo. *Diabetes* 57: 3307-3314, 2008.
129. Everaert BR, Van Craenenbroeck EM, Hoymans VY, Haine SE, Van Nassauw L, Conraads VM, Timmermans JP and Vrints CJ: Current perspective of pathophysiological and interventional effects on endothelial progenitor cell biology: Focus on PI3K/AKT/eNOS pathway. *Int J Cardiol* 144: 350-366, 2010.
130. Friese N, Gierschner MB, Schadzek P, Roger Y and Hoffmann A: Regeneration of damaged tendon-bone junctions (Entheses)-TAK1 as a potential node factor. *Int J Mol Sci* 21: 5177, 2020.
131. Hu JJ, Yin Z, Shen WL, Xie YB, Zhu T, Lu P, Cai YZ, Kong MJ, Heng BC, Zhou YT, *et al*: Pharmacological regulation of in situ tissue stem cells differentiation for soft tissue calcification treatment. *Stem Cells* 34: 1083-1096, 2016.
132. Sugimoto Y, Takimoto A, Akiyama H, Kist R, Scherer G, Nakamura T, Hiraki Y and Shikunami C: Scx+/Sox9+ progenitors contribute to the establishment of the junction between cartilage and tendon/ligament. *Development* 140: 2280-2288, 2013.
133. Spinetti G, Cordella D, Fortunato O, Sangalli E, Losa S, Gotti A, Carnelli F, Rosa F, Riboldi S, Sessa F, *et al*: Global remodeling of the vascular stem cell niche in bone marrow of diabetic patients. *Circ Res* 112: 510-522, 2013.

134. Semeraro F, Cancarini A, dell'Omo R, Rezzola S, Romano MR and Costagliola C: Diabetic retinopathy: Vascular and inflammatory disease. *J Diabetes Res* 2015: 582060, 2015.
135. Xu N, Liu H, Qu F, Fan J, Mao K, Yin Y, Liu J, Geng Z and Wang Y: Hypoxia inhibits the differentiation of mesenchymal stem cells into osteoblasts by activation of Notch signaling. *Exp Mol Pathol* 94: 33-39, 2013.
136. Hinoi E, Ochi H, Takarada T, Nakatani E, Iezaki T, Nakajima H, Fujita H, Takahata Y, Hidano S, Kobayashi T, *et al*: Positive regulation of osteoclastic differentiation by growth differentiation factor 15 upregulated in osteocytic cells under hypoxia. *J Bone Miner Res* 27: 938-949, 2012.
137. Wu YF, Wang HK, Chang HW, Sun J, Sun JS and Chao YH: High glucose alters tendon homeostasis through downregulation of the AMPK/Egr1 pathway. *Sci Rep* 7: 44199, 2017.
138. Guerquin MJ, Charvet B, Nourissat G, Havis E, Ronsin O, Bonnin MA, Ruggiu M, Olivera-Martinez I, Robert N, Lu Y, *et al*: Transcription factor EGR1 directs tendon differentiation and promotes tendon repair. *J Clin Invest* 123: 3564-3576, 2013.
139. Veidal SS, Karsdal MA, Vassiliadis E, Nawrocki A, Larsen MR, Nguyen QH, Hägglund P, Luo Y, Zheng Q, Vainer B and Leeming DJ: MMP mediated degradation of type VI collagen is highly associated with liver fibrosis-identification and validation of a novel biochemical marker assay. *PLoS One* 6: e24753, 2011.
140. Cui N, Hu M and Khalil RA: Biochemical and biological attributes of matrix metalloproteinases. *Prog Mol Biol Transl Sci* 147: 1-73, 2017.
141. Tsai WC, Liang FC, Cheng JW, Lin LP, Chang SC, Chen HH and Pang JH: High glucose concentration up-regulates the expression of matrix metalloproteinase-9 and -13 in tendon cells. *BMC Musculoskeletal Disorders* 14: 255, 2013.
142. Sharma P and Maffulli N: Tendon injury and tendinopathy. *J Bone Joint Surg Am* 87: 187-202, 2005.
143. Oshiro W, Lou J, Xing X, Tu Y and Manske PR: Flexor tendon healing in the rat: A histologic and gene expression study. *J Hand Surg Am* 28: 814-823, 2003.
144. Cabral-Pacheco GA, Garza-Veloz I, Castruita-De la Rosa C, Ramirez-Acuña JM, Perez-Romero BA, Guerrero-Rodriguez JF, Martinez-Avila N and Martinez-Fierro ML: The roles of matrix metalloproteinases and their inhibitors in human diseases. *Int J Mol Sci* 21: 9739, 2020.
145. Zhang J, Liu Z, Li Y, You Q, Yang J, Jin Y, Zou G, Tang J, Ge Z and Liu Y: FGF2: A key regulator augmenting tendon-to-bone healing and cartilage repair. *Regen Med* 15: 2129-2142, 2020.
146. Bedi A, Kovacevic D, Hettrich C, Gulotta LV, Ehteshami JR, Warren RF and Rodeo SA: The effect of matrix metalloproteinase inhibition on tendon-to-bone healing in a rotator cuff repair model. *J Shoulder Elbow Surg* 19: 384-391, 2010.
147. Lo IK, Marchuk LL, Hollinshead R, Hart DA and Frank CB: Matrix metalloproteinase and tissue inhibitor of matrix metalloproteinase mRNA levels are specifically altered in torn rotator cuff tendons. *Am J Sports Med* 32: 1223-1229, 2004.
148. Choi HR, Kondo S, Hirose K, Ishiguro N, Hasegawa Y and Iwata H: Expression and enzymatic activity of MMP-2 during healing process of the acute supraspinatus tendon tear in rabbits. *J Orthop Res* 20: 927-933, 2002.
149. Yoshihara Y, Hamada K, Nakajima T, Fujikawa K and Fukuda H: Biochemical markers in the synovial fluid of glenohumeral joints from patients with rotator cuff tear. *J Orthop Res* 19: 573-579, 2001.
150. Zhen EY, Brittain IJ, Laska DA, Mitchell PG, Sumer EU, Karsdal MA and Duffin KL: Characterization of metalloproteinase cleavage products of human articular cartilage. *Arthritis Rheum* 58: 2420-2431, 2008.
151. Huang Y and Kyriakides TR: The role of extracellular matrix in the pathophysiology of diabetic wounds. *Matrix Biol Plus* 6-7: 100037, 2020.
152. Uemura S, Matsushita H, Li W, Glassford AJ, Asagami T, Lee KH, Harrison DG and Tsao PS: Diabetes mellitus enhances vascular matrix metalloproteinase activity. *Circ Res* 88: 1291-1298, 2001.
153. Morey M, O'Gaora P, Pandit A and Hélaré C: Hyperglycemia acts in synergy with hypoxia to maintain the pro-inflammatory phenotype of macrophages. *PLoS One* 14: e0220577, 2019.
154. Bedi A, Fox AJ, Harris PE, Deng XH, Ying L, Warren RF and Rodeo SA: Diabetes mellitus impairs tendon-bone healing after rotator cuff repair. *J Shoulder Elbow Surg* 19: 978-988, 2010.
155. Boyle WJ, Simonet WS and Lacey DL: Osteoclast differentiation and activation. *Nature* 423: 337-342, 2003.
156. Teissier T, Temkin V, Pollak RD and Cox LS: Crosstalk between senescent bone cells and the bone tissue microenvironment influences bone fragility during chronological age and in diabetes. *Front Physiol* 13: 812157, 2022.
157. Relaix F and Zammit PS: Satellite cells are essential for skeletal muscle regeneration: The cell on the edge returns centre stage. *Development* 139: 2845-2856, 2012.
158. Hawke TJ and Garry DJ: Myogenic satellite cells: Physiology to molecular biology. *J Appl Physiol* (1985) 91: 534-551, 2001.
159. Espino-Gonzalez E, Dalbram E, Mounier R, Gondin J, Farup J, Jessen N and Treebak JT: Impaired skeletal muscle regeneration in diabetes: From cellular and molecular mechanisms to novel treatments. *Cell Metab* 36: 1204-1236, 2024.
160. Karla EM and Debbie CT: Role of skeletal muscle in insulin resistance and glucose uptake. *Compr Physiol* 10: 785-809, 2020.
161. Chen H, Huang X, Dong M, Wen S, Zhou L and Yuan X: The association between sarcopenia and diabetes: From pathophysiology mechanism to therapeutic strategy. *Diabetes Metab Syndr Obes* 16: 1541-1554, 2023.
162. Perry BD, Caldwell MK, Brennan-Speranza TC, Sbaraglia M, Jerums G, Garnham A, Wong C, Levinger P, Asrar Ul Haq M, Hare DL, *et al*: Muscle atrophy in patients with type 2 diabetes mellitus: Roles of inflammatory pathways, physical activity and exercise. *Exerc Immunol Rev* 22: 94-109, 2016.
163. Tiplunjong C, Khimmakong W, Pholpramool C and Surinlert P: Hyperglycaemia attenuated C2C12 myoblast proliferation and induced skeletal muscle atrophy via modulating myogenic regulatory factors genes expression in diabetic rats. *Sains Malaysiana* 52: 47-56, 2023.
164. Lu J, Chen H, Lyu K, Jiang L, Chen Y, Long L, Wang X, Shi H and Li S: The functions and mechanisms of tendon stem/progenitor cells in tendon healing. *Stem Cells Int* 2023: 1258024, 2023.
165. Walia B and Huang AH: Tendon stem progenitor cells: Understanding the biology to inform therapeutic strategies for tendon repair. *J Orthop Res* 37: 1270-1280, 2019.
166. Kohler J, Popov C, Klotz B, Alberton P, Prall WC, Haasters F, Müller-Deubert S, Ebert R, Klein-Hitpass L, Jakob F, *et al*: Uncovering the cellular and molecular changes in tendon stem/progenitor cells attributed to tendon aging and degeneration. *Aging Cell* 12: 988-999, 2013.
167. Shi L, Li YJ, Dai GC, Lin YC, Li G, Wang C, Chen H and Rui YF: Impaired function of tendon-derived stem cells in experimental diabetes mellitus rat tendons: Implications for cellular mechanism of diabetic tendon disorder. *Stem Cell Res Ther* 10: 27, 2019.
168. Zhang X, Lin YC, Rui YF, Xu HL, Chen H, Wang C and Teng GJ: Therapeutic roles of tendon stem/progenitor cells in tendinopathy. *Stem Cells Int* 2016: 4076578, 2016.
169. He Z, Zeng S, Qin B, Liu G, Liu H and Bao D: Investigation on the role of Icaritin in tendon injury repair: Focusing on tendon stem cell differentiation. *J Orthop Surg Res* 20: 379, 2025.
170. Guo J, Chan KM, Zhang JF and Li G: Tendon-derived stem cells undergo spontaneous tenogenic differentiation. *Exp Cell Res* 341: 1-7, 2016.
171. Qin S, Wang W, Liu Z, Hua X, Fu S, Dong F, Li A, Liu Z, Wang P, Dai L, *et al*: Fibrochondrogenic differentiation potential of tendon-derived stem/progenitor cells from human patellar tendon. *J Orthop Translat* 22: 101-108, 2019.
172. Shengnan Q, Fei D and Wen W: Mechanism research progress of tendon-derived stem cells in reconstruction of fibrocartilage zone at bone-tendon junction. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 31: 1006-1009, 2018 (In Chinese).
173. Shen S, Lin Y, Sun J, Liu Y, Chen Y and Lu J: A New tissue engineering strategy to promote tendon-bone healing: Regulation of osteogenic and chondrogenic differentiation of tendon-derived stem cells. *Orthop Surg* 16: 2311-2325, 2024.
174. Xu L, Xu K, Wu Z, Chen Z, He Y, Ma C, Moqbel SAA, Ran J, Zhang C, Wu L and Xiong Y: Pioglitazone attenuates advanced glycation end products-induced apoptosis and calcification by modulating autophagy in tendon-derived stem cells. *J Cell Mol Med* 24: 2240-2251, 2020.
175. Wongdee K and Charoenphandhu N: Update on type 2 diabetes-related osteoporosis. *World J Diabetes* 6: 673-678, 2015.
176. San-Millán I: The key role of mitochondrial function in health and disease. *Antioxidants* 12: 782, 2023.
177. Gao S and Hu J: Mitochondrial fusion: The machineries in and out. *Trends Cell Biol* 31: 62-74, 2021.

178. Rovira-Llopis S, Bañuls C, Diaz-Morales N, Hernandez-Mijares A, Rocha M and Victor VM: Mitochondrial dynamics in type 2 diabetes: Pathophysiological implications. *Redox Biol* 11: 637-645, 2017.
179. van der Blik AM, Shen Q and Kawajiri S: Mechanisms of mitochondrial fission and fusion. *Cold Spring Harb Perspect Biol* 5: a011072, 2013.
180. Van Huynh T, Rethi L, Rethi L, Chen CH, Chen YJ and Kao YH: The complex interplay between imbalanced mitochondrial dynamics and metabolic disorders in type 2 diabetes. *Cells* 12: 1223, 2023.
181. Yu T, Robotham JL and Yoon Y: Increased production of reactive oxygen species in hyperglycemic conditions requires dynamic change of mitochondrial morphology. *Proc Natl Acad Sci USA* 103: 2653-2658, 2006.
182. Shan Z, Fa WH, Tian CR, Yuan CS and Jie N: Mitophagy and mitochondrial dynamics in type 2 diabetes mellitus treatment. *Aging (Albany NY)* 14: 2902-2919, 2022.
183. Wang S, Zhao H, Lin S, Lv Y, Lin Y, Liu Y, Peng R and Jin H: New therapeutic directions in type II diabetes and its complications: mitochondrial dynamics. *Front Endocrinol (Lausanne)* 14: 1230168, 2023.
184. Wu MY, Yang GT, Lai TT and Li CJ: The oxidative stress and mitochondrial dysfunction during the pathogenesis of diabetic retinopathy. *Oxid Med Cell Longev* 2018: 3420187, 2018.
185. Kay L, Nicolay K, Wieringa B, Saks V and Wallimann T: Direct evidence for the control of mitochondrial respiration by mitochondrial creatine kinase in oxidative muscle cells in situ. *J Biol Chem* 275: 6937-6944, 2000.
186. Rizo-Roca D, Guimarães DSPSF, Pendergrast LA, Di LeoN, Chibalin AV, Maqdasy S, Rydén M, Näslund E, Zierath JR and Krook A: Decreased mitochondrial creatine kinase 2 impairs skeletal muscle mitochondrial function independently of insulin in type 2 diabetes. *Sci Transl Med* 16: eado3022, 2024.
187. Mogi M and Horiuchi M: Diabetes and sarcopenia. In: *Diabetes and Aging-related Complications*. Yamagishi Si (ed). Springer, Singapore, pp141-151, 2018.
188. Hickey MS, Carey JO, Azevedo JL, Houmard JA, Pories WJ, Israel RG and Dohm GL: Skeletal muscle fiber composition is related to adiposity and in vitro glucose transport rate in humans. *Am J Physiol* 268 (3 Pt 1): E453-E457, 1995.
189. Tanner CJ, Barakat HA, Dohm GL, Pories WJ, MacDonald KG, Cunningham PR, Swanson MS and Houmard JA: Muscle fiber type is associated with obesity and weight loss. *Am J Physiol Endocrinol Metab* 282: E1191-E1196, 2002.
190. Oberbach A, Bossenz Y, Lehmann S, Niebauer J, Adams V, Paschke R, Schön MR, Blüher M and Punkt K: Altered fiber distribution and fiber-specific glycolytic and oxidative enzyme activity in skeletal muscle of patients with type 2 diabetes. *Diabetes Care* 29: 895-900, 2006.
191. Giebelstein J, Poschmann G, Højlund K, Schechinger W, Dietrich JW, Levin K, Beck-Nielsen H, Podwojski K, Stühler K, Meyer HE and Klein HH: The proteomic signature of insulin-resistant human skeletal muscle reveals increased glycolytic and decreased mitochondrial enzymes. *Diabetologia* 55: 1114-1127, 2012.
192. Simoneau JA and Kelley DE: Altered glycolytic and oxidative capacities of skeletal muscle contribute to insulin resistance in NIDDM. *J Appl Physiol* (1985) 83: 166-171, 1997.
193. Albers PH, Pedersen AJT, Birk JB, Kristensen DE, Vind BF, Baba O, Nøhr J, Højlund K and Wojtaszewski JF: Human muscle fiber type-specific insulin signaling: impact of obesity and type 2 diabetes. *Diabetes* 64: 485-497, 2015.
194. Al Mamun A, Rahman MM, Zaman S, Munira MS, Uddin MS, Rauf A, Banu N and Ashraf GM: Molecular insight into the crosstalk of UPS components and Alzheimer's disease. *Curr Protein Pept Sci* 21: 1193-1201, 2020.
195. Wang W, Li M, Chen Z, Xu L, Chang M, Wang K, Deng C, Gu Y, Zhou S, Shen Y, *et al*: Biogenesis and function of extracellular vesicles in pathophysiological processes of skeletal muscle atrophy. *Biochem Pharmacol* 198: 114954, 2022.
196. Shen Y, Li M, Wang K, Qi G, Liu H, Wang W, Ji Y, Chang M, Deng C, Xu F, *et al*: Diabetic muscular atrophy: Molecular mechanisms and promising therapies. *Front Endocrinol (Lausanne)* 13: 917113, 2022.
197. Yin L, Li N, Jia W, Wang N, Liang M, Yang X and Du G: Skeletal muscle atrophy: From mechanisms to treatments. *Pharmacol Res* 172: 105807, 2021.
198. Teaney NA and Cyr NE: FoxO1 as a tissue-specific therapeutic target for type 2 diabetes. *Front Endocrinol (Lausanne)* 14: 1286838, 2023.
199. Kahn D, Macias E, Zarini S, Garfield A, Zemski Berry K, Gerszten R, Schoen J, Cree-Green M and Bergman BC: Quantifying the inflammatory secretome of human intermuscular adipose tissue. *Physiol Rep* 10: e15424, 2022.
200. Yang B, Yang X, Sun X, Shi J, Shen Y, Chen R and Sakuma K: IL-6 deficiency attenuates skeletal muscle atrophy by inhibiting mitochondrial ROS production through the upregulation of PGC-1 $\alpha$  in septic mice. *Oxid Med Cell Longev* 2022: 9148246, 2022.
201. Hardee JP, Fix DK, Wang X, Goldsmith EC, Koh HJ and Carson JA: Systemic IL-6 regulation of eccentric contraction-induced muscle protein synthesis. *Am J Physiol Cell Physiol* 315: C91-C103, 2018.
202. Haberecht-Müller S, Krüger E and Fielitz J: Out of control: The role of the ubiquitin proteasome system in skeletal muscle during inflammation. *Biomolecules* 11: 1327, 2021.
203. Taniguchi CM, Emanuelli B and Kahn CR: Critical nodes in signalling pathways: Insights into insulin action. *Nat Rev Mol Cell Biol* 7: 85-96, 2006.
204. Mahadev K, Motoshima H, Wu X, Ruddy JM, Arnold RS, Cheng G, Lambeth JD and Goldstein BJ: The NAD(P)H oxidase homolog nox4 modulates insulin-stimulated generation of H<sub>2</sub>O<sub>2</sub> and plays an integral role in insulin signal transduction. *Mol Cell Biol* 24: 1844-1854, 2004.
205. Zuo J, Zhang Z, Luo M, Zhou L, Nice EC, Zhang W, Wang C and Huang C: Redox signaling at the crossroads of human health and disease. *MedComm* (2020) 3: e127, 2022.
206. O'Reilly CL, Uranga S and Fluckey JD: Culprits or consequences: Understanding the metabolic dysregulation of muscle in diabetes. *World J Biol Chem* 12: 70-86, 2021.
207. Zellers JA, Eekhoff J, Walk R, Tang SY, Hastings MK and Lake S: Effects of advanced glycation endproducts on the mechanical properties of posterior tibialis in individuals with diabetes mellitus. *Foot Ankle Orthop* 4: 2473011419S00454, 2019.
208. Yammine K, Daccache E and Assi C: Morphological changes and pathological findings in the achilles tendons of diabetic patients: A meta-analysis of comparative clinical studies. *Muscles Ligaments Tendons J* 13: 201-210, 2026.
209. Muramatsu S, Nakamura W and Etô H: Chromosomal aberrations persisting in hematopoietic tissue of mice having been subjected to lethal x-ray doses following thrombocyte transfusion. *C R Seances Soc Biol Fil* 163: 243-248, 1969 (In French).
210. Khor BYC, Woodburn J, Newcombe L and Barn R: Plantar soft tissues and Achilles tendon thickness and stiffness in people with diabetes: A systematic review. *J Foot Ankle Res* 14: 35, 2021.
211. Reddy GK: Cross-linking in collagen by nonenzymatic glycation increases the matrix stiffness in rabbit achilles tendon. *Exp Diabetes Res* 5: 143-153, 2004.
212. Xu J, Wang J, Ji Y, Liu Y, Jiang J, Wang Y, Cui X, Wan Y, Guo B and Yu H: The impact of diabetes mellitus on tendon pathology: A review. *Front Pharmacol* 15: 1491633, 2024.
213. Luciano FB and Arntfield S: Use of transglutaminases in foods and potential utilization of plants as a transglutaminase source - Review. *Biotemas* 25: 1-11, 2012.
214. Rosenthal AK, Gohr CM, Mitton E, Monnier V and Burner T: Advanced glycation end products increase transglutaminase activity in primary porcine tenocytes. *J Investig Med* 57: 460-466, 2009.
215. Heinkel D, Gohr CM, Uzuki M and Rosenthal AK: Transglutaminase contributes to CPPD crystal formation in osteoarthritis. *Front Biosci* 9: 3257-3261, 2004.
216. Zellers JA, Eekhoff JD, Walk RE, Hastings MK, Tang SY and Lake SP: Human Achilles tendon mechanical behavior is more strongly related to collagen disorganization than advanced glycation end-products content. *Sci Rep* 11: 24147, 2021.
217. Hammond MA, Gallant MA, Burr DB and Wallace JM: Nanoscale changes in collagen are reflected in physical and mechanical properties of bone at the microscale in diabetic rats. *Bone* 60: 26-32, 2014.
218. Choi JH, Kim HR and Song KH: Musculoskeletal complications in patients with diabetes mellitus. *Korean J Intern Med* 37: 1099-1110, 2022.
219. Patel SH and Carroll CC: Impact of elevated serum advanced glycation end products and exercise on intact and injured murine tendons. *Connect Tissue Res* 64: 161-174, 2023.
220. Bayraktutan U: Free radicals, diabetes and endothelial dysfunction. *Diabetes Obes Metab* 4: 224-238, 2002.

221. Adela R, Nethi SK, Bagul PK, Barui AK, Mattapally S, Kuncha M, Patra CR, Reddy PN and Banerjee SK: Hyperglycaemia enhances nitric oxide production in diabetes: A study from South Indian patients. *PLoS One* 10: e0125270, 2015.
222. Xia W, Szomor Z, Wang Y and Murrell GA: Nitric oxide enhances collagen synthesis in cultured human tendon cells. *J Orthop Res* 24: 159-172, 2006.
223. Cazzaniga A, Scrimieri R, Giani E, Zuccotti GV and Maier JAM: Endothelial hyper-permeability induced by T1D sera can be reversed by iNOS inactivation. *Int J Mol Sci* 21: 2798, 2020.
224. Hwang MH and Kim S: Type 2 diabetes: Endothelial dysfunction and exercise. *J Exerc Nutrition Biochem* 18: 239-247, 2014.
225. Pandolfi A and De Filippis EA: Chronic hyperglycemia and nitric oxide bioavailability play a pivotal role in pro-atherogenic vascular modifications. *Genes Nutr* 2: 195-208, 2007.
226. Pitocco D, Tesaro M, Alessandro R, Ghirlanda G and Cardillo C: Oxidative Stress in Diabetes: Implications for Vascular and Other Complications. *Int J Mol Sci* 14: 21525-21550, 2013.
227. Mawhinney AE, Bertolet M, Minni M, Barsotti R, Young L and Chen Q: Effects of NOX-1 on Real-Time Blood Nitric Oxide and Hydrogen Peroxide in Acute Hyperglycemia. *FASEB J* 30: 734.10, 2016.
228. Singh R, Barden A, Mori T and Beilin L: Advanced glycation end-products: A review. *Diabetologia* 44: 129-146, 2001.
229. Nichols AEC, Best KT and Loiselle AE: The cellular basis of fibrotic tendon healing: Challenges and opportunities. *Transl Res* 209: 156-168, 2019.
230. Yang L, Song Y, Wang T, Cui Z, Wu J, Shi Y, Yu Z and Song B: Transcription factor c-Maf drives macrophages to promote hypertrophic scar formation. *J Cosmet Dermatol* 23: 639-647, 2023.
231. Teng C, Zhou C, Xu D and Bi F: Combination of platelet-rich plasma and bone marrow mesenchymal stem cells enhances tendon-bone healing in a rabbit model of anterior cruciate ligament reconstruction. *J Orthop Surg Res* 11: 96, 2016.
232. Nakase J, Kitaoka K, Matsumoto K and Tomita K: Facilitated tendon-bone healing by local delivery of recombinant hepatocyte growth factor in rabbits. *Arthroscopy* 26: 84-90, 2010.
233. Lu H, Liu F, Chen C, Wang Z, Chen H, Qu J, Zhang T, Xu D and Hu J: Low-Intensity pulsed ultrasound stimulation for tendon-bone healing. *Am J Phys Med Rehabil* 97: 270-277, 2018.
234. Font Tellado S, Balmayor ER and Van Griensven M: Strategies to engineer tendon/ligament-to-bone interface: Biomaterials, cells and growth factors. *Adv Drug Deliv Rev* 94: 126-140, 2015.
235. Qu D, Subramony SD, Boskey AL, Pleshko N, Doty SB and Lu HH: Compositional mapping of the mature anterior cruciate ligament-to-bone insertion. *J Orthop Res* 35: 2513-2523, 2017.
236. Spalazzi JP, Boskey AL, Pleshko N and Lu HH: Quantitative mapping of matrix content and distribution across the ligament-to-bone insertion. *PLoS One* 8: e74349, 2013.
237. Zhong S, Lan Y, Liu J, Seng Tam M, Hou Z, Zheng Q, Fu S and Bao D: Advances focusing on the application of decellularization methods in tendon-bone healing. *J Adv Res* 67: 361-372, 2025.
238. Patel S, Caldwell JM, Doty SB, Levine WN, Rodeo S, Soslowsky LJ, Thomopoulos S and Lu HH: Integrating soft and hard tissues via interface tissue engineering. *J Orthop Res* 36: 1069-1077, 2018.
239. Yang PJ and Temenoff JS: Engineering orthopedic tissue interfaces. *Tissue Eng Part B Rev* 15: 127-141, 2009.
240. Benjamin M, Toumi H, Ralphs JR, Bydder G, Best TM and Milz S: Where tendons and ligaments meet bone: attachment sites ('entheses') in relation to exercise and/or mechanical load. *J Anat* 208: 471-490, 2006.
241. Zitnay JL and Weiss JA: Load transfer, damage, and failure in ligaments and tendons. *J Orthop Res* 36: 3093-3104, 2018.
242. Killian ML: Growth and mechanobiology of the tendon-bone enthesis. *Semin Cell Dev Biol* 123: 64-73, 2022.
243. Barajaa MA, Nair LS and Laurencin CT: Bioinspired scaffold designs for regenerating musculoskeletal tissue interfaces. *Regen Eng Transl Med* 6: 451-483, 2020.
244. Apostolakis J, Durant TJ, Dwyer CR, Russell RP, Weinreb JH, Alaei F, Beitzel K, McCarthy MB, Cote MP and Mazzocca AD: The enthesis: A review of the tendon-to-bone insertion. *Muscles Ligaments Tendons J* 4: 333-342, 2014.
245. Wang N, Tytell JD and Ingber DE: Mechanotransduction at a distance: mechanically coupling the extracellular matrix with the nucleus. *Nat Rev Mol Cell Biol* 10: 75-82, 2009.
246. Humphrey JD, Dufresne ER and Schwartz MA: Mechanotransduction and extracellular matrix homeostasis. *Nat Rev Mol Cell Biol* 15: 802-812, 2014.
247. Xie W, Wei X, Kang H, Jiang H, Chu Z, Lin Y, Hou Y and Wei Q: Static and dynamic: Evolving biomaterial mechanical properties to control cellular mechanotransduction. *Adv Sci (Weinh)* 10: e2204594, 2023.
248. Xu Z, Xu W, Zhang T and Luo L: Mechanisms of tendon-bone interface healing: biomechanics, cell mechanics, and tissue engineering approaches. *J Orthop Surg Res* 19: 817, 2024.
249. Pastino AK, Greco TM, Mathias RA, Cristea IM and Schwarzbauer JE: Stimulatory effects of advanced glycation endproducts (AGEs) on fibronectin matrix assembly. *Matrix Biol* 59: 39-53, 2017.
250. Karamessinis PM, Tzinia AK, Kitsiou PV, Stetler-Stevenson WG, Michael AF, Fan WW, Zhou B, Margaritis LH and Tsilibary EC: Proximal tubular epithelial cell integrins respond to high glucose by altered cell-matrix interactions and differentially regulate matrixin expression. *Lab Invest* 82: 1081-1093, 2002.
251. Javed F and Romanos GE: Chronic hyperglycemia as a risk factor in implant therapy. *Periodontol* 2000 81: 57-63, 2019.
252. Wang S, Deng Z, Ma Y, Jin J, Qi F, Li S, Liu C, Lyu FJ and Zheng Q: The role of autophagy and mitophagy in bone metabolic disorders. *Int J Biol Sci* 16: 2675-2691, 2020.
253. Forner P and Sheu A: Bone health in patients with type 2 diabetes. *J Endocr Soc* 8: bvaa112, 2024.
254. Bonewald LF: The amazing osteocyte. *J Bone Miner Res* 26: 229-238, 2011.
255. Pathak JL, Bravenboer N and Klein-Nulend J: The osteocyte as the new discovery of therapeutic options in rare bone diseases. *Front Endocrinol (Lausanne)* 11: 405, 2020.
256. Zhu J, Tang Y, Wu Q, Ji YC and Kang FW: Mechanism of participation of osteocytes in the formation of osteoclasts under hypoxia. *Hua Xi Kou Qiang Yi Xue Za Zhi* 37: 463-468, 2019 (In Chinese).
257. Eckhardt BA, Rowsey JL, Thicke BS, Fraser DG, O'Grady KL, Bondar OP, Hines JM, Singh RJ, Thoreson AR, Rakshit K, *et al*: Accelerated osteocyte senescence and skeletal fragility in mice with type 2 diabetes. *JCI Insight* 5: e135236, 2020.
258. Villarino ME, Sánchez LM, Bozal CB and Ubios AM: Influence of short-term diabetes on osteocytic lacunae of alveolar bone. A histomorphometric study. *Acta Odontol Latinoam* 19: 23-28, 2006.
259. Tanaka K, Yamaguchi T, Kanazawa I and Sugimoto T: Effects of high glucose and advanced glycation end products on the expressions of sclerostin and RANKL as well as apoptosis in osteocyte-like MLO-Y4-A2 cells. *Biochem Biophys Res Commun* 461: 193-199, 2015.
260. Tchkonja T, Zhu Y, van Deursen J, Campisi J and Kirkland JL: Cellular senescence and the senescent secretory phenotype: Therapeutic opportunities. *J Clin Invest* 123: 966-972, 2013.
261. Moiseeva V, Cisneros A, Cobos AC, Tarrega AB, Oñate CS, Perdiguerro E, Serrano AL and Muñoz-Cánoves P: Context-dependent roles of cellular senescence in normal, aged, and disease states. *FEBS J* 290: 1161-1185, 2023.
262. Birch J and Gil J: Senescence and the SASP: Many therapeutic avenues. *Genes Dev* 34: 1565-1576, 2020.
263. Canalis E: Wnt signalling in osteoporosis: Mechanisms and novel therapeutic approaches. *Nat Rev Endocrinol* 9: 575-583, 2013.
264. Montgomery MK and Turner N: Mitochondrial dysfunction and insulin resistance: An update. *Endocr Connect* 4: R1-R15, 2015.
265. Galatz LM, Sandell LJ, Rothermich SY, Das R, Mastny A, Havlioglu N, Silva MJ and Thomopoulos S: Characteristics of the rat supraspinatus tendon during tendon-to-bone healing after acute injury. *J Orthop Res* 24: 541-550, 2006.
266. Rojas LB and Gomes MB: Metformin: An old but still the best treatment for type 2 diabetes. *Diabetol Metab Syndr* 5: 6, 2013.
267. Zhou Z, Tang Y, Jin X, Chen C, Lu Y, Liu L and Shen C: Metformin inhibits advanced glycation end products-induced inflammatory response in murine macrophages partly through AMPK activation and RAGE/NFκB pathway suppression. *J Diabetes Res* 2016: 4847812, 2016.
268. Ishibashi Y, Matsui T, Takeuchi M and Yamagishi S: Metformin inhibits advanced glycation end products (AGEs)-induced renal tubular cell injury by suppressing reactive oxygen species generation via reducing receptor for AGEs (RAGE) expression. *Horm Metab Res* 44: 891-895, 2012.
269. Pang R, Guan M, Zheng Z and Xue Y: Effects of metformin on apoptosis induced by advanced glycation end-products and expressions of caspase-3, Bax and Bcl-2 in human dermal fibroblasts in vitro. *Nan Fang Yi Ke Da Xue Xue Bao* 35: 898-902, 2015 (In Chinese).

270. Alikhani M, Alikhani Z, Boyd C, MacLellan CM, Raptis M, Liu R, Pischon N, Trackman PC, Gerstenfeld L and Graves DT: Advanced glycation end products stimulate osteoblast apoptosis via the MAP kinase and cytosolic apoptotic pathways. *Bone* 40: 345-353, 2007.
271. Neumann B, Baror R, Zhao C, Segel M, Dietmann S, Rawji KS, Foerster S, McClain CR, Chalut K, van Wijngaarden P and Franklin RJM: Metformin restores CNS remyelination capacity by rejuvenating aged stem cells. *Cell Stem Cell* 25: 473-485.e8, 2019.
272. Ma J, Zhang ZL, Hu XT, Wang XT and Chen AM: Metformin promotes differentiation of human bone marrow derived mesenchymal stem cells into osteoblast via GSK3 $\beta$  inhibition. *Eur Rev Med Pharmacol Sci* 22: 7962-7968, 2018.
273. Yang K, Cao F, Qiu S, Jiang W, Tao L and Zhu Y: Metformin promotes differentiation and attenuates H<sub>2</sub>O<sub>2</sub>-Induced oxidative damage of osteoblasts via the PI3K/AKT/Nrf2/HO-1 pathway. *Front Pharmacol* 13: 829830, 2022.
274. Liu Y, Wang X, Wu J, Wu A and Yu X: Metformin and bone metabolism: Unraveling their direct and indirect effects. *Ther Adv Endocrinol Metab* 16: 20420188251397207, 2025.
275. Zhou Q, Guan Z, Liu S, Xuan Y, Han G, Chen H, Jin X, Tao K and Guan Z: The effects of metformin and alendronate in attenuating bone loss and improving glucose metabolism in diabetes mellitus mice. *Aging (Albany NY)* 14: 272-285, 2022.
276. Cummings SR, San Martin J, McClung MR, Siris ES, Eastell R, Reid IR, Delmas P, Zoog HB, Austin M, Wang A, *et al*: Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N Engl J Med* 361: 756-765, 2009.
277. Cosman F, Crittenden DB, Adachi JD, Binkley N, Czerwinski E, Ferrari S, Hofbauer LC, Lau E, Lewiecki EM, Miyauchi A, *et al*: Romosozumab treatment in postmenopausal women with osteoporosis. *N Engl J Med* 375: 1532-1543, 2016.
278. Lawlor N, George J, Bolisetty M, Kursawe R, Sun L, Sivakamasundari V, Kycia I, Robson P and Stitzel ML: Single-cell transcriptomes identify human islet cell signatures and reveal cell-type-specific expression changes in type 2 diabetes. *Genome Res* 27: 208-222, 2017.
279. Brissova M, Fowler MJ, Nicholson WE, Chu A, Hirshberg B, Harlan DM and Powers AC: Assessment of human pancreatic islet architecture and composition by laser scanning confocal microscopy. *J Histochem Cytochem* 53: 1087-1097, 2005.
280. Blodgett DM, Nowosielska A, Afik S, Pechhold S, Cura AJ, Kennedy NJ, Kim S, Kucukural A, Davis RJ, Kent SC, *et al*: Novel observations from next-generation RNA sequencing of highly purified human adult and fetal islet cell subsets. *Diabetes* 64: 3172-3181, 2015.
281. Geach T: Transcriptomes reveal specific islet cell signatures. *Nat Rev Endocrinol* 12: 686, 2016.
282. Zadian SS, Jahanbin K, Nikooei S, Rostaminejad M, Rahimi A, Abdizadeh P and Alipoor B: Current trends in single-cell RNA sequencing applications in diabetes mellitus. *FEBS Open Bio* 15: 1738-1761, 2025.
283. Yang H, Luo J, Liu X, Luo Y, Lai X and Zou F: Unveiling cell subpopulations in T1D mouse islets using single-cell RNA sequencing. *Am J Physiol Endocrinol Metab* 326: E723-E734, 2024.
284. Honardoost MA, Adinatha A, Schmidt F, Ranjan B, Ghaeidamini M, Arul Rayan N, Gek Liang Lim M, Joanito I, Xiao Xuan Lin Q, Rajagopalan D, *et al*: Systematic immune cell dysregulation and molecular subtypes revealed by single-cell RNA-seq of subjects with type 1 diabetes. *Genome Med* 16: 45, 2024.
285. Segerstolpe Å, Palasantza A, Eliasson P, Andersson EM, Andréasson AC, Sun X, Picelli S, Sabirsh A, Clausen M, Bjursell MK, *et al*: Single-cell transcriptome profiling of human pancreatic islets in health and type 2 diabetes. *Cell Metab* 24: 593-607, 2016.
286. Armstrong DG, Ingelfinger JR, Boulton AJM and Bus SA: Diabetic foot ulcers and their recurrence. *N Engl J Med* 376: 2367-2375, 2017.
287. Wong R, Geyer S, Weninger W, Guimberteau JC and Wong JK: The dynamic anatomy and patterning of skin. *Exp Dermatol* 25: 92-98, 2016.
288. Theodoridis G, Tekkela S, Veves A, McGrath JA and Onoufriadis A: Single-cell transcriptomics in human skin research: Available technologies, technical considerations and disease applications. *Exp Dermatol* 31: 655-673, 2022.
289. Patel S, Srivastava S, Singh MR and Singh D: Mechanistic insight into diabetic wounds: Pathogenesis, molecular targets and treatment strategies to pace wound healing. *Biomed Pharmacother* 112: 108615, 2019.
290. Luo W, Li N, Liu J, Li D, Li Y, Ma Q, Lin C, Lu L and Lin S: Bulk and single-cell transcriptome analyses unravel gene signatures of mitochondria-associated programmed cell death in diabetic foot ulcer. *J Cell Mol Med* 28: e70319, 2024.
291. Wang Z, Wei D, Li S, Tang Q, Lu G, Gu S, Lu L, Liang F, Teng J, Lin J, *et al*: Healing mechanism of diabetic foot ulcers using single-cell RNA-sequencing. *Ann Transl Med* 11: 210, 2023.
292. Wang S, Sun ST, Zhang XY, Ding HR, Yuan Y, He JJ, Wang MS, Yang B and Li YB: The Evolution of Single-Cell RNA Sequencing Technology and Application: Progress and Perspectives. *Int J Mol Sci* 24: 2943, 2023.
293. Mo R, Jiang T, Di J, Tai W and Gu Z: Emerging micro- and nanotechnology based synthetic approaches for insulin delivery. *Chem Soc Rev* 43: 3595-3629, 2014.
294. Azevedo C, Pinto S, Benjakul S, Nilsen J, Santos HA, Traverso G, Andersen JT and Sarmiento B: Prevention of diabetes-associated fibrosis: Strategies in FcRn-targeted nanosystems for oral drug delivery. *Adv Drug Deliv Rev* 175: 113778, 2021.
295. Gualandi-Signorini AM and Giorgi G: Insulin formulations-a review. *Eur Rev Med Pharmacol Sci* 5: 73-83, 2001.
296. Kumar A, Zhang X and Liang XJ: Gold nanoparticles: Emerging paradigm for targeted drug delivery system. *Biotechnol Adv* 31: 593-606, 2013.
297. Ashique S, Sandhu NK, Chawla V and Chawla PA: Targeted drug delivery: Trends and perspectives. *Curr Drug Deliv* 18: 1435-1455, 2021.
298. Abbasi M, Heath B and McGinness L: Advances in metformin-delivery systems for diabetes and obesity management. *Diabetes Obes Metab* 26: 3513-3529, 2024.
299. Mansoor S, Kondiah PPD, Choonara YE and Pillay V: Polymer-based nanoparticle strategies for insulin delivery. *Polymers (Basel)* 11: 1380, 2019.
300. Pereira ASBF, de Souza Lima ML, da Silva-Junior AA, dos Santos Silva E, de Araújo Júnior RF, Martins AA, Alves JSF, Oliveira AS, De Santis Ferreira L, de Araújo Costa ECT, *et al*: In vitro-in vivo availability of metformin hydrochloride-PLGA nanoparticles in diabetic rats in a periodontal disease experimental model. *Pharm Biol* 59: 1574-1582, 2021.
301. Ghosh K, Kanapathipillai M, Korin N, McCarthy JR and Ingber DE: Polymeric nanomaterials for islet targeting and immunotherapeutic delivery. *Nano Lett* 12: 203-208, 2012.
302. He Z, Peng Q, Bin W, Zhao L, Chen Y, Zhang Y, Yang W, Yan X and Liu H: Nucleic acid aptamers in orthopedic diseases: promising therapeutic agents for bone disorders. *Bone Res* 13: 71, 2025.
303. Wu Z, Tang LJ, Zhang XB, Jiang JH and Tan W: Aptamer-modified nanodrug delivery systems. *ACS Nano* 5: 7696-7699, 2011.
304. Yan L, Zhang J, Lee CS and Chen X: Micro- and nanotechnologies for intracellular delivery. *Small* 10: 4487-4504, 2014.
305. Xiao Z and Farokhzad OC: Aptamer-Functionalized nanoparticles for medical applications: Challenges and opportunities. *ACS Nano* 6: 3670-3676, 2012.
306. Yamada Y, Tabata M, Yasuzaki Y, Nomura M, Shibata A, Bayashi Y, Taniguchi Y, Sasaki S and Harashima H: A nano-carrier system for the delivery of nucleic acids targeted to a pancreatic beta cell line. *Biomaterials* 35: 6430-6438, 2014.
307. Xia B, Xu F, Chen J, Shan S, Shen J, Zhang Y, Wang J, Zhou Z, Sun W, Jia Y, *et al*: Site-specific adaptive nanovesicles for oral insulin delivery. *Sci Adv* 11: eady6386, 2025.
308. Chu C, Wei M, Bian C, Bi X, Deng Y, Xiao P, Zhao J, Wang Y, He H, Gou J, *et al*: Oral dosed organo-silica nanoparticles restore glucose homeostasis and  $\beta$ -cell function in diabetes rats. *Adv Funct Mater* 36: e19628, 2025.
309. Ghasemiyeh P and Mohammadi-Samani S: Potential of nanoparticles as permeation enhancers and targeted delivery options for skin: Advantages and disadvantages. *Drug Des Devel Ther* 14: 3271-3289, 2020.
310. Montanheiro TLD, Schatkoski VM, de Menezes BRC, Pereira RM, Ribas RG, de Freitas ADM, Lemes AP, Fernandes MHFV and Thim GP: Recent progress on polymer scaffolds production: Methods, main results, advantages and disadvantages. *Express Polym Lett* 16: 197-219, 2022.
311. Papatheodorou K, Papanas N, Banach M, Papazoglou D and Edmonds M: Complications of diabetes 2016. *J Diabetes Res* 2016: 6989453, 2016.
312. Alowais SA, Alghamdi SS, Alsuhbany N, Alqahtani T, Alshaya AI, Almohareb SN, Aldairem A, Alrashed M, Bin Saleh K, Badreldin HA, *et al*: Revolutionizing healthcare: The role of artificial intelligence in clinical practice. *BMC Med Educ* 23: 689, 2023.

313. Dankwa-Mullan I, Rivo M, Sepulveda M, Park Y, Snowdon J and Rhee K: Transforming diabetes care through artificial intelligence: The future is here. *Popul Health Manag* 22: 229-242, 2019.
314. Ellahham S: Artificial intelligence: The future for diabetes care. *Am J Med* 133: 895-900, 2020.
315. Shu T, Zhang B and Yan Tang Y: An extensive analysis of various texture feature extractors to detect diabetes mellitus using facial specific regions. *Comput Biol Med* 83: 69-83, 2017.
316. Li J, Yuan P, Hu X, Huang J, Cui L, Cui J, Ma X, Jiang T, Yao X, Li J, *et al*: A tongue features fusion approach to predicting prediabetes and diabetes with machine learning. *J Biomed Inform* 115: 103693, 2021.
317. Bellemo V, Lim G, Rim TH, Tan GSW, Cheung CY, Sadda S, He MG, Tufail A, Lee ML, Hsu W, *et al*: Artificial intelligence screening for diabetic retinopathy: the real-world emerging application. *Curr Diab Rep* 19: 72, 2019.
318. American Diabetes Association: 11. Microvascular complications and foot care: Standards of medical care in diabetes-2020. *Diabetes Care* 43 (Suppl 1): S135-S151, 2020.
319. Keel S, Lee PY, Scheetz J, Li Z, Kotowicz MA, MacIsaac RJ and He M: Feasibility and patient acceptability of a novel artificial intelligence-based screening model for diabetic retinopathy at endocrinology outpatient services: A pilot study. *Sci Rep* 8: 4330, 2018.
320. Elafros MA, Andersen H, Bennett DL, Savelieff MG, Viswanathan V, Callaghan BC and Feldman EL: Towards prevention of diabetic peripheral neuropathy: Clinical presentation, pathogenesis, and new treatments. *Lancet Neurol* 21: 922-936, 2022.
321. Williams BM, Borroni D, Liu R, Zhao Y, Zhang J, Lim J, Ma B, Romano V, Qi H, Ferdousi M, *et al*: An artificial intelligence-based deep learning algorithm for the diagnosis of diabetic neuropathy using corneal confocal microscopy: A development and validation study. *Diabetologia* 63: 419-430, 2020.
322. Preston FG, Meng Y, Burgess J, Ferdousi M, Azmi S, Petropoulos IN, Kaye S, Malik RA, Zheng Y and Alam U: Artificial intelligence utilising corneal confocal microscopy for the diagnosis of peripheral neuropathy in diabetes mellitus and prediabetes. *Diabetologia* 65: 457-466, 2022.
323. Murphree DH, Arabmakki E, Ngufor C, Storlie CB and McCoy RG: Stacked classifiers for individualized prediction of glycemic control following initiation of metformin therapy in type 2 diabetes. *Comput Biol Med* 103: 109-115, 2018.
324. Tarumi S, Takeuchi W, Chalkidis G, Rodriguez-Loya S, Kuwata J, Flynn M, Turner KM, Sakaguchi FH, Weir C, Kramer H, *et al*: Leveraging artificial intelligence to improve chronic disease care: Methods and application to pharmacotherapy decision support for type-2 diabetes mellitus. *Methods Inf Med* 60(S 01): e32-e43, 2021.
325. Allalou A, Nalla A, Prentice KJ, Liu Y, Zhang M, Dai FF, Ning X, Osborne LR, Cox BJ, Gunderson EP and Wheeler MB: A predictive metabolic signature for the transition from gestational diabetes mellitus to type 2 diabetes. *Diabetes* 65: 2529-2539, 2016.
326. Huang YJ, Chen CH and Yang HC: AI-enhanced integration of genetic and medical imaging data for risk assessment of type 2 diabetes. *Nat Commun* 15: 4230, 2024.
327. Woldaregay AZ, Årsand E, Walderhaug S, Albers D, Mamykina L, Botsis T and Hartvigsen G: Data-driven modeling and prediction of blood glucose dynamics: Machine learning applications in type 1 diabetes. *Artif Intell Med* 98: 109-134, 2019.
328. Zheng Y, Ley SH and Hu FB: Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat Rev Endocrinol* 14: 88-98, 2018.
329. Deng Y, Lu L, Aponte L, Angelidi AM, Novak V, Karniadakis GE and Mantzoros CS: Deep transfer learning and data augmentation improve glucose levels prediction in type 2 diabetes patients. *NPJ Digit Med* 4: 109, 2021.
330. Wang G, Liu X, Ying Z, Yang G, Chen Z, Liu Z, Zhang M, Yan H, Lu Y, Gao Y, *et al*: Optimized glycemic control of type 2 diabetes with reinforcement learning: A proof-of-concept trial. *Nat Med* 29: 2633-2642, 2023.
331. Sheng B, Pushpanathan K, Guan Z, Lim QH, Lim ZW, Yew SME, Goh JHL, Bee YM, Sabanayagam C, Sevdalis N, *et al*: Artificial intelligence for diabetes care: Current and future prospects. *Lancet Diabetes Endocrinol* 12: 569-595, 2024.
332. The Lancet Diabetes Endocrinology: Diabetes care and AI: A looming threat or a necessary advancement? *Lancet Diabetes Endocrinol* 11: 441, 2023.



Copyright © 2026 He et al. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.