

1 α ,25-Dihydroxyvitamin D₃ enhances γ -glutamyl transpeptidase activity in LLC-PK1 porcine kidney epithelial cells

RITSU ITO, HIDEYUKI IHARA, TAKAHIRO OKADA and YOSHITAKA IKEDA

Division of Molecular Cell Biology, Department of Biomolecular Sciences,
Faculty of Medicine, Saga University, Saga 849-8501, Japan

Received September 27, 2013; Accepted November 15, 2013

DOI: 10.3892/mmr.2014.2436

Abstract. Mammalian γ -glutamyl transpeptidase (GGT) is expressed most highly in the kidney and serves to recover the constituent amino acids of glutathione in the proximal tubules. Serum GGT is used as a marker for obstructive jaundice and alcoholic liver disease and it has been reported that urinary GGT is a potential marker for bone resorption. The present study investigated the effect of derivatives of vitamin D₃ on GGT activity in LLC-PK1 porcine renal tubular epithelial cells in order to analyze the biochemical basis of bone metabolism-associated alterations in GGT activity in the kidney. In the presence of 1 α ,25-dihydroxyvitamin D₃ [1,25(OH)₂VD₃], GGT activity was observed to be significantly increased in LLC-PK1 cells, with an increase in GGT activity also found in the cell medium. While the stimulatory effect of 1-OH-VD₃ was similar to that of 1,25(OH)₂VD₃, vitamin D₃ and 25-OH-VD₃ had no effect on GGT activity. The increased GGT activity caused by 1,25(OH)₂VD₃ in LLC-PK1 cells was the result of long-term stimulation of the cells, in contrast to the GGT-induced increase in alkaline phosphatase, which is more transient. Polymerase chain reaction analysis revealed that the 1,25(OH)₂VD₃-induced increase in GGT activity was due to prolonged GGT turnover, rather than increased GGT expression, as no increase in GGT mRNA expression was observed. Thus, it is likely that the expression of GGT is not entirely constitutive in the kidney, but is altered under certain conditions, including under hormonal regulation.

Introduction

Mammalian γ -glutamyl transpeptidase (GGT) has a role in glutathione metabolism by catalyzing the hydrolysis of a γ -glutamyl moiety of glutathione and associated

compounds (1-3). GGT also catalyzes the transfer of γ -glutamyl groups from γ -glutamylated compounds, including that of glutathione to amino acids or dipeptides. The mammalian form of GGT is a membrane-bound glycoprotein with a typical type II membrane protein topology and is anchored to the extracellular surface of cell membranes through a non-cleavable N-terminal signal-anchor domain.

GGT activity is higher in the kidney, intestine and epididymis compared with other tissues and GGT is constitutively expressed in these tissues (4). A previous study using GGT-deficient mice revealed that GGT has an essential role in the kidney in the recovery of cysteine and cystine from glutathione that is excreted into the urine (5). However, while the activity of GGT is undetectable in the adult rat liver, it is relatively high in the fetal rat liver (6). GGT activity can be induced in the adult liver by various chemical compounds, including alcohol, xenobiotics and associated drugs. Furthermore, GGT expression is associated with hepatocarcinogenesis in rats (7-10). Thus, it is well established that GGT is of potential value in clinical chemistry for the diagnosis of certain types of hepatic disease.

Niida *et al* (11) reported that GGT is a bone-resorbing factor. This novel biological function of GGT does not depend on its enzymatic activity. GGT is capable of inducing osteoclasts by stimulating the expression of the receptor activator of nuclear factor- κ B, which has further been confirmed using specific antibodies *in vivo* (12). Furthermore, an investigation using transgenic mice indicated that GGT overexpression accelerates bone resorption, leading to the development of osteoporosis (13). It has also been reported that significant increases in urinary GGT activity are associated with elevated bone resorption (14), suggesting that urinary GGT activity has potential to be a marker for bone resorption.

It is well established that among the various animal tissues, GGT is most abundant in the kidney, and that this expression is observed exclusively in the brush border membrane of the proximal tubular epithelial cells (15,16). Thus, increased urinary GGT activity, which may be associated with bone resorption, is likely to be due to enhanced GGT release from epithelial cells into the luminal space of the proximal tubules. However, whether kidney GGT is affected in conjunction with an alteration in extracellular signal molecules and the systemic metabolic status or others, has yet to be elucidated. The present study aimed to investigate the effect of vitamin D₃ and its

Correspondence to: Dr Yoshitaka Ikeda, Division of Molecular Cell Biology, Department of Biomolecular Sciences, Faculty of Medicine, Saga University, 5-1-1 Nabeshima, Saga 849-8501, Japan
E-mail: yiked@med.saga-u.ac.jp

Key words: vitamin D₃, 1 α ,25-dihydroxyvitamin D₃, calcitriol, LLC-PK1 cells, γ -glutamyltranspeptidase

metabolites, which are factors involved in bone metabolism, on GGT activity in LLC-PK1 pig kidney cells, which are cells derived from the proximal tubular epithelium.

Materials and methods

Cell line and cell culture. LLC-PK1 cells were obtained from the Health Science Research Resources Bank (Osaka, Japan) and were maintained at 37°C in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, 100 μ g/ml streptomycin and 100 U/ml penicillin under 5% CO₂ in humidified air.

GGT activity assay. GGT activity was assessed according to the method described by Tate and Meister (17). Transpeptidation reactions were performed at 37°C using 1 mM γ -glutamyl-*p*-nitroanilide (donor substrate; Wako Pure Chemical Industries Ltd., Osaka, Japan) and 20 mM glycylglycine (acceptor substrate; Wako Pure Chemical Industries Ltd.), in 0.1 M Tris-HCl (pH 8.0). The release of *p*-nitroaniline was monitored spectrophotometrically at 410 nm and the activity was calculated using a molar extinction coefficient of 8.8.

Effect of vitamin D₃ and its derivatives on the enzyme activity of GGT in LLC-PK1 cells. In nearly confluent cultures of LLC-PK1 cells, the medium was changed immediately prior to the addition of the agent to be analyzed. Cells were cultured for the indicated durations in the presence of the agent. Vitamin D₃ and its derivatives (Sigma-Aldrich, St. Louis, MO, USA) were added to the cells subsequent to being dissolved in ethanol, which was used as the vehicle. Parathyroid hormone (PTH; American Research Products, Waltham, MA, USA) was dissolved in phosphate buffered saline (PBS). Following culture for the specified durations, the medium was collected and the cells were harvested and washed two or three times using PBS. Cells were centrifuged at 150 x g for 5 min, then resuspended in a small volume of PBS. The cells were lysed using sonication and were subjected to an enzyme activity assay.

Reverse transcription polymerase chain reaction (RT-PCR) analysis for GGT expression. Total RNA was extracted from the LLC-PK1 cells using an RNeasy® Mini Kit (Qiagen, Valencia, CA, USA) according to the manufacturer's instructions. First-strand cDNAs were synthesized from 0.5 μ g total RNA using oligo dT primers and a ReverTra Plus RT-PCR kit (Toyobo, Osaka, Japan). PCR amplification of the 495 bp cDNA fragments using specific primers for pig GGT was performed using a Program Temp Control System PC-708 (Astec, Fukuoka, Japan) in a 50 μ l reaction volume containing 1 μ l cDNA, 1.0 U KOD-Plus (Toyobo), 1 mM MgSO₄, 0.2 mM deoxynucleotide triphosphates and 2% dimethyl sulfoxide in the buffer supplied along with the enzyme according to the manufacturer's instructions. The RT-PCR products were electrophoresed on 1.5% agarose gel containing ethidium bromide.

Protein concentration determination. Protein concentration was determined using a Bradford protein assay kit (Pierce Chemical Company, Rockford, IL, USA) with bovine serum albumin as a standard.

Statistical analysis. Data were analyzed by a t-test using Prism statistical software (GraphPad Software, Inc., San Diego, CA, USA). Data are presented as the mean \pm standard deviation. $P < 0.01$ was considered to indicate a statistically significant difference.

Results

Effect of vitamin D₃ and its derivatives on GGT activity in LLC-PK1 renal tubular cells. LLC-PK1 cells are a well defined renal tubule-derived cell line which exhibit high GGT activity (18,19), similar to the level observed in the proximal tubules in the kidney. As shown in Fig. 1A, the biologically active form of vitamin D₃, 1 α ,25-dihydroxyvitamin D₃ [1,25(OH)₂VD₃], also known as calcitriol, was found to significantly enhance GGT activity in the LLC-PK1 cells, while the non-hydroxylated vitamin D₃ had no effect on GGT activity. The 1 α -monohydroxylated form of vitamin D₃ 1-OH-VD₃ also appeared to increase GGT activity to an extent similar to that induced by 1,25(OH)₂VD₃; however, this increase was not statistically significant due to experimental variations. After three days of culture, GGT activity was observed to be significantly higher in the medium of the cells treated with 1-OH-VD₃ and 1,25(OH)₂VD₃, compared with those treated with the vehicle or the control cells (Fig. 1B).

A precursor of the active form of vitamin D₃, 25(OH)VD₃, was found to have no effect on GGT activity in LLC-PK1 cells, although it is known that LLC-PK1 cells exhibit some 25(OH)VD₃ 1 α -hydroxylase activity (20). Thus, the levels of 1 α -hydroxylase may have been insufficient to cause an increase in the levels of 1,25(OH)₂VD₃ and stimulate GGT activity. Hydroxylase activity in the renal tubular epithelia is induced by PTH (23); however, the PTH receptor is not expressed in LLC-PK1 cells (21,22). In the present study, PTH was unable to stimulate GGT activity in the LLC-PK1 cells, even in the presence of 25(OH)VD₃ (data not shown).

Dose-dependent effect of 1,25(OH)₂VD₃ on GGT activity in LLC-PK1 renal tubular cells. LLC-PK1 cells were cultured for three days in the presence of various concentrations of 1,25(OH)₂VD₃. The GGT activity in the cells and medium was then assessed. As shown in Fig. 2, GGT activity was found to be significantly increased by the active form of vitamin D₃ 1,25(OH)₂VD₃ at all concentrations examined, compared with the cells treated with the vehicle. This increase in cellular GGT activity was associated with a significant increase in GGT secretion into the medium. Concentrations of 1,25(OH)₂VD₃ as low as 10 nM, were sufficient to stimulate GGT activity in the LLC-PK1 cells and to facilitate GGT secretion into the medium.

Time-dependent effect of 1,25(OH)₂VD₃ on GGT activity and mRNA expression in LLC-PK1 renal tubular cells. To further analyze the stimulation of GGT activity by 1,25(OH)₂VD₃, a time course for 1,25(OH)₂VD₃-induced GGT activity was performed. As shown in Fig. 3, 1,25(OH)₂VD₃ was observed to stimulate an increase in GGT activity in LLC-PK1 cells over a relatively long duration of time. Thus, the stimulatory effect of 1,25(OH)₂VD₃ was not transient, but appeared to be continuous, in contrast to the 1,25(OH)₂VD₃-induced alkaline phosphatase (ALP) activity reported in a previous study (24).




Figure 1. Stimulatory effect of 1,25(OH)₂VD₃ on GGT activity in LLC-PK1 cells. LLC-PK1 cells were cultured for three days in the presence of various derivatives of vitamin D₃. The vitamin D₃ derivatives were dissolved in ethanol as a vehicle and added to the cells at a final concentration of 100 μM. Ethanol concentrations were 1% (v/v) in all cultures except the control. GGT activities in (A) cell homogenates and (B) cell media were analyzed. Data are presented as the mean ± standard deviation (n=3/group). GGT, γ-glutamyl transpeptidase; 1,25(OH)₂VD₃, 1α,25-dihydroxyvitamin D₃.




Figure 2. Dose-dependent effect of 1,25(OH)₂VD₃ on GGT activity in LLC-PK1 cells. LLC-PK1 cells were cultured in the presence of various concentrations of 1,25(OH)₂VD₃ and GGT activity was assessed in the (A) cell homogenates and (B) cell media. *P<0.01 vs. vehicle. GGT, γ-glutamyl transpeptidase; 1,25(OH)₂VD₃, 1α,25-dihydroxyvitamin D₃.

Furthermore, PCR was used to analyze GGT mRNA expression in the LLC-PK1 cells, 24, 48 and 72 h after the addition of 1,25(OH)₂VD₃. Fig. 4 shows that after 72 h of culture with 1,25(OH)₂VD₃, no significant increase in GGT mRNA expression was observed compared with the control and vehicle cells. The same findings were observed for the cells that had been treated with 1,25(OH)₂VD₃ for 24 and 48 h (data not shown). Thus, it is likely that the increase in GGT activity caused by 1,25(OH)₂VD₃ is due to a prolonged GGT protein turnover rather than enhanced GGT biosynthesis.

Discussion

GGT expression, which is associated with carcinogenesis in the liver and other tissues, has been extensively investigated (1).

By contrast, alterations in GGT activity in the kidney have not been investigated, which is likely to be due to the fact that GGT expression is relatively high and stable in the kidney. It has been reported that urinary GGT may be a potential marker for enhanced bone resorption (14), but little is known about the biochemical basis of the mechanism involved in the stimulation of renal GGT activity and how GGT is secreted into the urine. Therefore, the present study aimed to investigate the effect of factors involved in bone metabolism on GGT activity in renal epithelial cells.

ALP is also expressed in the proximal tubules in the kidney (25,26). Do Thanh *et al* (24) reported that 1,25(OH)₂VD₃ causes a rapid and transient increase in ALP activity in LLC-PK1 cells, with an almost two-fold increase observed 6 h after the addition of 1,25(OH)₂VD₃. However,




Figure 3. Time course of the effect of 1,25(OH)₂VD₃ on GGT activity in LLC-PK1 cells. LLC-PK1 cells were cultured in the presence of 100 μ M 1,25(OH)₂VD₃ for the indicated durations and GGT activity was assessed in the (A) cell homogenates and (B) cell media. Open circles, control; closed circles, 1,25(OH)₂VD₃; and open triangles, vehicle. Data are presented as the mean \pm standard deviation (n=3/group). *P<0.01 vs. vehicle. GGT, γ -glutamyl transpeptidase; 1,25(OH)₂VD₃, 1 α ,25-dihydroxyvitamin D₃.

no significant increase was observed for GGT activity in the same time course (24). The findings of the present study contradict the lack of GGT activity induced by 1,25(OH)₂VD₃ reported by Do Thanh *et al.* (24). In the present study, GGT activity was found to be stimulated by long-term exposure to 1,25(OH)₂VD₃. However, the time courses for the enhanced activities induced by 1,25(OH)₂VD₃ vary between ALP and GGT, suggesting that GGT activity is stimulated by a different mechanism to ALP activity. Although both GGT and ALP are expressed in the proximal tubules of the kidney, their distributions differ (27). Thus, the tubules may be partitioned into four segments according to the distribution of ALP and GGT.

A similar effect of 1,25(OH)₂VD₃ on GGT activity has been reported in the rat brain (28). In rat astrocytes, 1,25(OH)₂VD₃ was found to increase GGT activity alone or through potentiating the stimulatory effect of lipopolysaccharides (29). The increased activity of GGT observed in rat astrocytes facilitated glutathione synthesis by supplying constituent amino acids. Thus, increases in GGT activity may have a role in astrocyte detoxification pathways against oxidative stress. Extracellular




Figure 4. PCR analysis of GGT expression in LLC-PK1 cells. The 495 bp DNA fragments for the coding region of porcine GGT were amplified using PCR with specific primers. The left lane is a DNA size marker. PCR, polymerase chain reaction; GGT, γ -glutamyl transpeptidase; 1,25(OH)₂D₃, 1 α ,25-dihydroxyvitamin D₃.

glutathione is degraded by a series of reactions that are initiated by the hydrolysis of the γ -glutamyl group of glutathione by GGT. The resultant free amino acids are recovered by cells and utilized for the resynthesis of glutathione (1-3). In rat astrocytes, it is likely that GGT activity was stimulated by 1,25(OH)₂VD₃ either in a short- or long-term manner (28-29).

A previous study analyzed the biomarker potential of urinary GGT using a bone resorption mouse model and revealed that intravenous administration of PTH significantly increased urinary GGT activity (14). This *in vivo* effect of PTH may result from the direct induction of GGT in renal tubules; however, such an effect was not observed in the present study due to the lack of PTH receptor in LLC-PK1 cells (21,22). At present, whether the effect of PTH on GGT activity depends on a mechanism that involves tissues or organs other than the proximal tubular cells has yet to be elucidated. In addition, the specific origin of urinary GGT is unknown, but it is possible that some urinary GGT is derived from circulating GGT in the blood.

The findings of the present study suggested that 1,25(OH)₂VD₃ stimulates GGT activity in renal proximal tubular cells. This increased activity may, at least in part, modulate glutathione metabolism in the kidney. Furthermore, in the present study, the 1,25(OH)₂VD₃-induced increases in cellular GGT were associated with increases in GGT secretion into the cell medium. The findings of the present study and those of previous reports suggest that urinary or renal GGT activity may be responsive to the status of bone metabolism. Further investigations are required to provide further evidence for the use of GGT as a biomarker, which may contribute to the diagnosis and monitoring of certain bone diseases, including osteoporosis and abnormal bone resorption.

Acknowledgements

The authors would like to thank Dr Shumpei Niida for the advice. This study was supported by a Grant-in-Aid for Comprehensive Research on Aging and Health from the Ministry of Health, Labour and Welfare of Japan.

References

- Ikeda Y and Taniguchi N: Gene expression of gamma-glutamyl-transpeptidase. *Methods Enzymol* 401: 408-425, 2005.
- Taniguchi N and Ikeda Y: gamma-Glutamyl transpeptidase: catalytic mechanism and gene expression. *Adv Enzymol Relat Areas Mol Biol* 72: 239-278, 1998.
- Meister A and Tate SS: Glutathione and related gamma-glutamyl compounds: biosynthesis and utilization. *Annu Rev Biochem* 45: 559-604, 1976.
- Meister A, Tate SS and Griffith OW: Gamma-glutamyl transpeptidase. *Methods Enzymol* 77: 237-253, 1981.
- Lieberman MW, Wiseman AL, Shi ZZ, Carter BZ, Barrios R, Ou CN, Chévez-Barrios P, Wang Y, Habib GM, Goodman JC, Huang SL, Lebovitz RM and Matzuk MM: Growth retardation and cysteine deficiency in gamma-glutamyl transpeptidase-deficient mice. *Proc Natl Acad Sci USA* 93: 7923-7926, 1996.
- Iannaccone PM and Koizumi J: Pattern and rate of disappearance of gamma-glutamyl transpeptidase activity in fetal and neonatal rat liver. *J Histochem Cytochem* 31: 1312-1316, 1983.
- Teschke R and Petrides AS: Hepatic gamma-glutamyltransferase activity: its increase following chronic alcohol consumption and the role of carbohydrates. *Biochem Pharmacol* 31: 3751-3756, 1982.
- Barouki R, Chobert MN, Finidori J, Aggerbeck M, Nalpas B and Hanoune J: Ethanol effects in a rat hepatoma cell line: induction of gamma-glutamyltransferase. *Hepatology* 3: 323-329, 1983.
- Solt DB, Medline A and Farber E: Rapid emergence of carcinogen-induced hyperplastic lesions in a new model for the sequential analysis of liver carcinogenesis. *Am J Pathol* 88: 595-618, 1977.
- Farber E: Cellular biochemistry of the stepwise development of cancer with chemicals: G. H. A. Clowes memorial lecture. *Cancer Res* 44: 5463-5474, 1984.
- Niida S, Kawahara M, Ishizuka Y, Ikeda Y, Kondo T, Hibi T, Suzuki Y, Ikeda K and Taniguchi N: Gamma-Glutamyltranspeptidase stimulates receptor activator of nuclear factor-kappaB ligand expression independent of its enzymatic activity and serves as a pathological bone-resorbing factor. *J Biol Chem* 279: 5752-5756, 2004.
- Ishizuka Y, Moriwaki S, Kawahara-Hanaoka M, Uemura Y, Serizawa I, Miyauchi M, Shibata S, Kanaya T, Takata T, Taniguchi N and Niida S: Treatment with anti-gamma-glutamyl transpeptidase antibody attenuates osteolysis in collagen-induced arthritis mice. *J Bone Miner Res* 22: 1933-1942, 2007.
- Hiramatsu K, Asaba Y, Takeshita S, Nimura Y, Tatsumi S, Katagiri N, Niida S, Nakajima T, Tanaka S, Ito M, Karsenty G and Ikeda K: Overexpression of gamma-glutamyltransferase in transgenic mice accelerates bone resorption and causes osteoporosis. *Endocrinology* 148: 2708-2715, 2007.
- Asaba Y, Hiramatsu K, Matsui Y, Harada A, Nimura Y, Katagiri N, Kobayashi T, Takewaka T, Ito M, Niida S and Ikeda K: Urinary gamma-glutamyltransferase (GGT) as a potential marker of bone resorption. *Bone* 39: 1276-1282, 2006.
- Scherberich JE, Kleemann B and Mondorf W: Isolation of kidney brush border gamma-glutamyl transpeptidase from urine by specific antibody gel chromatography. *Clin Chim Acta* 93: 35-41, 1979.
- Shiozawa M, Yamashita S, Aiso S and Yasuda K: A monoclonal antibody against human kidney gamma-glutamyl transpeptidase: preparation, immunochemical, and immuno-histochemical characterization. *J Histochem Cytochem* 37: 1053-1061, 1989.
- Tate SS and Meister A: gamma-Glutamyl transpeptidase from kidney. *Methods Enzymol* 113: 400-419, 1985.
- Rabito CA, Kreisberg JJ and Wight D: Alkaline phosphatase and gamma-glutamyl transpeptidase as polarization markers during the organization of LLC-PK1 cells into an epithelial membrane. *J Biol Chem* 259: 574-582, 1984.
- Altman RA, Orr AV, Lagenaur CF, Curthoys NP and Hughey RP: Expression of rat renal gamma-glutamyltranspeptidase in LLC-PK1 cells as a model for apical targeting. *Biochemistry* 32: 3822-3828, 1993.
- Yoshida T, Yoshida N, Nakamura A, Monkawa T, Hayashi M and Saruta T: Cloning of porcine 25-hydroxyvitamin D3 1alpha-hydroxylase and its regulation by cAMP in LLC-PK1 cells. *J Am Soc Nephrol* 10: 963-970, 1999.
- Nakahama H, Kakihara M, Fukuhara Y, Ueda N, Orita Y and Kamada T: Parathyroid hormone enhances gentamicin uptake by opossum kidney cells but not by LLC-PK1 cells. *Nephron* 58: 85-89, 1991.
- Bringhurst FR, Juppner H, Guo J, Urena P, Potts JT Jr, Kronenberg HM, Abou-Samra AB and Segre GV: Cloned, stably expressed parathyroid hormone (PTH)/PTH-related peptide receptors activate multiple messenger signals and biological responses in LLC-PK1 kidney cells. *Endocrinology* 132: 2090-2098, 1993.
- Siegel N, Wongsurawat N and Armbrecht HJ: Parathyroid hormone stimulates dephosphorylation of the renoredoxin component of the 25-hydroxyvitamin D3-1alpha-hydroxylase from rat renal cortex. *J Biol Chem* 261: 16998-17003, 1986.
- Do Thanh X, Massicot F, Do B, Breget R, Nivet V, Durand D, Warnet JM, Claude JR and Clot JP: 1 alpha,25-dihydroxyvitamin D3 stimulated alkaline phosphatase activity in cultured pig kidney epithelial LLC-PK1 cells. *Acta Physiol Scand* 158: 107-111, 1996.
- Petitclerc C and Plante GE: Renal transport of phosphate: role of alkaline phosphatase. *Can J Physiol Pharmacol* 59: 311-323, 1981.
- Letellier M, Plante GE, Brière N and Petitclerc C: Participation of alkaline phosphatase in the active transport of phosphates in brush border membrane vesicles. *Biochem Biophys Res Commun* 108: 1394-1400, 1982.
- Brière N, Martel M, Plante G and Petitclerc C: Heterogeneous distribution of alkaline phosphatase and gamma-glutamyl transpeptidase in the mouse nephron. *Acta Histochem* 74: 103-108, 1984.
- Garcion E, Thanh XD, Bled F, Teissier E, Dehouck MP, Rigault F, Brachet P, Girault A, Torpier G and Darcy F: 1,25-Dihydroxyvitamin D3 regulates gamma 1 transpeptidase activity in rat brain. *Neurosci Lett* 216: 183-186, 1996.
- Garcion E, Sindji L, Leblondel G, Brachet P and Darcy F: 1,25-dihydroxyvitamin D3 regulates the synthesis of gamma-glutamyl transpeptidase and glutathione levels in rat primary astrocytes. *J Neurochem* 73: 859-866, 1999.