Expression of protein kinase C isoforms in cultured human Tenon's capsule fibroblast cells

PING MA¹, QIANYING GAO², ZHICHONG WANG² and KEMING YU^2

¹Department of Ophthalmology, Shandong Provincial Hospital Affiliated to Shandong University, Jinan, Shandong 250021; ²Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, Guangdong 510060, P.R. China

Received October 20, 2014; Accepted July 17, 2015

DOI: 10.3892/mmr.2015.4205

Abstract. Members of the protein kinase C (PKC) family are involved in physiological and pathophysiological processes, and exert an important role in signal transduction. The aim of the present study was to determine which of the 12 protein kinase C (PKC) isoforms (PKC_{α}, PKC_{β I}, PKC_{β II}, PKC_{γ}, PKC_{δ} , PKC_{ϵ} , PKC_{η} , PKC_{θ} , PKC_{μ} , PKC_{ζ} , PKC_{λ} and PKC_{λ} were expressed in vitro in cultured human Tenon's capsule fibroblasts (HTFs). HTFs from cell passages three to five were investigated for the presence of the 12 PKC isoforms at the cellular, mRNA and protein levels using laser scanning confocal microscopy (LSCM), reverse transcription-polymerase chain reaction (RT-PCR) and western blot analysis, respectively. These analyses yielded similar results for several of the PKC isoforms (PKC_{α}, PKC_{δ}, PKC_{ϵ}, PKC_{μ}, PKC_{μ}, PKC_{ξ}, PKC_{λ} and PKC_{λ} ; however, $PKC_{\beta II}$, $PKC_{\beta II}$, PKC_{γ} and PKC_{θ} were not expressed in the cultured HTFs. LSCM revealed that eight PKC isoforms, PKC_{α} , PKC_{δ} , PKC_{ε} , PKC_{μ} , PKC_{μ} , PKC_{ζ} , PKC_{λ} and PKC₁, appeared almost exclusively in the cytoplasm of the cells. Notably, PKC_{δ} was expressed particularly well in the cytoskeleton. The present study revealed that all 12 PKC isoforms were expressed and that eight of the isoforms were present in cultured HTFs. These results will be of value when determining specific roles for the PKC isoforms in HTF proliferation, which may provide a novel therapeutic target for bleb scarring in glaucoma filtering surgery.

Introduction

Glaucoma, characterized by progressive optic neuropathy, is the second most commonly occurring disease causing blind-

E-mail: yukeming66@126.com

ness worldwide. Glaucoma filtration surgery (GFS) provides the gold standard for the management of intraocular pressure (IOP), after medication and laser surgery have failed to control IOP adequately. However, scar tissue, which can form under the conjunctiva, obstructs aqueous flow and causes the filter to fail. Previous studies have demonstrated that human Tenon's capsule fibroblasts (HTFs) located in the incision area exert a major role in scar formation by promoting the proliferation, migration and synthesis of the extracellular matrix (1). Several antimetabolites, including mitomycin C and 5-fluorouracil, have been used to prevent post-operative ocular scar tissue formation. However, a disadvantage is that these antimetabolites are associated with marked side-effects, including hypotony, endophthalmitis, bleb leakage and loss of vision (2,3). Several alternative methods have been demonstrated to reduce scar formation through the inhibition of the proliferation of the HTFs (4,5).

Protein kinase C (PKC) comprises a family of protein isoforms, which occupy a central role in cellular processes, including proliferation, differentiation, mitosis and inflammation. To date, at least 12 isoforms of PKC have been cloned, and these are divided into three major groups: Classical PKCs $(PKC_{\alpha}, PKC_{\beta I}, PKC_{\beta II} \text{ and } PKC_{\gamma}), \text{ novel } PKCs (PKC_{\delta}, PKC_{\epsilon},$ PKC_{η} , PKC_{θ} and PKC_{μ}) and atypical PKCs (PKC_{ξ} , PKC_{λ} and PKC₁) (6). The differences in functionality among the specific PKC isoforms are predominantly due to their subcellular localization, activation or inhibition by different stimuli, and transcriptional regulation. It was reported that tranilast inhibits the proliferation and migration of HTFs in vitro, at least in part, by downregulating the expression of PKC (7,8). Alkylphosphocholines were identified as effective inhibitors of HTF proliferation and migration, and cell-mediated contraction of collagen gels at non-toxic concentrations. A previous study indicated that the mechanism of action appeared to involve the inhibition of the PKC pathway (9). It was therefore feasible that several of the PKC isoforms may exert a role in HTF proliferation, and the present study was focused on analyzing the expression of the 12 PKC isoforms in cultured HTFs with a view towards elucidating their role(s) in HTF proliferation.

Materials and methods

Culture of HTFs. Fresh human Tenon's capsule tissues (48 h post-mortem) from donors, were obtained from the Eye Bank

Correspondence to: Dr Keming Yu, Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, 54 South Xianlie Road, Guangzhou, Guangdong 510060, P.R. China

Key words: protein kinase C, fibroblast, scarring, glaucoma, filtering surgery

6026

of Zhongshan Ophthalmic Center (Guangzhou, China). The HTFs were cultured in 6-well plates (BD Biosciences, Lincoln Park, NJ, USA) with explants in Dulbecco's modified Eagle's medium/F12 nutrient mixture (Invitrogen Life Technologies, Grand Island, NY, USA), containing 5% FBS (Hyclone, Logan, UT, USA), 100 U/ml penicillin G (Gibco Life Technologies, Carlsbad, CA, USA), 100 mg/ml streptomycin sulfate (Gibco Life Technologies) and L-glutamate (Invitrogen Life Technologies), as previously described (10). HTFs from cell passage 3 to 5 were subsequently used in the experiments. The cells were lysed for total RNA extraction to assess the mRNA expression levels, or to extract proteins and assess the protein expression levels.

Immunofluorescence analysis. HTFs (1x10⁶/liter) grown on coverslips were fixed in 4% paraformaldehyde for 15 min at room temperature, prior to rinsing three times in phosphate buffered saline (PBS). The cell cultures were permeabilized with 0.2% Triton X-100 (Sigma-Aldrich, St. Louis, MO, USA) in PBS at room temperature for 10 min. Indirect immunostaining was performed, as described previously (11). The primary antibodies, rabbit anti-PKCa polyclonal antibody (1:200; cat. no. sc-208), rabbit anti-PKCγ polyclonal antibody (1:100; cat. no. sc-211) and mouse anti-PKCn monoclonal antibody (1:100; cat. no. sc-136036) were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA); mouse anti-PKCδ monoclonal antibody (1:100; cat. no. 610397), mouse anti-PKCe monoclonal antibody (1:200; cat. no. 610086), mouse anti-PKC0 monoclonal antibody (1:50; cat. no. 612734), mouse anti-PKC1 monoclonal antibody (1:50; cat. no. 610175) and mouse anti-PKCλ monoclonal antibody (1:50; cat. no. 610208) were purchased from BD Biosciences; rabbit anti-PKCBI polyclonal antibody (1:200; cat. no. p3078), rabbit anti-PKCβII polyclonal antibody (1:100; cat. no. p8371), rabbit anti-PKCζ polyclonal antibody (1:30; cat. no. SAB2104776) and rabbit anti-PKCµ polyclonal antibody (1:50; cat. no. SAB1306354) were from Sigma-Aldrich. The cells were incubated with the antibodies overnight in a solution of PBS at a temperature of 4°C. Following washing with PBS, Alexa Fluor 488-conjugated secondary antibodies [1:200 (cat. nos. A28175 and A27034); Invitrogen Life Technologies] were applied for 1 h and $1 \mu g/ml$ Hoechst 33342 (Sigma-Aldrich) was used for nuclear counterstaining. Either secondary antibody alone, without primary antibodies, or isoform immunoglobulin G (BD Biosciences) served as the negative control. The staining was imaged using a laser-scanning confocal microscope (LSCM510META; Zeiss, Thornwood, NY, USA). Each antibody was used in a minimum of three separate experiments.

Total RNA extraction and reverse transcription-polymerase chain reaction (RT-PCR). The total RNA was isolated from the cells using an RNeasy Mini kit (Qiagen, Valencia, CA, USA), according to the manufacturer's instructions. The RNA concentration was quantified spectrophotometry, using an ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA) prior to storage at -80°C. The RNA (5 μ g) was reverse-transcribed using the SuperScriptTM first-strand synthesis system, according to the manufacturer's instructions (Invitrogen Life Technologies). cDNAs (2 μ g) encoding the PKC isoform genes were amplified by PCR as follows: Denaturation for 30 sec, followed by annealing (56°C for PKC_a, PKC_{βII}, PKC_γ, PKC_e, PKC_η, PKC_θ, PKC₄, PKC₅ and PKC_β; 60°C for PKC₈; 52°C for PKC_{βI}) for 30 sec, and elongation at 72°C for 60 sec, for 30 cycles. The resulting PCR products were analyzed by 2% gel electrophoresis. Primer sequences for 11 PKC isoforms and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were published previously (12), and are listed in Table 1 (PKC λ cannot be detected due to the absence of its human PKC cDNA in GeneBank; http://www.ncbi.nlm.nih.gov/genbank/). Each PCR experiment was performed a minimum of three times with each set of primers.

Western blot analysis. The culture medium was removed and washed twice with ice-cold PBS. The HTFs were lysed using sample buffer [60 mM Tris/HCl (pH 6.8), 2% (w/v) sodium dodecyl sulfate (SDS), 100 mM 2-mercaptoethanol and 0.01% (w/v) Bromophenol Blue] (13). The lysates were incubated on ice for 30 min and the extracts were harvested using a cell scraper, and boiled for 5 min prior to storage at -20°C.

Subsequently, western blotting was performed, as previously described (14). Briefly, 40 μ g protein per well, measured using a bicinchoninic acid protein assay kit (Bio-Rad Laboratories, Inc., Hercules, CA, USA), was loaded onto a 12% SDS-polyacrylamide gel for SDS-polyacrylamide gel electrophoresis. The proteins were separated and electrotransferred onto polyvinylidene fluoride membranes (EMD Millipore, Bedford, MA, USA) for 1 h at 350 mA. The membranes were blocked with 5% non-fat milk disolved in TTBS buffer, containing, 50 mM Tris/HCl (pH 7.5), 0.9% NaCl and 0.1% Tween-20, for 1 h at room temperature and incubated with primary antibodies overnight at 4°C, prior to subsequent incubation with horseradish peroxidase-conjugated secondary antibody for 1 h at room temperature. The signals were detected using an enhanced chemiluminescence kit (GE Healthcare, Inc., Piscataway, NJ, USA), according to the manufacturer's instructions. Each PKC isoform was detected in a minimum of three independent experiments. Rat brain lysate (BD Biosciences) served as a positive control, as recommended by the manufacturer's instructions for the primary antibody.

Results

Immunofluorescence analysis of the PKC isoforms in HTFs. Using laser scanning confocal microscopy (LSCM), eight PKC isoforms (PKC_{α}, PKC_{δ}, PKC_{ϵ}, PKC_{ζ}, PKC_{η}, PKC_{ι}, PKC_{λ} and PKC_{μ}) were identified in the cultured HTFs, predominantly localized in the cytoplasm of the cells, as revealed by immunofluorescence staining. In particular, PKC_{δ} was expressed in the cytoskeleton. However, no staining was identified for the PKC_{β I}, PKC_{β II}, PKC_{γ} or PKC_{θ} isoforms in the HTFs (Fig. 1). Note that the primary antibody replaced with β -actin, and the primary antibody replaced with IgG isotype (positive and negative controls, respectively), are not shown in Fig. 1.

mRNA expression levels of the PKC isoforms in HTFs. The results from the 2% agarose gel electrophoresis experiment revealed the presence of mRNAs coding for seven of the

	1 1	1		r -	D	•		1			•	. •		1		1	•		. •		1	
	0	٦I	0		ν_r	imer	C 01	nd	reverce	trance	rin	tion	no	1.	marace	ch	10111	reac	tion	conc	11117	and
1	a	л		ι.	11	muci	s ai	IU.		uanse	11D	uon	-00	1 1	morase	UI.	ıam	ruau	uon	COIR	uu	JIIS.

Primer	Sequence $(5' \rightarrow 3')$	Amplicon (bp)
РКСа	Forward: ATCCGCAGTGGAATGAGTCCTTTACAT Reverse: TTGGAAGGTTGTTTCCTGTCTTCAGAG	327
РКСβІ	Forward: CTGTGGAACTGACTCCCACTG Reverse: ATACTGAAGCATTTTGGTATC	404
РКСβІІ	Forward: GACCGTTTTTCACCCGCCA Reverse: CCATCTCATAGAGATGCTCC	309
РКСү	Forward: CACGAAGTCAAGAGCCACAA Reverse: TAGCTATGCAGGCGGAACTT	233
РКСб	Forward: CAACTACATGAGCCCCACCT Reverse: GAGGCTCTCTGGGTGACTTG	189
ΡΚCε	Forward: GATGCAGAAGGTCACTGCAA Reverse: GTCGTCATGGAGGATGGACT	249
РКСҲ	Forward: GTTATCGATGGGATGGATGG Reverse: GCACCAGCTCTTTCTTCACC	166
РКСη	Forward: GAACAGAGGTTCGGGATCAA Reverse: ATATTTCCGGGTTGGAGACC	239
РКСө	Forward: ACAAACAGGGCTACCAGTGC Reverse: ATGCCACATGCATCACACTT	250
PKCı	Forward: TACGGCCAGGAGATACAACC Reverse: TCGGAGCTCCCAACAATATC	169
РКСµ	Forward: ACGGCACTATTGGAGATTGG Reverse: TGACCACATTTTCTCCCACA	206
GAPDH	Forward: ACCCAGAAGACTGTGGATGG Reverse: TGCTGTAGCCAAATTCGTTG	415

PKC, protein kinase C.



Figure 1. Immunofluorescence staining of the 12 PKC isoforms in HTFs using light-scanning confocal microscopy (magnification, x100). The nuclei of the HTFs were labeled with Hoechst 33342 and appear blue. A green signal indicates the positive staining of the PKC isoforms. PKC, protein kinase C; HTF, human Tenon's capsule fibroblasts. Scale bar, 50 μ m.



Figure 2. Reverse transcription-polymerase chain reaction analysis of 11 protein kinase C (PKC) isoforms in human Tenon's fibroblasts. GAPDH (415 bp) was used as a control. The molecular size markers are indicated on the left. GADPH, glyceraldehyde-3-phosphate dehydrogenase.



Figure 3. Western blot analysis of the 12 PKC isoforms in HTFs. Each gel was loaded with 40 μ g protein. Lanes 2-5 were HTF protein samples derived from independent cell lines. Lane 6 was a positive control. The markers are indicated on the left. β -actin (42 kDa) was used for protein normalization. PKC, protein kinase C; HTF, human Tenon's capsule fibroblast.

PKC isoforms. PKC_a (327 bp), PKC_b (189 bp), PKC_e (249 bp), PKC_q (239 bp), PKC_t (169 bp) and PKC₄ (206 bp) were present at a higher level compared with PKC₅ (166 bp), which only

produced a weak signal. $PKC_{\beta I}$, $PKC_{\beta II}$, PKC_{γ} and PKC_{θ} were not detected at the mRNA level (Fig. 2). GAPDH was used as a positive control.

Protein expression levels of the PKC isoforms in HTFs. An analysis of the protein expression levels of the PKC isoforms in HTFs was also performed. Using western blotting, seven PKC isoforms [PKC_α (80 kDa), PKC_δ (78 kDa), PKC_ε (90 kDa), PKC_ζ (78 kDa), PKC_η (82 kDa), PKC₁ (74 kDa) and PKC_μ (110 kDa)] were observed to be expressed in the HTFs, corroborating the results of the mRNA expression level analysis. In addition, PKC_λ (74 kDa) was also expressed in the HTFs. However, the other four isoforms, PKC_{βI}, PKC_{βII}, PKC_γ and PKC_η, were not detected (Fig. 3). Lanes 2-5 feature HTF protein lysates derived from independent cell lines. Rat brain lysate was used as a positive control (lane 6) and β-actin (42 kDa) was used for protein normalization.

Discussion

Scarring is the predominant reason for the failure of GFS. Successful filtration surgery depends directly on an individual's wound-healing response. HTFs are crucially important in this process. Previous studies have revealed that subconjunctival scarring of the filtering bleb site is predominantly mediated by HTF proliferation, migration and contraction (15,16). In order to assess the role of PKCs in the biological function of HTFs, the specific expression of the 12 PKC isoforms were characterized. The present study assessed for the first time, to the best of our knowledge, the expression levels of the 12 PKC isoforms in HTFs at the cellular, mRNA and protein level, using LSCM, RT-PCR and western blotting, respectively.

A similar expression pattern of the PKC isoforms was identified by each of the three methods. Eight of the PKC isoforms, PKC_a, PKC_b, PKC_c, PKC_c, PKC_u, PKC_a and PKC_µ, were expressed in cultured HTFs; however, no expression was observed for the four other isoforms, PKC_{βI}, PKC_{βII}, PKC_γ and PKC_θ. With the exception of PKC_b, the protein expression levels of the other 11 isoforms were consistent with their gene expression levels. A previous report revealed that 8 of 10 of the PKC isoforms were expressed in rat subconjunctival fibroblasts, as determined using western blot analysis, including PKC_α, PKC_β, PKC_γ, PKC_b, PKC_c, PKC_γ, PKC₁ and PKC_λ. PKC_ζ was not examined (17). The differences identified between the two studies are likely to be species-specific.

It has been suggested that isoform-specific functions may be conferred by the subcellular localization of the PKCs. Using immunofluorescence staining, the present study revealed that the subcellular localization of the PKC isoforms occurs predominantly in the cytoplasm of the cells. In addition, PKC_{δ} was localized specifically to the cytoskeleton, which may be associated with its function in HTFs.

Although the PKC isoforms exhibit very few differences in terms of their structures, substrate preferences, expression and localization, individual PKC isoforms still appear to be tissue- and cell-specific. Epidermal growth factor activates the PKC_{α} phosphorylation pathway to stimulate goblet cell proliferation (18). Hepatocyte growth factor induces the migration of retinal pigment epithelial cells by enhancing the activation of PKC_{δ} and the phosphorylation of ERK (19). However, the majority of studies to date have focused on the activation of PKC in general (17). Therefore it is necessary to elucidate which PKC isoforms exert specific roles in HTFs proliferation. In conclusion, the present study demonstrated that 8 of the 12 PKC isoforms were expressed in HTFs at both the protein and the mRNA level. This represents an important step in understanding their precise physiological role, and how they are regulated during the process of proliferation. To elucidate whether one or several of the isoforms are involved in HTF proliferation, due to their role in cellular signal transduction of scarring post-GFS, remains to be elucidated by further experiments.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (no. 81100643).

References

- 1. Occleston NL, Daniels JT, Tarnuzzer RW, Sethi KK, Alexander RA, Bhattacharya SS, Schultz GS and Khaw PT: Single exposures to antiproliferatives: Long-term effects on ocular fibroblast wound-healing behavior. Invest Ophthalmol Vis Sci 38: 1998-2007, 1997.
- Anand N, Arora S and Clowes M: Mitomycin C augmented glaucoma surgery: Evolution of filtering bleb avascularity, transconjunctival oozing, and leaks. Br J Ophthalmol 90: 175-180, 2006.
- Beckers HJ, Kinders KC and Webers CA: Five-year results of trabeculectomy with mitomycin C. Graefes Arch Clin Exp Ophthalmol 241: 106-110, 2003.
- Meyer-Ter-Vehn T, Katzenberger B, Han H, Grehn F and Schlunck G: Lovastatin inhibits TGF-β-induced myofibroblast transdifferentiation in human tenon fibroblasts. Invest Ophthalmol Vis Sci 49: 3955-3960, 2008.
- Xiao YQ, Liu K, Shen JF, Xu GT and Ye W: SB-431542 inhibition of scar formation after filtration surgery and its potential mechanism. Invest Ophthalmol Vis Sci 50: 1698-1706, 2009.
- 6. Johannes FJ, Prestle J, Eis S, Oberhagemann P and Pfizenmaier K: PKCu is a novel, atypical member of the protein kinase C family. J Biol Chem 269: 6140-6148, 1994.
- Ji CN, Hu YZ, Ding ZP and Li GG: The investigation of tranilast on the proliferation and migration of human Tenon's capsule fibroblasts. Zhonghua Yan Ke Za Zhi 40: 165-169, 2004. (In Chinese).
- Chihara E, Dong J, Ochiai H and Hamada S: Effects of tranilast on filtering blebs: A pilot study. J Glaucoma 11: 127-133, 2002.
- Eibl KH, Banas B, Kook D, Ohlmann AV, Priglinger S, Kampik A and Welge-Luessen UC: Alkylphosphocholines: A new therapeutic option in glaucoma filtration surgery. Invest Ophthalmol Vis Sci. 45: 2619-2624, 2004.
- 10. Huang SS, Ge J, Wang LN, Yin XB, Wei YT and Ma P: Preliminary study of inhibition of human Tenon's capsule fibroblasts *in vitro* by RNA interference targeting IKK-beta. Zhonghua Yan Ke Za Zhi 141: 1076-1081, 2005. (In Chinese).
- Ma P, Wang Z, Pflugfelder SC and Li DQ: Toll-like receptors mediate induction of peptidoglycan recognition proteins in human corneal epithelial cells. Exp Eye Res 90: 130-136, 2010.
- Yu K, Ma P, Ge J, Willey CD, Yang P, Wang Z and Gao Q: Expression of protein kinase C isoforms in cultured human retinal pigment epithelial cells. Graefes Arch Clin Exp Ophthalmol 245: 993-999, 2007.
- Campa MJ, Wang MZ, Howard B, Fitzgerald MC and Patz EF Jr: Protein expression profiling identifies macrophage migration inhibitory factor and cyclophilin a as potential molecular targets in non-small cell lung cancer. Cancer Res 63: 1652-1656, 2003.
- 14. Ma P, Bian F, Wang Z, Zheng X, Chotikavanich S, Pflugfelder SC and Li DQ: Human corneal epithelium-derived thymic stromal lymphopoietin links the innate and adaptive immune responses via TLRs and Th2 cytokines. Invest Ophthalmol Vis Sci 50: 2702-2709, 2009.
- 15. Van Bergen T, Vandewalle E, Van de Veire S, Dewerchin M, Stassen JM, Moons L and Stalmans I: The role of different VEGF isoforms in scar formation after glaucoma filtration surgery. Exp Eye Res 93: 689-699, 2011.

6030

- 16. Ye H, Qian Y, Lin M, Duan Y, Sun X, Zhuo Y and Ge J: Cationic nano-copolymers mediated IKKβ targeting siRNA to modulate wound healing in a monkey model of glaucoma filtration surgery. Mol Vis 16: 2502-2510, 2010.
- Nomura N, Nomura M, Takahira M and Sugiyama K: Phorbol 12-myristate 13-acetate-activated protein kinase C increased migratory activity of subconjunctival fibroblasts via stress-activated protein kinase pathways. Mol Vis 13: 2320-2327, 2007.
- Li D, Shatos MA, Hodges RR and Dartt DA: Role of PKCα activation of Src, PI-3K/AKT, and ERK in EGF-stimulated proliferation of rat and human conjunctival goblet cells. Invest Ophthalmol Vis Sci. 54: 5661-5674, 2013.
 Chen YJ, Tsai RK, Wu WC, He MS, Kao YH and Wu WS:
- 19. Chen YJ, Tsai RK, Wu WC, He MS, Kao YH and Wu WS: Enhanced PKCδ and ERK signaling mediate cell migration of retinal pigment epithelial cells synergistically induced by HGF and EGF. PLoS One 7: e44937, 2012.