

A novel splice-site mutation of *WRN* (c.IVS28+2T>C) identified in a consanguineous family with Werner Syndrome

PAN-FENG WU^{1,2*}, JIE-YUAN JIN^{2*}, JING-JING LI², JI-QIANG HE¹, LIANG-LIANG FAN², MIN JIN¹, HAO HUANG², KUN XIA², JU-YU TANG¹ and RONG XIANG^{1,2}

¹Department of Orthopedics, Xiangya Hospital of Central South University, Changsha, Hunan 410078;

²School of Life Sciences, The State Key Laboratory of Medical Genetics, Central South University, Changsha, Hunan 410013, P.R. China

Received January 18, 2016; Accepted January 16, 2017

DOI: 10.3892/mmr.2017.6465

Abstract. Werner Syndrome (WS) is a rare, adult-onset progeroid syndrome that is associated with multiple age-associated complications and relatively short life expectancy. The characteristics of WS include a 'bird-like' appearance, canities, cataracts and ulcerations around the ankles. In addition, certain patients develop hypogonadism with atrophic genitalia and infertility. The average life span of affected individuals is 54 years. Previous studies have demonstrated that mutations in the Werner syndrome RecQ like helicase gene (*WRN*) may contribute to WS. The present study investigated a consanguineous family with WS, comprising of 4 generations from Northwest China (Gansu province). A novel homozygous splice-site mutation in *WRN* (c.IVS28+2T>C) was identified in this family and was predicted to be deleterious. No further relevant mutations were identified by direct sequencing of the genes lamin A/C, barrier to autointegration factor 1, zinc metalloproteinase STE24 and DNA polymerase $\Delta 1$. cDNA sequencing and alignments were performed to further confirm the pathogenicity of this mutation. The results support the important role of *WRN* in WS and expand the spectrum of known *WRN* mutations. In addition, it may provide novel approaches in genetic diagnosis and counseling of families with WS.

Introduction

Werner Syndrome (WS; OMIM entry no. 277700; <https://omim.org/entry/277700>) is a rare progeroid syndrome associated with a number of aging phenotypes. WS often occurs in consanguineous families and affects ~1 in 1 million individuals in the general population (1-3). Patients with WS often exhibit adult-onset progeria and have an increased risk of developing cancer (4). At birth, patients with WS do not present any clinical symptoms; the lack of pubertal growth spurts are generally the first symptom identified (5). Over time, the typical characteristics of WS become evident, including a 'bird-like' appearance, cataracts, canities, ulcerations around the ankles and certain patients develop hypogonadism with atrophic genitalia and infertility (6). The average life span of affected individuals is 54 years (5).

WS is caused by mutations in the Werner syndrome RecQ like helicase gene (*WRN*), which was first cloned in 1996 (7). *WRN* is located on chromosome 8p11-p12, spanning ~250 kb and consists of 35 exons, 34 of which are protein coding (7). *WRN* is generally considered to follow an autosomal-recessive pattern of inheritance (8). *WRN*, coding a 180 kDa multifunctional nuclear protein, belongs to the family of RecQ type helicases (9). Sequence analysis and subsequent biochemical analysis have revealed that human *WRN* possesses helicase and exonuclease functions (10). It serves a role in DNA replication, transcription, repair, recombination and heterochromatin maintenance (including telomere maintenance), indicating that one of the major causes of WS pathogenesis may be associated with genomic instability (11). In the absence of functioning *WRN*, cells accumulate potentially toxic DNA intermediates or critically short telomeres, which induce genetic instability, misexpression and mutagenesis (12-14). In addition, they may drive cell loss and produce tissue-specific defects (15). Compromised cell or tissue structure and function leads to two seemingly divergent outcomes: Senescence and neoplasms (16,17). To date, the majority of disease-inducing mutations in *WRN* are truncating mutations (5).

In addition to *WRN*, a small number of heterozygous mutations in the lamin A/C gene (*LMNA*) have been revealed to produce similar phenotypes to WS, which suggests that *LMNA* may be an underlying disease-inducing gene in

Correspondence to: Professor Rong Xiang, School of Life Sciences, The State Key Laboratory of Medical Genetics, Central South University, 172 Tongzipo Road, Changsha, Hunan 410013, P.R. China
E-mail: shirlesmile@csu.edu.cn

*Contributed equally

Key words: Werner syndrome, Werner syndrome RecQ like helicase, splice-site mutation, consanguineous family

WS (3,18,19). *LMNA* encodes nuclear intermediate filaments, lamin A and lamin C (1). A previous study demonstrated that WS patients with *LMNA* mutations exhibited a younger onset when compared to patients with classical WS (5). In addition, a number of studies have suggested that barrier to autointegration factor 1 (*BANF1*), zinc metalloproteinase STE24 (*ZMPSTE24*) and DNA polymerase $\Delta 1$ (*POLD1*) may be involved in progeroid syndrome (15,20,21).

The present study investigated the potential causative gene in a consanguineous family with WS from Northwest China. A novel homozygous splice-site mutation (c.IVS28+2T>C) was identified in intron 28 of *WRN* in the proband and co-segregated with the affected WS family members. To the best of our knowledge, this mutation has not been reported in previous studies, nor was it identified in our previous control cohorts or the single nucleotide polymorphism (dbSNP) database (<https://www.ncbi.nlm.nih.gov/projects/SNP/>) and the Exome Variant Server database (<http://evs.gs.washington.edu/EVS/>).

Materials and methods

Patients. A consanguineous family from Northwest China (Gansu province) consisting of 11 living members across four generations participated in the present study (Fig. 1A). The proband, family member 2 from the 4th generation (IV:2), was diagnosed with WS. The remaining 10 members (II:2, II:4, III:1, III:2, III:3, III:4, IV:1, IV:3, IV:4 and V1) were phenotypically normal. The proband was admitted to Xiangya Hospital in April 2015 (Changsha, China) for treatment of an ankle ulcer. The 38-year-old patient (gender, male) had a 'bird-like' face, gray hair, a husky voice and a recurrent ulceration around his ankle which first presented 12 years previously (Fig. 1B). The Review Board of Xiangya Hospital of the Central South University (Hunan, China) approved this research and all family members involved gave written informed consent.

DNA extraction. Genomic DNA was extracted from the peripheral blood of the patient and the other family members using a DNeasy Blood & Tissue kit (Qiagen, Inc., Valencia, CA, USA) on the QIAcube automated DNA extraction robot (Qiagen, Inc.).

Mutation sequencing. The entire coding regions, including the flanking intronic sequences of *WRN* [Refseq (<https://www.ncbi.nlm.nih.gov/refseq/>), NM_000553], *LMNA* (NM_170,707), *BANF1* (NM_003860), *ZMPSTE24* (NM_005857) and *POLD1* (NM_001256849) were amplified by polymerase chain reaction (PCR; primer sequences are available upon request). PCR product sequences were determined using the ABI 3100 Genetic Analyzer (Thermo Fisher Scientific, Inc., Waltham, MA, USA) as previously described (22).

Multiple sequence alignments and bioinformatic prediction of mutation. The multiple *WRN* protein sequences across mammals were aligned using the multiple sequence comparison by log-expectation program (version 3.6; <https://www.ncbi.nlm.nih.gov>, and the MUSCLE software) (23,24).

RNA extraction and reverse transcription for verification. Total RNA was extracted from mononuclear cells from the

peripheral blood of the patient using the Nucleospin RNA II kit (Macherey-Nagel GmbH, Düren, Germany) and DNase (DNase I, RNase-free (1 U/ μ l); Thermo Fisher Scientific, Inc.) treated (25). Reverse transcription (RT)-qPCR was performed to convert extracted total RNA into cDNA using the PrimeScript RT reagent kit (Takara Bio, Inc., Otsu, Japan) according to the manufacturer's instructions (26). cDNA was then sequenced following amplification by PCR. The PCR was conducted in a 25 μ l reaction mixture, which consisted of 0.3 mM deoxyribonucleotide triphosphates, 1X PCR buffer (10 mM Tris-HCl pH 9.0, 50 mM KCl, 0.1% Triton X-100, and 0.01% w/v gelatin), 2.0 mM MgCl₂, 0.5 μ M of each primer (forward and reverse), 1.5 U of Taq polymerase, and 50 ng genomic DNA. The thermal cycling consisted of an initial denaturation at 95°C for 4 min, followed by 35 cycles of amplification consisting of denaturation at 95°C for 1 min, primer annealing at 55–61°C for 30 sec and primer extension at 72°C for 1 min. A final extension step was performed at 72°C for 7 min. The results were compared to the normal control for variant analysis via multiple sequence alignment using the MUSCLE software (24).

Results

The present study investigated the case of a patient with WS (IV:2) born to consanguineous parents (family members III:2 and III:3) in a family comprised of four generations with 11 living members. The proband, with a 'bird-like' face, husky voice, canities and ankle ulcers, conforms to the typical phenotypes associated with WS. The present study investigated the potential causative genes among all family members. Sequence analysis of *WRN*, *LMNA*, *BANF1*, *ZMPSTE24* and *POLD1*, identified a previously unreported homozygous splice-site mutation in intron 28 (c.IVS28+2T>C) of the *WRN* gene in the proband (IV:2) and co-segregated with the affected family members (Fig. 1C). No further relevant mutations were identified by direct sequencing of the genes for *LMNA*, *BANF1*, *ZMPSTE24* and *POLD1*. In addition, the cDNA sequencing results from the proband identified deletion of *WRN* exon 28 (74 nucleotides; Fig. 1D). This newly discovered c.IVS28+2T>C mutation was not identified in the 200 control cohorts that our group studied previously (23). In addition, this mutation was not present in the dbSNP (<https://www.ncbi.nlm.nih.gov/projects/SNP/>) or Exome Variant Server databases (<http://evs.gs.washington.edu/EVS/>).

Discussion

The *WRN* gene encodes a nuclear protein which belongs to the family of RecQ type helicases and comprises of five functional domains including an exonuclease region, a helicase region, a RecQ C-terminus consensus region, an RNase D consensus region and a nuclear localization signal region (NLS) (27). According to previous studies, homology-dependent recombination repair (HDR) may be used to repair DNA damage while suppressing gene loss or rearrangement. In addition, *WRN* appears to serve a role late in the HDR process when recombinant molecules are topologically disentangled for segregation to daughter cells (28,29), as well as in the maintenance of telomere length and the suppression




Figure 1. (A) Ancestry of the family affected with WS. Family members are identified by their generation (indicated by roman numerals) and a number. Squares represent male family members and circles, female members; the triangle represents a fetus. The black symbol represents a member with WS, the white symbols represent unaffected members and the half black-half white symbols represent carriers. The arrow indicates the proband. (B) Phenotypes of the proband. The proband has a 'bird-like' face, canities, a husky voice and a recurrent ulceration around the ankle. (C) Sequencing results of the *WRN* mutation. Sequence chromatograms indicate a homozygous splice-site mutation (c.IVS28+2T>C) in the proband. (D) The reverse sequencing results of the cDNA with the *WRN* mutation. The sequence chromatogram reveals a deletion of exon 28 following a homozygous splice-site mutation (c.IVS28+2T>C). The green rectangular box represents exon 29 and the corresponding sequence and the blue rectangular box represents exon 27 and corresponding sequence. The red letters indicate deleted nucleotides in exon 28 and the red 'Y' shape indicates the normal location of exon 28. WS, Werner syndrome; *WRN*, Werner syndrome RecQ like helicase.




Figure 2. *WRN* mutations identified in Werner syndrome patients. The rectangular box represents the *WRN* protein with the N-terminus on the left and C-terminus on the right. Known functional domains include an exonuclease region, a helicase region, a RQC region, a HRDC region and a NLS. Mutations are grouped based on the mutation site. The red words represent the previous splice-site mutation. *Indicates a termination codon. *WRN*, Werner syndrome RecQ like helicase; RQC, RecQ C-terminus consensus region; HRDC, RNase D consensus region; NLS, nuclear localization signal.

of telomere sister-chromatid exchanges (30,31). *WRN* may also be involved in non-homologous DNA-end joining, base-excision repair, DNA-damage signaling and transcription (15). Zhang *et al* (11) revealed that the progressive heterochromatin disorganization observed in *WRN*-deficient mesenchymal stem cells underlies cellular aging (32). Thus, loss of *WRN* may disrupt genetic stability, lead to cell aging and death. Therefore, it may be pivotal in human aging and the development of WS.

The present study revealed that the novel mutation (c.IVS28+2T>C) causes a change in the splicing pattern, leading to a deletion of 74 nucleotides in the mRNA. It suggested that the substitution and subsequent skipping of exon 28 may produce a frameshift transcript, resulting in the absence of the NLS domain. The NLS domain is essential for *WRN* protein targeting to the nucleus via the nuclear pore complex (33). In addition, the majority of previously reported

causative WRN mutations also give rise to a lack of NLS at the C-terminus of the protein (Fig. 2) (5,34).

During this research, the wife of the proband (IV:1) was pregnant. It was speculated that the infant (V:2) may be a carrier without any pathological phenotype. However, amniocentesis or shotgun sequencing of maternal plasma DNA is required to produce accurate and convincing results to verify this. As cancer predisposition is a key feature in WS, the proband may have a higher risk of cancer development and should therefore have regular health examinations.

In conclusion, the present study identified a novel homozygous splice-site mutation (c.IVS28+2T>C) in a four generation family with WS. The present identification of a novel mutation expands the spectrum of known WRN mutations (only 87 mutations were identified as of February 2015; <http://www.hgmd.cf.ac.uk/ac/search.php>) and it may contribute to novel approaches to genetic diagnosis and counseling of families with WS.

Acknowledgements

The authors would like to thank the State Key Laboratory of Medical Genetics of China (Hunan, China) for their technical assistance. The present study was supported by the National Natural Science Foundation of China (Beijing, China; grant no. 81370394) and the National Basic Research Program of China (973 Program; Beijing, China; grant no. 2012CB517900).

References

- Masala MV, Scapaticci S, Olivieri C, Pirodda C, Montesu MA, Cuccuru MA, Pruneddu S, Danesino C and Cerimele D: Epidemiology and clinical aspects of Werner's syndrome in North Sardinia: Description of a cluster. *Eur J Dermatol* 17: 213-216, 2007.
- Hasty P, Campisi J, Hoeijmakers J, van Steeg H and Vijg J: Aging and genome maintenance: Lessons from the mouse? *Science* 299: 1355-1359, 2003.
- Chen L, Lee L, Kudlow BA, Dos Santos HG, Sletvold O, Shafeghati Y, Botha EG, Garg A, Hanson NB, Martin GM, *et al*: LMNA mutations in atypical Werner's syndrome. *Lancet* 362: 440-445, 2003.
- Muftuoglu M, Oshima J, von Kobbe C, Cheng WH, Leistriz DF and Bohr VA: The clinical characteristics of Werner syndrome: Molecular and biochemical diagnosis. *Hum Genet* 124: 369-377, 2008.
- Oshima J and Hisama FM: Search and insights into novel genetic alterations leading to classical and atypical Werner syndrome. *Gerontology* 60: 239-246, 2014.
- Friedrich K, Lee L, Leistriz DF, Nürnberg G, Saha B, Hisama FM, Eyman DK, Lessel D, Nürnberg P, Li C, *et al*: WRN mutations in Werner syndrome patients: Genomic rearrangements, unusual intronic mutations and ethnic-specific alterations. *Hum Genet* 128: 103-111, 2010.
- Yu CE, Oshima J, Fu YH, Wijsman EM, Hisama F, Alisch R, Matthews S, Nakura J, Miki T, Ouais S, *et al*: Positional cloning of the Werner's syndrome gene. *Science* 272: 258-262, 1996.
- Sinha JK, Ghosh S and Raghunath M: Progeria: A rare genetic premature ageing disorder. *Indian J Med Res* 139: 667-674, 2014.
- Choudhary S, Sommers JA and Brosh RM Jr: Biochemical and kinetic characterization of the DNA helicase and exonuclease activities of werner syndrome protein. *J Biol Chem* 279: 34603-34613, 2004.
- Orren DK, Theodore S and Machwe A: The Werner syndrome helicase/exonuclease (WRN) disrupts and degrades D-loops in vitro. *Biochemistry* 41: 13483-13488, 2002.
- Zhang W, Li J, Suzuki K, Qu J, Wang P, Zhou J, Liu X, Ren R, Xu X, Ocampo A, *et al*: Aging stem cells. A Werner syndrome stem cell model unveils heterochromatin alterations as a driver of human aging. *Science* 348: 1160-1163, 2015.
- Laud PR, Multani AS, Bailey SM, Wu L, Ma J, Kingsley C, Lebel M, Pathak S, De Pindo RA and Chang S: Elevated telomere-telomere recombination in WRN-deficient, telomere dysfunctional cells promotes escape from senescence and engagement of the ALT pathway. *Gen Dev* 19: 2560-2570, 2005.
- Eller MS, Liao X, Liu S, Hanna K, Bäckvall H, Oprea PL, Bohr VA and Gilchrist BA: A role for WRN in telomere-based DNA damage responses. *Proc Natl Acad Sci USA* 103: 15073-15078, 2006.
- Dhillon KK, Sidorova J, Saintigny Y, Poot M, Gollahon K, Rabinovitch PS and Monnat RJ Jr: Functional role of the Werner syndrome RecQ helicase in human fibroblasts. *Aging Cell* 6: 53-61, 2007.
- Kudlow BA, Kennedy BK and Monnat RJ Jr: Werner and Hutchinson-Gilford progeria syndromes: Mechanistic basis of human progeroid diseases. *Nat Rev Mol Cell Biol* 8: 394-404, 2007.
- Monnat RJ Jr and Saintigny Y: Werner syndrome protein-unwinding function to explain disease. *Sci Aging Knowledge Environ* 2004: re3, 2004.
- Kipling D, Davis T, Ostler EL and Faragher RG: What can progeroid syndromes tell us about human aging? *Science* 305: 1426-1431, 2004.
- Csoka AB, Cao H, Sammak PJ, Constantinescu D, Schatten GP and Hegele RA: Novel lamin A/C gene (LMNA) mutations in atypical progeroid syndromes. *J Med Genet* 41: 304-308, 2004.
- Renard D, Fourcade G, Milhau D, Bessis D, Esteves-Vieira V, Boyer A, Roll P, Bourgeois P, Levy N and De Sandre-Giovannoli A: Novel LMNA mutation in atypical Werner syndrome presenting with ischemic disease. *Stroke* 40: e11-e14, 2009.
- Lessel D, Hisama FM, Szakszon K, Saha B, Sanjuanelo AB, Salbert BA, Steele PD, Baldwin J, Brown WT, Piussan C, *et al*: POLD1 germline mutations in patients initially diagnosed with Werner syndrome. *Human mutation* 36: 1070-1079, 2015.
- Barcena C, Osorio FG and Freije JM: Detection of nuclear envelope alterations in senescence. *Methods Mol Biol* 965: 243-251, 2013.
- Tan ZP, Huang C, Xu ZB, Yang JF and Yang YF: Novel ZFPM2/FOG2 variants in patients with double outlet right ventricle. *Clin Genet* 82: 466-471, 2012.
- Xiang R, Fan LL, Huang H, Cao BB, Li XP, Peng DQ and Xia K: A novel mutation of GATA4 (K319E) is responsible for familial atrial septal defect and pulmonary valve stenosis. *Gene*: Oct 26, 2013 (Epub ahead of print).
- Edgar RC: MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32: 1792-1797, 2004.
- Kadari A, Mekala S, Wagner N, Malan D, Köth J, Doll K, Stappert L, Eckert D, Peitz M, Matthes J, *et al*: Robust generation of cardiomyocytes from human iPS cells requires precise modulation of BMP and WNT signaling. *Stem Cell Rev* 11: 560-569, 2015.
- Gotoh A, Hamada Y, Shiobara N, Kumagai K, Seto K, Horikawa T and Suzuki R: Skew in T cell receptor usage with polyclonal expansion in lesions of oral lichen planus without hepatitis C virus infection. *Clin Exp Immunol* 154: 192-201, 2008.
- Oshitari T, Kitahashi M, Mizuno S, Baba T, Kubota-Taniai M, Takemoto M, Yokote K, Yamamoto S and Roy S: Werner syndrome with refractory cystoid macular edema and immunohistochemical analysis of WRN proteins in human retinas. *BMC Ophthalmol* 14: 31, 2014.
- Prince PR, Emond MJ and Monnat RJ Jr: Loss of Werner syndrome protein function promotes aberrant mitotic recombination. *Genes Dev* 15: 933-938, 2001.
- Saintigny Y, Makienko K, Swanson C, Emond MJ and Monnat RJ Jr: Homologous recombination resolution defect in werner syndrome. *Mol Cell Biol* 22: 6971-6978, 2002.
- Chang S, Multani AS, Cabrera NG, Naylor ML, Laud P, Lombard D, Pathak S, Guarente L and DePinho RA: Essential role of limiting telomeres in the pathogenesis of Werner syndrome. *Nat Genet* 36: 877-882, 2004.
- Crabbe L, Verdun RE, Hagglblom CI and Karlseder J: Defective telomere lagging strand synthesis in cells lacking WRN helicase activity. *Science* 306: 1951-1953, 2004.
- Jones B: Ageing: Heterochromatin disorganization associated with premature ageing. *Nat Rev Genet* 16: 318, 2015.
- Floch AG, Tareste D, Fuchs PF, Chadrin A, Naciri I, Léger T, Schlenstedt G, Palancade B and Doye V: Nuclear pore targeting of the yeast Pom33 nucleoporin depends on karyopherin and lipid binding. *J Cell Sci* 128: 305-316, 2015.
- Oshima J, Yu CE, Piussan C, Klein G, Jabkowski J, Balci S, Miki T, Nakura J, Ogihara T, Ellis J, *et al*: Homozygous and compound heterozygous mutations at the Werner syndrome locus. *Hum Mol Genet* 5: 1909-1913, 1996.