# Altered gene expression of hepatic cytochrome P450 in a rat model of intermittent hypoxia with emphysema

HONGZHI YU<sup>1\*</sup>, HONGXIA SHAO<sup>1\*</sup>, QI WU<sup>1</sup>, XIN SUN<sup>2</sup>, LI LI<sup>1</sup>, KUAN LI<sup>3</sup>, XUE LI<sup>3</sup>, YU LI<sup>3</sup>, QIUYANG ZHANG<sup>3</sup>, JUNPING WU<sup>1</sup> and HUAIYONG CHEN<sup>3</sup>

<sup>1</sup>Department of Respiration; <sup>2</sup>Key Research Laboratory for Infectious Disease Prevention for State Administration of Traditional Chinese Medicine; <sup>3</sup>Department of Basic Medicine, Tianjin Institute of Respiratory Diseases, Tianjin Haihe Hospital, Tianjin Medical University, Tianjin 300350, P.R. China

Received May 10, 2016; Accepted March 28, 2017

DOI: 10.3892/mmr.2017.6642

Abstract. Patients with respiratory overlap syndrome (OS), defined as concomitant chronic obstructive pulmonary disease and obstructive sleep apnea syndrome, may exhibit an increased blood concentration of ingested drugs. This poor elimination of drugs is primarily attributed to downregulated gene expression of the drug-metabolizing cytochrome P450 enzymes (CYPs) in the liver. However, the underlying mechanisms of the decreased expression of CYPs in OS are poorly understood. In order to address this, a rat model of intermittent hypoxia with emphysema was evaluated in the present study, by analyzing liver gene expression using the reverse transcription-quantitative polymerase chain reaction. Intermittent hypoxia and cigarette smoke exposure caused upregulation of hepatic inflammatory cytokines, while CYPs were downregulated. This downregulation of CYPs was associated with an increase in nuclear factor (NF)-KB expression and a decrease in the expression of nuclear receptors pregnane X receptor, constitutive androstane receptor and glucocorticoid receptor, which are the upstream regulatory molecules of CYPs. The results of the present study indicated that, during the development of OS, systematic inflammatory reactions may downregulate hepatic CYP gene expression via the NF-κB signaling pathway.

E-mail: wq572004@163.com

#### \*Contributed equally

### Introduction

Chronic obstructive pulmonary disease (COPD) and obstructive sleep apnea syndrome (OSAS) represent two of the most prevalent chronic respiratory disorders in clinical practice. Respiratory overlap syndrome (OS), defined as the coexistence of these two diseases, occurs in ~1% of adults and leads to the development of increased nocturnal oxygen desaturation compared with mono-COPD or OSAS (1-3). The possible coexistence of COPD with OSAS is notable, as systemic inflammation develops in each disorder (3,4) and may contribute to the pathogenesis of associated comorbidities. Hypoxia induces systemic inflammatory reactions and acts as an aggravating factor of liver injury (5,6). It has been hypothesized that, in patients with mono-COPD or OSAS, inflammatory mediators generated in the lung may 'spill over' into the bloodstream and promote the release of inflammatory proteins from the liver, which may cause damage to target organs and act as a source of systemic inflammation (7,8).

Inflammation modulates the expression of hepatic metabolizing enzymes. Cytochrome p450 enzymes (CYPs) catalyze the oxidative metabolism of numerous drugs, and the level of CYP expression may affect drug efficacy, toxicity and consequently, therapeutic outcome (9). Previous studies demonstrated that the blood concentration of theophylline was above the normal range when the expression and activity of CYP1A2 was inhibited (10,11). A further previous study demonstrated that sleep hypoxia combined with emphysema synergistically enhanced hepatic inflammation and produced a more apparent liver-derived inflammatory state (12). It has been reported that inflammatory challenges may suppress the expression of major CYPs (13). Underexpression of the nuclear receptors pregnane X receptor (PXR) and constitutive androstane receptor (CAR), as upstream regulatory molecules, may inhibit the transcription of CYPs (11-13) by attenuating nuclear translocation (14,15). It is hypothesized that severe inflammation in the liver tissues of patients with OS alters hepatic metabolism via nuclear receptors.

In the present study, a previously-published rat model of OS was developed by exposing rats to intermittent hypoxia (IH) and cigarette smoke (CS). Gene expression analysis of liver samples with reverse transcription-quantitative polymerase

*Correspondence to:* Professor Qi Wu, Department of Respiration, Tianjin Institute of Respiratory Diseases, Tianjin Haihe Hospital, Tianjin Medical University, 890 Jingu Road, Jinnan, Tianjin 300350, P.R. China

*Key words:* intermittent hypoxia, emphysema, cytochrome P450, hepatic

chain reaction (RT-qPCR) was used to evaluate the effect of IH with CS on inflammatory cytokines and CYPs. The present study may provide a molecular mechanism to explain adverse drug reactions in patients with OS.

#### Materials and methods

*Ethics statement*. Rats were used in accordance with the protocol approved by the Animal Care Committee of Tianjin Medical University (permit no. 2010-0002).

Animals and treatments. Rats were provided by the Model Animal Center of the Radiological Medicine Research Institute, Chinese Academy of Medical Science (Beijing, China), and housed in standard laboratory cages (5 rats/cage) with food and water available ad libitum. As described previously (12), a total of 30 male Wistar rats weighing 180±20 g at age 6 weeks were divided into two groups of 15 according to exposure condition, as follows: i) Control group; ii) IH with CS experimental group. For IH, the rats were treated in a 120 sec cycle, comprising 30 sec nitrogen followed by 90 sec air, between 9:00 a.m. and 5:00 p.m. daily. For CS exposure, the rats underwent whole-body exposure to the smoke of five unfiltered cigarettes (Dagianmen, Yunnan, China;  $\leq 15 \text{ mg tar}$ ,  $\leq 1.1 \text{ mg nicotine and } \leq 13 \text{ mg CO}$ ) for 30 min twice-daily (before 9:00 a.m. and after 5:00 p.m.), 7 days/week for 14 weeks, inside a 0.6 m<sup>3</sup> custom-made plexiglas chamber (16,17).

*Measurement of serum liver enzymes.* Blood samples from control and experimental rats were centrifuged at 600 g for 15 min at 4°C, and the serum was stored at -80°C prior to being assayed. The serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) concentrations were quantified using a transaminase CII test, according to the manufacturer's protocol (Wako Pure Chemical Industries, Ltd., Osaka, Japan).

*Liver tissue sampling.* Following treatment, 10 rats from the control and experimental groups were anesthetized with 10% chloral hydrate (0.3 ml/100 g body weight) and sacrificed. For gene expression analysis using reverse transcription-quantitative polymerase chain reaction (RT-qPCR), liver tissues were excised, rinsed in ice-cold PBS, frozen in liquid nitrogen and stored at -80°C prior to analysis. For hematoxylin and eosin staining, liver tissues were fixed in 10% formalin at 4°C overnight, embedded in paraffin, and sliced into  $5-\mu$ m-thick sections. Sections were then stained with 1% hematoxylin and eosin solution for 5 min at room temperature, according to the manufacturer's protocol.

Preparation of RNA from tissue samples. RNA was extracted from liver tissues using TRIzol<sup>®</sup> reagent (Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA). The extract yield and quality were determined by measuring the absorbance at 260 and 280 nm using the Maestro Nano Micro-Volume Spectrophotometer (Maestrogen, Inc., Hsinchu, Taiwan). The absorbance ratio at 260:280 nm was between 1.8 and 2.0. The RNA was subsequently reverse-transcribed into cDNA. RT-qPCR. mRNA (3  $\mu$ g) was reverse-transcribed into cDNA using oligo(dT) primers for 1 h at 50°C, with the TIAN Script RT kit (Tiangen Biotech Co., Ltd., Beijing, China), according to the manufacturer's protocol. The cDNA served as a template for qPCR, which was performed using SYBR Green PCR core reagents (Bio-Rad Laboratories, Inc., Hercules, CA, USA). Specific gene primers were designed using the PrimerQuest SM software (http://sg.idtdna.com/Primerguest/Home/Index; Integrated DNA Technologies, Inc., Coralville, IA, USA) and commercially produced (BGI Tech, Shenzen, China) (Table I). DNA amplification was performed using a CFX96 Real-Time System (Bio-Rad Laboratories, Inc.) with the following reaction conditions: An initial heating cycle of 95°C for 2 min; 40 cycles, alternating between denaturation at 95°C for 25 sec and primer annealing at 60°C for 25 sec; and final extension at 72°C for 20 sec. Melt curves were used to clarify the identity of the amplicons and the housekeeping gene GAPDH served as an internal control. The relative mRNA expression of targeted genes was calculated using the comparative Cq (threshold cycle) method and normalized to GAPDH mRNA in the same sample (18). The specific  $\Delta Cq$  was calculated as follows:  $[\Delta Cq=(Cq_{GAPDH})-(Cq_{target})]$ ; relative expression was defined as  $2^{-\Delta\Delta Cq}$ 

Statistical analysis. The numerical data are presented as the mean  $\pm$  standard error of the mean. The statistical significance of the differences between the two groups was assessed using Student's t-test. P<0.05 was considered to indicate a statistically significant difference. Statistical analysis was performed using Microsoft Excel software version 2007 (Microsoft Corporation, Redmond, WA, USA).

# Results

IH with CS exposure causes elevated expression of liver enzymes, upregulated mRNA expression of inflammatory cytokines and hepatocyte damage. Consistent with a previous study (12), IH with CS exposure resulted in apparent emphysemic alterations in rat lungs and decreased blood gas concentration, indicating that the rat model of IH with emphysema was successfully established (data not presented). Hematoxylin and eosin staining of the liver demonstrated that various hepatic lesions were observed in the IH with CS group, and not in the control group (Fig. 1A and B). Compared with the control group, hepatic lobules in the IH with CS group exhibited partial clarity and integrity, sinusoids were broadened, inflammatory cell infiltration was observed in the periportal space and various foci of lobular inflammatory cell accumulation were noted. Inflammatory cell infiltrates were observed in the portal area and light staining of the cytoplasm of liver cells suggested cytoplasmic loss (Fig. 1C).

The concentration of serum ALT and AST in the experimental group was increased compared with the control group (P<0.05; Table II), which suggested a more severe impairment of hepatocyte function. Additionally, the mRNA expression levels of interleukin (IL)-1 $\beta$ , IL-6, and tumor necrosis factor (TNF)- $\alpha$  in the livers of the experimental rats were significantly increased compared with the control group (P<0.05; Fig. 2). The results of the present study suggested that early-phase inflammation and mild hepatocyte damage had occurred.

Table I. Primer sequences for the present study.

Genes	Forward (5'-3')	Reverse (5'-3')
IL-1β	TCCCTGAACTCAACTGTGAAATA	GGCTTGGAAGCAATCCTTAATC
IL-6	GAAGTTAGAGTCACAGAAGGAGTG	GTTTGCCGAGTAGACCTCATAG
TNF-α	ACCTTATCTACTCCCAGGTTCT	GGCTGACTTTCTCCTGGTATG
NF-κB	AGACATCCTTCCGCAAACTC	TAGGTCCATCCTGCCCATAA
PXR	GAAGATCATGGCTGTCCTCAC	CGTCCGTGCTGCTGAATAA
CAR	GAGACCATGACCAGTGAAGAAG	AGTCAGGGCATGGAAATGATAG
GR	CAGCAGTGAAATGGGCAAAG	GGGCAAATGCCATGAGAAAC
CYP1A2	GACAAGACCCTGAGTGAGAAG	GAGGATGGCTAAGAAGAGGAAG
CYP2C9	CCCAAGGGCACAACCATATTA	CTTTCTGGATGAAGGTGGCA
CYP2C19	CCCAAGGGCACAACCATATTA	TTTGACCCTCGTCACTTTCTG
CYP2D4	CCTTTCAGCCCTAACACTCTAC	ATGAAGCGTGGGTCATTGT
CYP3A2	GGAAACCCGTCTGGATTCTAAG	GAAGTGTCTCATAAAGCCCTGT
GAPDH	ACTCCCATTCTTCCACCTTTG	AATATGGCTACAGCAACAGGG

IL, interleukin; TNF, tumor necrosis factor; NF, nuclear factor; PXR, pregnane X receptor; CAR, constitutive androstane receptor; GR, glucocorticoid receptor; CYP, cytochrome P450.



Figure 1. Liver pathology of rat exposed to IH and CS. Control rats or rats exposed to IH+CS were sacrificed and liver tissues were processed to conduct hematoxylin and eosin staining. Compared with the integrity and clarity of the liver tissues in (A) the control group, the liver tissues from (B) the IH+CS group demonstrated hepatocytes which were arranged in rows around central veins, and were visible as polygons in the sinusoid; magnification, x200. (C) Further magnification of a section of IH+CS liver tissue; magnification, x400. IH, intermittent hypoxia; CS, cigarette smoke.

Underexpressed mRNA levels of CYPs in the injured liver. Expression of each CYP is influenced by a unique combination of mechanisms and factors, including genetic polymorphisms, xenobiotic induction, regulation by cytokines, hormones and disease states, in addition to sex, age, and others (13). Compared with the control group, the IH with CS group exhibited markedly increased mRNA expression of CYP1A2, CYP2C9, CYP2C19, CYP2D4, and CYP3A2 (Fig. 3). The results of the present study demonstrated that liver inflammation and hepatocyte damage affect the transcription of major CYPs in a rat model of IH with emphysema, which was consistent with the previous literature suggesting inflammation may induce downregulation of CYP expression (19,20).

Upregulation of nuclear factor (NF)- $\kappa B$  and subsequent downregulation of nuclear receptors in the liver. Transcription factor NF- $\kappa B$  serves a role in inflammatory reactions and oxidative stress (21,22). CYP3A expression is modulated by nuclear receptors, including PXR and CAR (23-25). If the nuclear translocation of these receptors is decreased, CYP3A expression will consequently decrease. In addition, the synthesis and nuclear translocation of these receptors is negatively-associated with NF- $\kappa$ B nuclear translocation (14,26). Consequently, NF- $\kappa$ B expression was significantly increased in the IH with CS rats, compared with the control group (P<0.05; Fig. 4). Additionally, in the IH with CS group, the hepatic mRNA expression of PXR, CAR and the glucocorticoid receptor (GR) was significantly decreased compared with that in the control group (P<0.05; Fig. 5). The results of the present study demonstrated that upregulated NF- $\kappa$ B may be involved in reduced hepatic CYP expression, by negatively impacting the synthesis and nuclear translocation of PXR, CAR and GR.

## Discussion

The emerging pathophysiology of OS encompasses intermittent hypoxia and emphysema (27). In the present study, a rat model of intermittent hypoxia with emphysema was developed. The results of the present study suggested that an early phase of inflammation and mild hepatocyte damage had occurred in

Table II. Serum ALT and AST levels in the liver.
--

Group	n	ALT, U/I	AST, U/I
Contro	15	21.0±5.31	17.8±3.0
IH+CS	15	50.5±2.1ª	26.0±3.2ª

<sup>a</sup>P<0.05. IH, intermittent hypoxia; CS, cigarette smoke; ALT, alanine aminotransferase; AST, aspartate aminotransferase.



Figure 2. mRNA expression levels of inflammatory cytokines in the liver. The mRNA expression levels of the inflammatory cytokines (A) IL-1 $\beta$ , (B) IL-6, and (C) TNF- $\alpha$  in the liver were measured using the reverse transcription-quantitative polymerase chain reaction. GAPDH was used as an internal reference. The data are presented as the mean ± standard error of the mean, and were obtained from 10 rats/group. Student's t-test. \*P<0.05 vs. the control. IL, interleukin; TNF, tumor necrosis factor; IH, intermittent hypoxia.



Figure 3. mRNA expression levels of CYPs in the liver. The mRNA expression levels of (A) CYP1A2, (B) CYP2C9, (C) CYP2C19, (D) CYP2D4 and (E) CYP3A2 were measured using the reverse transcription-quantitative polymerase chain reaction. GAPDH was used as an internal reference. The data are presented as the mean  $\pm$  standard error of the mean, and were obtained from 10 rats/group. Student's t-test; \*P<0.05, \*\*P<0.01 vs. the control. CYP, cytochrome P450; IH, intermittent hypoxia.

the liver. The results of the present study were consistent with previous studies, which stated that hypoxia contributes to liver injury (28,29). Alterations in hepatic CYP expression, and the mechanisms underlying these alterations, were analyzed in the present study using the IH with CS rat model. In the IH with CS group, the hepatic mRNA expression levels of a number of CYP molecular species, specifically CYP1A2, CYP2C9, CYP2C19, CYP2D4 and CYP3A2, were markedly decreased compared with the control group.



Figure 4. mRNA expression level of NF- $\kappa$ B in the liver. Following the establishment of the rat model, the liver was removed and the mRNA expression level of NF- $\kappa$ B was measured using the reverse transcription-quantitative polymerase chain reaction. GAPDH was used as an internal reference. The data are presented as the mean  $\pm$  standard error of the mean, and were obtained from ten rats per group. Student's t-test; \*\*P<0.01 vs. the control. NF- $\kappa$ B, nuclear factor- $\kappa$ B; IH, intermittent hypoxia.



Figure 5. mRNA expression levels of (A) PXR, (B) CAR, and (C) GR in the liver. After established successfully model, the liver was removed, and the mRNA expression levels of PXR, CAR, and GR were measured using the reverse transcription-quantitative polymerase chain reaction. GAPDH was used as an internal reference. The data are presented as the mean ± standard error of the mean, and were obtained from ten rats per group. Student's t-test; \*P<0.05, \*\*P<0.01 vs. the control. PXR, pregnane X receptor; CAR, constitutive androstane receptor; GR, glucocorticoid receptor; IH, intermittent hypoxia.

The present study sought to analyze why the expression of hepatic CYPs decreased in the rat model of intermittent hypoxia with emphysema. It is known that CYPs are generated in damaged hepatocytes, where the degree of damage may affect the quantity and quality of CYPs. Previous studies have demonstrated that the severity of liver injury is positively-associated with levels of hypoxia and inflammation (30,31). Compared with patients with only COPD or OSAS, nocturnal hypoxemia, hypoxia and hypercapnia are more severe in patients with OS (32), and the present study demonstrated that intermittent hypoxia and emphysema interact synergistically. The mechanism of the decrease in CYP expression was analyzed by focusing on hepatic inflammation, as the pathological conditions of early-phase hepatic inflammation and mild hepatocyte damage were detected in the present experimental model. Hepatic inflammation is promoted by the binding of cyclic pro-inflammatory factors to Toll-like receptor 4 in hepatic Kupffer cells. The activated Kupffer cells subsequently initiate the secretion of inflammatory cytokines, which promote the activity of NF-KB (21). Activated NF-KB dissociates from the inhibitor of NF-kB and translocates to the nucleus (33). Nuclear NF- $\kappa$ B forms a complex with the transcription factor GR and prevents the binding of GR to GR-responsive elements, thereby inhibiting the transcription of PXR and CAR (14,15). The decrease in PXR and CAR expression causes a decrease in nuclear translocations (26,34). Additionally, the decrease in the expression of PXR and CAR, which bind to the responsive elements of DNA, results in the inhibition of CYP3A transcription and a subsequent decrease in its expression (35-37). Hepatic NF-kB nuclear translocation was examined in the present study, and was observed to be increased in the experimental group compared with the control group. By contrast, the hepatic mRNA expression of PXR, CAR and GR was decreased in the experimental group. The results of the present study demonstrated that, in a rat model of intermittent hypoxia with emphysema, hepatic inflammatory cytokines activate NF-kB, which inhibits the transcription of PXR and CAR. This inhibited transcription subsequently leads to a decrease in nuclear translocation, which inhibits the transcription of CYP3A.

The transcription of CYP2C is modified by CAR (38,39). Therefore, it is hypothesized that the mechanism underlying the decrease in the expression level of CYP2C is identical to that underlying the decrease in the CYP3A expression level. However, the transcription of CYP1A and CYP2D is not primarily regulated by PXR and CAR (40,41). Therefore, the decrease in the nuclear translocation of these CYP molecular species cannot be explained by the decrease in the nuclear translocation of PXR and CAR. However, it has been previously demonstrated that the expression levels of these CYP molecular species are downregulated by inflammatory cytokines (42). It is therefore hypothesized that the decrease in the hepatic expression levels of CYP1A and CYP2D in the IH with CS group, in the present study, may be triggered by inflammatory cytokines, as is the case with other CYP molecular species.

In conclusion, early-phase hepatic inflammation and mild hepatocyte damage was detected in the rat model of intermittent hypoxia with emphysema generated in the present study; additionally, hepatic CYP expression was markedly decreased. Based on the results of the present study, it is hypothesized that, during the development of OS, systematic inflammatory reactions may downregulate hepatic CYP gene expression via the NF- $\kappa$ B signaling pathway. The results of the present study may provide an important molecular mechanism to explain adverse drug reactions in patients with OS.

### Acknowledgements

The present study was supported by the Natural Science Foundation of Tianjin City (grant nos. 13JCYBJC22400, 14JCYBJC25700 and 13JCYBJC40000) and the National Natural Science Foundation of China (grant nos. 31471121 and 81270144).

# References

- McNicholas WT: Chronic obstructive pulmonary disease and obstructive sleep apnea: Overlaps in pathophysiology, systemic inflammation and cardiovascular disease. Am J Respir Crit Care Med 180: 692-700, 2009.
- Ioachimescu OC and Teodorescu M: Integrating the overlap of obstructive lung disease and obstructive sleep apnoea: OLDOSA syndrome. Respirology 18: 421-431, 2013.
- 3. Wouters EF: Local and systemic inflammation in chronic obstructive pulmonary disease. Proc Am Thorac Soc 2: 26-33, 2005.

- 4. Thunström E, Glantz H, Fu M, Yucel-Lindberg T, Petzold M, Lindberg K and Peker Y: Increased inflammatory activity in nonobese patients with coronary artery disease and obstructive sleep apnea. Sleep 38: 463-471, 2015.
- Gonzalez NC and Wood JG: Alveolar hypoxia-induced systemic inflammation: What low PO (2) does and does not do. Adv Exp Med Biol 662: 27-32, 2010.
- Fuhrmann V, Kneidinger N, Herkner H, Heinz G, Nikfardjam M, Bojic A, Schellongowski P, Angermayr B, Kitzberger R, Warszawska J, et al: Hypoxic hepatitis: Underlying conditions and risk factors for mortality in critically ill patients. Intensive Care Med 35: 1397-1405, 2009.
- van Eeden SF and Sin DD: Chronic obstructive pulmonary disease: A chronic systemic inflammatory disease. Respiration 75: 224-238, 2008.
- Dopp JM, Moran JJ, Abel NJ, Wiegert NA, Cowgill JB, Olson EB and Sims JJ: Influence of intermittent hypoxia on myocardial and hepatic P-glycoprotein expression in a rodent model. Pharmacotherapy 29: 365-372, 2009.
- 9. Wrighton SA, Schuetz EG, Thummel KE, Shen DD, Korzekwa KR and Watkins PB: The human CYP3A subfamily: Practical considerations. Drug Metab Rev 32: 339-361, 2000.
- Pan X, Wang P, Hu N, Liu L, Liu X, Xie L and Wang G: A physiologically based pharmacokinetic model characterizing mechanism-based inhibition of CYP1A2 for predicting theophylline/antofloxacin interaction in both rats and humans. Drug Metab Pharmacokinet 26: 387-398, 2011.
  Li Y, Wang Y, Yang Y, Luo Y and Chen P: Effect of chronic
- 11. Li Y, Wang Y, Yang Y, Luo Y and Chen P: Effect of chronic intermittent hypoxia on the expression of fractalkine in rat liver. Zhong Nan Da Xue Xue Bao Yi Xue Ban 38: 984-990, 2013 (In Chinese).
- Feng J, Wang QS, Chiang A and Chen BY: The effects of sleep hypoxia on coagulant factors and hepatic inflammation in emphysematous rats. PLoS One 5: e13201, 2010.
- Zanger UM and Schwab M: Cytochrome P450 enzymes in drug metabolism: Regulation of gene expression, enzyme activities, and impact of genetic variation. Pharmacol Ther 138: 103-141, 2013.
- Assenat E, Gerbal-Chaloin S, Larrey D, Saric J, Fabre JM, Maurel P, Vilarem MJ and Pascussi JM: Interleukin lbeta inhibits CAR-induced expression of hepatic genes involved in drug and bilirubin clearance. Hepatology 40: 951-960, 2004.
  Widén C, Gustafsson JA and Wikström AC: Cytosolic glucocor-
- Widén C, Gustafsson JA and Wikström AC: Cytosolic glucocorticoid receptor interaction with nuclear factor-kappa B proteins in rat liver cells. Biochem J 373: 211-220, 2003.
- Magalang UJ, Cruff JP, Rajappan R, Hunter MG, Patel T, Marsh CB, Raman SV and Parinandi NL: Intermittent hypoxia suppresses adiponectin secretion by adipocytes. Exp Clin Endocrinol Diabetes 117: 129-134, 2009.
- Feng J, Chiang AA, Wu Q, Chen BY, Cui LY, Liang DC, Zhang ZL and Yao W: Sleep-related hypoxemia aggravates systematic inflammation in emphysematous rats. Chin Med J (Engl) 123: 2392-2399, 2010.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. Methods 25: 402-408, 2001.
- 19. Slaviero KA, Clarke SJ and Rivory LP: Inflammatory response: An unrecognised source of variability in the pharmacokinetics and pharmacodynamics of cancer chemotherapy. Lancet Oncol 4: 224-232, 2003.
- Aitken AE, Richardson TA and Morgan ET: Regulation of drug-metabolizing enzymes and transporters in inflammation. Annu Rev Pharmacol Toxicol 46: 123-149, 2006.
- Ivanenkov YA, Balakin KV and Lavrovsky Y: Small molecule inhibitors of NF-kB and JAK/STAT signal transduction pathways as promising anti-inflammatory therapeutics. Mini Rev Med Chem 11: 55-78, 2011.
- 22. Sandireddy R, Yerra VG, Areti A, Komirishetty P and Kumar A: Neuroinflammation and oxidative stress in diabetic neuropathy: Futuristic strategies based on these targets. Int J Endocrinol 2014: 674987, 2014.
- Pondugula SR, Dong H and Chen T: Phosphorylation and protein-protein interactions in PXR-mediated CYP3A repression. Expert Opin Drug Metab Toxicol 5: 861-873, 2009.
- 24. Chai X, Zeng S and Xie W: Nuclear receptors PXR and CAR: Implications for drug metabolism regulation, pharmacogenomics and beyond. Expert Opin Drug Metab Toxicol 9: 253-266, 2013.
- 25. Shah P, Guo T, Moore DD and Ghose R: Role of constitutive androstane receptor in Toll-like receptor-mediated regulation of gene expression of hepatic drug-metabolizing enzymes and transporters. Drug Metab Dispos 42: 172-181, 2014.

- 26. Ghose R, Omoluabi O, Gandhi A, Shah P, Strohacker K, Carpenter KC, McFarlin B and Guo T: Role of high-fat diet in regulation of gene expression of drug metabolizing enzymes and transporters. Life Sci 89: 57-64, 2011.
- 27. Chaouat A, Weitzenblum E, Krieger J, Ifoundza T, Oswald M and Kessler R: Association of chronic obstructive pulmonary disease and sleep apnea syndrome. Am J Respir Crit Care Med 151: 82-86, 1995.
- Tanné F, Gagnadoux F, Chazouillères O, Fleury B, Wendum D, Lasnier E, Lebeau B, Poupon R and Serfaty L: Chronic liver injury during obstructive sleep apnea. Hepatology 41: 1290-1296, 2005.
- Savransky V, Nanayakkara A, Vivero A, Li J, Bevans S, Smith PL, Torbenson MS and Polotsky VY: Chronic intermittent hypoxia predisposes to liver injury. Hepatology 45: 1007-1013, 2007.
- Ebert EC: Hypoxic liver injury. Mayo Clin Proc 81: 1232-1236, 2006.
- 31. Tian JL, Zhang Y and Chen BY: Sleep apnea hypopnea syndrome and liver injury. Chin Med J (Engl) 123: 89-94, 2010.
- Verbraecken J and McNicholas WT: Respiratory mechanics and ventilatory control in overlap syndrome and obesity hypoventilation. Respir Res 14: 132, 2013.
- Baeuerle PA and Baltimore D: I kappa B: A specific inhibitor of the NF-kappa B transcription factor. Science 242: 540-546, 1988.
- 34. Ogura J, Terada Y, Tsujimoto T, Koizumi T, Kuwayama K, Maruyama H, Fujikawa A, Takaya A, Kobayashi M, Itagaki S, *et al*: The decrease in farnesoid X receptor, pregnane X receptor and constitutive androstane receptor in the liver after intestinal ischemia-reperfusion. J Pharm Pharm Sci 15: 616-631 2012.
- 35. Beigneux AP, Moser AH, Shigenaga JK, Grunfeld C and Feingold KR: Reduction in cytochrome P-450 enzyme expression is associated with repression of CAR (constitutive androstane receptor) and PXR (pregnane X receptor) in mouse liver during the acute phase response. Biochem Biophys Res Commun 293: 145-149, 2002.

- 36. Gu X, Ke S, Liu D, Sheng T, Thomas PE, Rabson AB, Gallo MA, Xie W and Tian Y: Role of NF-kappaB in regulation of PXR-mediated gene expression: A mechanism for the suppression of cytochrome P-450 3A4 by proinflammatory agents. J Biol Chem 281: 17882-17889, 2006.
- 37. Wang YM, Ong SS, Chai SC and Chen T: Role of CAR and PXR in xenobiotic sensing and metabolism. Expert Opin Drug Metab Toxicol 8: 803-817, 2012.
- Chen Y, Ferguson SS, Negishi M and Goldstein JA: Identification of constitutive androstane receptor and glucocorticoid receptor binding sites in the CYP2C19 promoter. Mol Pharmacol 64: 316-324, 2003.
- 39. Wortham M, Czerwinski M, He L, Parkinson A and Wan YJ: Expression of constitutive and rostane receptor, hepatic nuclear factor 4 alpha and P450 oxidoreductase genes determines interindividual variability in basal expression and activity of a broad scope of xenobiotic metabolism genes in the human liver. Drug Metab Dispos 35: 1700-1710, 2007.
- 40. Aleksunes LM and Klaassen CD: Coordinated regulation of hepatic phase I and II drug-metabolizing genes and transporters using AhR-, CAR-, PXR-, PPARα- and Nrf2-null mice. Drug Metab Dispos 40: 1366-1379, 2012.
- 41. Cai Y, Konishi T, Han G, Campwala KH, French SW and Wan YJ: The role of hepatocyte RXR alpha in xenobiotic-sensing nuclear receptor-mediated pathways. Eur J Pharm Sci 15: 89-96, 2002.
- 42. Abdulla D, Goralski KB, Del Busto Cano EG and Renton KW: The signal transduction pathways involved in hepatic cytochrome P450 regulation in the rat during a lipopolysaccharide-induced model of central nervous system inflammation. Drug Metab Dispos 33: 1521-1531, 2005.