

The protective effect of berberine hydrochloride on LPS-induced osteoclastogenesis through inhibiting TRAF6-Ca²⁺-calcineurin-NFATc1 signaling pathway

FEI YE, QINGZHONG ZHOU, LONG TIAN, FEI LEI and DAXIONG FENG

Department of Orthopaedic, The Affiliated Hospital of Southwest Medical University,
Luzhou, Sichuan 646000, P.R. China

Received March 1, 2017; Accepted July 20, 2017

DOI: 10.3892/mmr.2017.7338

Abstract. The present study investigated the protective effect of berberine hydrochloride on lipopolysaccharide (LPS)-induced acute bone destruction through inhibition of the TNF receptor associated factor 6 (TRAF6)-Ca²⁺-calcineurin-nuclear factor of activated T-cell 1 (NFATc1) signaling pathway. An osteoclast culture system of RAW264.7 cells induced by LPS *in vitro* was established. A polymerase chain reaction (PCR) assay was applied to determine the effect of berberine hydrochloride on the mRNA expression levels of fos-related antigen 2 (Fra-2), tartrate-resistant acid phosphatase (TRAP), β 3-integrin, cathepsin K, dendritic cell-specific transmembrane protein (DC-STAMP), V-type proton ATPase subunit d 2 (Atp6v0d2) and NFATc1. An ELISA assay was performed to measure the release of tumor necrosis factor- α (TNF- α). Western blot analysis was used to measure the effect of berberine hydrochloride on the expression of calcineurin in the LPS-induced NFATc1 signaling pathway, as well as the expression levels of phosphoinositide phospholipase C- γ 1 (PLC- γ 1), toll like receptor 4 (TLR4) and TRAF6. The effect of berberine hydrochloride on Ca²⁺ concentration was detected using a confocal technique with a Flou-3/acetoxymethyl ester Ca²⁺ probe. The PCR results demonstrated that berberine hydrochloride inhibited the mRNA expression levels of Fra-2, TRAP, β 3-integrin, cathepsin K, DC-STAMP, Atp6v0d2 and NFATc1. Furthermore, the ELISA results demonstrated that TNF- α expression was decreased. The western blot analysis revealed that berberine hydrochloride treatment results in decreased expression levels of PLC- γ 1, TLR4 and TRAF6, and inhibition of Ca²⁺ influx. In conclusion, the results of the present study suggest that berberine hydrochloride targets TRAF6 and

NFATc1, thus inhibiting osteoclastogenesis and bone destruction via inhibition of the TRAF6-Ca²⁺-calcineurin-NFATc1 signaling pathway.

Introduction

As a dynamic organ, the bone undergoes continuous resorption and generation process, of which the process is called remodeling. Under normal circumstances, there is a balance between osteoclast-mediated resorption and osteoblast-mediated formation, which maintains metabolism and homeostasis. However, the loss of balance may lead to skeletal diseases (1). Osteoclasts are the only cells that are responsible for bone resorption. Overexpression of osteoclasts leads to excessive bone resorption and bone destruction, thus causing osteoporosis, osteoarthritis and other common joint diseases (2-4).

Berberine hydrochloride, as an isoquinoline alkaloid, with molecular formula [C₂₀H₁₈NO₄]⁺, also known as berberine, presents in many plants of the Berberidaceae families. Berberine can be precipitated in ether as yellow needle-like crystals, with melting point at 145°C, soluble in water, antibacterial to hemolytic *Streptococcus*, *Staphylococcus aureus*, *Neisseria gonorrhoeae*, and Freund's *Shigella*, and also can enhance leukocyte phagocytosis. Berberine hydrochloride has been widely used in the treatment of gastroenteritis, bacillary dysentery, pulmonary tuberculosis, scarlet fever, acute tonsillitis and respiratory infections. However, it has been reported neither whether the drug can affect osteoclastogenesis, nor osteoclast-associated bone destruction can be improved (5). Therefore, it is of certain significance to be investigated (6).

Materials and methods

RAW264.7 cell culture and the effect of berberine on lipopolysaccharide (LPS)-induced osteoclastogenesis. The 3rd generation passaged RAW264.7 cells (from Sun Yat-Sen University) were inoculated into 96-well plates at a density of 1x10⁴/well and cultured in DMEM medium containing 10% FBS and 1% double antibody, 200 μ l/well, and then put in an incubator at 37°C with 5% CO₂. RAW264.7 cells were treated with LPS (100 ng/ml) except control group; normal

Correspondence to: Dr Daxiong Feng, Department of Orthopaedic, The Affiliated Hospital of Southwest Medical University, 25 Taiping Street, Luzhou, Sichuan 646000, P.R. China
E-mail: fengdaxiong0606@163.com

Key words: berberine hydrochloride, osteoclasts, Ca²⁺, NFATc1, LPS, bone destruction

saline was added to the model group, and Berberine hydrochloride (5, 10 and 20 μM ; Manst Biological Technology Co., Ltd.) was added into the Drug group simultaneously. Each group was set with three multiple holes. Tartrate-resistant acid phosphatase (TRAP) staining was performed after 5 days' culture. The staining method was carried out according to the kit instructions. The cells were observed under an inverted microscope.

ELISA method to detect tumor necrosis factor- α (TNF- α) levels. After 24 h culture, the cells were put in the ELISA plate and added with diluted samples or standardized samples. The Control group holes were set at 100 μl . Biotinylated antibody were added into each hole at 50 μl , mixed, and sealed with closure plates. The cells were incubated at 37°C for 90 min, washed for 4 times, then added with streptavidin-HRP, with 100 μl per hole. Subsequently, the cells were incubated at 37°C for 30 min, and washed for 4 times. Color reagent TMB was added at 100 μl per hole. The cells were colored at 37°C for 10 min. Stop solution was added at 50 μl per hole. Absorbance at a wavelength of 450 nm was read by a microplate reader, and TNF- α density was calculated.

Quantitative polymerase chain reaction (PCR) assay to detect the mRNA expressions of fos-related antigen 2 (Fra-2), TRAP, β 3-integrin, cathepsin K, dendritic cell-specific transmembrane protein (DC-STAMP), V-type proton ATPase subunit d 2 (Atp6v0d2) and NFATc1. The cells were incubated for 24 h and added with 1 ml TRIzol reagent, then agitated sufficiently. A total of 0.2 ml chloroform was added, mixed, agitated, and then the cells were ice-bathed standing for 5 min. The cells were centrifuged at 12,000 rpm for 20 min under 4°C. The supernatant was replaced with an equal volume of isopropanol, then ice-bathed standing for 5 min. The cells were centrifuged at 12,000 rpm for 20 min under 4°C. The supernatant was discarded and a total of 1 ml ethanol at 75% was added. The solution was agitated to dissolve the RNA thoroughly. The cells were centrifuged at 10,000 rpm for 5 min under 4°C. The supernatant was carefully removed, and the EP tube was turned upside down, then dried under room temperature for 15 min. A total of 20 μl RNase-free water was added to dissolve the precipitate, and 1 μl solution was suctioned into an Eppendorf, and the rest was stored in -70°C refrigerator. The 1 μl solution was diluted to 80 μl . OD260 and OD280 ratio was measured by a spectrophotometer, then total RNAs were calculated.

The cDNA synthesis and reverse transcription were carried out using the PrimeScript RT reagent kit according to the instructions, as follows: 5 times PrimeScript buffer 2 μl , PrimeScript RT Enzyme Mix 0.5 μl , Total RNA 2 μl , RNase Free dH₂O 5 μl , and Total 10 μl . The reaction was carried out in water bath at 37°C for 15 min, and then at 85°C for 15 sec. After the reaction, an appropriate amount of cDNA was quantified by real-time PCR, and the remaining samples were stored under -20°C. Fluorescence real-time PCR was performed according to the instructions of the SYBR Premix Ex Taq™ II (Perfect Real Time) kit: SYBR Premix Ex Taq 12.5 μl , PCR Forward Primer 1 μl , PCR Reverse Primer 1 μl , DNA template 2 μl , d H₂O 8.5 μl , and Total 25 μl . Fluorescence real-time PCR was performed as follows: Amplification curve: 5 min

pre-denaturation at 95°C; 20 sec denaturation at 95°C; 30 sec annealing at 60°C; and 20 sec extending at 72°C. This stage was used for fluorescence signal acquisition, 40 cycles in total. The dissolution curve: 60°C-95°C, 0.5°C in each incremental, for 20 sec. This stage is for fluorescence signal acquisition, a total of 71 cycles. The size of PCR product was verified and a total of 10 μl was obtained to perform electrophoresis in 1% agar gel.

PCR primers were as follows: Fra-2, 5'-CCAACACGTAGT TTGAAGAC-3' and 5'-TCCTGCCGCAGTTACACCG-3'; TRAP, 5'-TGACATCGAGCAGGTGAAAG-3' and 5'-GAG TAGCAAGGAATGAGC-3'; β 3-integrin, 5'-AGCGGACAT TCTGGAAATG-3' and 5'-TCGTTTCATGCACACTGCTGA-3'; cathepsin K, 5'-ACCGACAACAGCAGCATT-3' and 5'-AGC TTGCTGTGCTTCAGT-3'; DC-STAMP, 5'-AACTGTTGT GGCCTGAATC-3' and 5'-CGGTAAATGCAGGCGTAT-3'; Atp6v0d2, 5'-CATCCCATCACCATCTTCC-3' and 5'-TCA CACATGACGAAGGCA-3'; NFATc1, 5'-TCCTCCATGAAC AACAG-3' and 5'-AGACGTGGTTTAGGAATGCAG-3'; and GAPDH, 5'-AACTTTGGCATGTGGAAGG-3' and 5'-ACACATTGGGGGTAGGAAC-3'.

Western blotting to detect the expressions of calcineurin in PLC γ , toll like receptor 4 (TLR4), TNF receptor associated factor 6 (TRAF6) and nuclear factor of activated T-cell 1 (NFATc1) signaling pathways. The above cells were cultured for 24 h, harvested, and washed with PBS twice. Each bottle was added with 400 μl of cell lysate, followed by 40 μl PMSF, agitated. The cells were placed on ice 10 min for sufficient lysis. After being repeatedly aspirated by a sterile syringe, the cells were put into an EP tube, which was ice-bathed for 30 min, and subsequently centrifuged at 12,000 x g for 15 min. After the supernatant was transferred to a new EP tube, 20 μl of protein sample buffer was added into each 100 μl tube, boiled for 5 min, mixed, and then stored under -80°C. The protein from the above samples was separated by 12% SDS-PAGE electrophoresis. The separated protein bands were transferred to a PVDF membrane by wet method and closed under room temperature for 1 h. Subsequently, the primary antibody was added (concentration 1:1,000) for 4°C overnight incubation. The primary antibody was discarded, and the secondary antibody (BA1026) was added (concentration 1:1,000) for 1 h incubation. The secondary antibody was abandoned, and color development and fixation were carried out after chemiluminescence. The expressions of calcineurin in PLC γ , TLR4, TRAF6 and NFATc1 signaling pathways were measured.

Confocal assay to detect the effect of berberine hydrochloride on intracellular Ca²⁺ concentration. After 24 h culture, the supernatant was discarded. The cells were washed with HESS, following this, HBSS containing 5Flou-3/AM and 0.05% Pluronic F127 was added. The cells were incubated for 30 min under 37°C. The supernatant was abandoned, and the cells were washed with HEPES buffer saline for 3 times. Subsequently, a sum of 2 ml HEPES buffer saline was added. At the excitation wavelength of 488 nm and the emission wavelength of 500-530 nm, the intracellular Ca²⁺ concentration was measured according to the intensity changes of Fluo-3/AM by using AIR laser scanning confocal microscopy.




Figure 1. Effect of berberine on LPS-induced osteoclastogenesis by TRAP staining (x100). LPS, lipopolysaccharide; TRAP, tartrate-resistant acid phosphatase.




Figure 2. Effect of berberine hydrochloride on LPS-induced TNF- α production in RAW264.7 cells. **P<0.01, *P<0.05 vs. Model; ##P<0.01, #P<0.05 vs. 5 μ M; &P<0.05 vs. 10 μ M. LPS, lipopolysaccharide; TNF- α , tumor necrosis factor- α .

Statistical methods. All data were expressed as mean \pm SD. Comparisons between the two groups were carried out using the t-test, and multiple sets of data (>2) comparison using the one-way ANOVA, where P<0.05 denoted statistical significance. The data were analyzed by GraphPad Prism 5.0 (GraphPad Software Inc., San Diego, CA, USA).

Results

Effects of berberine on LPS-induced osteoclastogenesis. TRAP staining showed that RAW264.7 could be differentiated into mature osteoclasts after 5 days of LPS induction, while osteoclastogenesis was not seen in RAW264.7 cells in the negative Control group. The results showed that LPS-induced RAW264.7 cells' differentiation into osteoclasts can be significantly inhibited by Berberine hydrochloride of high concentration. See Fig. 1.

The inhibition of berberine on LPS-induced osteoclast TNF- α secretion. In the supernatant, LPS significantly induced TNF- α secretion, while the inhibition of TNF- α by berberine exhibited in a dose-dependent manner compared with that of the Control group, suggesting that berberine may inhibit osteoclastogenesis through inhibiting TNF- α production. See Fig. 2.

PCR assay to detect the mRNA expressions of Fra-2, TRAP, β 3-integrin, cathepsin K, DC-STAMP, Atp6v0d2 and NFATc1. The mRNA expressions of Fra-2, TRAP, DC-STAMP and Atp6v0d2 in RAW264.7 cells induced by LPS were significantly inhibited by berberine hydrochloride, suggesting that

berberine could inhibit LPS-induced osteoclasts-related gene expression, thereby inhibiting the differentiation of RAW264.7 cells to osteoclasts.

During the osteoclastogenesis, NFATc1 was amplified and NFATc1 mRNA was upregulated. The results showed that berberine can inhibit the increase of NFATc1 gene expression induced by LPS in a concentration-dependent manner. The experimental results confirmed the hypothesis that the inhibition of osteoclastogenesis by berberine may be related to the activation of nuclear transcription factor NFATc1. See Fig. 3.

Western blot to detect the calcineurin expressions of PLC γ , p-PLC γ , TLR4, TRAF6 and NFATc1, p-NFATc1 signaling pathways. WB results showed that berberine hydrochloride significantly inhibited the expression of NFATc1 in the nucleus. The results further confirmed the hypothesis that berberine inhibited the osteoclastogenesis possibly by inhibiting the activation of nuclear transcription factor NFATc1.

After stimulated by LPS, PLC γ was activated and increased intracellular Ca²⁺ concentration. The results showed that the activation of PLC γ , or the p-PLC γ protein, was significantly upregulated by LPS which was inhibited by berberine, suggesting that berberine could inhibit the activation of PLC γ , thereby inhibiting Ca²⁺ influx, which reduced intracellular Ca²⁺ concentration.

In order to further confirm the effect of berberine hydrochloride on LPS signal transduction pathway, the effect of berberine on the expression of LPS-induced TRAF6 protein was investigated. The expression of TRAF6 protein was significantly increased in RAW264.7 cells after LPS stimulation, but was significantly inhibited by berberine. The results were consistent with the detection of PCR. See Fig. 4.

Berberine to reduce the intracellular Ca²⁺ concentration. The intracellular Ca²⁺ concentration was measured by the Confocal technique. The results showed that LPS significantly increased intracellular Ca²⁺ concentration. However, berberine significantly reduced intracellular Ca²⁺ concentration, in particular, the high concentration group reduced intracellular Ca²⁺ concentration the most, even close to the normal group, suggesting that berberine hydrochloride can strongly inhibit Ca²⁺ influx. As shown in Fig. 5.

Discussion

'Bone immunology' is more and more a hot topic, in which the osteoclast is the only cell responsible for bone resorption. Under the category of 'Bone immunology', there are excessive




Figure 3. PCR detection of berberine on the expression of various genes. **P<0.01, *P<0.05 vs. Model; #P<0.01, #P<0.05 vs. 5 μ M; &P<0.01, &P<0.05 vs. 10 μ M; !P<0.01 vs. 20 μ M. PCR, polymerase chain reaction; LPS, lipopolysaccharide; TRAP, tartrate-resistant acid phosphatase; Fra-2, fos-related antigen 2; Atp6v0d2, V-type proton ATPase subunit d 2; DC-STAMP, dendritic cell-specific transmembrane protein.

activations of osteoclasts in degenerative diseases such as osteoporosis, inflammatory arthritis, and tumor-induced bone

destruction (7). Activated osteoclasts can absorb bones, resulting in bone loss, causing fractures, reactive bone




Figure 4. Effect of berberine hydrochloride on the expressions of various proteins. **P<0.01, *P<0.05 vs. Control; ##P<0.01, #P<0.05 vs. 5 μM; &&P<0.01, &P<0.05 vs. 10 μM; †P<0.01, †P<0.05 vs. 20 μM. PLC-γ1, phosphoinositide phospholipase C-γ1; TLR4, toll like receptor 4; TRAF6, TNF receptor associated factor 6; NFATc1, nuclear factor of activated T-cell 1.




Figure 5. Effect of berberine hydrochloride on Fluo-3/AM fluorescence intensity at 200 s (x100).

proliferation, and subsequent pain and disability. Drugs that whip the osteoclast differentiation can be applied to the treatment of osteoporosis, and in recent years expanding to the diseases such as tumor bone metastasis and rheumatoid arthritis (8).

LPS is a component of the G-cell wall, which can induce bone destruction through chronic infection. *In vitro* TRAP staining showed that berberine inhibited LPS-induced osteoclastogenesis. Studies have shown that inflammatory factors such as TNF-α, IL-1, and IL-6 can promote osteoclast-mediated bone resorption, posing a key regulatory factor in osteoclast activity (9,10). These inflammatory factors contribute to the formation of strong osteoclastogenesis stimulating molecules PGE2 by increasing the expression of COX-2 in osteoblasts and stromal cells. Increased TNF-α and other inflammatory factors are associated with the fracture of the femoral head in older women (11,12).

The present study found that berberine can reduce the TNF-α levels in LPS stimulated supernatant, suggesting that berberine may have the potential anti-inflammatory and anti-osteoclastogenesis effect by downregulating the production of TNF-α. In addition, it is found that berberine significantly inhibited the expression of TRAP gene in RAW264.7 cells. Inhibition of TNF-α activity may be the reason for the inhibition of TRAP activity.

Intracellular Ca²⁺ influx can activate calcineurin, which can de-phosphorylate the inactivated p-NFATc1, and promote NFATc1 into the nucleus, thereby activating osteoclast-specific gene expression. The present study found that berberine inhibited calcineurin expression, thereby inhibiting cytoplasmic

p-NFATc1 de-phosphorylation and NFATc1 nuclear translocation, indicating that drugs can inhibit the activation of NFATc1, and thus inhibit osteoclast-associated gene expression. And molecules such as DC-STAMP, Atp6v0d2 and Fra-2 are all related to osteoclastogenesis. During the osteoclastogenesis, the osteoclast precursors fuse into large multinucleated macrophages under the intervention of DC-STAMP and Atp6v0d2 (13,14).

Fra-2 can regulate the size and survival of osteoclasts (15). β3-integrin, cathepsin K and MMP-9 are necessary for osteoclasts to carry out bone resorption. Both β3-integrin and Cathepsin K played important roles in the bone resorption by mediating migration and adsorption of osteoclasts (16,17). Cathepsin K exhibited high expression in osteoclasts, making a key enzyme in degradation of organic bone matrix, thus facilitated the bone resorption of osteoclasts (18-20).

In conclusion, berberine hydrochloride may target through the TRAF6 and NFATc1, and inhibit osteoclastogenesis and bone destruction probably through inhibiting TRAF6-Ca²⁺-calcineurin-NFATc1 signaling pathways.

References

1. Geusens P and Lems WF: Osteoimmimology and osteoporosis. *Arthritis Res Ther* 13: 242, 2011.
2. Deal C: Bone loss in rheumatoid arthritis: Systemic, periarticular, and focal. *Curr Rheumatol Rep* 14: 231-237, 2012.
3. McClung M, Harris ST, Miller PD, Bauer DC, Davison KS, Dian L, Hanley DA, Kendler DL, Yuen CK and Lewiecki EM: Bisphosphonate therapy for osteoporosis: Benefits, risks, and drug holiday. *Am J Med* 126: 13-20, 2013.
4. Schett G and Gravallese E: Bone erosion in rheumatoid arthritis: Mechanisms, diagnosis and treatment. *J Nat Rev Rheumatol* 8: 656-664, 2012.

5. Tao K, Xiao D, Weng J, Xiong A, Kang B and Zeng H: Berberine promotes bone marrow-derived mesenchymal stem cells osteogenic differentiation via canonical Wnt/ β -catenin signaling pathway. *Toxicol Lett* 240: 68-80, 2016.
6. Banks L and Takayanagi H: Immunology and bone. *J Biochem* 154: 29-39, 2013.
7. Stickeler E and Fehm T: Targeted and osteo-oncologic: Treatment in early breast cancer: What is state of the art and what might become so within the next 5 years. *Breast Care (Basel)* 9: 161-167, 2014.
8. D'Amico AV: US Food and drug administration approval of drugs for the treatment of prostate cancer: A new era has begun. *J Clin Oncol* 32: 362-364, 2014.
9. Manolagas SC and Jilka RL: Bone marrow, cytokines, and bone remodeling. Emerging insights into the pathophysiology of osteoporosis. *N Engl J Med* 332: 305-311, 1995.
10. Braun T and Schett G: Pathways for bone loss in inflammatory disease. *Curr Osteoporos Rep* 10: 101-108, 2012.
11. Okada Y, Lorenzo JA, Freeman AM, Tomita M, Morham SG, Raisz LG and Pilbeam CC: Prostaglandin G/H synthase-2 is required for maximal formation of osteoclast-like cells in culture. *J Clin Invest* 105: 823-832, 2000.
12. Barbour KE, Boudreau R, Danielson ME, Youk AO, Wactawski-Wende J, Greep NC, LaCroix AZ, Jackson RD, Wallace RB, Bauer DC, *et al*: Inflammatory markers and the risk of hip fracture: The women's health initiative. *J Bone Miner Res* 27: 1167-1176, 2012.
13. Kim K, Lee SH, Ha Kim J, Choi Y and Kim N: NFATc1 induces osteoclast fusion via up-regulation of Atp6v0d2 and the dendritic cell specific transmembrane protein (DC-STAMP). *Mol Endocrinol* 22: H6-85, 2008.
14. Yagi M, Miyamoto T, Sawatani Y, Iwamoto K, Hosogane N, Fujita N, Morita K, Ninomiya K, Suzuki T, Miyamoto K, *et al*: DC-STAMP is essential for cell-cell fusion in osteoclasts and foreign body giant cells. *J Exp Med* 202: 345-351, 2005.
15. Bozec A, Bakiri L, Hoebertz A, Eferl R, Schilling AF, Kommenovic V, Scheuch H, Priemel M, Stewart CL, Amling M and Wagner EF: Osteoclast size is controlled by Fra-2 through LIF/LIF-receptor signalling and hypoxia. *Nature* 454: 221-225, 2008.
16. Song I, Kim R, Kim K, Jin HM, Youn BU and Kim N: Regulatory mechanism of NFATc1 in RANKL induced osteoclast activation. *FEBS Lett* 583: 2435-2440, 2009.
17. Wei ZF, Tong B, Xia YF, Lu Q, Chou GX, Wang ZT and Dai Y: Norisoboldine suppresses osteoclast differentiation through preventing the accumulation of TRAF6-TAK1 complexes and activation of MAPKs/NF- κ B/c-Fos/NFATc1 pathways. *PLoS One* 8: e59171, 2013.
18. Asagiri M and Takayanagi H: The molecular understanding of osteoclast differentiation. *Bone* 40: 251-264, 2007.
19. Costa AG, Cusano NE, Silva BC, Cremers S and Bilezikian JP: Cathepsin K: Its skeletal actions and role as a therapeutic target in osteoporosis. *Nat Rev Rheumatol* 7: 447-456, 2011.
20. Broadhead ML, Clark JC, Dass CR, Choong PF and Myers DE: The apertic targeting of osteoclast function and pathways. *Expert Opin Ther Targets* 15: 169-181, 2011.