ADME/toxicity prediction and antitumor activity of novel nitrogenous heterocyclic compounds designed by computer targeting of alkylglycerone phosphate synthase

YU ZHU^{1*}, YUAN HAN^{1*}, YING MA^{2*} and PING YANG¹

¹Department of Clinical Laboratory, Tianjin Key Laboratory of Cerebral Vessels and Neural Degeneration, Tianjin Huanhu Hospital, Tianjin 300350; ²Tianjin Key Laboratory on Technologies Enabling Development of Clinical Therapeutics and Diagnostics (Theranostics), School of Pharmacy, Tianjin Medical University, Tianjin 300070, P.R. China

Received September 12, 2017; Accepted February 15, 2018

DOI: 10.3892/ol.2018.8873

Abstract. Alkylglycerone phosphate synthase (AGPS) is an oncogene and can be considered as an antitumor drug target. The aim of the present study was to design novel nitrogenous heterocyclic compound improving targetability by computer-aided drug design technology targeting AGPS. A total of 12 nitrogenous heterocyclic compounds were designed and predicted the absorption, distribution, metabolism and excretion parameters/toxicity. Their activity in terms of proliferation inhibition, cell cycle arrest and apoptosis induction was then measured using an MTS assay and a high-content screening system in U251 cells. The results showed that anti-glioma activity was present in compounds N4, N5, N6, N7, N8 and N12, which was in accordance with the computer prediction. Therefore, these compounds may be suitable for the development of a novel glioma therapeutic drug.

Introduction

Malignant glioma is a disease that severely impairs human health. In our previous study, alkylglycerone phosphate synthase (AGPS) was found to be an oncogene, and the hypothesis was made that AGPS may be a target of glioma inhibitors (1,2). The aim of the present study was to design nitrogenous heterocyclic compounds via computer-aided design and determine the molecular biological differences between tumor cells and normal cells, in order to improve

Correspondence to: Dr Yu Zhu, Department of Clinical Laboratory, Tianjin Key Laboratory of Cerebral Vessels and Neural Degeneration, Tianjin Huanhu Hospital, 6 Jizhao Road, Tianjin 300350, P.R. China

E-mail: zhuyutj@126.com

*Contributed equally

Key words: computer-aided drug design, alkylglycerone phosphate synthase, nitrogenous heterocyclic compound

selectivity and enhance the current research on targetability of the nitrogenous heterocyclic compounds. AGPS was selected as the target around which to structure a two-dimensional (2D) and 3D structure-activity model via computer-aided drug design (CADD) technology to make macromolecular docking. Nitrogenous heterocyclic compounds were designed and a database was established in an effort to obtain a comprehensive understanding of the structure-activity relationship among candidate compounds.

It has been found that the onset and progression course of a tumor is often accompanied with lipid metabolic confusion (3). The inactivation of key enzymes synthesized by AGPS can lower the ether ester level of tumor cells and reduce the pathogenicity of the cancer, while their overexpression can raise the ether ester level of the tumor cells and enhance cell proliferation and movement ability, thereby promoting growth and invasion of tumor (4). This indicates the potential of AGPS to become a novel target for antineoplastic agents, and its specific inhibitor will provide advantages which surpass those provided by traditional chemotherapy. Our previous study also validated the use of RNA interference technology to silence the expression of AGPS in glioma and hepatoma cells in order to inhibit their proliferation and invasion, and to improve the sensitivity of cells that are resistant to medicine (5).

Based on the aforementioned study (6), AGPS was selected as target to structure 2D and 3D structure-activity association models via CADD technology in the present study. A total of 12 nitrogenous heterocyclic compounds were designed and a database was established, in order to obtain a comprehensive understanding of the structure-activity associations among the candidate compounds and then further optimize the lead compound structure.

Materials and methods

Protein structure and the database. A 3D structure model [Protein Data Bank (PDB) ID: 2UUV] of human AGPS was downloaded from PDB (7). The amino acid residue in the crystal structure did not mutate. Crystal structure resolution was <2.6.

Ligand preparation. Ligand preparation was designed by chemical drug experience and utilized ChemBioDraw Ultra 11.0 in the ChemOffice 2010 software package (PerkinElmer, Inc., Waltham, MA, USA) to plot the micromolecule planar construction. The plotted planar construction was then imported into ChemBio3D Ultra 11.0 to generate a spatial structure, which was then saved in *.mol2 format. The file in *.mol2 format was imported into Discovery Studio 3.5 software (BIOVIA, San Diego, CA, USA) for docking and Erwin Schrodinger 2009 software (Schrodinger, LLC, Portland, OR, USA) for absorption, distribution, metabolism and excretion (ADME) analysis, and the LigPrep module was adopted to optimize the micromolecule, desalt and add an electric charge to produce ionization consistent with human body pH. The force field of the optimization molecule was OPLS_2005, which is the same as that of the optimization receptor, thereby forming a tautomeride of the molecule. The alloisomerism treatment adopted the input spatial structure to perform the chiral method of molecular docking.

Molecular docking. Molecular docking was conducted using the 'docking' module in the Discovery Studio 3.5 software, with 'receptor box producing' selected and centered on the original ligand micromolecule to automatically generate the receptor box file. Flexible docking with standard accuracy was adopted. All calculations were conducted using the Dell Precision T5500 workstation (Dell Inc., Round Rock, TX, USA) with the Red Hat Enterprise Linux 6.0 operating system (Red Hat, Raleigh, NC, USA).

The compound structure was imported into Discovery Studio 3.5, and the preparation work for the micromolecule prior to docking was completed using the prepare ligands tool module in the software package, including restoring valence, generating 3D conformation and minimizing the energy of the micromolecule. Molecular docking was performed by the CDOCKER module in Discovery Studio 3.5, with the original ligand molecule as the center and 9Å as the radius, to automatically generate docking region. The result used 'CDOCKER ENERGY' as the scoring index. The larger the value was, the larger the affinity of receptor docking.

ADME analysis. ADME was an important part in the medicine discovery process to select and optimize the compound. The downloaded AGPS target crystal structure (2UUV) was imported into Schrodinger 2009 for ADME analysis in order to first correct the chemical bond in the protein structure, treat the metallic ions, hydrogenate, make the amino acid residue at end N and end C neutral, reserve the original ligand and protein, and remove impurity atoms and water molecules. The OPLS_2005 position was selected to optimize the protein structure, and sample water orientations served as the parameters to optimize the hydrogen bond, with the convergence standard being RMSD0.5Å.

Toxicity analysis. Toxicity prediction was also important in the selection of a potential drug. Toxicity analysis was conducted using the 'TOPKAT' module in Discovery Studio 3.5 software. Toxicity parameters included rodent carcinogenicity, mutagenicity, the ames test, skin irritancy, ocular irritation, aerobic biodegradability, and whether the molecule was a non-carcinogen, non-mutagen and non-degradable.

Cell culture. AGPS was demonstrated to be overexpressed in glioma in a previous study (8), so the glioma U251 cell line was used to explored the activity of compounds at first, provided by the Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). The cells were cultured in Dulbecco's modified Eagle's medium with 10% fetal bovine serum (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA) in a cell incubator with 5% CO₂ at 37°C.

MTS assay. A total of $5x10^3$ U251 cells in suspension with Dulbecco's modified Eagle's medium with 10% fetal bovine serum were added to 96-well plates at 37°C for 24 h, and N1-N12 (designed by the School of Pharmacy, Tianjin Medical University, Tianjin, China; synthesized by Werian Biotech Co., Jinan, China) at different concentrations (0, 2, 5, 10, 20, 50, 100 and 200 µM) were added at 37°C for 72 h. Then MTS (Promega Corporation, Madison, WI, USA) solvent was added and incubated at 37°C for 1 h and the OD value was measured using a Multiskan FC Microplate Reader (Thermo Fisher Scientific Inc.). The cell proliferation was then measured by MTS assay. Inhibition rate was calculated as follows: Inhibition rate (%)=(1-optical density value_{treatment group}/optical density value_{control group}) x100. The half maximal inhibitory concentration (IC₅₀) was calculated by Graphpad Prism 6.0 (GraphPad Software, Inc., La Jolla, CA, USA).

High-content screening (HCS) assay. The cell cycle and apoptosis were measured by HCS platform CellInsight CX5 (Thermo Fisher Scientific, Inc.) via fluorescence intensity. For the cell cycle measurement, $8x10^3$ U251 cells in suspension were added to 96-well plates with Premo Cdt1-red fluorescent protein and Premo geminin-green fluorescent protein (both Thermo Fisher Scientific, Inc.) at 37°C for 24 h. N1-N12 (100 μ M) [with 100 μ l phosphate-buffered saline (PBS) as the negative control] were added for 24 h, and then the cell cycle was measured and analyzed by HCS platform at 488 nm excitation.

For the cell apoptosis measurements, 8×10^3 U251 cells in suspension were added to 96-well plates for 24 h, and 100 μ M compounds (with PBS as the negative control) were added for 24 h. The cells were then incubated with Annexin V and Hoechst 33258 (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) in the dark at room temperature for 15 min, and the cell apoptosis was measured and analyzed by HCS platform at 488 nm excitation.

Statistical analysis. The experimental data were statistically analyzed with SPSS 11.0 statistical software (SPSS, Inc., Chicago, IL, USA) and are expressed as the mean \pm standard deviation. The statistical analysis was performed using one-way analysis of variance with the Tukey-Kramer post hoc test. P<0.05 was considered to indicate a statistically significant difference.

Results

Analysis on affinity. A total of 12 compounds were designed for the analysis on affinity, which is represented by the docking score (a high docking score indicates a high affinity) (Fig. 1). The 2D combined mode with amino acid residue (Fig. 2) and the 3D combined mode (Fig. 3) with AGPS show the



143	3
-----	---

Table I. Absorpt	tion, distribution.	metabolism an	d excretion	prediction of	nitrogenous	heterocyclic com	pounds
					0	~	

Compound	MW (Da)	donorHB	accptHB	QPlogPo/w	QPlogS	PSA
N1	353.299	4.75	13.75	-2.350	-2.809	215.170
N2	394.349	5.50	14.00	-2.247	-2.459	230.122
N3	270.250	3.50	10.50	-1.464	-2.210	148.217
N4	323.310	4.75	8.75	-0.409	-3.568	170.861
N5	329.315	4.75	9.50	-0.617	-3.307	178.267
N6	350.336	4.75	8.75	0.100	-3.810	172.853
N7	362.390	3.50	9.50	0.008	-2.991	144.227
N8	376.417	3.50	9.50	0.259	-3.435	143.247
N9	376.417	3.50	10.50	0.488	-2.838	142.237
N10	376.417	3.50	9.50	0.505	-2.489	135.360
N11	376.417	4.00	10.50	0.340	-2.651	139.990
N12	380.449	3.50	9.50	0.506	-2.749	139.842

MW, molecular weight; donorHB, hydrogen bond donor; accptHB, hydrogen bond receptor; QPlogPo/w, lipid water partition coefficient; QPlogS, water solubility partition coefficient; PSA, polar surface area.

Name	Structure	Docking score	Name	Structure	Docking score
N1		-10.0112	N7	NH2 N N N N O S	-10.5824
N2		-9.1720	N8		-9.9654
N3		-8.0454	N9		-9.8299
N4		-8.6013	N10		-9.7717
N5		-7.4427	N11		-10.1660
N6		-9.6292	N12		-9.9439

Figure 1. Structure and docking score of nitrogenous heterocyclic compounds N1-N12.



Figure 2. 2D combined mode of nitrogenous heterocyclic compounds N1-N12. The green circle and green line represent the hydrophobic bond, the pink circle and blue line represent the hydrogen bond, and the orange line represents the pi bond (a covalent bond) in interactions of the compounds. 2D, two-dimensional.

hydrophobic bond (green circle and green line), hydrogen (pink circle and blue line) and pi bond (a covalent bond, orange line) interactions of the compounds (the dotted lines in Fig. 2).

ADME prediction. ADME properties are an important index to check whether clinical candidates can reach the required standard (9). Bad properties imply a high risk of failure for this candidate, which may become a less than ideal drug. The ADME of an ideal medicine is as follows: Hydrogen bond donor, \leq 5; hydrogen bond receptor, <10; molecular weight, <500 Da; lipid water partition coefficient, <5; water-solubility partition coefficient, -6.5<logS<-0.5; and polar surface area, 7.0-200.0 (10).

Table I shows the ADME results of the nitrogenous heterocyclic compounds, with some that satisfied the aforementioned qualifications. N4, N5, N6, N7, N8 and N12 were considered to exhibit potential drug properties. *Toxicity prediction*. Table II shows the toxicity prediction results of the nitrogenous heterocyclic compounds. All compounds were found to exhibit suitable toxicity for drug development.

Effect of nitrogenous heterocyclic compounds on the proliferation of U251 cells. Table III and Fig. 4 show the IC_{50} of the nitrogenous heterocyclic compounds for 72 h. It was found that compounds N4, N5, N6, N7, N8 and N12, which had IC_{50} values of <100 μ M, were suitable for drug development.

Effect of nitrogenous heterocyclic compounds on the cell apoptosis of U251 cells. Table IV shows the effect of the nitrogenous heterocyclic compounds on cell apoptosis for 24 h. Compounds N4, N5, N6, N7, N8 and N12 induced a >10% cell apoptosis rate at a 100 μ M concentration (Fig. 5).





Figure 3. 3D combined mode of nitrogenous heterocyclic compounds N1-N12. 3D, three-dimensional.

Effect of nitrogenous heterocyclic compounds on the cell cycle of U251 cells. Table V shows the effect of nitrogenous heterocyclic compounds on the cell cycle for 24 h. It was found that compounds N2, N3, N4, N5, N6, N7, N8, N11 and N12 induced a >10% cell cycle arrest at a 100 μ M concentration (Fig. 6).

Discussion

Malignant glioma is a disease that seriously damages human health; it accounts for ~70% of primary malignant brain tumors, with an annual incidence of ~5/100,000 individuals (11,12). The number of new cases every year exceeds 14,000. As for glioblastoma (World Health Organization stage IV) patients, median survival time is only between 14.6 and 17 months, and the annual number of associated mortalities reaches 30,000 (13-15). The chemical features comprise: The hydrogen bond receptor, hydrogen bond donor, hydrophobic center, positive charge center, negative charge center and aromatic ring center (16-18). In the present study, compounds N1-N12 were first docked, resulting in high docking scores with the target AGPS. This implies their potential ability to inhibit AGPS.

In previous years, the medicine discovery phase mainly focused on the discovery of active compounds, and other problems, including pharmacokinetics, toxicity, solubility and stability, were not considered until the development phase. Thus, solely analyzing the affinity of the compound and target cannot ascertain the possibility of these compounds becoming potential medicines (19). ADME properties are an important index to check whether clinical candidates can reach the required standard. Bad properties imply a high risk of failure for this candidate, which may then become a less than ideal drug. According to a previous study, ~40% of failures to develop a medicine in the development phase are due to poor biopharmaceutical properties (PK and bioavailability) (20). High development expenses make such failures a primary

Compound	Mouse female NTP prediction	Mouse male NTP prediction	Rat female NTP prediction	Rat male NTP prediction	Ames prediction	Skin irritancy	Ocular irritancy	Aerobic biodegradability prediction
N1	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Mutagen	None	Mild	Non-degradable
N2	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
N3	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
N4	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
N5	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
N6	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Mutagen	None	Mild	Non-degradable
N7	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Mutagen	None	Mild	Non-degradable
N8	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Mutagen	None	Mild	Non-degradable
N9	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Mutagen	None	Mild	Non-degradable
N10	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
N11	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
N12	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-carcinogen	Non-mutagen	None	Mild	Non-degradable
NTP, National	Toxicology Program.							

Table III. IC ₅₀ of nitrogenous heterocyclic compounds	in U251
cells.	

Compound	$IC_{50}, \mu M$
N1	251.7
N2	128.5
N3	106.8
N4	86.7
N5	73.5
N6	97.5
N7	88.2
N8	92.8
N9	168.7
N10	205.6
N11	153.8
N12	71.7

IC₅₀, half maximal inhibitory concentration.

Table IV. Apoptosis rate of nitrogenous heterocyclic compounds in U251 cells.

Compound	Apoptosis rate, %
PBS control	1.1
N1	8.7
N2	6.2
N3	6.3
N4	18.7
N5	24.5
N6	15.6
N7	21.8
N8	17.2
N9	6.3
N10	8.7
N11	5.5
N12	23.7

economic loss in medicine development. Thus, ADME has become an indispensable part of the medicine discovery process, and it can be used to supervise the selection and optimization of precious lead compounds. The QikProp module (in part of the Erwin Schrodinger 2009 software) can predict the following properties of compounds: the logarithm value of brain to plasma concentration ratio, water solubility, liposolubility, MDCK cell line and Caco-2 cell line for permeability, general central nervous system activity, K ion channel blocking of human ether-a-go-go related gene log IC₅₀ and the bonding activity of serum albumin (log Khsa). The present study experiments predicted water solubility, polar surface area, MDCK cell permeability, and oil and water partition coefficient. It was found that N4, N5, N6, N7, N8 and N12 exhibited potential drug properties. The toxicity predictions of the compounds were also investigated with Discovery Studio using the TOPKAT protocol, and the toxicity

Table II. Toxicity prediction of nitrogenous heterocyclic compounds.



Table V. Effect of nitrog	cenous heter	ocyclic comp	ounds on the	e cell cycle o	f U251 cells								
Cell cycle stage (%)	PBS	N1	N2	N3	N4	N5	N6	N7	N8	6N	N10	N11	N12
G ₀ /G ₁	43.7	38.7	41.2	39.7	37.1	39.5	41.2	36.7	37.5	41.8	43.2	41.2	34.8
S	51.6	51.3	46.8	48.3	45.7	43.6	44.6	48.6	49.8	50.6	50.7	46.3	42.3
G ₂ /M	4.7	10.0	12.0	12.0	17.2	16.9	14.2	14.7	12.7	7.6	6.1	12.5	22.9
PBS, phosphate-buffered s	aline.												



Figure 4. MTS assay showed that nitrogenous heterocyclic compounds suppressed the proliferation of U251 cells and the IC_{50} was determined. IC_{50} , half maximal inhibitory concentration.



Figure 5. Effect of nitrogenous heterocyclic compounds on the cell apoptosis of U251 cells. HCS assay showed that nitrogenous heterocyclic compounds induced cell apoptosis of U251 cells. *P<0.05 compared with the PBS group. PBS, phosphate-buffered saline.



Figure 6. Effect of nitrogenous heterocyclic compounds on the cell cycle of U251 cells. HCS assay showed that nitrogenous heterocyclic compounds induced cell cycle arrest in the G2/M phase of U251 cells. $^{\circ}P<0.05$ compared with the PBS group. PBS, phosphate-buffered saline.

of the compounds was found to be suitable for the development into a medical drug.

HCS was the novel technology used for the drug activity screening in the present study, and it was able to analyze the effect on cell proliferation, cell apoptosis and the cell cycle by high throughput and quick screening using the HCS platform. The effect on proliferation, apoptosis and the cell cycle by 12 nitrogenous heterocyclic compounds was analyzed, and N4, N5, N6, N7, N8 and N12 were shown to be most active *in vitro* according the ADME prediction, showing the accuracy of the CADD screening model.

In conclusion, 12 nitrogenous heterocyclic compounds were designed in the present study, 6 of which presented with suitable ADME parameters and toxicity predictions; these 6 may be suitable for development into novel glioma therapeutic drugs. This study provides a foundation for the study into novel nitrogenous heterocyclic anti-glioma drugs and future studies will investigate the toxicity in animal models.

Acknowledgements

Not applicable.

Funding

The present study was supported by the National Natural Science Foundation of China (grant nos. 31501159 and 81601047), the Tianjin Public Health Key Research Project (grant no. 15KG108), the Tianjin Science and Technology Key Project on Chronic Disease Prevention and Treatment (grant no. 16ZXMJSY00020), the Special Program of Talent Development for Excellent Youth Scholars in Tianjin, China (grant no. TJTZJH-QNBJRC-2-9) and the Tianjin 131 Creative Talents Cultivation Project (1st Class, 2016).

Availability of data and materials

All data and materials relevant to the present study are described in this published article or available from the corresponding author on reasonable request.

Authors' contributions

YZ performed the statistical analysis, YZ and YH designed the study and performed the cell experiments, YM designed and synthesized the computer-aided drug, and PY performed the statistical analysis.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

 Ames H, Halushka MK and Rodriguez FJ: miRNA regulation in gliomas: Usual suspects in glial tumorigenesis and evolving clinical applications. J Neuropathol Exp Neurol 76: 246-254, 2017.

- Cives M, Ghayouri M, Morse B, Brelsford M, Black M, Rizzo A, Meeker A and Strosberg J: Analysis of potential response predictors to capecitabine/temozolomide in metastatic pancreatic neuroendocrine tumors. Endocr Relat Cancer 23: 759-767, 2016.
- Dilillo M, Ait-Belkacem R, Esteve C, Pellegrini D, Nicolardi S, Costa M, Vannini E, Graaf EL, Caleo M and McDonnell LA: Ultra-high mass resolution MALDI imaging mass spectrometry of proteins and metabolites in a mouse model of glioblastoma. Sci Rep 7: 603, 2017.
- 4. Piano V, Benjamin DI, Valente S, Nenci S, Marrocco B, Mai A, Aliverti A, Nomura DK and Mattevi A: Discovery of inhibitors for the ether lipid-generating enzyme AGPS as anti-cancer agents. ACS Chem Biol 10: 2589-2597, 2015.
- Zhu Y, Liu XJ, Yang P, Zhao M, Lv LX, Zhang GD, Wang Q and Zhang L: Alkylglyceronephosphate synthase (AGPS) alters lipid signaling pathways and supports chemotherapy resistance of glioma and hepatic carcinoma cell lines. Asian Pac J Cancer Prev 15: 3219-3226, 2014.
- 6. Zhu Y, Liu A, Zhang X, Qi L, Zhang L, Xue J, Liu Y and Yang P: The effect of benzyl isothiocyanate and its computer-aided design derivants targeting alkylglycerone phosphate synthase on the inhibition of human glioma U87MG cell line. Tumour Biol 36: 3499-3509, 2015.
- 7. Razeto A, Mattiroli F, Carpanelli E, Aliverti A, Pandini V, Coda A and Mattevi A: The crucial step in ether phospholipid biosynthesis: Structural basis of a noncanonical reaction associated with a peroxisomal disorder. Structure 15: 683-692, 2007.
- Zhu Y, Zhu L, Lu L, Zhang L, Zhang G, Wang Q and Yang P: Role and mechanism of the alkylglycerone phosphate synthase in suppressing the invasion potential of human glioma and hepatic carcinoma cells *in vitro*. Oncol Rep 32: 431-436, 2014.
- 9. Battu MB, Chandra AM, Sriram D and Yogeeswari P: Pharmacophore-based 3DQSAR and molecular docking studies to identify new non-peptidic inhibitors of cathepsin S. Curr Med Chem 21: 1910-1921, 2014.
- 10. Verma SK and Thareja S: Structure based comprehensive modelling, spatial fingerprints mapping and ADME screening of curcumin analogues as novel ALR2 inhibitors. PLoS One 12: e0175318, 2017.
- 11. Silginer M, Weller M, Stupp R and Roth P: Biological activity of tumor-treating fields in preclinical glioma models. Cell Death Dis 8: e2753, 2017.
- 12. Lin CY and Huang HM: Unilateral malignant optic glioma following glioblastoma multiforme in the young: A case report and literature review. BMC Ophthalmol 17: 21, 2017.
- 13. Louis DN, Perry A, Reifenberger G, *et al*: The 2016 World Health Organization classification of tumors of the central nervous system: A summary. Acta Neuropathol 131: 803-820, 2016.
- 14. Wang N, Zhang Q, Ning B, Luo L and Fang Y: β-Asarone promotes Temozolomide's entry into glioma cells and decreases the expression of P-glycoprotein and MDR1. Biomed Pharmacother 90: 368-374, 2017.
- 15. Holdhoff M, Ye X, Supko JG, Nabors LB, Desai AS, Walbert T, Lesser GJ, Read WL, Lieberman FS, Lodge MA, *et al*: Timed sequential therapy of the selective T-type calcium channel blocker mibefradil and temozolomide in patients with recurrent high-grade gliomas. Neuro Oncol 19: 845-852, 2017.
- Cai Z, Zhang G, Zhang X, Liu Y and Fu X: Current insights into computer-aided immunotherapeutic design strategies. Int J Immunopathol Pharmacol 28: 278-285, 2015.
- Medeiros Turra K, Pineda Rivelli D, Berlanga de Moraes Barros S and Mesquita Pasqualoto KF: Constructing and validating 3D-pharmacophore models to a set of MMP-9 inhibitors for designing novel anti-melanoma agents. Mol Inform 35: 238-252, 2016.
- Valasani KR, Chaney MO, Day VW and Shidu Yan S: Acetylcholinesterase inhibitors: Structure based design, synthesis, pharmacophore modeling, and virtual screening. J Chem Inf Model 53: 2033-2046, 2013.
- Cao D, Wang J, Zhou R, Li Y, Yu H and Hou T: ADMET evaluation in drug discovery. 11. PharmacoKinetics Knowledge Base (PKKB): A comprehensive database of pharmacokinetic and toxic properties for drugs. J Chem Inf Model 52: 1132-1137, 2012.
- 20. Gomeni R, Bani M, D'Angeli C, Corsi M and Bye A: Computer-assisted drug development (CADD): An emerging technology for designing first-time-in-man and proof-of-concept studies from preclinical experiments. Eur J Pharm Sci 13: 261-270, 2001.
 - This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.