

# The expression and functional characterization associated with cell apoptosis and proteomic analysis of the novel gene MLAA-34 in U937 cells

WEN-JUAN ZHANG, WANG-GANG ZHANG, PENG-YU ZHANG,  
XING-MEI CAO, AI-LI HE, YIN-XIA CHEN and LIU-FANG GU

Department of Clinical Hematology, Affiliated No. 2 Hospital, Xi'an Jiaotong University  
College of Medicine, Xi'an, Shaanxi 710004, P.R. China

Received August 12, 2012; Accepted September 25, 2012

DOI: 10.3892/or.2012.2129

**Abstract.** MLAA-34 is a novel acute monocytic leukemia (M5)-associated antigen (MLAA) that plays a role in the apoptosis of U937 cells. However, the expression and molecular mechanism of MLAA-34 in U937 cells remain largely unclear. Here, we utilized three strategies to gain insight into the expression and molecular functions of MLAA-34 and to identify its interacting proteins and pathways involved in the fine-tuning of the MLAA-34 response. Western blot analysis was performed to assess the expression of MLAA-34 in 41 cell lines and five mixed cell types, which revealed that MLAA-34 is most strongly expressed in U937 cells. Immunostaining indicated that MLAA-34 is localized in the cytoplasm and cell membrane. Furthermore, lentivirus-mediated overexpression of MLAA-34 in the U937 cell line led to significant suppression of apoptosis and increased the potential of tumorigenicity. Co-immunoprecipitation (Co-IP), shotgun and bioinformatic analysis identified 256 proteins and 225 of them were annotated by gene ontology categories. This analysis revealed 71 proteins involved in cell apoptosis or proliferation of biological processes and signaling pathways. Moreover, the effect of MLAA-34 apoptosis may be through interaction with the Ras, Wnt, calcium and chemokine signaling pathways and thirteen of the annotated proteins may interact with MLAA-34 and participate in carcinogenesis directly. This study provides a basis for a better understanding of the molecular mechanism and proteomics in the inhibition of apoptosis by MLAA-34 in U937 cells and indicates that MLAA-34 may be a potential candidate for the early diagnosis and therapeutic application of M5.

## Introduction

Leukemia is the leading cause of mortality worldwide in patients with malignant tumors under the age of 35 years. Patients with acute myeloid leukemia (AML) who have relapsed or are refractory to conventional chemotherapy have a poorer prognosis and response to chemotherapy than those with *de novo* AML, which remains a formidable therapeutic challenge even with the introduction of several new therapeutic strategies (1-3). M5 is largely incurable with high relapse rates, infiltration and a median remission duration of only six months, approximately (4). Moreover, M5 has been reported to have a worse prognosis than other subtypes of AML (5). Thus, a vaccine or a new drug against M5 is required as a strategic tool for the control of this disease, but none are currently available for practical use.

The MLAA-34 gene (GenBank no. AY288977.2) has been confirmed to be a novel splice variant of CAB39L (calcium binding protein 39-like). MLAA-34 was first discovered in M5 in an effort to identify monocytic leukemia-associated antigens by serologic analysis of a recombinant cDNA expression library (SEREX) that reacted exclusively with sera from allogeneic leukemia patients but not with normal donor sera (6,7). The 1671 kb gene is located on 13q14.2 and was initially cloned in our laboratory from U937 cells (7). CAB39L has three alternative transcripts and has been predicted to encode a 337 aa protein. The three alternative transcripts of CAB39L have been recognized to encode the same protein, differing only in their 5' untranslated regions [GenBank nos. BC010993 (1482 bp), BX647518 (2371 bp) and AY288977.2].

In our previous study, MLAA-34 and CAB39L were identified with RNA interference (RNAi) in the U937 cell line as novel anti-apoptotic factors that are closely related to carcinogenesis or progression of M5 (7). Clinical research has shown that MLAA-34 mRNA expression is upregulated in refractory/relapsed M5 patients compared with newly diagnosed, healthy donors and AML patients in complete remission; high expression of MLAA-34 is more prominent in the M5 subtype than in other AML patients; MLAA-34 overexpression has been found to be associated with unfavorable clinical features at diagnosis and has been shown to be an independent prognostic factor (8).

---

*Correspondence to:* Professor Wang-Gang Zhang, Department of Clinical Hematology, Affiliated No. 2 Hospital, Xi'an Jiaotong University College of Medicine, 157 West 5 Road, Xi'an, Shaanxi 710004, P.R. China  
E-mail: juner0705@163.com

**Key words:** MLAA-34, U937, acute monocytic leukemia, lentivirus, anti-apoptosis, proteomics

However, for MLAA-34, there are no exact reports regarding its cellular localization and expression in manifold cell lines; the anti-apoptotic mechanism of MLAA-34 remains unclear.

The purpose of this study was to conduct an in-depth search for the expression and anti-apoptotic mechanism of MLAA-34 through the lentivirus-mediated overexpression in the U937 cell line, and to then apply proteomics to identify its correlated proteins or pathways that might perform functions important for the apoptosis and proliferation of U937 cells.

## Materials and methods

**Cell culture.** U937, HL60, K562, RPMI-8226, HepG2, Hep3B, MHCC97-H, RC-K8, SGC-7901, Eca109, BGC823, MKN45, GES-1, BxPC-3, A375, T24, HUVEC, BMSCs, LO2, HeLa, 293T, 293, RD, RT4, 5637, EJ, UM-UC-3, 2537, J82, Tsu-Prl, MAH, LiBr, Hut-78, HCT116<sup>+</sup>, FBL-3, C6, astrocyte, 3T3-L1, NIH3T3, Vero and MDCK cell lines were all maintained in our laboratory and cultured in RPMI-1640 or DMEM supplemented with 10% fetal calf serum. The medium for cell lines expressing the neomycin resistance gene was supplemented with 0.5 mg/ml G418. Human epithelial tissue, normal human peripheral blood mononuclear cells (PBMCs), M5 patient and non-M5 acute leukemia patient PBMCs were all obtained from over 30 cases of patients or healthy young individuals. Mouse splenocytes were obtained from 30 mice.

**Antibodies and reagents.** CAB39L and MLAA-34 share the same open reading frame (ORF), the CAB39L antibody was used in this report. Antibodies specific for CAB39L (sc-100390),  $\beta$ -catenin (sc-133240), Rab-3D (sc-26559), Rap-1B (sc-1481) and PGK1 (sc-130335) were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA). A monoclonal mouse antibody against  $\beta$ -actin was obtained from Sigma-Aldrich (St. Louis, MO, USA). The SAP kit and AP-Red kit were provided by Zhongshan Co. Beijing, China (SAP-9102, ZLI-9042). The lentivirus packaging system and enhanced infection solution (ENi.S) were purchased from GeneChem Limited Company (Shanghai, China). The SYBR Green PCR kit and SYBR Master Mixture were purchased from Takara Bio, Inc. (Dalian, China). The Endo-free Plasmid Mini kit was purchased from Qiagen, USA (12163). M-PER<sup>®</sup> Mammalian Protein Extraction Reagent was purchased from Pierce, Rockford, IL, USA (78503).

**Western blot analysis.** Cells were collected at a concentration of  $2 \times 10^7$ /ml. Following sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), the proteins were transferred to polyvinylidene fluoride membranes, which were incubated with the primary antibody CAB39L (1:200). Western blot analyses were performed according to standard methods. The protein bands were visualized by applying SuperSignal West Pico Chemiluminescent Substrate (34079; Pierce). The exposed film was then analyzed using a densitometer.

**Immunohistochemistry and immunofluorescence.** For analysis of the subcellular localization of MLAA-34, U937 cells were washed with ice-cold PBS, blocked with 10% normal goat serum and incubated with a primary antibody against

CAB39L at a dilution of 1:50 for 2 h at 37°C. Next, the cells were washed again and incubated with the appropriate biotinylated secondary antibody (goat anti-mouse IgG antibody) for 20 min at 37°C. Incubation with serum alkaline phosphatase (SAP; ALP) was then performed at 37°C for 20 min, and the immunolabeling was visualized with a mixture of AP-Red solution. Counterstaining with hematoxylin was performed. For immunofluorescence, the cell samples were incubated with the monoclonal antibody CAB39L (diluted 1:50) and fluorescein isothiocyanate (FITC)-labeled or rhodamine-labeled goat anti-mouse IgG as the primary and secondary antibodies, respectively. The mounted cells were visualized with a fluorescent microscope.

**Construction and identification of the MLAA-34 lentivirus vector and upregulated MLAA-34 stably transfected cell line.** The full-length MLAA-34 cDNA sequence was assembled by searching the NCBI database and amplified by RT-PCR from U937 cells. First-strand cDNA synthesis was performed using a commercial kit (Boehringer Mannheim, Milan, Italy). The restriction enzyme site for AgeI (ACCGGT) was introduced into the 5' and 3' PCR primers. To generate cDNA coding for full-length MLAA-34 by PCR, the following primers were designed using plasmid MLAA-34 as the template: MLAA-34-Age, I-F, GAGGATCCCCGGGTACCGGTCGCCACCATGAAAAAATGCCTTTG and MLAA-34-Age, I-R, TCACCATGGTGGCGACCGGAGGGGCCGTTTCTTCAAG. The PCR conditions consisted of 30 cycles, and the cycle parameters were: 94°C for 5 min, then 30 cycles of 94°C for 30 sec, 55°C for 30 sec, 68°C for 1 min, followed by a final extension of 68°C for 10 min. The PCR product was purified using an Agarose Gel DNA Purification kit (Takara Bio, Inc.). The two recovered products were ligated using an In-Fusion kit (631774; Becton, Dickinson and Co., USA). To confirm that the ligation was correct, MLAA-34-SEQF, GACAGATAGGCACTCGGAG; Ubi-F, GGGTCAATATGTAATTTTCAGTG; and EGFP-N-R, CGTCGCCGTCCAGCTCGACCAG primers were designed. The cycle parameters were: 30 cycles of 94°C for 30 sec, 94°C for 30 sec, 60°C for 30 sec, 72°C for 50 sec, followed by a final extension of 72°C for 6 min. For detection of MLAA-34 expressed by recombinant lentivirus *in vitro*, purified pGC-FU-MLAA-34 vectors were transfected into 293T cells using Lipofectamine 2000 reagent (11668-019; Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. This vector was termed MLAA-34-Lentivirus, and the vector without MLAA-34 cDNA was pGC-FU-GFP-LV. The titer of the recombinant lentivirus was determined by real-time qPCR on 293T cells. For identification of the recombinant MLAA-34 lentivirus vector, the virus was added to targeted U937 cells at multiplicity of infections (MOIs) of 10, 20, 50, 80, 100, 120 and 200 with ENi.S and 5  $\mu$ g/ml polybrene. MLAA-34-Lentivirus and pGC-FU-GFP-LV transfected U937 cells were used as the test, and non-transfected cells were used as the control. The expression level of MLAA-34 was detected by western blot analysis and RT-PCR. The best MOI was chosen.

Cells were grown in selective media (containing G418) for two weeks, expanded and grown as independent clones for at least two weeks. Resistant colonies were counted, and the expression of GFP was confirmed by fluorescence microscopy, RT-PCR and western blot analysis.

**Fluorescence microscopy, MTT, flow cytometry and DNA ladder.** To determine the effect of upregulation of MLAA-34 by the MLAA-34-Lentivirus, non-transfected cells and cells transfected with pGC-FU-GFP-LV and MLAA-34-Lentivirus were examined. Cells were seeded in 96-well plates at a density of  $1 \times 10^4$  cells/well. Cellular proliferation was measured once per day during a seven-day period. In brief, 20  $\mu$ l of sterile MTT (Sigma) dye (5 mg/ml) was added to the cells, which were then incubated for another 4 h at 37°C. Then, 150  $\mu$ l of dimethylsulfoxide was added to each well. The spectrophotometric absorbance was measured at a wavelength of 490 nm on an enzyme immunoassay analyzer.

Fixed cells were stained with 2.5 g/ml of DAPI (4',6-diamidino-2-phenylindole) solution to detect apoptotic nuclei. Quantification of apoptosis was determined by counting the number of apoptotic cells. The cells were stained using an Annexin V-PE/7-AAD apoptosis detection kit (KGA1015; Nanjing KeyGen Biotech. Co., Ltd.) according to the manufacturer's instructions and were analyzed by flow cytometry using a Beckman Coulter flow cytometer.

For cell cycle analysis, the cells were fixed in 70% ethanol and stained with propidium iodide (PI; Biosea Biotechnology Co., Beijing, China) at a final concentration of 20  $\mu$ g/ml in Triton X-100 containing 10 mg/ml RNase. Following incubation, the samples were analyzed on a flow cytometer.

Fragmented DNA was isolated using a DNA extraction kit (C0008; Beyotime) according to the manufacturer's instructions. The eluants containing DNA pellets were electrophoresed on a 1% agarose gel at 80 V for 1.5 h. The gel was examined and photographed using an ultraviolet gel documentation system.

**Co-immunoprecipitation (Co-IP) and SDS-PAGE.** Co-IP was performed using a Profound™ Mammalian Co-IP kit (23605; Pierce). Transfected U937 cells ( $2 \times 10^7$ /ml) were washed, centrifuged and resuspended in lysis buffer for incubation. The cell lysates were centrifuged to remove the supernatant material, and the CAB39L antibody was cross-linked to the antibody coupling resin. The lysed cell sample was then applied to the antibody support to form immune complexes. Then, unbound proteins were washed away three times. The samples were then eluted, and coupling buffer was added to obtain the immunoprecipitated protein. Finally, the Co-IP protein concentrations were determined using a BCA Protein Assay kit (23225; Pierce). The proteins were analyzed by SDS-PAGE, and the gel was stained with Coomassie Blue.

**Mass spectrometry analysis (MS, shotgun) and protein identification.** After separation by SDS-PAGE, discrete bands were excised from and subjected to in-gel tryptic digestion. The extracted peptides were analyzed using shotgun HPLC-ESI-MS proteomics approach (LTQ; Thermo Finnigan, San Jose, CA, USA). High-performance liquid chromatography (HPLC) separation was performed with a capillary LC pump. The mobile phases used for the reverse phase were i) 0.1% formic acid in water, pH 3.0; ii) 0.1% formic acid in ACN. The collision energy was set automatically by the LTQ system. Following acquisition of full scan mass spectrum, three MS/MS scans were acquired for the next three most intense ions using dynamic exclusion. Peptides and proteins were identified using




Figure 1. Expression of MLAA-34 by western blot analysis. A, commixture of M5 patient PBMCs; B, commixture of non-M5 acute leukemia patient PBMCs; C, normal human PBMCs; D, human epithelial tissue.

Bioworks Browser 3.1 software (Thermo Finnigan), which uses the MS and MS/MS spectra of peptide ions to search against the NCBI human protein database. The protein identification criteria that we used were based on Delta CN ( $\geq 0.1$ ) and Xcorr (one charge  $\geq 1.9$ , two charges  $\geq 2.2$ , three charges  $\geq 3.75$ ). The protein identification results were extracted from the SEQUEST out file with in-house software (BuildSummary). The cellular localization, molecular function and biologic process were determined using the gene ontology annotation DAVID (<http://david.abcc.ncifcrf.gov/>). For pathway analysis, the KEGG database was searched. To identify the corresponding proteins in mixed protein obtained by Co-IP, western blot analysis was performed as previously described.

**Statistical analysis.** The RT-PCR results were analyzed by the self-contained software of iQ5 (Bio-Rad Co.). Statistical analyses were performed using an analysis of variance (ANOVA). All results are expressed as the means  $\pm$  standard deviations from at least three experiments.  $P < 0.05$  was considered to indicate statistically significant differences.

## Results

**Expression of human MLAA-34 protein.** With western blot analysis, a strong specific band of ~39 kDa was observed in U937 and MHCC97-H cells, and reduced expression was observed in other leukemia or lymphoma cell lines and PBMCs from leukemia patients. Much fainter bands were observed in solid tumor cell lines, and no expression was detected in normal human cell lines or primary animal cells (Fig. 1).

**Identification and cellular localization of MLAA-34.** Immunohistochemical staining confirmed the presence of MLAA-34 in U937 cells and the subcellular localization was detected primarily in the cytomembrane and cytoplasm (Fig. 2).




Figure 2. The subcellular localization of MLAA-34 protein in U937 cells (x40). The MLAA-34 protein was distributed predominantly in the cytomembrane and cytoplasm (red or green). The nuclei were stained by DAPI (x40).




Figure 3. Lentivirus-mediated gene transfer system for overexpression of MLAA-34 was constructed successfully and markedly decreases apoptosis and promotes proliferation in U937 cells. (A) Morphological observation of MLAA-34-Lentivirus transfected U937 cells at different MOIs. The transfection efficiency was higher at MOI=50 and 80, but the growth condition was the best at MOI=50. (B) MLAA-34-Lentivirus upregulated the expression of MLAA-34 in U937 cells, estimated by western blot analysis (left) and RT-PCR (right). (C) Morphological changes in the morphology of the cell nucleus were observed by DAPI staining (blue). (D) Cell viability was measured by MTT assays. (E) The cells were stained with Annexin V-PE and 7-AAD for flow cytometry analysis.

*MLAA-34 is upregulated by the lentiviral vector.* A human MLAA-34 lentivirus gene transfer vector encoding the green fluorescent protein (GFP) sequence was constructed. The pGC-

FU-MLAA-34-GFP plasmid has an insert of ~771 bp, which is in accord with the MLAA-34 cDNA [identities, 1009/1012 (99%)]. The pilot experiments showed that 293T cells could




Figure 4. Proteomics analysis of MLAA-34 in stably transfected U937 cells. (A) SDS-PAGE gel images after Co-IP and the cutting sites in the gel band. Four spaced sections were excised. From left to right were cell lysate, elutriant after Co-IP, sediment after Co-IP and marker. (B) Western blot analysis. The major bands that migrated at ~21, 25, 92 and 45 kDa corresponding to Rap-1B, Rab-3D, β-catenin and PGK1. (C-E) Numbers and percentages of the annotated proteins with molecular function, biological process and cellular localization. (F and G) Distributions of theoretical molecular mass and PI for all of the annotated proteins.

be successfully infected by the packaged virus; the virus titer reached higher than  $2 \times 10^8$  TU/ml, indicating that a high-titer lentiviral packaging platform was preliminarily established. The pGC-FU-MLAA-34-GFP plasmid was confirmed by

western blot analysis. MLAA-34-Lentivirus and control pGC-FU-GFP-LV virus were produced. After obtaining ideal U937 cells, we transfected the cells with the MLAA-34-Lentivirus and pGC-FU-GFP-LV viruses at different MOIs. The transfection




Figure 5. Categories of related pathways in 225 annotated proteins according to KEGG pathway taxonomy.

tion efficiency was ~95% or higher on Day 5 or later at the MOI of 50 (Fig. 3A). Five days after transfection, the recombinant MLAA-34-Lentivirus caused a pronounced increase in the expression of MLAA-34 compared with non-transfected U937 cells (Fig. 3B).

**Establishment of U937 cell line stably overexpressing MLAA-34.** In preliminary studies, 400  $\mu\text{g/ml}$  of G418 were found to maintain adequate selection pressure. The expression of GFP and MLAA-34 were observed. After the cells had been frozen in liquid nitrogen for six months and revived monthly, the U937 cells expressed higher levels of MLAA-34 in ~400  $\mu\text{g/ml}$  of G418, and ~95% of the lentivirus-transfected U937 cells overexpressed MLAA-34. These results suggested that the stably transfected U937 cell line was successfully established by lentivirus and that the expression of MLAA-34 can be long lasting even after passage.

**Effect of upregulating MLAA-34 on apoptosis and growth of U937 cells.** Observations of morphology revealed increasing cell shrinkage, nuclear condensation and fragmentation in non-transfected and pGC-FU-GFP-LV transfected cells. By contrast, cells transfected with MLAA-34-Lentivirus predominantly appeared uniformly stained without condensation (Fig. 3C). These results further support the findings that anti-apoptotic changes in the cell and nuclear morphology are induced by MLAA-34 overexpression. MTT assays suggested that the lentiviral overexpression of MLAA-34 induces anti-apoptotic effects

that result in a promotion effect on U937 cells; these data suggest that MLAA-34 might accelerate cell proliferation (Fig. 3D). In agreement with the anti-apoptotic effects of MLAA-34, cells overexpressing MLAA-34 accumulated in the S-phase (~67.63% compared with ~49.6% of cells in the S-phase in the control) and showed a corresponding increase in cell numbers in the G2/M phase. The percentages of early (lower right) and late apoptotic (upper right) cells were markedly reduced in U937 cells after transfection with MLAA-34-Lentivirus (Fig. 3E). These results are in agreement with the DNA ladder assay and are even more evident at the MOI=50, in which the cells transfected with MLAA-34-Lentivirus showed a further increase. All of these results suggest that MLAA-34 inhibits apoptosis in U937 cells.

**Co-IP, shotgun and western blot analysis.** Protein extracts with Co-IP were separated by SDS-PAGE and the gel was cut into four pieces for shotgun ESI-MS analysis (Fig. 4A). A total of 256 proteins were identified by the LC ESI-MS analysis and BIOWORKS in the NCBI HUMAN protein databases, of which 225 (87.9%) proteins were annotated by DAVID and the remaining 31 (12.1%) proteins have no DAVID terms (Table I). The expression of Rap-1B, Rab-3D,  $\beta$ -catenin and PGK1 was verified by western blot analysis (Fig. 4B).

Classification of the 225 annotated proteins in terms of molecular function, biological process and cellular localization was performed according to the DAVID. Molecular function was clustered and the protein binding (140, 41.8%) and nucleotide binding (60, 17.9%) groups were the majority (Fig. 4C).

Table I. The 225 annotated proteins identified by the shotgun approach.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
1	IP100022434	ALB albumin	Apoptosis	
2	IP100023598	TUBB4 tubulin $\beta$ -4 chain		
3	IP100013475	TUBB2A tubulin $\beta$ -2A chain		
4	IP100180675	TUBA1A tubulin $\alpha$ -1A chain		
5	IP100387144	TUBA1B tubulin $\alpha$ -1B chain		
6	IP100013683	TUBB3 tubulin $\beta$ -3 chain		
7	IP100007752	TUBB2C tubulin $\beta$ -2C chain		
8	IP100218343	TUBA1C tubulin $\alpha$ -1C chain	Apoptosis	
9	IP100021439	ACTB actin, cytoplasmic 1		
10	IP100646909	TUBA8 tubulin $\alpha$ -8 chain		
11	IP100646779	TUBB6 tubulin $\beta$ -6 chain		
12	IP100257508	DPYSL2 dihydropyrimidinase-related protein 2		
13	IP100908469	cDNA FLJ52712, highly similar to Tubulin $\beta$ -6 chain		
14	IP100410714	HBA1; HBA2 hemoglobin, $\alpha$ -2; hemoglobin, $\alpha$ -1		
15	IP100021428	ACTA1 actin, $\alpha$ skeletal muscle	Cell proliferation, Ras protein signal transduction	Chemokine signaling pathway, Attenuation of GPCR signaling, Erk1/Erk2 MAPK signaling pathway, CXCR4 signaling pathway
16	IP100026268	GNB1 guanine nucleotide-binding protein G(i)/G(s)/G(t) subunit $\beta$ -1		
17	IP100220281	GNAO1 isoform A-1 of Guanine nucleotide-binding protein G(o) subunit $\alpha$	Regulation of calcium ion transport, G-protein coupled receptor protein signaling pathway	
18	IP100021907	MBP myelin basic protein		
19	IP100024067	CLTC clathrin heavy chain 1	Regulation of calcium ion transport, G-protein coupled receptor protein signaling pathway	
20	IP100216171	ENO2 enolase 2 ( $\gamma$ , neuronal)		
21	IP100465248	ENO1 isoform $\alpha$ -enolase of A-enolase		
22	IP100220706	HBB1 hemoglobin subunit $\gamma$ -1		
23	IP100219018	GAPDH glyceraldehyde-3-phosphate dehydrogenase		
24	IP100398700	GNAO1 isoform A-2 of Guanine nucleotide-binding protein G(o) subunit $\alpha$		
25	IP100025363	GFAP glial fibrillary acidic protein		
26	IP100303476	ATP5B ATP synthase, H <sup>+</sup> transporting, mitochondrial F1 complex, $\beta$ polypeptide	Regulation of calcium-mediated signaling	Calcium signaling pathway Calcium signaling pathway Chemokine signaling pathway
27	IP100022977	CKB creatine kinase B-type		
28	IP100413140	DNM1 dynamin 1		
29	IP100154742	IGL $\lambda$ protein		
30	IP100022463	TF serotransferrin		
31	IP100220737	NCAM1 neural cell adhesion molecule 1		
32	IP100022891	SLC25A4 ADP/ATP translocase 1		
33	IP100007188	SLC25A5 ADP/ATP translocase 2		
34	IP100009532	ABAT 4-aminobutyrate aminotransferase, mitochondrial		
35	IP100012451	GNB4 guanine nucleotide-binding protein subunit $\beta$ -4		
36	IP100291006	MDH2 malate dehydrogenase 2, NAD (mitochondrial)		



Table I. Continued.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
37	IP100298497	FGB fibrinogen $\beta$ chain		
38	IP100220993	CNP 2',3'-cyclic nucleotide 3' phosphodiesterase		
39	IP100027547	DCD dermcidin		
40	IP100219446	PEBP1 phosphatidylethanolamine-binding protein 1	Regulation of cAMP-mediated signaling, regulation of MAPKKK cascade	Calcium signaling pathway
41	IP100414123	CRMP1 collapsin response mediator protein 1		
42	IP100217507	NEFM neurofilament, medium polypeptide		
43	IP100465439	ALDOA aldolase A, fructose-bisphosphate		
44	IP100029111	DPYSL3 dihydropyrimidinase-like 3		
45	IP100219813	RTN1 reticulon-1		
46	IP100001453	INA $\alpha$ -internexin		
47	IP100237671	NEFL neurofilament light polypeptide	Apoptosis	
48	IP100743576	ATP6V0A1 ATPase, H <sup>+</sup> transporting, lysosomal V0 subunit a1		
49	IP100418262	ALDOC fructose-bisphosphate aldolase C	Apoptosis	
50	IP100029751	CNTN1 contactin-1	Notch signaling pathway	
51	IP100549543	NCDN neurochondrin		
52	IP100024975	KIF15 kinesin family member 15		
53	IP100027497	GPI glucose-6-phosphate isomerase		
54	IP100010154	GDI1 GDP dissociation inhibitor 1		
55	IP100554752	PRKAR2B protein kinase, cAMP-dependent, regulatory, type II, $\beta$	Small GTPase mediated signal transduction	Apoptosis, Insulin signaling pathway
56	IP100028888	HNRNPd heterogeneous nuclear ribonucleoprotein D0		
57	IP100033025	SEPT7 septin 7		
58	IP100784156	AP2B1 adaptor-related protein complex 2, $\beta$ 1 subunit		
59	IP100026272	HIST1H2AB; HIST1H2AE histone cluster 1, H2ae; histone cluster 1, H2ab		
60	IP100219661	PLP1 proteolipid protein 1		
61	IP100015671	TUBAL3 tubulin $\alpha$ chain-like 3		
62	IP100216298	TXN thioredoxin	Cell proliferation	
63	IP100215715	CAMK2A calcium/calmodulin-dependent protein kinase II $\alpha$	Regulation of NF- $\kappa$ B transcription factor activity	ErbB signaling pathway, Calcium signaling pathway, Wnt signaling pathway
64	IP100020926	HOXA4 homeobox A4		
65	IP100022314	SOD2 superoxide dismutase 2, mitochondrial		
66	IP100024266	MGST3 microsomal glutathione S-transferase 3	Cell proliferation, apoptosis	
67	IP100382470	HSP90AA1 heat shock protein 90 kDa $\alpha$ (cytosolic), class A member 1 isoform 1		NOD-like receptor signaling pathway, pathways in cancer, Ahr signal transduction pathway, AKT signaling pathway
68	IP100019971	STXBP2 syntaxin-binding protein 2		
69	IP100289861	ZCCHC11 zinc finger CCHC domain-containing protein 11		
70	IP100013508	ACTN1 $\alpha$ -actinin-1		
71	IP100007682	ATP6V1A V-type proton ATPase catalytic subunit A		
72	IP100003925	PDHB pyruvate dehydrogenase E1 component subunit $\beta$ , mitochondrial		
73	IP100910290	Aryl hydrocarbon receptor nuclear translocator	Cell proliferation	Pathways in cancer, Ahr signal transduction pathway



Table I. Continued.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
74	IP100022488	HPX hemopexin	Regulation of protein kinase cascade, interferon- $\gamma$ -mediated signaling pathway, regulation of JAK-STAT cascade	
75	IP100023302	SYN2 synapsin-2		
76	IP100902614	USP24 ubiquitin carboxyl-terminal hydrolase 24		
77	IP100220644	PKM2 pyruvate kinase isozymes M1/M2		
78	IP100414676	HSP90AB1 heat shock protein HSP90- $\beta$	Interferon- $\gamma$ -mediated signaling pathway, type I interferon-mediated signaling pathway	NOD-like receptor signaling pathway, pathways in cancer
79	IP100647704	IGHA1 immunoglobulin heavy constant $\alpha$ 1	Cell proliferation	PPAR signaling pathway
80	IP100215747	FABP7 fatty acid-binding protein, brain		
81	IP100026053	CLDN11 claudin-11		
82	IP100025447	EEF1A1 elongation factor 1- $\alpha$		ErbB signaling pathway, Calcium signaling pathway, Wnt signaling pathway
83	IP100182944	CAMK2B calcium/calmodulin-dependent protein kinase type II $\beta$ chain		
84	IP100411486	OPALIN opalin		
85	IP100299608	PSMD1 proteasome (prosome, macropain)		
		26S subunit, non-ATPase, 1		
86	IP100299399	S100B protein S100-B	Cell proliferation	Calcium signaling pathway
87	IP100175169	ARFGAP1 ADP-ribosylation factor GTPase-activating protein 1	Small GTPase mediated signal transduction, Ras protein signal transduction	
88	IP100005614	SPTBN1 spectrin $\beta$ chain, brain 1		
89	IP100017597	MAPRE3 microtubule-associated protein RP/EB family member 3		
90	IP100175092	RNF149 ring finger protein 149		
91	IP100293613	TBK1 TANK-binding kinase 1	Regulation of protein kinase cascade, regulation of I- $\kappa$ B kinase/NF- $\kappa$ B cascade	Toll-like receptor signaling pathway, RIG-I-like receptor signaling pathway
92	IP100015029	PTGES3 prostaglandin E synthase 3		
93	IP100169383	PGK1 phosphoglycerate kinase 1		
94	IP100015148	RAP1B ras-related protein Rap-1b	Cell proliferation, small GTPase mediated signal transduction	MAPK signaling pathway, Chemokine signaling pathway
95	IP100028946	RTN3 reticulon-3	Apoptosis	
96	IP100163849	EPS15L1 epidermal growth factor receptor substrate 15-like 1	Calcium ion binding	
97	IP100645078	UBA1 ubiquitin-like modifier-activating enzyme 1		
98	IP100005705	PPP1CC $\gamma$ -1 of serine/threonine-protein phosphatase PP1- $\gamma$ catalytic subunit		
99	IP100159927	NCAN neurocan core protein	Calcium ion binding	
100	IP100003420	MAPRE2 microtubule-associated protein, RP/EB family, member 2	Cell proliferation	
101	IP100017566	FABXL7 F-box/LRR-repeat protein 7		
102	IP100027252	PHB2 prohibitin-2		
103	IP100015141	CKMT2 creatine kinase, sarcomeric mitochondrial		
104	IP100027770	SYP synaptophysin		Ubiquitin mediated proteolysis
105	IP100290035	PCDH15 protocadherin-15		Insulin signaling pathway
106	IP100027462	S100A9 S100 calcium binding protein A9	Calcium ion binding Calcium ion binding	

Table I. Continued.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
107	IP100022462	TFRC transferrin receptor protein 1	Apoptosis	Ras signaling pathway
108	IP100300020	SLC1A2 excitatory amino acid transporter 2		
109	IP100019884	ACTN2 $\alpha$ -actinin-2		
1110	IP100000875	EEF1G cDNA FLJ56389, highly similar to Elongation factor 1- $\gamma$		
1111	IP100435928	RASGRF1 Ras protein-specific guanine nucleotide-releasing factor 1		
1112	IP100790581	MPRIIP protein		
1113	IP100383660	ZNF530 zinc finger protein 530		
1114	IP100218896	ADH1A alcohol dehydrogenase 1A		
1115	IP100300341	TCEB1 transcription elongation factor B polypeptide 1		Ubiquitin mediated proteolysis, pathways in cancer
1116	IP100021891	FGG $\Gamma$ -B of Fibrinogen $\gamma$ chain		
1117	IP100747180	WDR52 WD repeat protein 52		
1118	IP100642126	KIAA1618 isoform 1 of protein ALO17		
1119	IP100164441	UNC13A unc-13 homolog A		
120	IP100027820	ESPN isoform 1 of Espin		
121	IP100185659	CCDC62 isoform 2 of Coiled-coil domain-containing protein 60		
122	IP100000816	YWHAE 14-3-3 protein epsilon		
123	IP100160552	TNR isoform 1 of Tenascin-R		
124	IP100164347	CNGB1 cyclic nucleotide gated channel $\beta$ 1 isoform b		
125	IP100166979	KIAA1239 Leucine-rich repeat and WD repeat-containing protein KIAA1239		
126	IP100217240	WDR75 WD repeat-containing protein 75		
127	IP100017704	COTL1 coactosin-like protein	Cell proliferation	
128	IP100008305	HPCAL4 hippocalcin-like protein 4		
129	IP100440493	ATP5A1 ATP synthase subunit $\alpha$ , mitochondrial		
130	IP100024547	C2orf25 chromosome 2 open reading frame 25		
131	IP100074962	ANK2 isoform 4 of Ankyrin-2		
132	IP100395663	ANKS1A ankyrin repeat and SAM domain-containing protein 1A		
133	IP100029769	HCK isoform p59-HCK of Tyrosine-protein kinase HCK		Chemokine signaling pathway, GPCR signaling
134	IP100216592	HNRNPC isoform C1 of Heterogeneous nuclear ribonucleoproteins C1/C2		
135	IP100000792	CRYZ quinone oxidoreductase		
136	IP100219806	S100A7 S100 calcium binding protein A7	S100/CaBP-9k-type, calcium binding	
137	IP100216856	ANKMY2 ankyrin repeat and MYND domain-containing protein 2		
138	IP100027434	RHOC rho-related GTP-binding protein RhoC	Small GTPase mediated signal transduction, regulation of I- $\kappa$ B kinase/NF- $\kappa$ B cascade	
139	IP100396341	C2orf67 chromosome 2 open reading frame 67		
140	IP100015785	CRB1 crumbs homolog 1	Calcium ion binding	
141	IP100893234	OBSL1 obscurin-like 1		
142	IP100028277	FTO isoform 1 of Protein fto	Anti-apoptosis	
143	IP100060800	LOC124220 uncharacterized protein UNQ773/PRO1567		
144	IP100007765	HSPA9 stress-70 protein, mitochondrial		
145	IP100852669	ZNF516 zinc finger protein 516		
146	IP100024994	TULP4 tubby-related protein 4		

Table I. Continued.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
147	IP100010466	PRKCB isoform B-I of protein kinase C $\beta$ type		MAPK signaling pathway, ErbB signaling pathway, Calcium signaling pathway, Chemokine signaling pathway, Phosphatidylinositol signaling system, Wnt signaling pathway, VEGF signaling pathway, pathways in cancer
148	IP100815811	ZNF235 zinc finger protein 235	Cell proliferation	
149	IP100163187	FSCN1 fascin homolog 1		
150	IP100793780	TMCO5B transmembrane and coiled-coil domain-containing protein 5B		
151	IP100011088	CLDN12 claudin-12		
152	IP100011986	C5orf42 chromosome 5 open reading frame 42		
153	IP100061780	ITCH itchy E3 ubiquitin protein ligase homolog	Cell proliferation	Ubiquitin mediated proteolysis
154	IP100152653	DNAH5 dynein heavy chain 5, axonemal		
155	IP100175416	PLCH11-phosphatidylinositol-4,5-bisphosphate phosphodiesterase $\beta$ -1	Calcium ion binding	
156	IP100218352	ESR1 estrogen receptor1	Estrogen receptor signaling pathway	
157	IP100791536	MCF.2 cell line derived transforming sequence-like 2	Regulation of Rho protein signal transduction, regulation of Ras protein signal transduction, regulation of small GTPase mediated signal transduction	
158	IP100009619	CADM3 isoform 2 of cell adhesion molecule 3	Apoptosis	
159	IP100179330	UBC; RPS27A; UBB ubiquitin and ribosomal protein S27a precursor		
160	IP100217776	ICK intestinal cell (MAK-like) kinase	Calcium ion binding	
161	IP100009439	SYT1 synaptotagmin-1		
162	IP100784869	DNAH10 isoform 1 of Dynein heavy chain 10, axonemal		
163	IP100020265	ANKRD20A1 ankyrin repeat domain-containing protein 20A1		MAPK signaling pathway, Chemokine signaling pathway, VEGF signaling pathway, Pathways in cancer, Ras signaling pathway
164	IP100007189	CDC42 isoform 1 of cell division control protein 42 homolog		
165	IP100032325	CSTA cystatin-A	Small GTPase mediated signal transduction	
166	IP100032808	RAB3D ras-related protein Rab-3D	Apoptosis	Ras signaling pathway
167	IP100216308	VDAC1 voltage-dependent anion-selective channel protein 1	Cell proliferation, small GTPase mediated signal transduction, Ras protein signal transduction, Rho protein signal transduction, Cdc42 protein signal transduction, G-protein coupled receptor protein signaling pathway	Calcium signaling pathway
168	IP100021841	APOA1 apolipoprotein A-1		PPAR signaling pathway
169	IP100645906	CXorf39 isoform 1 of uncharacterized protein CXorf39		
170	IP100220032	CTNND2 isoform 2 of Catenin $\delta$ -2		
171	IP100002459	ANXA6 Annexin VI isoform 2		
172	IP100184119	DNAJC6 isoform 2 of putative tyrosine-protein phosphatase auxilin	Calcium ion transport	

Table I. Continued.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
173	IP100152949	TMEM168 transmembrane protein 168	Apoptosis, MAPKKK cascade, second-messenger-mediated signaling, regulation of JAK-STAT cascade	Wnt signaling pathway, TGF- $\beta$ signaling pathway, AKT signaling pathway, Erk1/Erk2 MAPK signaling pathway
174	IP100032402	ATP8A1 isoform long of probable phospholipid-transporting ATPase 1A		
175	IP100008380	PPP2CA serine/threonine-protein phosphatase 2A catalytic subunit $\alpha$ isoform		
176	IP100219217	LDHB L-lactate dehydrogenase B chain	G-protein coupled receptor protein signaling pathway G-protein coupled receptor protein signaling pathway, activation of protein kinase C activity by G-protein coupled receptor protein signaling pathway	Phosphatidylinositol signaling system
177	IP100394855	C12orf63 chromosome 12 open reading frame 63		
178	IP100025366	CS citrate synthase, mitochondrial		
179	IP100218570	PGAM2 phosphoglycerate mutase 2		
180	IP100303484	OR52K2 olfactory receptor 52K2		
181	IP100005565	DGKQ diacylglycerol kinase $\theta$	Rab protein signal transduction, Ras protein signal transduction, small GTPase mediated signal transduction Apoptosis, regulation of protein kinase cascade, regulation of phosphoinositide 3-kinase cascade, regulation of NF- $\kappa$ B transcription factor activity	
182	IP100168218	DOK7 isoform 2 of protein Dok-7		
183	IP100154645	TBC1D15 isoform 1 of TBC1 domain family member 15		
184	IP100386494	SPPL2B isoform 1 of signal peptide peptidase-like 2B		
185	IP100465436	CAT catalase		
186	IP100007612	KCNJ1 isoform 1 of ATP-sensitive inward rectifier potassium channel 1	Small GTPase mediated signal transduction, Ras protein signal transduction	
187	IP100020153	BSN protein bassoon		
188	IP100298547	PARK7 protein DJ-1	G-protein coupled receptor protein signaling pathway, second-messenger-mediated signaling, cAMP-mediated signaling Calcium ion binding	MAPK signaling pathway, Calcium signaling pathway, Chemokine signaling pathway, Apoptosis, Wnt signaling pathway, Hedgehog signaling pathway, Insulin signaling pathway
189	IP100456969	DYNC1H1 cytoplasmic dynein 1 heavy chain 1		
190	IP100030144	PPIAL4C; PPIAL4A; PPIAL4G; PPIAL4B		
		Peptidylprolyl cis-trans isomerase A-like 4B		
191	IP100376119	PRKACB isoform 2 of cAMP-dependent protein kinase catalytic subunit $\beta$		
192	IP100025753	DSG1 desmoglein-1		
193	IP100292934	USP53 inactive ubiquitin carboxyl-terminal hydrolase 53		
194	IP100024684	MX2 interferon-induced GTP-binding protein Mx2		
195	IP100384998	NFASC isoform 7 of Neurofascin		
196	IP100217494	SMG7 sng-7 homolog, nonsense mediated mRNA decay factor ( <i>C. elegans</i> )		

Table I. Continued.

No.	Accession no.	Protein description	Biological processes (partly)	KEGG pathways (partly)
197	IP00171594	DACT1 dapper homolog 1	Wnt receptor signaling pathway	
198	IP00006612	SNAP91 isoform 1 of Clathrin coat assembly protein AP180		
199	IP00291922	PSMA5 proteasome subunit $\alpha$ type-5		
200	IP00056040	NRN1L neuritin-like protein		
201	IP00013421	GPM6B isoform 1 of neuronal membrane glycoprotein M6-b		
202	IP000738216	KIAA0947 isoform 1 of uncharacterized protein KIAA0947	Small GTPase mediated signal transduction, cAMP-mediated signaling	MAPK signaling pathway
203	IP00022133	EPB41L2 4.1G protein		
204	IP000853219	RAPGEF2 rap guanine nucleotide exchange factor 2		
205	IP00375609	JAKMIP3 janus kinase and microtubule-interacting protein 3		
206	IP00178185	BICD2 isoform 1 of protein bicaudal D homolog 2	Calcium ion binding	
207	IP00006146	SAA2 serum amyloid A2		
208	IP00014843	LRRC16A isoform 1 of Leucine-rich repeat-containing protein 16A		
209	IP00183368	SMG1 phosphatidylinositol 3-kinase-related kinase ( <i>C. elegans</i> )		
210	IP00019038	LYZ lysozyme C		
211	IP00397801	FLG2 filaggrin-2	Wnt receptor signaling pathway, calcium ion binding Apoptosis, cell proliferation, regulation of MAPKK cascade	Wnt signaling pathway, pathways in cancer
212	IP00186290	EEF2 elongation factor 2		
213	IP00783097	GARS glycyl-tRNA synthetase		
214	IP00029468	ACTR1A $\alpha$ -centractin		
215	IP00010845	NDUFS8 NADH dehydrogenase (ubiquinone) iron-sulfur protein 8, mitochondrial		
216	IP00256861	MACF1 microtubule-actin crosslinking factor 1	Apoptosis, ER-nuclear signaling pathway	
217	IP00017292	CTNNB1 Catenin $\beta$ -1		
218	IP00005966	NDUFA12 13 kDa differentiation-associated protein variant (Fragment)		
219	IP00022229	APOB apolipoprotein B-100		
220	IP00307259	DNAJC13 dnaJ homolog subfamily C member 13		
221	IP00216085	COX6B1 cytochrome c oxidase subunit VIb isoform 1	Apoptosis, ER-nuclear signaling pathway	
222	IP00022774	VCP valosin-containing protein		
223	IP00479640	Clorf113 chromosome 1 open reading frame 113		
224	IP00033019	KCNB1 potassium voltage-gated channel subfamily B member 1		
225	IP00455876	RING1 isoform 2 of E3 ubiquitin-protein ligase RING1		

Seventy-one of the proteins are correlated with cell proliferation or apoptosis according to biological processes and KEGG pathways (in the right two tiers). PEBP1, GNB1, ARFGAP1, RAPIB, RASGRF1, RHOC, PRKCB, MCF2, CDC42, RAB3D, APOA1, PPP2CA, TBC1D15, PARK7, PRKACB, RAPGEF2 and CTNNB1 are concerned with the Ras signaling pathway; CAMK2A, CAMK2B, PRKCB, PPP2CA, CTNNB1, MACF1 and DACT1 are concerned with the Wnt signaling pathway; SLC25A4, SLC25A5, PEBP1, CAMK2A, CAMK2B, S100B, PRKCB, VDACL1 and PRKACB participate in the Calcium signaling pathway; GNB1, GNB4, RAPIB, HCK, PRKCB, CDC42 and PRKACB are involved in the Chemokine signaling pathway; 15 proteins are concerned with calcium-mediated biological processes; 10 proteins are concerned with small GTPase mediated signal transduction and 8 proteins are concerned with the G-protein coupled receptor protein signaling pathway.

For biological processes, annotated proteins are particularly involved in the cell process (167, 27.5%) and the multicellular organismal process (75, 12.4%) (Fig. 4D). Most (172, 18.6%) of the annotated proteins were localized in the intracellular (Fig. 4E). Distribution of molecular mass and isoelectric points (PI) of the annotated proteins was analyzed. Molecular mass ranged between 10.19 and 620.42 kDa in size, most of them were between 10 and 60 kDa (Fig. 4F). PI of the proteins ranged between 4.35 and 11.05 with the most PIs between four and ten (Fig. 4G). To uncover the signaling pathways of the 225 annotated proteins, the protein sequences were searched against the KEGG reference pathway database. The pathways were ascribed to metabolism, genetic information processing, environmental information processing, cellular processes, organismal systems and human diseases (Fig. 5). Among them, the immune system, cancer and signal transduction were more than others. On the other hand, the specific expressed proteins related pathways displayed more differences and 71 proteins were involved in cell apoptosis or proliferation biological processes and KEGG pathways (Table I).

## Discussion

In this study, to evaluate the function of MLAA-34 in M5 cells, we used the well-characterized cell line U937. In our previous research, we reported that the MLAA-34 protein is probably a cytoplasmic protein predicted by the amino acid sequence analysis of the encoded protein (7). Here, we verified that MLAA-34 is localized in the cytoplasm and cell membrane. Western blot analysis showed that the expression of MLAA-34 differed between different cell types and was observed to be stronger in U937. Although U937 cells are generally difficult to transfect, the U937 cells were transfected with MLAA-34-Lentivirus and pGC-FU-GFP-LV. A stably transfected U937 cell line was successfully established and expressed MLAA-34 at a high level, which aided in the study exploring the effect of MLAA-34 on M5 and will be critical for further research using U937 cells and animal models. In addition, an analysis of the cell morphology, apoptosis, proliferation and cell cycle revealed that the overexpression of MLAA-34 markedly inhibited apoptosis of U937 cells. These results suggested that MLAA-34 maybe a novel anti-apoptotic factor of M5, which is consistent with the RNAi in our previous study.

The proteins that interact with MLAA-34 or CAB39L remain unclear. To analyze complex mixtures of proteins, shotgun is considered the most powerful (9,10). Using the MLAA-34 protein as bait, 256 proteins were identified and 225 of them have DAVID terms. Among these proteins, 71 proteins correlated with cell apoptosis or proliferation biological processes and KEGG pathways. Twenty-eight proteins are involved in cell apoptosis or proliferation; nine proteins are associated with the calcium signaling pathway and seven proteins participate in the chemokine signaling pathway; 17 proteins are concerned with the Ras signaling transduction pathway and 8 proteins are concerned with Wnt signaling pathway. The Ras, Wnt, calcium and chemokine signaling pathways may be involved in anti-apoptosis with MLAA-34 in U937 cells. As is known, the Ras family plays an important role in the molecular pathogenesis of myeloid leukemia, and Ras mutations have been preferentially associated with monocytic

subtypes in AML (11). The Ras and Wnt signaling pathways are known to be key anti-apoptosis pathways in AML-M5 (12). Understanding the molecular genetics of leukemia has led to an appreciation that particular molecular abnormalities give rise to specific subtypes of the disease. For example, in myeloid leukemogenesis, PML-RAR- $\alpha$  and BCR-ABL are defining features of acute promyelocytic leukemia and chronic myeloid leukemia, respectively (13). In this case, MLAA-34 may either play an important role in leukemogenesis or play a dual role in subsequent differentiation, as in the case of PML/RAR. The results suggest that MLAA-34 might be an important agent for subtype diagnosis in AML. However, an understanding of how these identified proteins or pathways interact with MLAA-34 requires further study.

In addition to the typical pathways such as pathways in cancer and apoptosis, there were several notable pathways such as the GPCR signaling, the insulin signaling pathway, the ErbB signaling pathway, the NOD-like receptor signaling pathway, the Ahr signal transduction pathway, the AKT signaling pathway, the Toll-like receptor signaling pathway, the RIG-I-like receptor signaling pathway, the ubiquitin mediated proteolysis, the hedgehog signaling pathway, the phosphatidylinositol signaling system, the PPAR signaling pathway, the VEGF signaling pathway and the TGF- $\beta$  signaling pathway worthy of further validation (Table I). Otherwise, there are some proteins mainly involved in tumorigenesis concerned with MLAA-34 as discussed below. PGK1 is secreted by tumor cells and may play a role in inhibiting tumor angiogenesis (14). GAPDH has been shown to be upregulated in several types of cancer and downregulated by chemotherapeutic drugs, and could be considered a potential target to observe the effects of bisphosphonates on cancer cells (15). In addition, GAPDH was the best control gene in the apoptosis pattern on the myeloid cell lines (16). CRMP1 is a suppressor of tumor cell invasion of the local stroma and might be a functional modulator of the Wnt signaling pathway *in vivo* (17,18). As the trigger of TBK-1 pathway, TBK1 is important for tumor angiogenesis and tumor-associated microvascular inflammation and expressed at significant levels in many solid tumors (19,20). A recent study has demonstrated that SEPT7 could function in gliomagenesis and in the suppression of glioma cell proliferation (21).

Markedly, some p53 or caspase-related proteins were also identified, such as CLTC, PPP2CA, SOD2, PARK7, HSPA9, TXN, ESR1 and YWHA. CLTC associates with p53 not only in nuclei but also in cytosol, and co-localizes with p53 at the plasma membrane in human cancer cells (22). CLTC expression enhances p53-dependent transactivation (23). As a downstream mediator of the antiproliferative effects of PPP2CA, p53 plays an important role in PPP2CA-directed cell cycle arrest and apoptosis (24). The SOD2 growth-retarding functions are at least partially due to triggering of a p53-dependent cellular senescence program (25). DJ-1 (PARK7) bound to p53 *in vitro* and *in vivo* and they were found colocalized. DJ-1 positively regulates p53 through Topors-mediated sumoylation (26). Previous studies indicated that HSPA9 could bind to p53 and sequesters it in the cytoplasm, thus providing a mechanism of inactivation of wild-type p53 and contributing to human carcinogenesis (27,28). Additional studies have shown that TXN induces p53 DNA binding activity *in vitro* and enhances p53-dependent expression of its target gene p21 and DNA repair

genes (29). Additional studies also indicated that caspases could be activated by TXN due to its disulfide reducing properties (30). ESR1 might activate caspases-8, -9 and -3 and induce tumor cell apoptosis, it also showed the downregulation of  $\beta$ -catenin signaling implicating the suppression of proliferation and metastasis of tumor cells (31,32). The cleavage of YWHAE by caspase-3 during apoptosis might contribute to cell death by preventing the association of YWHAE with Bad (33). The key event during apoptosis that is common to all pathways is the activation of caspases. P53 is a well-known tumor suppressor gene, and mutational inactivation of p53 function or deletion of the gene increases susceptibility to cancer (34-37). On the basis of these findings, we will further study the interaction between MLAA-34 and caspases or p53 to investigate the anti-apoptotic mechanisms of MLAA-34 in U937 cells.

To our knowledge, this is the first report showing the cellular localization and expression of MLAA-34 in U937 cells. We have demonstrated for the first time that the overexpression of MLAA-34 by lentivirus can significantly suppress the apoptosis of U937 cells, and a cell line stably overexpressing MLAA-34 was successfully established. Another key finding of this study is the information from proteomics evidence that MLAA-34 may be a tumor-correlated gene, and this is the first time it is revealed that the preliminary framework of proteins and pathways interlink with MLAA-34 in U937. Furthermore, it will be essential to integrate data from many different sources to obtain an accurate understanding of MLAA-34 protein networks.

Gene therapy remains the most promising, if not the only, approach to treating genetic diseases. An example of this is the use of rituximab for the treatment of lymphoma and other types of cancer. Rituximab is a mouse/human chimeric IgG(1)- $\kappa$  monoclonal antibody that targets the CD20 antigen found on the surface of malignant and normal B lymphocytes (38). Most cellular processes are performed by multiprotein complexes. The identification and analysis of their components provides insight into how the ensemble of expressed proteins (the proteome) is organized into functional units (39). Nevertheless, for a viable clinical approach, extensive research is needed in the future to regulate the expression of the target gene and improve its safety.

In conclusion, our current results provide new evidence that MLAA-34 may be a novel anti-apoptotic factor *in vitro*, and the data presented here show a strong correlation between anti-apoptosis with the upregulation of MLAA-34. In addition, preliminary proteomic analysis suggests that a number of genes belonging to different signaling pathways may be involved in apoptosis in U937 cells in association with MLAA-34, which would disclose a novel cross-link between MLAA-34 and the Ras, Wnt, calcium and chemokine signaling pathways. Findings of the present study will lead to a better understanding of the mechanisms involved in M5, and MLAA-34 may serve as a potential novel marker for the early diagnosis and gene therapy of M5.

## Acknowledgements

This study was supported by the National Natural Science Foundation of China under award nos. 30971284, 81000219 and 18110021.

## References

- Ossenkoppele GJ, Graveland WJ, Sonneveld P, Daenen SM, Biesma DH, Verdonck LF, Schaafsma MR, Westveer PH, Peters GJ, Noordhuis P, Muus P, Selleslag D, van der Holt B, Delforge M, Lowenberg B and Verhoef GE: The value of fludarabine in addition to ARA-C and G-CSF in the treatment of patients with high-risk myelodysplastic syndromes and AML in elderly patients. *Blood* 103: 2908-2913, 2004.
- Faderl S, Gandhi V, O'Brien S, Bonate P, Cortes J, Estey E, Beran M, Wierda W, Garcia-Manero G, Ferrajoli A, Estrov Z, Giles FJ, Du M, Kwari M, Keating M, Plunkett W and Kantarjian H: Results of a phase 1-2 study of clofarabine in combination with cytarabine (ara-C) in relapsed and refractory acute leukemias. *Blood* 105: 940-947, 2005.
- Fiedler W, Serve H, Döhner H, Schwittay M, Ottmann OG, O'Farrell AM, Bello CL, Allred R, Manning WC, Cherrington JM, Louie SG, Hong W, Brega NM, Massimini G, Scigalla P, Berdel WE and Hossfeld DK: A phase 1 study of SU11248 in the treatment of patients with refractory or resistant acute myeloid leukemia (AML) or not amenable to conventional therapy for the disease. *Blood* 105: 986-993, 2005.
- Niitsu N, Yamamoto-Yamaguchi Y, Kasukabe T, Okabe-Kado J, Umeda M and Honma Y: Antileukemic efficacy of 2-deoxycorformycin in monocytic leukemia cells. *Blood* 96: 1512-1516, 2000.
- Tallman MS, Kim HT, Paietta E, Bennett JM, Dewald G, Cassileth PA, Wiernik PH and Rowe JM: Acute monocytic leukemia (French-American-British classification M5) does not have a worse prognosis than other subtypes of acute myeloid leukemia: a report from the Eastern Cooperative Oncology Group. *J Clin Oncol* 22: 1276-1286, 2004.
- Chen G, Zhang W, Cao X, Li F, Liu X and Yao L: Serological identification of immunogenic antigens in acute monocytic leukemia. *Leuk Res* 29: 503-509, 2005.
- Zhang PY, Zhang WG, He AL, Wang JL and Li WB: Identification and functional characterization of the novel acute monocytic leukemia associated antigen MLAA-34. *Cancer Immunol Immunother* 58: 281-290, 2009.
- Zhao J, He A, Zhang W, Meng X and Gu L: Quantitative assessment of MLAA-34 expression in diagnosis and prognosis of acute monocytic leukemia. *Cancer Immunol Immunother* 60: 587-597, 2011.
- Lu B, McClatchy DB, Kim JY and Yates JR III: Strategies for shotgun identification of integral membrane proteins by tandem mass spectrometry. *Proteomics* 8: 3947-3955, 2008.
- He P, He HZ, Dai J, Wang Y, Sheng QH, Zhou LP, Zhang ZS, Sun YL, Liu F, Wang K, Zhang JS, Wang HX, Song ZM, Zhang HR, Zeng R and Zhao X: The human plasma proteome: analysis of Chinese serum using shotgun strategy. *Proteomics* 5: 3442-3453, 2005.
- Bowen DT, Frew ME, Hills R, Gale RE, Wheatley K, Groves MJ, Langabeer SE, Kottaridis PD, Moorman AV, Burnett AK and Linch DC: RAS mutation in acute myeloid leukemia is associated with distinct cytogenetic subgroups but does not influence outcome in patients younger than 60 years. *Blood* 106: 2113-2119, 2005.
- Morgan MA, Dolp O and Reuter CW: Cell-cycle-dependent activation of mitogen-activated protein kinase kinase (MEK-1/2) in myeloid leukemia cell lines and induction of growth inhibition and apoptosis by inhibitors of RAS signaling. *Blood* 97: 1823-1834, 2001.
- Pearn L, Fisher J, Burnett AK and Darley RL: The role of PKC and PDK1 in monocyte lineage specification by Ras. *Blood* 109: 4461-4469, 2007.
- Lay AJ, Jiang XM, Kisker O, Flynn E, Underwood A, Condron R and Hogg PJ: Phosphoglycerate kinase acts in tumour angiogenesis as a disulphide reductase. *Nature* 408: 869-873, 2000.
- Valenti MT, Bertoldo F, Dalle Carbonare L, Azzarello G, Zenari S, Zanatta M, Balducci E, Vinante O and Lo Cascio V: The effect of bisphosphonates on gene expression: GAPDH as a housekeeping or a new target gene? *BMC Cancer* 6: 49, 2006.
- Ullmannová V and Haskovec C: The use of housekeeping genes (HKG) as an internal control for the detection of gene expression by quantitative real-time RT-PCR. *Folia Biol (Praha)* 49: 211-216, 2003.
- Pan SH, Chao YC, Hung PF, Chen HY, Yang SC, Chang YL, Wu CT, Chang CC, Wang WL, Chan WK, Wu YY, Che TF, Wang LK, Lin CY, Lee YC, Kuo ML, Lee CH, Chen JJ, Hong TM and Yang PC: The ability of LCRMP-1 to promote cancer invasion by enhancing filopodia formation is antagonized by CRMP-1. *J Clin Invest* 121: 3189-3205, 2011.



18. Stelzl U, Worm U, Lalowski M, Haenig C, Brembeck FH, Goehler H, Stroedicke M, Zenkner M, Schoenherr A, Koepfen S, Timm J, Mintzlaff S, Abraham C, Bock N, Kietzmann S, Goedde A, Toksöz E, Droege A, Krobitsch S, Korn B, Birchmeier W, Lehrach H and Wanker EE: A human protein-protein interaction network: a resource for annotating the proteome. *Cell* 122: 957-968, 2005.
19. Korherr C, Gille H, Schäfer R, Koenig-Hoffmann K, Dixelius J, Egland KA, Pastan I and Brinkmann U: Identification of proangiogenic genes and pathways by high-throughput functional genomics: TBK1 and the IRF3 pathway. *Proc Natl Acad Sci USA* 103: 4240-4245, 2006.
20. Czabanka M, Korherr C, Brinkmann U and Vajkoczy P: Influence of TBK-1 on tumor angiogenesis and microvascular inflammation. *Front Biosci* 13: 7243-7249, 2008.
21. Jia ZF, Huang Q, Kang CS, Yang WD, Wang GX, Yu SZ, Jiang H and Pu PY: Overexpression of septin 7 suppresses glioma cell growth. *J Neurooncol* 98: 329-340, 2010.
22. Endo Y, Sugiyama A, Li SA, Ohmori K, Ohata H, Yoshida Y, Shibuya M, Takei K, Enari M and Taya Y: Regulation of clathrin-mediated endocytosis by p53. *Genes Cells* 13: 375-386, 2008.
23. Enari M, Ohmori K, Kitabayashi I and Taya Y: Requirement of clathrin heavy chain for p53-mediated transcription. *Genes Dev* 20: 1087-1099, 2006.
24. Ofek P, Ben-Meir D, Kariv-Inbal Z, Oren M and Lavi S: Cell cycle regulation and p53 activation by protein phosphatase 2C alpha. *J Biol Chem* 278: 14299-14305, 2003.
25. Behrend L, Mohr A, Dick T and Zwacka RM: Manganese superoxide dismutase induces p53-dependent senescence in colorectal cancer cells. *Mol Cell Biol* 25: 7758-7769, 2005.
26. Shinbo Y, Taira T, Niki T, Iguchi-Ariga SM and Ariga H: DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3. *Int J Oncol* 26: 641-648, 2005.
27. Grover A, Priyandoko D, Gao R, Shandilya A, Widodo N, Bisaria VS, Kaul SC, Wadhwa R and Sundar D: Withanone binds to mortalin and abrogates mortalin-p53 complex: Computational and experimental evidence. *Int J Biochem Cell Biol* 44: 496-504, 2012.
28. Wadhwa R, Yaguchi T, Hasan MK, Mitsui Y, Reddel RR and Kaul SC: Hsp70 family member, mot-2/mthsp70/GRP75, binds to the cytoplasmic sequestration domain of the p53 protein. *Exp Cell Res* 274: 246-253, 2002.
29. Ueno M, Masutani H, Arai RJ, Yamauchi A, Hirota K, Sakai T, Inamoto T, Yamaoka Y, Yodoi J and Nikaido T: Thioredoxin-dependent redox regulation of p53-mediated p21 activation. *J Biol Chem* 274: 35809-35815, 1999.
30. Ueda S, Nakamura H, Masutani H, Sasada T, Yonehara S, Takabayashi A, Yamaoka Y and Yodoi J: Redox regulation of caspase-3(-like) protease activity: regulatory roles of thioredoxin and cytochrome c. *J Immunol* 161: 6689-6695, 1998.
31. Hsu HH, Cheng SF, Chen LM, Liu JY, Chu CH, Weng YJ, Li ZY, Lin CS, Lee SD, Kuo WW and Huang CY: Over-expressed estrogen receptor- $\alpha$  upregulates hTNF- $\alpha$  gene expression and downregulates  $\beta$ -catenin signaling activity to induce the apoptosis and inhibit proliferation of LoVo colon cancer cells. *Mol Cell Biochem* 289: 101-109, 2006.
32. Kouzmenko AP, Takeyama K, Ito S, Furutani T, Sawatsubashi S, Maki A, Suzuki E, Kawasaki Y, Akiyama T, Tabata T and Kato S: Wnt/ $\beta$ -catenin and estrogen signaling converge in vivo. *J Biol Chem* 279: 40255-40258, 2004.
33. Won J, Kim DY, La M, Kim D, Meadows GG and Joe CO: Cleavage of 14-3-3 protein by caspase-3 facilitates bad interaction with Bcl-x(L) during apoptosis. *J Biol Chem* 278: 19347-19351, 2003.
34. Vousden KH and Prives C: p53 and prognosis: new insights and further complexity. *Cell* 120: 7-10, 2005.
35. Vousden KH and Lane DP: p53 in health and disease. *Nat Rev Mol Cell Biol* 8: 275-283, 2007.
36. Pietsch EC, Sykes SM, McMahon SB and Murphy ME: The p53 family and programmed cell death. *Oncogene* 27: 6507-6521, 2008.
37. Kruse JP and Gu W: Modes of p53 regulation. *Cell* 137: 609-622, 2009.
38. Plosker GL and Figgitt DP: Rituximab: a review of its use in non-Hodgkin's lymphoma and chronic lymphocytic leukaemia. *Drugs* 63: 803-843, 2003.
39. Gavin AC, Bösch M, Krause R, Grandi P, Marzioch M, Bauer A, Schultz J, Rick JM, Michon AM, Cruciat CM, Remor M, Höfert C, Schelder M, Brajenovic M, Ruffner H, Merino A, Klein K, Hudak M, Dickson D, Rudi T, Gnau V, Bauch A, Bastuck S, Huhse B, Leutwein C, Heurtier MA, Copley RR, Edelmann A, Querfurth E, Rybin V, Drewes G, Raida M, Bouwmeester T, Bork P, Seraphin B, Kuster B, Neubauer G and Superti-Furga G: Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* 415: 141-147, 2002.