

Table SI. Association between *H. pylori* and IM.

<i>H. pylori</i>				
Group	Positive (%)	Negative (%)	OR (95%CI)	P-value
Non-IM	18,801 (31.2)	41,389 (68.8)		
IM	10,258 (37.3)	17,251 (62.7)	1.31 (1.27-1.35)	<0.001

IM, intestinal metaplasia; *H. pylori*, *Helicobacter pylori*; OR, odds ratio; CI, confidence interval.

Table SII. The average age of the patients who obtained *H. pylori* status by histopathological methods in each calendar year.

Year	n	<i>H. pylori</i> positive	<i>H. pylori</i> negative	Average age
2009	3,819	1,366	2,453	52.41±14.56
2010	3,844	1,312	2,532	52.96±14.16
2011	8,572	3,183	5,389	52.84±14.51
2012	8,274	2,891	5,383	53.13±14.55
2013	8,835	3,042	5,793	53.06±14.08
2014	8,294	2,933	5,361	52.65±13.63
2015	8,973	3,049	5,924	52.70±13.72
2016	10,589	3,247	7,342	52.89±13.65
2017	12,445	3,532	8,913	52.91±13.66
2018	14,054	4,504	9,550	53.01±13.49

One-way analysis of variance test, $F=1.53$, $P=0.132$; post-hoc multiple comparisons (Student-Newman-Keuls test), all $P>0.05$. Values are expressed as the mean ± standard deviation. *H. pylori*, *Helicobacter pylori*.