

Improvement of metabolic syndrome by irbesartan via the PPAR γ /HGF pathway in apolipoprotein E knockout mice

AMARNATH CHATTERJEE^{1*}, HIROSHI KUSUNOKI^{1,2*}, YOSHIAKI TANIYAMA^{1,2},
HIROMI RAKUGI² and RYUICHI MORISHITA¹

Departments of ¹Clinical Gene Therapy and ²Geriatric Medicine and Nephrology,
Osaka University, Graduate School of Medicine, Suita 565-0871, Japan

Received July 28, 2012; Accepted October 4, 2012

DOI: 10.3892/br.2012.28

Abstract. Irbesartan, a partial agonist of peroxisome proliferators activated receptor- γ (PPAR γ), has been reported to improve insulin resistance and lipid profile in patients with diabetes mellitus or metabolic syndrome (MS). However, the down effectors of PPAR γ have yet to be elucidated. Thus, in this study, we focused on the role of the hepatocyte growth factor (HGF) in the anti-metabolic effects of irbesartan, using apolipoprotein E (ApoE) knockout (KO) mice. ApoE KO mice placed on a high-fat diet (HFD) for 12 weeks were divided into four groups: i) the control (HFD only), ii) the HFD + irbesartan (5 mg/kg/day), iii) the HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day) and iv) the HFD + irbesartan + anti-HGF neutralizing antibody (200 μ g/week). The liver and epididymal adipose tissues were evaluated histologically. Serum adiponectin and HGF levels were also measured by ELISA. Fatty liver (as detected by oil-red O staining) and macrophage infiltration were markedly reduced by irbesartan. Irbesartan treatment also reduced macrophage infiltration into epididymal adipose tissue and hypertrophy of adipocytes. However, these effects of irbesartan were attenuated by GW9662 as well as by anti-HGF neutralizing antibody. Serum and hepatic HGF levels were also markedly increased by irbesartan, whereas GW9662 decreased the HGF level. In conclusion, irbesartan, an angiotensin (Ang) receptor blocker (ARB) and partial agonist of PPAR γ (metabosartan), demonstrated a reduction in fatty liver and chronic inflammation, such as macrophage infiltration, beyond its blood pressure-lowering effect. These favorable characteristics of irbesartan

might be due to local HGF activation through its partial PPAR γ agonistic action, in addition to Ang II blockade. Upregulation of local HGF by irbesartan might provide a novel advantage in a strategy for the prevention and treatment of cardiovascular diseases (CVDs).

Introduction

Metabolic syndrome (MS), an extremely frequent condition characterized by dyslipidemia, insulin resistance, abdominal obesity and hypertension, is associated with an elevated risk of cardiovascular events. Of the risk factors for MS, high blood pressure is the most significant, since hypertension is closely associated with insulin resistance and dyslipidemia. Angiotensin (Ang) type I receptor (AT1R) blockers (ARBs) have been widely used in the treatment of hypertension and hypertension-related cardiovascular end-organ damage (1,2), since treatment with ARBs is known to improve the clinical manifestations of MS. However, new antihypertensive drugs should be designed to affect cell and biochemical mechanisms contributing to increased blood pressure and to also address disordered lipid metabolism in a more favorable manner.

To this end, peroxisome proliferator-activated receptor- γ (PPAR γ) is in the center of interest, since ligands for PPAR γ improve insulin sensitivity, reduce triglyceride levels and decrease the risk for atherosclerosis (3-5). Notably, among the approved ARBs, irbesartan and telmisartan were demonstrated to constitute a unique subset of ARBs that are also capable of activating PPAR γ (2,6). Findings by Schupp *et al* (2) demonstrated that irbesartan and telmisartan could be considered partial and selective PPAR modulators (SPPARMs) (2,7). The SPPARM approach has also been suggested to be a method to avoid unwanted complications of PPAR γ ligands, such as obesity and edema (7). In patients with MS, irbesartan markedly reduced blood pressure, and was associated with a reduction in cardiovascular risk factors, such as high-density lipoprotein (HDL) cholesterol, serum triglyceride, fasting blood glucose and waist circumference (8). Beneficial therapeutic effects of irbesartan in hypertensive patients with MS might be mediated via the AT1 receptor antagonistic and partial PPAR γ agonistic actions. Based upon these observations, ARBs with partial PPAR γ agonistic activity, such as

Correspondence to: Professor Ryuichi Morishita or Dr Yoshiaki Taniyama, Department of Clinical Gene Therapy, Osaka University Graduate School of Medicine, 2-2 Yamada-oka, Suita 565-0871, Japan

E-mail: morishit@cgt.med.osaka-u.ac.jp

E-mail: taniyama@cgt.med.osaka-u.ac.jp

*Contributed equally

Key words: irbesartan, proliferator-activated receptor, metabolic syndrome, angiotensin receptor blocker

irbesartan are now termed ‘metabosartans’. Hepatocyte growth factor (HGF) was previously shown to be a downstream effector of PPAR γ agonists (9). Thus, in this study, we investigated the effects of a metabosartan, irbesartan, on fatty liver and the hypertrophy of adipocytes in apolipoprotein E (ApoE) knockout (KO) mice.

Materials and methods

Animals, diets and drug treatment. ApoE KO mice were used as a murine model exhibiting fatty liver and hyperlipidemia, which are typical phenotypes of MS. The experiments carried out in animals were performed in accordance with guidelines laid down by the Ethics Committee for Animal Experiments of the Osaka University Graduate School of Medicine (Osaka, Japan). Male ApoE KO mice with a C57/BL6 background were obtained from the Jackson Laboratory (Bar Harbor, ME, USA). Irbesartan was donated by Shionogi Pharma, Inc. (Osaka, Japan). HGF neutralizing antibody was purchased from Kringle Pharma, Inc. (Osaka, Japan). GW9662 was purchased from Cayman Chemical Company (Ann Arbor, MI, USA).

Ten-week old mice ($n=6$ per group) were fed a high-fat diet (HFD) (MF plus 0.5% (wt/wt) cholesterol and 10% yashi oil; Oriental Yeast Co., Ltd., Tokyo, Japan). Mice were divided into four groups: i) the control group, ApoE KO mice with HFD only; ii) the irbesartan group, ApoE KO mice with HFD and 5 mg/kg/day irbesartan; iii) the HGF antibody group, ApoE KO mice with HFD, 5 mg/kg/day irbesartan and 200 μ g/week HGF neutralizing antibody and iv) the GW9662 group, ApoE KO mice with HFD, 5 mg/kg/day irbesartan and 0.5 mg/kg/day GW9662 (Fig. 1). Drugs were dissolved in water and administered *ad libitum*. There were 6 mice/group, which were housed in the animal facilities of the Osaka University (Osaka, Japan). The mice had free access to water and food during the experimental period. Following three months of drug and HFD treatment, the mice were sacrificed. Samples of epididymal adipose tissue and liver were evaluated.

Measurement of HGF and adiponectin. HGF concentration was measured by an enzyme-linked immunosorbent assay (ELISA), using an IMMUNIS mouse HGF ELISA kit (Institute of Immunology Co., Ltd., Tokyo, Japan). Mouse liver samples were disintegrated with IMMUNIS HGF extraction buffer, using a Multi-beads shocker (Yasui Kikai, Osaka, Japan) at 2000 x g for 15 sec. Homogenates were centrifuged at 14,000 x g for 30 min. The supernatant was used for the HGF assay, according to the manufacturer's instructions. Serum adiponectin level was also measured using an ELISA kit (Otsuka Pharmaceuticals, Co., Ltd., Tokyo, Japan).

Immunohistochemical staining. Tissues fixed with 4% formalin and embedded in paraffin were subjected to immunohistochemical staining, as described previously (10). Immunostaining of F4/80 was performed using anti-mouse F4/80 antibody (ab6640; Abcam, Cambridge, UK). Immunostained images were quantified using the NIH Image J software (<http://rsb.info.nih.gov/ij/>) and then analyzed visually under a light microscope by two investigators blinded to treatment.




Figure 1. Experimental protocol is shown. ApoE KO mice placed on a high-fat diet (HFD) for 12 weeks were divided into four groups: i) control (HFD only); ii) HFD + irbesartan (5 mg/kg/day); iii) HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day) and iv) HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week). ApoE, apolipoprotein E; KO, knockout; PPAR γ , peroxisome proliferators activated receptor- γ .

Statistical analysis. Data were expressed as the mean \pm standard error of the mean (SEM). Comparisons were made using analysis of variance (ANOVA) followed by Tukey's simultaneous multiple comparisons. $P<0.05$ was considered to indicate a statistically significant difference.

Results

Effects of irbesartan on adipose tissue in ApoE KO mice. Administration of irbesartan at 5 mg/kg/day for 12 weeks lowered the epididymal adipose tissue weight. Histological examination showed a smaller adipocyte size and crown-like structures (formed by the gathering of macrophages), as detected by F4/80 (a macrophage marker)-positive areas in the irbesartan-treated group (Fig. 2A). Consistent with the histological findings, adipocyte diameter was markedly decreased by irbesartan (Fig. 2B) and the F4/80-positive areas were also markedly decreased by irbesartan (Fig. 2C). These beneficial effects of irbesartan were attenuated by treatment with anti-HGF antibody or an inhibitor of PPAR γ , GW9662.

Irbesartan also reduced the weight of epididymal adipose tissue (Fig. 3A). This effect was also attenuated by anti-HGF antibody or GW9662 treatment (Fig. 3A). Notably, irbesartan significantly increased the serum adiponectin level, while anti-HGF antibody or GW9662 treatment decreased the serum adiponectin level (Fig. 3B).

Effect of irbesartan on fatty liver in ApoE KO mice. In this model, the control group demonstrated findings of severe fatty liver, such as marked deposition of lipid, as assessed by oil-red O staining (Fig. 4A). Notably, irbesartan markedly reduced lipid accumulation, while anti-HGF antibody or GW9662 treatment reversed the beneficial effect of irbesartan (Fig. 4A and B). Irbesartan also decreased serum aspartate transaminase (AST) as compared to the control (Table I), whereas anti-HGF antibody or GW9662 treatment attenuated the changes induced by irbesartan. Although no significant changes were detected in the low-density lipoprotein (LDL)

Table I. Serum parameters in each group.

Parameter	Irbesartan			
	Control	(-)	GW9662	HGF-Ab
AST (U/l)	182.0±10.3	133.3±5.7 ^a	168.8±5.7 ^b	161.5±4.8 ^b
ALT (U/l)	30.5±2.3	29.6±1.7	30.9±2.5	31.4±2.9
LDL-chol (mg/dl)	1034±56	925±16	991±73	1006±29
T-chol (mg/dl)	1248±21	1069±48	1095±89	1194±47
FFA (mEq/l)	1.58±0.19	1.13±0.02 ^a	1.29±0.05	1.43±0.08 ^b
HDL-chol (mg/dl)	62.2±3.7	88.2±4.4 ^a	58.0±5.7 ^a	66.3±6.6

^aP<0.05 vs. control. ^bP<0.05 vs. irbesartan only. Data are shown as the mean ± standard error of the mean. HGF-Ab, hepatocyte growth factor neutralizing antibody; AST, aspartate transaminase; ALT, alanine aminotransferase; LDL-chol, low-density lipoprotein cholesterol; T-chol, total cholesterol; FFA, free fatty acid; HDL-chol, high-density lipoprotein cholesterol.




Figure 2. Effects of irbesartan, GW9662 and hepatocyte growth factor neutralizing antibody (HGF-Ab) on F4/80 staining and adipocyte diameter in adipose tissue. (A) Typical micrographs of immunostaining for F4/80 in periodic cross-sections of epididymal adipose tissue after 12 weeks of high-fat diet (HFD). Brown-stained area shows F4/80 protein-positive area; (a) control (HFD only), (b) HFD + irbesartan (5 mg/kg/day), (c) HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day) and (d) HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week); bar, 100 μ m. (B) Adipocyte diameter (μ m). (C) Quantitative data of immunohistochemical staining for macrophages (F4/80) (%). Control, HFD only; (-), HFD + irbesartan (5 mg/kg/day); GW9662, HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day); HGF-Ab, HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week). *P<0.01 vs. control, †P<0.05 vs. irbesartan only. Data are shown as the mean ± standard error of the mean. PPAR γ ; peroxisome proliferators activated receptor- γ .




Figure 3. Effects of irbesartan, GW9662 and hepatocyte growth factor neutralizing antibody (HGF-Ab) on serum adiponectin and epididymal adipose tissue weight. (A) Serum concentration of adiponectin (mg/ml). (B) Epididymal adipose tissue weight [g/100 g of body weight (BW)] after 12 weeks of high-fat diet (HFD). Control, HFD only; (-), HFD + irbesartan (5 mg/kg/day); GW9662, HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day); HGF-Ab, HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week). *P<0.01 vs. control, †P<0.05 vs. irbesartan only. PPAR γ ; peroxisome proliferators activated receptor- γ .

Table II. Body weight, blood pressure and food intake in each group.

	Irbesartan			
	Control	(-)	GW9662	HGF-Ab
Body weight (g)				
PreHFD	27.6 \pm 0.9	27.8 \pm 0.9	27.4 \pm 0.6	27.6 \pm 0.7
PostHFD	35.7 \pm 1.6	31.1 \pm 0.8	33.3 \pm 1.6	33.2 \pm 1.3
Systolic BP (mmHg)				
PreHFD	103.8 \pm 2.0	104.3 \pm 2.2	108.2 \pm 0.9	105.5 \pm 3.4
PostHFD	104.7 \pm 2.1	101.3 \pm 2.3	103.5 \pm 2.1	107.7 \pm 3.1
Diastolic BP (mmHg)				
PreHFD	70.3 \pm 1.6	68.5 \pm 1.9	69.8 \pm 2.0	68.7 \pm 3.3
PostHFD	67.3 \pm 2.9	65.0 \pm 2.3	67.5 \pm 1.5	66.5 \pm 2.1
Food intake (g/day)	2.06 \pm 0.11	2.05 \pm 0.10	2.09 \pm 0.11	2.08 \pm 0.09

HGF-Ab, hepatocyte growth factor neutralizing antibody; HFD, high-fat diet; BP, blood pressure.




Figure 4. Effects of irbesartan, GW9662 and hepatocyte growth factor neutralizing antibody (HGF-Ab) on liver. (A) Upper panels: typical micrographs of periodic cross-sections of liver with oil-red O staining after 12 weeks of high-fat diet (HFD). (a) Control (HFD only), (b) HFD + irbesartan (5 mg/kg/day), (c) HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day) and (d) HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week); bar, 100 μ m. Lower panels: typical micrographs of immunostaining for macrophages (F4/80) in periodic cross-sections of liver tissue after 12 weeks of HFD. Brown-stained area shows F4/80 protein-positive area; (e) control (HFD only), (f) HFD + irbesartan (5 mg/kg/day), (g) HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day) and (h) HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week); bar, 100 μ m. (B) Quantitative data for percentage of oil deposits (oil-red O-positive area) in liver. (C) Quantitative data of immunohistochemical staining for macrophages (F4/80) in liver (%). (D) Liver weight [g/100g of body weight (BW)]. Control, HFD only; (-), HFD + irbesartan (5 mg/kg/day); GW9662, HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day); HGF-Ab, HFD + irbesartan + neutralizing antibody against HGF (200 μ g/week). * P <0.01 vs. control, † P <0.05 vs. irbesartan only. PPAR γ ; peroxisome proliferators activated receptor- γ .

cholesterol and total cholesterol levels, the free fatty acid (FFA) level was markedly decreased and HDL cholesterol level was markedly increased by irbesartan (Table I). The changes were also reversed by anti-HGF antibody or GW9662 treatment. No statistically significant changes were observed in body weight

(BW), blood pressure or food intake in the groups (Table II). Therefore, this beneficial effect of irbesartan was not due to blood pressure lowering or a decrease in food intake.

As resident macrophages are crucially involved in the progression of fatty liver, immunostaining with F4/80 was




Figure 5. Effect of irbesartan and GW9662 on hepatocyte growth factor (HGF) protein levels in serum and liver. (A) Serum HGF protein concentration (ng/ml). (B) HGF protein concentration in liver (ng/mg tissue protein). Control, HFD only, irbesartan, HFD + irbesartan (5 mg/kg/day); irbesartan + GW9662, HFD + irbesartan + GW9662, a PPAR γ antagonist (0.5 mg/kg/day). *P<0.01 vs. control, †P<0.01 vs. irbesartan only. HFD, high-fat diet; PPAR γ ; peroxisome proliferators activated receptor- γ .




Figure 6. Summary of findings. First, irbesartan activates PPAR γ and then hepatocyte growth factor (HGF) is upregulated and exerts favorable effects against pathological characteristics of metabolic syndrome (MS). When this signaling is blocked, these beneficial effects are reduced. PPAR γ ; peroxisome proliferators activated receptor- γ .

performed. As shown in Fig. 4C, irbesartan markedly reduced the infiltration of macrophages in the liver, as detected by F4/80 staining. Irbesartan also reduced the liver weight (Fig. 4D). Consistently, these effects of irbesartan were also reversed by anti-HGF antibody or GW9662 treatment (Fig. 4C and D).

Effect of irbesartan on HGF protein levels in serum and liver. To confirm the role of HGF in the reduction of fatty liver by irbesartan, we measured the tissue HGF protein level in the liver. As shown in Fig. 5A, irbesartan markedly increased hepatic HGF protein as compared to the control, while GW9662 treatment markedly inhibited the increase in the HGF level induced by irbesartan. Similar changes were observed in the serum HGF level. As shown in Fig. 5B, the serum HGF level was markedly increased by irbesartan, while GW9662 treatment reduced the increase.

Discussion

The present study has demonstrated that irbesartan significantly reduced fatty liver and chronic inflammation, such as macrophage infiltration through the PPAR γ -HGF pathway, beyond

its blood pressure-lowering effect. In general, adiponectin is a well-known downstream effector of PPAR γ (12,13). The present study demonstrated that irbesartan markedly increased adiponectin expression. However, another downstream effector of PPAR γ , HGF, is also likely to act as a ‘guardian’ in MS, in addition to adiponectin. Previous studies suggested that HGF mediates multiple various biological effects in various cells including anti-fibrotic, anti-inflammatory and anti-apoptotic activities (14). The present study clearly demonstrated that the anti-metabolic effects of irbesartan were largely mediated by the PPAR γ -HGF pathway, with the exception of Ang II blockade.

Recent data have suggested that the presence of NAFLD in type 2 diabetes mellitus is linked to an increased risk of cardiovascular disease (CVD) independent of MS. In addition, the prevalence of carotid plaques has been reported to be higher in patients with NAFLD compared to the normal controls, regardless of classical cardiovascular risk factors (15). Therefore, NAFLD should be considered an independent risk factor for CVD. However, currently there are limited therapeutic options to treat NAFLD. To this end, the present findings suggesting that irbesartan reduced fatty liver and adipocyte hypertrophy may provide a new therapeutic option to treat NAFLD.

The manner in which PPAR γ activates local HGF has yet to be elucidated. PPAR γ has been reported to bind to the putative peroxisome proliferator response element (PPRE) in the promoter region of the HGF gene, resulting in an increase in HGF gene transcription, mRNA expression and protein secretion (9). Our group of investigators have reported that telmisartan (another PPAR γ agonistic ARB), but not losartan (a classical ARB), improved endothelial dysfunction and fatty liver, due to an increased tissue HGF level (16). More directly, using AT1R KO mice, telmisartan, but not losartan, exhibited renal protective effects in a unilateral ureteral obstruction model (10). The partial PPAR γ agonistic effect of irbesartan might provide an additional advantage as a strategy for the prevention and treatment of CVD, beyond its blood pressure-lowering effect through Ang II blockade. Notably, although irbesartan as well as telmisartan selectively induced PPAR γ target genes, distinctive gene expression profiles have also been reported in 3T3-L1 adipocytes (2). The X-ray crystal structure exhibited various binding modes to PPAR γ between

irbesartan and telmisartan (6). In addition, irbesartan is also known to activate PPAR α . Irbesartan has been reported to improve hepatic steatosis and hepatic fibrosis by activating PPAR α , in NASH model FLS ob/ob mice (17). Another study also demonstrated that irbesartan upregulated PPAR α in obese Koletsky (*fa^k/fa^k*) rats and improved their metabolic disorders (18). In our model, PPAR α activation may also act to improve the pathological characteristics of the MS.

Overall, irbesartan, an ARB with partial PPAR γ agonistic activity (metabosartan), demonstrated a reduction in fatty liver and chronic inflammation, such as macrophage infiltration, beyond its blood pressure-lowering effect (Fig. 6). These favorable characteristics of irbesartan might be due to local HGF activation through partial PPAR γ agonistic activity, in addition to Ang II blockade. Upregulation of local HGF by irbesartan might provide a novel advantage in a strategy for the prevention and treatment of CVDs.

Acknowledgements

This study was partially supported by a Grant-in-Aid from the Organization for Pharmaceutical Safety and Research, a Grant-in-Aid from the Ministry of Public Health and Welfare, a Grant-in-Aid from the Japan Promotion of Science and by special coordination funds of the Ministry of Education, Culture, Sports, Science and Technology of the Japanese Government.

References

- de Gasparo M, Catt KJ, Inagami T, Wright JW and Unger T: International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol Rev* 52: 415-472, 2000.
- Schupp M, Janke J, Clasen R, Unger T and Kintscher U: Angiotensin type I receptor blockers induce peroxisome proliferator-activated receptor-gamma activity. *Circulation* 109: 2054-2057, 2004.
- Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM and Kliewer SA: An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 270: 12953-12956, 1995.
- Rosen ED and Spiegelman BM: PPARgamma: a nuclear regulator of metabolism, differentiation, and cell growth. *J Biol Chem* 276: 37731-37734, 2001.
- Picard F and Auwerx J: PPAR(gamma) and glucose homeostasis. *Annu Rev Nutr* 22: 167-197, 2002.
- Benson SC, Pershadsingh HA, Ho CI, Chittiboyina A, Desai P, Pravenec M, Qi N, Wang J, Avery MA and Kurtz TW: Identification of telmisartan as a unique angiotensin II receptor antagonist with selective PPARgamma-modulating activity. *Hypertension* 43: 993-1002, 2004.
- Schupp M, Clemenz M, Gineste R, Witt H, Janke J, Helleboid S, Hennuyer N, Ruiz P, Unger T, Staels B and Kintscher U: Molecular characterization of new selective peroxisome proliferator-activated receptor gamma modulators with angiotensin receptor blocking activity. *Diabetes* 54: 3442-3452, 2005.
- Parhofer KG, Münzel F and Krekler M: Effect of the angiotensin receptor blocker irbesartan on metabolic parameters in clinical practice: the DO-IT prospective observational study. *Cardiovasc Diabetol* 6: 36, 2007.
- Li Y, Wen X, Spataro BC, Hu K, Dai C and Liu Y: Hepatocyte growth factor is a downstream effector that mediates the antifibrotic action of peroxisome proliferator-activated receptor-gamma agonists. *J Am Soc Nephrol* 17: 54-65, 2006.
- Kusunoki H, Taniyama Y, Azuma J, Iekushi K, Sanada F, Otsu R, Iwabayashi M, Okayama K, Rakugi H and Morishita R: Telmisartan exerts renoprotective actions via peroxisome proliferator-activated receptor- γ /hepatocyte growth factor pathway independent of angiotensin II type 1 receptor blockade. *Hypertension* 59: 308-316, 2012.
- Bieghs V, Rensen PC, Hofker MH and Shiri-Sverdlov R: NASH and atherosclerosis are two aspects of a shared disease: central role for macrophages. *Atherosclerosis* 220: 287-293, 2012.
- Janke J, Schupp M, Engeli S, Gorzelniak K, Boschmann M, Sauma L, Nystrom FH, Jordan J, Luft FC and Sharma AM: Angiotensin type I receptor antagonists induce human in-vitro adipogenesis through peroxisome proliferator-activated receptor-gamma activation. *J Hypertens* 24: 1809-1816, 2006.
- Clasen R, Schupp M, Foryst-Ludwig A, Sprang C, Clemenz M, Krikov M, Thöne-Reineke C, Unger T and Kintscher U: PPARgamma-activating angiotensin type-1 receptor blockers induce adiponectin. *Hypertension* 46: 137-143, 2005.
- Nakamura T and Mizuno S: The discovery of hepatocyte growth factor (HGF) and its significance for cell biology, life sciences and clinical medicine. *Proc Jpn Acad Ser B Phys Biol Sci* 86: 588-610, 2010.
- Choi SY, Kim D, Kang JH, Park MJ, Kim YS, Lim SH, Kim CH and Lee HS: Nonalcoholic fatty liver disease as a risk factor of cardiovascular disease: relation of non-alcoholic fatty liver disease to carotid atherosclerosis. *Korean J Hepatol* 14: 77-88, 2008 (In Korean).
- Nakagami H, Osako MK, Takami Y, Hanayama R, Koriyama H, Mori M, Hayashi H, Shimizu H and Morishita R: Differential response of vascular hepatocyte growth factor concentration and lipid accumulation between telmisartan and losartan in ApoE-deficient mice. *Mol Med Rep* 1: 657-661, 2008.
- Kato J, Koda M, Kishina M, Tokunaga S, Matono T, Sugihara T, Ueki M and Murawaki Y: Therapeutic effects of angiotensin II type 1 receptor blocker, irbesartan, on non-alcoholic steatohepatitis using FLS-ob/ob male mice. *Int J Mol Med* 30: 107-113, 2012.
- Rong X, Li Y, Ebihara K, Zhao M, Kusakabe T, Tomita T, Murray M and Nakao K: Irbesartan treatment up-regulates hepatic expression of PPARalpha and its target genes in obese Koletsky (*fa(k)/fa(k)*) rats: a link to amelioration of hypertriglyceridaemia. *Br J Pharmacol* 160: 1796-1807, 2010.