

α -Tocopherol protects keratinocytes against ultraviolet A irradiation by suppressing glutathione depletion, lipid peroxidation and reactive oxygen species generation

CHI-MING WU, YA-LI CHENG, YOU-HUA DAI, MEI-FEI CHEN and CHEE-CHAN WANG

Department of Cosmetic Science, Vanung University, Tao-Yuan 32061, Taiwan, R.O.C.

Received December 09, 2013; Accepted January 30, 2014

DOI: 10.3892/br.2014.236

Abstract. This study aimed to investigate whether α -tocopherol is able to protect keratinocytes against ultraviolet A (UVA) radiation by increasing glutathione (γ -glutamylcysteinylglycine; GSH) levels or decreasing lipid peroxidation and reactive oxygen species (ROS) generation. The cell survival fraction was 43.6% when keratinocytes were irradiated with UVA at a dose of 8 J/cm². α -Tocopherol was added prior to UVA irradiation and the cell viability was assayed. The cell survival fractions were 60.2, 77.1, 89.0, 92.9 and 96.2% when α -tocopherol was added at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml, respectively. These results suggested that α -tocopherol is capable of protecting keratinocytes against UVA irradiation. Furthermore, the levels of GSH, lipid peroxidation and ROS were measured. The levels of GSH were 0.354 and 0.600 mmol/g protein in keratinocytes irradiated with UVA (8 J/cm²) and in non-irradiated cells, respectively, whereas they were 0.364, 0.420, 0.525, 0.540 and 0.545 mmol/g protein when α -tocopherol was added at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml, respectively. The levels of lipid peroxidation were 20.401 or 5.328 μ mol/g [malondialdehyde (MDA)/protein] in keratinocytes irradiated with UVA (8 J/cm²) and in non-irradiated cells, respectively, whereas they were 11.685, 6.544, 5.847, 4.390 and 2.164 μ mol/g (MDA/protein) when α -tocopherol was added at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml, respectively. The levels of ROS were 3,952.17 or 111.87 1/mg protein in keratinocytes irradiated with UVA (8 J/cm²) and in non-irradiated cells, respectively, whereas they were 742.48, 579.36, 358.16, 285.63 and 199.82 1/mg protein when α -tocopherol was added at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml, respectively. These findings suggested that α -tocopherol protects keratinocytes against UVA irradiation, possibly through

increasing the levels of GSH or decreasing the levels of lipid peroxidation and ROS generation.

Introduction

Ultraviolet (UV) radiation (200-400 nm) is associated with a number of acute and chronic effects on the skin, which may result in inflammation, immunosuppression, premature aging and the development of malignancies (1). Ultraviolet A (UVA) radiation (320-400 nm), which is not absorbed by the ozone layer, comprises >95% of the UV light that reaches the surface of the earth. UVA penetrates the epidermis and affects the epidermal and dermal layers of the skin. At the cellular level, UVA exposure causes significant oxidative stress via generation of reactive oxygen species (ROS), such as singlet oxygen, hydroxyl radical, superoxide anion and hydrogen peroxide (2). ROS are rapidly removed by non-enzymatic antioxidants, particularly glutathione (γ -glutamylcysteinylglycine; GSH), as well as enzymatic antioxidants, such as catalase, superoxide dismutase, GSH peroxidase and GSH reductase, which maintain the pro-/antioxidant balance resulting in cell and tissue stabilization. However, a surplus of ROS may overwhelm the skin antioxidant defense mechanisms causing a pro-/antioxidant disequilibrium. The overproduction of ROS induces oxidation of nucleic acids, proteins and membrane lipids, which may result in intracellular GSH and NADH/NADPH depletion and, therefore, energy loss from the cell. UV-generated ROS also affects the regulation of the gene expression of signaling molecules/cascades, such as mitogen-activated protein kinases, interrelated inflammatory cytokines, nuclear factor- κ B and activator protein-1 (3).

α -Tocopherol is the most active form of vitamin E, a fat-soluble vitamin that exists in 8 different forms in humans, and is a powerful biological antioxidant. α -Tocopherol is considered to be the major membrane-bound antioxidant employed by cells (4). α -Tocopherol scavenges acylperoxyl radicals and, thus, exerts a protective effect against cell membrane lipid peroxidation induced by UVA (5-8). Moreover, α -tocopherol and ascorbic acid work together in a cyclic process (9). During the antioxidant reaction, α -tocopherol is converted to an α -tocopherol radical by the donation of a labile hydrogen to a lipid or lipid peroxyl radical. The α -tocopherol radical is thus reduced to the original α -tocopherol form by ascorbic acid (9).

Correspondence to: Professor Chi-Ming Wu, Department of Cosmetic Science, Vanung University, 1 Van-Nung Road, Tao-Yuan 320, Taiwan R.O.C.
E-mail: chimingwu@mail.vnu.edu.tw

Key words: α -tocopherol, keratinocytes, ultraviolet A radiation, glutathione, lipid peroxidation

Among the cutaneous antioxidants, the tripeptide GSH, plays a pivotal role in protecting skin cells against oxidative damage by directly scavenging ROS or by acting as a coenzyme in GSH peroxidase- or GSH S-transferase-catalyzed reactions (10,11). It was demonstrated that GSH is also involved in DNA repair and cell apoptosis (12,13). Moreover, GSH participates in a number of biological processes, such as mitochondrial respiration, inflammatory response, signal transduction, regulation of gene expression and cell proliferation (14). GSH is able to regenerate the most essential antioxidants, vitamins C and E, back to their active forms and reduce the tocopherol radical of vitamin E directly or indirectly, via the reduction of semidehydroascorbate to ascorbate.

This study aimed to investigate the protective activity of α -tocopherol against UVA in an experimental model of HaCaT human keratinocytes. The effects of α -tocopherol on UVA-induced cellular oxidative stress, particularly on the intracellular levels of GSH, lipid peroxidation and ROS were investigated.

Materials and methods

Materials. The HaCaT human keratinocyte cell line was obtained from the Food Industry Research and Development Institute (Hsinchu, Taiwan). Dulbecco's modified Eagle's medium (DMEM), heat-inactivated fetal calf serum (FCS), penicillin-streptomycin and trypsin-EDTA solutions were purchased from Gibco-BRL (Carlsbad, CA, USA). Malondialdehyde (MDA) was purchased from Merck KGaA (Darmstadt, Germany). Sterile dimethylsulfoxide (DMSO), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB), thiobarbituric acid (TBA), dichlorodihydrofluorescein diacetate (H_2DCFDA) and α -tocopherol were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Cell culture. The HaCaT cells were grown in DMEM supplemented with heat-inactivated FCS (10%, v/v), streptomycin (100 U/ml) and penicillin (0.1 mg/ml) in a humidified atmosphere of 5% CO_2 at 37°C. The culture medium was changed 3 times per week. The cells were subcultured following trypsinization and seeded in 6-well plates at a density of 1×10^5 cells/cm².

UVA irradiation and pretreatment with α -tocopherol. The keratinocytes were pretreated with α -tocopherol (2.9-14.7 IU/ml) at 37°C for 24 h, irradiated and incubated in serum-free medium at 37°C for an additional 24 h. Stock solutions of α -tocopherol were dissolved in DMSO, with a final concentration of DMSO in the medium of 1% (v/v). Irradiated and non-irradiated control cells were treated with serum-free medium. Prior to UVA irradiation, the cells were washed with phosphate-buffered saline (PBS) and covered with a layer of PBS. Dishes with keratinocytes were irradiated (8 J/cm² UVA) on ice-cold plates to eliminate UVA thermal stimulation. Non-irradiated cells were treated similarly and were incubated in the dark. For the irradiation, a solar simulator system (Bio-Sun; Vilber Lourmat, Marne-la-Vallée, France) with a fixed wavelength of 365 nm was used.

MTT assay. Cell viability was monitored following UVA irradiation and pretreatment with α -tocopherol. MTT was used to quantify the metabolically active living cells. Mitochondrial dehydrogenases metabolize MTT to a purple formazan dye, which is measured photometrically at 570 nm using a spectrophotometer (15).

Intracellular GSH levels. Intracellular GSH was measured using a reaction with DTNB (16). The keratinocytes were rinsed with PBS, scraped into cooled perchloric acid (1%, v/v) and sonicated. The aliquots were frozen for protein determination using the Bradford assay. The suspension was centrifuged (13,000 rpm for 10 min at 4°C) and the supernatant was used to measure GSH in the reaction with the reaction mixture [800 mmol/l Tris-HCl, 20 mmol/l EDTA (pH 8.2) and 20 mg/ml DTNB]. A dilution series of GSH (0.15-1 mM) were prepared as standard. The absorbance was measured on a microplate reader at 412 nm.

Lipid peroxidation. The levels of lipid peroxidation were determined by measuring MDA in the cell extract. MDA is the end product of lipid peroxidation, which is induced by free radicals or activated oxygen species when cells are irradiated by UVA. The amount of MDA was measured using the TBA assay (17). TBA reacts with MDA, yielding red complexes, which absorb at 535 nm. A dilution series of MDA (0.1-10 μ M) were prepared as standard. The cells were washed with cooled PBS, scraped into trichloroacetic acid (2.8%, w/v), sonicated and aliquots were used for protein determination with the Bradford assay. The suspension was mixed with TBA (1%, w/v) in a ratio of 2:1, heated for 30 min at 95°C and centrifuged (13,000 rpm for 10 min at 4°C). The amount of MDA was determined spectrophotometrically.

ROS generation. The radical-scavenging efficacy of α -tocopherol in irradiated cells was monitored by the dichlorodihydrofluorescein assay. The polar, pre-fluorescent dichlorodihydrofluorescein diacetate (H_2DCFDA) undergoes deacetylation by cytosolic esterases to form dichlorodihydrofluorescein, which reacts with ROS to produce fluorescein. The fluorescence is monitored at specific excitation/emission wavelengths of 488/525 nm. Following irradiation and pretreatment with α -tocopherol, the cells were incubated with H_2DCFDA (5 nmol/l) for 15 min at 37°C. The cells were then washed with PBS, scraped into 2 ml of PBS and sonicated (18). The fluorescence was measured using a spectrophotometer (LS 50B; PerkinElmer, Waltham, MA, USA) and the protein concentration was determined with the Bradford assay.

Statistical analysis. The means \pm standard error of the mean were calculated from at least 3 repeated groups in all the experiments. Statistical significance between groups was determined with the Student's t-test. $P < 0.05$ was considered to indicate a statistically significant difference between two groups.

Results

Modulation of cell viability in UVA-irradiated cells by α -tocopherol. The cell survival fraction was 43.6% when keratinocytes were irradiated with UVA at a dose of 8 J/cm². The




Figure 1. Modulation of cell viability in UVA-irradiated cells by α -tocopherol. The cell survival fractions were measured with the MTT assay when α -tocopherol was added at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml prior to UVA irradiation (8 J/cm²). Data are expressed as the means \pm SD. *P<0.05 compared to irradiated cells. UVA, ultraviolet A; SD, standard deviation.




Figure 2. Prevention of UVA-induced GSH depletion by α -tocopherol. The keratinocytes were pretreated with α -tocopherol at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml and irradiated with UVA (8 J/cm²). The intracellular GSH levels were assayed. Data are expressed as the means \pm SD. *P<0.05 compared to irradiated cells. UVA, ultraviolet A; GSH, glutathione; SD, standard deviation.

keratinocytes were pretreated with α -tocopherol (2.9-14.7 IU/ml) prior to UVA irradiation. The cell survival fractions were 60.2, 77.1, 89.0, 92.9 and 96.2% following addition of α -tocopherol at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml, respectively (Fig. 1). α -Tocopherol pretreatment suppressed the UVA-induced decrease in cell viability in a concentration-dependent manner. These findings suggest that α -tocopherol is capable of protecting keratinocytes against UVA irradiation.

Prevention of UVA-induced GSH depletion by α -tocopherol. As illustrated in Fig. 2, the GSH level in UVA-irradiated HaCaT cells (8 J/cm²) was decreased to ~50% of that in control cells (0.600→0.354 mmol/g protein). When α -tocopherol was added prior to UVA irradiation, the GSH levels in the cells




Figure 3. Modulation of UVA-induced lipid peroxidation by α -tocopherol. The keratinocytes were pretreated with α -tocopherol at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml and irradiated with UVA (8 J/cm²). The intracellular lipid peroxidation levels were assayed. Data are expressed as the means \pm SD. *P<0.05 compared to irradiated cells. UVA, ultraviolet A; MDA, malondialdehyde; SD, standard deviation.




Figure 4. Modulation of UVA-induced ROS generation by α -tocopherol. The keratinocytes were pretreated with α -tocopherol at concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml and irradiated with UVA (8 J/cm²). The intracellular ROS generation was assayed. Data are expressed as the means \pm SD. *P<0.05 compared to irradiated cells. UVA, ultraviolet A; ROS, reactive oxygen species; SD, standard deviation.

were 0.364, 0.420, 0.525, 0.540 and 0.545 mmol/g protein at α -tocopherol concentrations of 2.9, 5.9, 8.8, 11.8 and 14.7 IU/ml, respectively. Therefore, the application of α -tocopherol to UVA-irradiated keratinocytes led to a dose-dependent prevention of GSH depletion.

Modulation of UVA-induced lipid peroxidation by α -tocopherol. The development of lipid peroxidation in irradiated HaCaT cells was measured using the TBA assay. Under our experimental conditions, there was a distinct increase in lipid peroxidation in the UVA-irradiated cells compared to the control cells (5.238→20.401 MDA/protein, μ mol/g). As illustrated in Fig. 3, the amount of MDA was significantly reduced by α -tocopherol pretreatment.

Modulation of UVA-induced ROS generation by α -tocopherol. To investigate whether the addition of α -tocopherol affects UVA-induced ROS generation, a fluorescence assay using H₂DCFDA was performed. α -Tocopherol significantly reduced ROS generation in UVA-irradiated keratinocytes in a dose-dependent manner (Fig. 4).

Discussion

UV radiation is the principal factor causing skin cancer in humans. Several studies demonstrated that supplementation with antioxidants decreases UV-induced skin damage *in vitro* and *in vivo* (19). In this study, we demonstrated the ability of α -tocopherol to prevent and reduce UVA-related damage at the cellular level in human keratinocytes. Treatment of HaCaT cells with α -tocopherol prior to UVA exposure increased cell viability and suppressed intracellular GSH depletion, lipid peroxidation and ROS generation. The cell viability assay demonstrated that α -tocopherol protects HaCaT human keratinocytes against UVA-induced apoptosis. It is well known that during and following UVA irradiation, the generation of ROS is significantly increased in exposed cells (20,21). As UVA-related biological effects are primarily mediated by ROS, their elimination is essential for protection against UVA damage. The application of α -tocopherol led to a significant increase in cell survival in irradiated HaCaT cells. α -Tocopherol pretreatment exhibited maximal protection at the highest concentration tested.

Pretreatment of cells with α -tocopherol resulted in a concentration-dependent reduction in GSH depletion. The role of GSH in protecting the skin from oxidative damage caused by various chemicals and UV exposure has been well documented. Among non-enzymatic antioxidants, GSH is considered to be the most important, as it serves as a substrate for 2 major antioxidant enzymes, GSH-peroxidase and GSH-transferase, and it also participates in vitamin C and E regeneration (22). The GSH level is directly associated with the degree of lipid peroxidation in the cell membrane (23), as GSH participates in eliminating lipid peroxidation products, including 4-hydroxynonenal, by forming a GSH conjugate (24).

The cutaneous antioxidant system is complex and incompletely understood. We previously demonstrated that magnesium ascorbyl phosphate and coenzyme Q₁₀ increased intracellular GSH levels (25). Kagan *et al* (26) reported that vitamin C is able to regenerate vitamin E from the α -tocopheroxyl radical. Furthermore, α -tocopherol and ascorbic acid work together in a cyclic process. During the antioxidant reaction, α -tocopherol is converted to an α -tocopherol radical through the donation of a labile hydrogen to a lipid or lipid peroxyl radical. The α -tocopherol radical is thus reduced to the original α -tocopherol form by ascorbic acid (9). α -Lipoic acid was previously shown to elevate intracellular GSH levels *in vitro* by increasing *de novo* synthesis (27), an effect dependent upon the metabolic reduction of lipoic to dihydrolipoic acid. Dihydrolipoic acid is then released into the culture medium, where it reduces cystine to cysteine. Cysteine is readily taken up by the neutral amino acid transport system and utilized for GSH synthesis. Through this mechanism, lipoic acid enables cysteine to bypass the Xc⁻ transport system, which is weakly expressed in lymphocytes and inhibited by

glutamate. Thereby, lipoic acid enables the key enzyme of GSH synthesis, γ -glutamylcysteine synthetase, which is regulated by an uptake-limited cysteine supply, to work at optimum conditions. However, the precise mechanism underlying the α -tocopherol-induced increase in intracellular GSH levels requires elucidation by further studies.

α -Tocopherol is the most active form of vitamin E in humans and is a powerful biological antioxidant, considered to be the major membrane-bound antioxidant employed by cells (4). The main antioxidant function of α -tocopherol is protection against lipid peroxidation (28). The overall process of lipid peroxidation includes three stages: initiation, propagation and termination. Once formed, peroxyl radicals are rearranged via a cyclization reaction to endoperoxides (precursors of MDA), with MDA as the final product (29). Our findings demonstrated that the amount of MDA was markedly reduced by treatment with α -tocopherol.

In conclusion, α -tocopherol protects keratinocytes against UVA irradiation, possibly through increasing the levels of GSH or decreasing the levels of lipid peroxidation and ROS generation.

Acknowledgements

This study was supported by grants from the National Science Council (no. 98-2314-B-238-001) and the Vanung University (no. VIT-98-CM-01), Taiwan, R.O.C..

References

1. Melnikova VO and Ananthaswamy HN: Cellular and molecular events leading to the development of skin cancer. *Mutat Res* 571: 91-106, 2005.
2. Svobodova A, Psotova J and Walterova D: Natural phenolics in the prevention of UV-induced skin damage. A review. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 147: 137-145, 2003.
3. Svobodova A, Walterova D and Vostalova J: Ultraviolet light induced alteration to the skin. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 150: 25-38, 2006.
4. Burton GW and Ingold KU: Vitamin E as an *in vitro* and *in vivo* antioxidant. *Ann NY Acad Sci* 570: 7-22, 1989.
5. Burton GW, Page YJ, Gabe EJ and Ingold KU: Antioxidant activity of vitamin E and related phenols. Importance of stereo-electronic factors. *J Am Chem Soc* 102: 7792-7794, 1980.
6. Fryer MJ: Evidence for the photoprotective effects of vitamin E. *Photochem Photobiol* 58: 304-312, 1993.
7. Nachbar F and Korting HC: The role of vitamin E in normal and damaged skin. *J Mol Med (Berl)* 73: 7-17, 1995.
8. Norkus EP, Bryce GF and Bhagavan HN: Uptake and bioconversion of alpha-tocopheryl acetate to alpha-tocopherol in skin of hairless mice. *Photochem Photobiol* 57: 613-615, 1993.
9. Kojo S: Vitamin C: basic metabolism and its function as an index of oxidative stress. *Curr Med Chem* 11: 1041-1064, 2004.
10. Afaq F and Mukhtar H: Effects of solar radiation on cutaneous detoxification pathways. *J Photochem Photobiol B* 63: 61-69, 2001.
11. Hayes JD and McLellan LI: Glutathione and glutathione-dependent enzymes represent a co-ordinately regulated defence against oxidative stress. *Free Radic Res* 31: 273-300, 1999.
12. Fonnum F and Lock EA: The contributions of excitotoxicity, glutathione depletion and DNA repair in chemically induced injury to neurones: exemplified with toxic effects on cerebellar granule cells. *J Neurochem* 88: 513-531, 2004.
13. Hall AG: Review: The role of glutathione in the regulation of apoptosis. *Eur J Clin Invest* 29: 238-245, 1999.
14. Sies H: Glutathione and its role in cellular functions. *Free Radic Biol Med* 27: 916-921, 1999.
15. Green LM, Reade JL and Ware CF: Rapid colorimetric assay for cell viability: application to the quantitation of cytotoxic and growth inhibitory lymphokines. *J Immunol Methods* 70: 257-268, 1984.

16. Sedlak J and Lindsay RH: Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem* 25: 192-205, 1968.
17. Buege JA and Aust SD: Microsomal lipid peroxidation. *Methods Enzymol* 52: 302-310, 1978.
18. D'Angelo S, Ingrosso D, Migliardi V, *et al*: Hydroxytyrosol, a natural antioxidant from olive oil, prevents protein damage induced by long-wave ultraviolet radiation in melanoma cells. *Free Radic Biol Med* 38: 908-919, 2005.
19. Swindells K and Rhodes LE: Influence of oral antioxidants on ultraviolet radiation-induced skin damage in humans. *Photodermatol Photoimmunol Photomed* 20: 297-304, 2004.
20. Tyrrell RM: The molecular and cellular pathology of solar ultraviolet radiation. *Mol Aspects Med* 15: 1-77, 1994.
21. Morita A and Krutmann J: Ultraviolet A radiation-induced apoptosis. *Methods Enzymol* 319: 302-309, 2000.
22. Svobodova A, Rambouskova J, Walterova D and Vostalova J: Protective effects of phenolic fraction of blue honeysuckle fruits against UVA-induced damage to human keratinocytes. *Arch Dermatol Res* 300: 225-233, 2008.
23. Schneider LA, Dissemond J, Brenneisen P, Hainzl A, Briviba K, Wlaschek M and Scharffetter-Kochanek K: Adaptive cellular protection against UVA-1-induced lipid peroxidation in human dermal fibroblasts shows donor-to-donor variability and is glutathione dependent. *Arch Dermatol Res* 297: 324-328, 2006.
24. Tarozzi A, Marchesi A, Hrelia S, Angeloni C, Andrisano V, Fiori J, Cantelli-Forti G and Hrelia P: Protective effects of cyanidin-3-*O*-beta-glucopyranoside against UVA-induced oxidative stress in human keratinocytes. *Photochem Photobiol* 81: 623-629, 2005.
25. Hwang TL, Tsai CJ, Chen JL, Changchien TT, Wang CC and Wu CM: Magnesium ascorbyl phosphate and coenzyme Q₁₀ protect the keratinocytes against UVA irradiation by suppressing glutathione depletion. *Mol Med Rep* 6: 375-378, 2012.
26. Kagan V, Witt E, Goldman R, Scita G and Packer L: Ultraviolet light-induced generation of vitamin E radicals and their recycling. A possible photosensitizing effect of vitamin E in skin. *Free Radic Res Commun* 16: 51-64, 1992.
27. Han D, Handelman G, Marcocci L, Sen CK, Roy S, Kobuchi H, Tritschler HJ, Flohe L and Packer L: Lipoic acid increases de novo synthesis of cellular glutathione by improving cystine utilization. *Biofactors* 6: 321-338, 1997.
28. Pryor WA: Vitamin E and heart disease: basic science to clinical intervention trials. *Free Radic Biol Med* 28: 141-164, 2000.
29. Marnett LJ: Lipid peroxidation-DNA damage by malondialdehyde. *Mutat Res* 424: 83-95, 1999.