

# Association of the interleukin-18 receptor 1 and interleukin-18 receptor accessory protein polymorphisms with the risk of esophageal cancer

JINGFENG ZHU<sup>1\*</sup>, CHAO LIU<sup>1\*</sup>, XIAO TENG<sup>2</sup>, JUN YIN<sup>1</sup>, LIANG ZHENG<sup>3</sup>, LIMING WANG<sup>4</sup>, WEIFENG TANG<sup>1</sup>, HAIYONG GU<sup>1</sup>, BING GU<sup>5</sup> and LIANG CHEN<sup>6</sup>

<sup>1</sup>Department of Cardiothoracic Surgery, Affiliated People's Hospital of Jiangsu University, Zhenjiang, Jiangsu 212002; <sup>2</sup>State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100037; <sup>3</sup>Department of Cardiothoracic Surgery, The First People's Hospital of Changzhou and The Third Affiliated Hospital of Suzhou University, Changzhou, Jiangsu 213003; <sup>4</sup>Cancer Institute, Department of Chemotherapy, People's Hospital Affiliated to Jiangsu University, Zhenjiang, Jiangsu 212002; <sup>5</sup>Department of Laboratory Medicine, The Affiliated Hospital of Xuzhou Medical College, Xuzhou, Jiangsu 221000; <sup>6</sup>Department of Thoracic and Cardiac Surgery, The First Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu 210009, P.R. China

Received May 6, 2015; Accepted June 24, 2015

DOI: 10.3892/br.2015.552

**Abstract.** Esophageal cancer is the fourth leading cause of cancer-associated fatalities and the fourth most commonly diagnosed cancer. In addition to environmental risk factors, genetic factors may have a significant role in esophageal cancer carcinogenesis. A hospital-based case-control study was conducted to evaluate the genetic effects of functional single-nucleotide polymorphisms in the interleukin-18 (*IL-18*), IL-18 receptor 1 protein (*IL-18R1*), IL-18 receptor accessory protein (*IL-18RAP*) and *IL-28B* on the development of esophageal cancer. In total, 380 esophageal squamous cell carcinoma (ESCC) cases and 380 controls were recruited for the present study. The *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T

and *IL-28B* rs8099917 T>G genotypes were determined. No association was observed between the *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms and the risk of ESCC. However, in stratification analyses, a significantly decreased risk of ESCC associated with the *IL-18R1* rs13015714 G>T polymorphism and a significantly increased risk of ESCC associated with the *IL-18RAP* rs917997 C>T polymorphism was evident among male patients and patients who smoked or consumed alcohol. These findings highlighted that functional polymorphisms *IL-18R1* rs13015714 G>T and *IL-18RAP* rs917997 C>T may contribute to ESCC susceptibility among these subgroups. However, the present results were obtained with a limited sample size and further epidemiological studies are warranted to clarify the role of *IL-18R1* and *IL-18RAP* variants in the development of ESCC.

**Correspondence to:** Dr Bing Gu, Department of Laboratory Medicine, The Affiliated Hospital of Xuzhou Medical College, 99 Huaihai Road West, Xuzhou, Jiangsu 221000, P.R. China  
E-mail: gb20031129@163.com

Professor Liang Chen, Department of Thoracic and Cardiac Surgery, The First Affiliated Hospital of Nanjing Medical University, 300 Guangzhou Road, Nanjing, Jiangsu 210009, P.R. China  
E-mail: clbright0909@hotmail.com

\*Contributed equally

**Abbreviations:** CI, confidence interval; IL-18, interleukin-18; OR, odds ratio; SNP, single-nucleotide polymorphism

**Key words:** interleukin-18, interleukin-18 receptor 1, interleukin-18 receptor accessory protein, polymorphisms, esophageal cancer, molecular epidemiology

## Introduction

Esophageal carcinoma is the fourth leading cause of cancer-associated fatalities and the fourth most commonly diagnosed cancer in China in 2010 (1). For esophageal cancer patients, the prior study reported that the 5-year survival rate is extremely poor and accounts for only 12.3% (2). In the highest-risk area, esophageal squamous cell carcinoma (ESCC) accounts for >90% of esophageal cancers (3,4). ESCC carcinogenesis is multifactorial, and in addition to the established environmental risk factors, such as heavy drinking and smoking (5), genetic aberrations, such as single-nucleotide polymorphisms (SNPs), may have significant roles (6).

Interleukin-18 (IL-18), an interferon- $\gamma$  (IFN- $\gamma$ )-inducing factor, upregulates several cytokines, including IL-1 $\beta$  and tumor necrosis factor- $\alpha$ , and IFN- $\gamma$  promotes T helper cell type 1 (Th1) differentiation (7). IL-18 is also one of the main

cytokines of the inflammasomes and has been confirmed to effect carcinogenesis and tumor progression significantly (8). The IL-18 receptor is comprised of IL-18 receptor accessory protein (IL-18RAP) and IL-18 receptor 1 (IL-18R1) protein (9,10). Upon binding to its receptor, IL-18R1 protein, IL-18 triggers the recruitment of IL-18RAP, which initiates signaling. IL-18 has a critical role in MyD88-mediated signaling to prevent colon adenocarcinoma development (11). IL-18RAP forms the signaling chain of this receptor complex and has been shown to be crucial for signaling of IL-18, resulting in the production of IFN- $\gamma$  (12). These two subunits of the IL-18R are mainly expressed on Th1 cells in response to IL-12 and/or IFN- $\alpha$  (13). IL-18RAP was also correlated with inflammatory bowel disease (14). A previous study reported that the *IL-18R1* and *IL-18RAP* genes were associated with atherosclerosis and its cardiovascular complications (15).

*IL-18* is located at the 11q22.2-22.3 chromosome and its promoter region is relatively unique and is comprised of several transcription initiation sites. The *IL-18* rs360719 polymorphism (A>G mutation) leads to loss of the octamer transcription factor-1 (OCT-1) transcription factor binding site. OCT-1 is identified as a ubiquitously expressed factor and has an important role in the regulation of certain genes. OCT-1 can also downregulate the expression of specific cytokines (16). Allele A of rs917997, an SNP that is 1.5 kb downstream of *IL-18RAP*, was strongly associated with coeliac disease susceptibility (17). This allele is also correlated with lower mRNA levels of *IL-18RAP* in whole blood. Furthermore, the *IL-18RAP* GA haplotype of rs13015714 and rs917997 showed the strongest association with coeliac disease (17).

The *IL-28B* rs8099917 T>G polymorphism has been demonstrated with the response to IFN- $\gamma$ -based antiviral therapy in the natural course of hepatitis C and following liver transplantation (18,19).

The biological and pathological significance of *IL-18*, *IL-18R1*, *IL-18RAP* and *IL-28B* suggested that the functional genetic variations in these genes may contribute to the development of ESCC. The objective of the present study was to evaluate the association between *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G genotypes and ESCC susceptibility in a hospital-based case-control study. Genotyping analyses were performed for these 4 SNPs with 380 ESCC cases and 380 cancer-free controls in a Chinese Han population.

## Patients and methods

*Ethical approval of the study protocol.* The study was approved by the Review Board of Jiangsu University (Zhenjiang, China). The study complied with the World Medical Association Declaration of Helsinki regarding ethical conduct of research involving human subjects and/or animals. All the subjects provided written informed consent for inclusion in the study.

*Study subjects.* A total of 380 subjects with esophageal cancer were consecutively recruited from the Affiliated People's Hospital of Jiangsu University and Affiliated Hospital of Jiangsu University (Zhenjiang, China) between

October 2008 and November 2009. By pathological analysis, all the esophageal cancer cases were diagnosed as ESCC. The exclusion criteria were patients who previously had: Cancer; any metastasized cancer; radiotherapy or chemotherapy. Frequency-matched to the cases with regards to age ( $\pm 5$  years) and gender, the controls were patients without cancer that were recruited from the two hospitals during the same time period. The majority of the control subjects were admitted to these two hospitals for the treatment of trauma.

Using a pre-tested questionnaire, demographic data (such as age and gender) and the associated ESCC risk factors were collected by trained interviewers. The definition of 'smokers' was individuals who smoked one cigarette per day for >1 year. The definition of 'alcohol drinkers' was subjects who consumed  $\geq 3$  alcoholic drinks a week for >6 months.

*Isolation of DNA and genotyping by a custom-by-design 48-Plex SNPscan™ kit.* In total, 2-ml blood samples were collected from each subject using vacutainers and transferred to tubes lined with ethylenediamine-N,N,N',N'-tetraacetic acid. Genomic DNA was isolated from whole blood with the QIAamp DNA Blood Mini kit (Qiagen, Berlin, Germany). Sample DNA (10 ng) was amplified by polymerase chain reaction according to the manufacturer's instructions. For *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G SNPs, genotyping was performed using a custom-by-design 48-Plex SNPscan™ kit (Genesky Biotechnologies Inc., Shanghai, China), as previously described (20). For quality control, repeated analyses were conducted for 4% of randomly selected samples with high DNA quality.

*Statistical analysis.* Differences in the distributions of demographic characteristics, selected variables and genotypes of the *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G variants between the cases and controls were evaluated using the  $\chi^2$  test. The associations between the 4 SNPs and risk of ESCC were estimated by computing the odds ratios (ORs) and their 95% confidence intervals (CIs) using logistic regression analyses for crude and adjusted ORs when adjusting for age, gender, smoking and drinking status. The Hardy-Weinberg equilibrium (HWE) was tested by a goodness-of-fit  $\chi^2$  test to compare the observed genotype frequencies to the expected ones among the control subjects. All the statistical analyses were performed with SAS 9.1.3 (SAS Institute Inc., Cary, NC, USA).  $P < 0.05$  was considered to indicate a statistically significant difference.

## Results

*Characteristics of the study population.* Characteristics of cases and controls included in the study are provided in Table I. The cases and controls appeared to be adequately matched on age and gender, as suggested by the  $\chi^2$  tests ( $P = 0.056$  and  $P = 0.346$ , respectively). As shown in Table I, no significant difference was detected on drinking status between the cases and the controls ( $P = 0.183$ ), however, smoking rate was higher in ESCC patients compared with the control subjects ( $P = 0.014$ ). In Table II, the primary information for

Table I. Distribution of the selected demographic variables and risk factors in ESCC cases and controls.

Variables	Cases (n=380), n (%)	Controls (n=380), n (%)	P-value <sup>a</sup>
Age, years			
<60	142 (37.4)	117 (30.8)	0.056
≥60	238 (62.6)	263 (69.2)	
Gender			
Male	269 (70.8)	257 (67.6)	0.346
Female	111 (29.2)	123 (32.4)	
Tobacco use			
Never	220 (57.9)	253 (66.6)	0.014
Ever	160 (42.1)	127 (33.4)	
Alcohol use			
Never	253 (66.6)	270 (71.1)	0.183
Ever	127 (33.4)	110 (28.9)	

<sup>a</sup>Two-sided  $\chi^2$  test. ESCC, esophageal squamous cell carcinoma.

these 4 genotyped SNPs is listed. The genotyping success rate was 96.97% for *IL-18* rs360719 A>G, 96.45% for *IL-18R1* rs13015714 G>T, 96.32% for *IL-18RAP* rs917997 C>T and 97.11% for *IL-28B* rs8099917 T>G in all 760 samples. For all the SNPs, the concordance rates of repeated analyses were 100%. Minor allele frequency (MAF) in the controls was similar to MAF for Chinese subjects in a database for all 4 SNPs (Table II). The observed genotype frequencies for *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms in the controls were in HWE (P=0.774, P=0.249, P=0.465 and P=0.325, respectively) (Table II).

Table II. Primary information for *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms.

Genotyped SNPs	<i>IL-18</i> rs360719 A>G	<i>IL-18R1</i> rs13015714 G>T	<i>IL-18RAP</i> rs917997 C>T	<i>IL-28B</i> rs8099917 T>G
Chromosome	11	2	2	19
Location	5'-Flanking	5'-Flanking	3'-Flanking	5'-Flanking
Chr Pos (genome build 36.3)	111541359	102338297	102437000	44435005
Regulome DB score <sup>a</sup>	2b	6	No data	4
TFBS <sup>b</sup>	Y	-	-	-
MAF <sup>c</sup> for Chinese population	0.142	0.547	0.488	0.035
MAF in the controls (n=380)	0.134	0.511	0.496	0.049
P-value for HWE <sup>d</sup> test in our controls	0.774	0.249	0.465	0.325
Genotyping value, %	96.97	96.45	96.32	97.11

<sup>a</sup><http://www.regulomedb.org/>; <sup>b</sup>TFBS, transcription factor binding site (<http://snpinfo.niehs.nih.gov/snpinfo/snpfunc.htm>); <sup>c</sup>MAF, minor allele frequency; *IL-18* rs360719 A>G MAF information was available for the Chinese Han and Japanese populations; <sup>d</sup>HWE, Hardy-Weinberg equilibrium. SNPs, single-nucleotide polymorphisms; *IL-18*, interleukin-18; *IL-18R1*, IL-18 receptor 1; *IL-18RAP*, IL-18 receptor accessory protein; Chr Pos, chromosome position; Y, yes.

*Associations between IL-18 rs360719 A>G, IL-18R1 rs13015714 G>T, IL-18RAP rs917997 C>T and IL-28B rs8099917 T>G polymorphisms and the risk of ESCC.* The genotype distributions of *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G in the cases and the controls are shown in Table III. In the single locus analyses, the genotype frequencies of *IL-18* rs360719 A>G were 72.6 (AA), 26.4 (AG) and 1.1% (GG) in the case patients and 74.8 (AA), 23.6 (AG) and 1.6% (GG) in the control subjects, and the difference was not statistically significant (P=0.579). When the *IL-18* rs360719 AA was used as the reference, the AG genotype was not associated with the risk for ESCC (AG vs. AA: Adjusted OR=1.15; 95% CI, 0.82-1.62; P=0.411); the GG genotype was not associated with the risk for ESCC (GG vs. AA: Adjusted OR=0.68; 95% CI, 0.19-2.46; P=0.553). In the dominant model, the *IL-18* rs360719 AG/GG variants were not associated with the risk for ESCC, compared with the *IL-18* rs360719 AA genotype (adjusted OR=1.12; 95% CI, 0.81-1.56; P=0.498). In the recessive model, when the *IL-18R1* rs13015714 AA/AG genotypes were used as the reference group, the GG homozygote genotype was not associated with the risk for ESCC (adjusted OR=0.65; 95% CI, 0.18-2.37; P=0.517) (Table III).

No association was observed between *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms and the risk of ESCC (Table III).

*Stratification analyses of IL-18 rs360719 A>G, IL-18R1 rs13015714 G>T, IL-18RAP rs917997 C>T and IL-28B rs8099917 T>G polymorphisms and the risk of ESCC.* To evaluate the effects of *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G genotypes on ESCC risk according to different age, gender, smoking and alcohol drinking status, the stratification analyses were performed. A significantly decreased risk of ESCC associated with the *IL-18R1* rs13015714 G>T polymorphism was evident among male patients (GT+TT vs. GG:

Table III. Logistic regression analyses of the associations between *IL-18* rs360719 A>G, *IL-18R1* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms and the risk of ESCC.

Genotype	Cases (n=380), n (%)	Controls (n=380), n (%)	Crude OR (95% CI)	P-value	Adjusted OR <sup>a</sup> (95% CI)	P-value
<i>IL-18</i> rs360719 A>G						
AA	267 (72.6)	276 (74.8)	1.00		1.00	
AG	97 (26.4)	87 (23.6)	1.15 (0.83-1.61)	0.406	1.15 (0.82-1.62)	0.411
GG	4 (1.1)	6 (1.6)	0.69 (0.19-2.47)	0.568	0.68 (0.19-2.46)	0.553
GG vs. AG vs. AA						0.579
AG+GG	101 (27.4)	93 (25.2)	1.12 (0.81-1.56)	0.490	1.12 (0.81-1.56)	0.498
AA+AG	364 (98.9)	363 (98.4)	1.00		1.00	
GG	4 (1.1)	6 (1.6)	0.67 (0.19-2.38)	0.530	0.65 (0.18-2.37)	0.517
G allele	105 (14.3)	99 (13.4)				
<i>IL-18R1</i> rs13015714 G>T						
GG	100 (27.5)	83 (22.4)	1.00		1.00	
GT	173 (47.7)	196 (53.0)	0.73 (0.51-1.05)	0.086	0.71 (0.49-1.02)	0.060
TT	90 (24.8)	91 (24.6)	0.82 (0.54-1.24)	0.348	0.83 (0.54-1.25)	0.364
TT vs. GT vs. GG						0.229
GT+TT	263 (72.5)	287 (77.6)	0.76 (0.54-1.06)	0.110	0.75 (0.53-1.05)	0.089
GG+GT	273 (75.2)	279 (75.4)	1.00		1.00	
TT	90 (24.8)	91 (24.6)	1.01 (0.72-1.41)	0.950	1.04 (0.74-1.46)	0.829
T allele	353 (48.6)	378 (51.1)				
<i>IL-18RAP</i> rs917997 C>T						
CC	91 (25.0)	90 (24.5)	1.00		1.00	
CT	167 (45.9)	191 (51.9)	0.87 (0.61-1.24)	0.426	0.84 (0.58-1.20)	0.334
TT	106 (29.1)	87 (23.6)	1.21 (0.80-1.81)	0.369	1.19 (0.79-1.80)	0.404
TT vs. CT vs. CC						0.177
CT+TT	273 (75.0)	278 (75.5)	0.97 (0.69-1.36)	0.865	0.95 (0.67-1.33)	0.753
CC+CT	258 (70.9)	281 (76.4)	1.00		1.00	
TT	106 (29.1)	87 (23.6)	1.33 (0.95-1.85)	0.093	1.34 (0.96-1.87)	0.085
T allele	379 (52.1)	365 (49.6)				
<i>IL-28B</i> rs8099917 T>G						
TT	335 (91.0)	334 (90.3)	1.00		1.00	
TG	31 (8.4)	36 (9.7)	0.86 (0.52-1.42)	0.553	0.87 (0.52-1.44)	0.585
GG	2 (0.5)	0 (0.0)	-	0.980	-	0.980
GG vs. TG vs. TT						0.306
TG+GG	33 (9.0)	36 (9.7)	0.91 (0.56-1.50)	0.722	0.92 (0.56-1.52)	0.741
TT+TG	366 (99.5)	370 (100)	1.00		1.00	
GG	2 (0.5)	0 (0.0)	-	0.980	-	0.980
G allele	35 (4.8)	36 (4.9)				

<sup>a</sup>Adjusted for age, gender, smoking status and alcohol consumption. *IL-18*, interleukin-18; *IL-18R1*, IL-18 receptor 1; *IL-18RAP*, IL-18 receptor accessory protein; ESCC, esophageal squamous cell carcinoma; OR, odds ratio; CI, confidence interval.

Adjusted OR=0.65; 95% CI, 0.43-0.98; P=0.040), patients who smoked (GT+TT vs. GG: Adjusted OR=0.43; 95% CI, 0.23-0.79; P=0.007) or were alcohol drinkers (GT+TT vs. GG: Adjusted OR=0.38; 95% CI, 0.19-0.76; P=0.006) (Table IV). A significantly increased risk of ESCC associated with the *IL-18RAP* rs917997 C>T polymorphism was evident among male patients (TT vs. CT+CC: Adjusted

OR=1.58; 95% CI, 1.05-2.38; P=0.029), patients who smoked (TT vs. CT+CC: Adjusted OR=2.36; 95% CI, 1.29-4.30; P=0.005) or were alcohol drinkers (TT vs. CT+CC: Adjusted OR=3.01; 95% CI, 1.52-5.97; P=0.002) (Table V). For *IL-18* rs360719 A>G and *IL-28B* rs8099917 T>G polymorphisms, no association was observed following stratification (data not shown).

Table IV. Stratified analyses between the *IL-18R1* rs13015714 G>T polymorphism and ESCC risk by gender, age, smoking status and alcohol consumption.

Variables	<i>IL18R1</i> rs13015714 G>T, case/control <sup>a</sup>				Adjusted OR <sup>b</sup> (95% CI); P-value				
	GG	GT	TT	GT+TT	GG	GT	TT	GT+TT	TT vs. (GT+GG)
Gender									
Male	72/52	124/136	62/65	186/201	1.00	0.63 (0.40-0.97); 0.037 <sup>c</sup>	0.69 (0.42-1.15); 0.157	0.65 (0.43-0.98); 0.040 <sup>c</sup>	0.95 (0.63-1.43); 0.816
Female	28/31	49/60	28/26	77/86	1.00	0.90 (0.48-1.71); 0.753	1.26 (0.60-2.65); 0.546	1.01 (0.55-1.83); 0.983	1.34 (0.72-2.50); 0.350
Age, years									
<60	34/22	65/65	37/26	102/91	1.00	0.57 (0.29-1.11); 0.349	0.91 (0.42-1.96); 0.204	0.66 (0.35-1.25); 0.803	1.34 (0.73-2.47); 0.098
≥60	66/61	108/131	53/65	161/196	1.00	0.77 (0.50-1.19); 0.245	0.75 (0.45-1.24); 0.261	0.77 (0.51-1.15); 0.198	0.88 (0.58-1.35); 0.566
Smoking status									
Never	53/62	96/122	58/62	154/184	1.00	0.98 (0.62-1.56); 0.938	1.16 (0.69-1.95); 0.583	1.04 (0.68-1.61); 0.857	1.17 (0.77-1.79); 0.464
Ever	47/21	77/74	32/29	109/103	1.00	0.42 (0.22-0.80); 0.008 <sup>c</sup>	0.45 (0.21-0.95); 0.036 <sup>c</sup>	0.43 (0.23-0.79); 0.007 <sup>c</sup>	0.81 (0.45-1.48); 0.499
Alcohol consumption									
Never	62/66	113/130	63/66	176/196	1.00	0.96 (0.62-1.48); 0.846	1.08 (0.66-1.78); 0.753	1.00 (0.66-1.51); 0.998	1.16 (0.74-1.68); 0.601
Ever	38/17	60/66	27/25	87/91	1.00	0.38 (0.18-0.78); 0.008 <sup>c</sup>	0.39 (0.17-0.93); 0.033 <sup>c</sup>	0.38 (0.19-0.76); 0.006 <sup>c</sup>	0.79 (0.40-1.54); 0.485

<sup>a</sup>Genotyping was successful in 363 (95.5%) ESCC cases and 370 (97.4%) controls for *IL-18R1* rs13015714 G>T; <sup>b</sup>adjusted for age, gender, smoking status and alcohol consumption (in addition to stratified factors accordingly) in a logistic regression model; <sup>c</sup>statistically significant (P<0.05). ESCC, esophageal squamous cell carcinoma; *IL-18R1*, interleukin-18 receptor 1; OR, odds ratio.

Table V. Stratified analyses between the *IL-18RAP* rs917997 C>T polymorphism and ESCC risk by gender, age, smoking status and alcohol consumption.

Variables	<i>IL18RAP</i> rs917997 C>T, case/control <sup>a</sup>				Adjusted OR <sup>b</sup> (95% CI); P-value				
	CC	CT	TT	CT+TT	CC	CT	TT	CT+TT	TT vs. (CT+CC)
Gender									
Male	61/65	120/132	77/54	197/186	1.00	0.92 (0.59-1.42); 0.703	1.49 (0.90-2.46); 0.119	1.08 (0.72-1.63); 0.701	1.58 (1.05-2.38); 0.029 <sup>c</sup>
Female	30/25	47/59	29/33	76/92	1.00	0.63 (0.32-1.21); 0.166	0.69 (0.33-1.44); 0.327	0.65 (0.35-1.21); 0.172	0.94 (0.52-1.70); 0.848
Age, years									
<60	37/25	62/62	37/24	99/86	1.00	0.59 (0.31-1.15); 0.122	1.01 (0.47-2.16); 0.984	0.71 (0.38-1.32); 0.277	1.43 (0.77-2.65); 0.262
≥60	54/65	105/129	69/63	174/192	1.00	1.01 (0.64-1.57); 0.978	1.34 (0.81-2.20); 0.257	1.12 (0.73-1.70); 0.609	1.33 (0.89-1.99); 0.166
Smoking status									
Never	59/61	94/120	55/65	149/185	1.00	0.80 (0.51-1.25); 0.326	0.82 (0.49-1.37); 0.441	0.80 (0.53-1.23); 0.312	0.94 (0.62-1.45); 0.791
Ever	32/29	73/71	51/22	124/93	1.00	0.94 (0.50-1.76); 0.835	2.25 (1.07-4.74); 0.033 <sup>c</sup>	1.24 (0.68-2.26); 0.479	2.36 (1.29-4.30); 0.005 <sup>c</sup>
Alcohol consumption									
Never	65/65	110/126	64/70	174/196	1.00	0.85 (0.55-1.31); 0.462	0.85 (0.52-1.39); 0.523	0.85 (0.57-1.28); 0.434	0.95 (0.63-1.42); 0.790
Ever	26/25	57/65	42/17	99/82	1.00	0.90 (0.44-1.83); 0.766	2.79 (1.19-6.54); 0.019 <sup>c</sup>	1.27 (0.64-2.50); 0.493	3.01 (1.52-5.97); 0.002 <sup>c</sup>

<sup>a</sup>Genotyping was successful in 364 (95.8%) ESCC cases and 368 (96.8%) controls for *IL-18RAP* rs917997 C>T; <sup>b</sup>adjusted for age, gender, smoking status and alcohol consumption (in addition to stratified factors accordingly) in a logistic regression model; <sup>c</sup>statistically significant (P<0.05). ESCC, esophageal squamous cell carcinoma; *IL-18RAP*, interleukin-18 receptor accessory protein; OR, odds ratio.

Table VI. Logistic regression analyses of associations between *IL-18* rs360719 A>G, *IL-18RI* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms and risk of esophageal cancer lymph node metastasis.

Genotype	LN meta (+) (n=85), n (%)	LN meta (-) (n=270), n (%)	Crude OR (95% CI)	P-value	Adjusted OR <sup>a</sup> (95% CI)	P-value
<i>IL-18</i> rs360719 A>G						
AA	63 (75.9)	185 (70.9)	1.00		1.00	
AG	18 (21.7)	74 (28.4)	0.71 (0.40-1.29)	0.263	0.70 (0.39-1.27)	0.241
GG	2 (2.4)	2 (0.8)	2.94 (0.41-21.28)	0.287	2.66 (0.35-20.11)	0.343
GG vs. AG vs. AA						0.255
AG+GG	20 (24.1)	76 (29.1)	0.77 (0.44-1.37)	0.375	0.76 (0.43-1.34)	0.339
AA+AG	81 (97.6)	259 (99.2)	1.00		1.00	
GG	2 (2.4)	2 (0.8)	3.20 (0.44-23.06)	0.249	2.91 (0.39-21.86)	0.299
G allele	22 (13.3)	78 (14.9)				
<i>IL-18RI</i> rs13015714 G>T						
GG	22 (26.8)	73 (28.4)	1.00		1.00	
GT	41 (50.0)	123 (47.9)	1.11 (0.61-2.00)	0.739	1.09 (0.60-1.98)	0.782
TT	19 (23.2)	61 (23.7)	1.03 (0.51-2.09)	0.927	1.04 (0.51-2.11)	0.917
TT vs. GT vs. GG						0.941
GT+TT	60 (73.2)	184 (71.6)	1.08 (0.62-1.89)	0.782	1.07 (0.61-1.88)	0.809
GG+GT	63 (76.8)	196 (76.3)	1.00		1.00	
TT	19 (23.2)	61 (23.7)	0.97 (0.54-1.75)	0.917	0.98 (0.54-1.78)	0.957
T allele	79 (48.2)	245 (47.7)				
<i>IL-18RAP</i> rs917997 C>T						
CC	20 (24.4)	61 (23.6)	1.00		1.00	
CT	40 (48.8)	120 (46.3)	1.02 (0.55-1.89)	0.958	0.99 (0.53-1.86)	0.984
TT	22 (26.8)	78 (30.1)	0.86 (0.43-1.72)	0.670	0.84 (0.42-1.70)	0.635
TT vs. CT vs. CC						0.849
CT+TT	62 (75.6)	198 (76.4)	0.96 (0.54-1.71)	0.876	0.94 (0.52-1.68)	0.824
CC+CT	60 (73.2)	181 (69.9)	1.00		1.00	
TT	22 (26.8)	78 (30.1)	0.85 (0.49-1.48)	0.569	0.85 (0.48-1.49)	0.563
T allele	84 (51.2)	276 (53.3)				
<i>IL-28B</i> rs8099917 T>G						
TT	75 (90.4)	238 (91.2)	1.00		1.00	
TG	8 (9.6)	21 (8.0)	1.21 (0.51-2.84)	0.664	1.20 (0.51-2.84)	0.680
GG	0 (0.0)	2 (0.8)	—	0.990	—	0.990
GG vs. TG vs. TT						0.660
TG+GG	8 (9.6)	23 (8.8)	1.10 (0.47-2.57)	0.819	1.07 (0.46-2.51)	0.872
TT+TG	83 (100)	259 (99.2)	1.00		1.00	
GG	0 (0.0)	2 (0.8)	—	0.990	—	0.990
G allele	8 (4.8)	25 (4.8)				

<sup>a</sup>Adjusted for age, gender, smoking status and alcohol consumption. LN meta, lymph node metastasis; *IL-18*, interleukin-18; *IL-18RI*, IL-18 receptor 1; *IL-18RAP*, IL-18 receptor accessory protein; OR, odds ratio; CI, confidence interval.

Associations between *IL-18* rs360719 A>G, *IL-18RI* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms and the risk of esophageal cancer lymph node metastasis. Analyses between *IL-18* rs360719 A>G, *IL-18RI* rs13015714 G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G polymorphisms and risk of esophageal cancer lymph node metastasis were further

conducted. No association was observed between the 4 polymorphisms and lymph node metastasis (Table VI).

## Discussion

In the present hospital-based case-control study of ESCC, the associations of *IL-18* rs360719 A>G, *IL-18RI* rs13015714

G>T, *IL-18RAP* rs917997 C>T and *IL-28B* rs8099917 T>G were associated with the risk of ESCC in a high-risk Chinese population. The multivariable logistic analysis revealed that a significantly decreased risk of ESCC associated with the *IL-18R1* rs13015714 G>T polymorphism was evident among male patients and patients who were smokers or alcohol drinkers. In addition, a significantly increased risk of ESCC associated with the *IL-18RAP* rs917997 C>T polymorphism was evident among male patients and patients who were smokers or alcohol drinkers.

*IL-18R* is comprised of *IL-18RAP* and *IL-18R1*. Allele A of *IL-18RAP* rs917997 was strongly correlated with coeliac disease susceptibility (17). The *IL-18RAP* rs917997 A allele has a significant effect on the level of *IL-18RAP* mRNA expression. The *IL-18RAP* rs917997 C allele is strongly associated with a protective effect in Barrett's esophagus and esophageal adenocarcinoma (21). The CC genotype at the *IL-18RAP* locus rs917997 was associated with a protective effect against esophageal disease, which is in accordance with the present findings in the stratification analyses (21).

The frequencies of genetic polymorphisms often vary between different ethnic groups (22). In the present study, the MAF of *IL-18R1* rs13015714 T was 0.511 among 380 control subjects, which is consistent with that of the Chinese Han population (0.547); however, this was significantly lower than that of the European population (0.796) and sub-Saharan African population (0.894) in the SNP database (<http://www.ncbi.nlm.nih.gov/SNP>). Similarly, the MAF of *IL-18RAP* rs917997 T was 0.496 among 380 control subjects, which is consistent with that of the Chinese Han population (0.488); however, this was significantly higher than that of the European population (0.208) and sub-Saharan African population (0.049) in the SNP database (<http://www.ncbi.nlm.nih.gov/SNP>).

Considering *IL-18R1* rs13015714 G>T mutant alleles in the control group, OR, ESCC samples and control samples, the power of the present analysis ( $\alpha=0.05$ ) was 0.679 in 258 ESCC cases and 253 controls with adjusted OR=0.65 for *IL-18R1* rs13015714 G>T in the male subgroup. The power of the analysis ( $\alpha=0.05$ ) was 0.932 in 156 ESCC cases and 124 controls with adjusted OR=0.43 in the smoking subgroup and 0.950 in 125 ESCC cases and 108 controls with adjusted OR=0.38 in the drinking subgroup.

For *IL-18RAP* rs917997 C>T, the power of the analysis ( $\alpha=0.05$ ) was 0.728 in 258 ESCC cases and 251 controls with adjusted OR=1.58 in the male subgroup, 0.937 in 156 ESCC cases and 122 controls with adjusted OR=2.36 in the smoking subgroup and 0.983 in 125 ESCC cases and 107 controls with adjusted OR=3.01 in the drinking subgroup.

Several weaknesses should be addressed in the case-control study. Firstly, all the subjects were selected from hospitals, which may have led to a bias and consequently affected the validity of the findings. Secondly, the relatively small sample size restricted the statistical power of the study, particularly in the stratification analyses. Further larger sample size studies with a well-designed two-stage fine-mapping strategy are warranted to confirm the present findings. Thirdly, due to the lack of detailed information on cancer metastasis and survival, further investigations on the potential role of *IL-18R1* rs13015714 G>T and *IL-18RAP* rs917997 C>T polymorphisms in ESCC progression and prognosis were not performed.

In conclusion, the present study provides evidence that functional *IL-18R1* rs13015714 G>T and *IL-18RAP* rs917997 C>T polymorphisms may contribute to the development of ESCC. However, the power of the present analysis was relatively low with a limited sample size, particularly in the subgroup analyses. Therefore, only preliminary conclusions were drawn. Larger studies with other ethnic populations are required to confirm the present findings.

### Acknowledgements

The present study was supported in part by the National Natural Science Foundation of China (grant nos. 81101889 and 81000028), Jiangsu Province Natural Science Foundation (grant nos. BK2010333 and BK2011481), Social Development Foundation of Zhenjiang (grant no. SH2010017) and Changzhou Young Talents and Science-Technology Foundation of Health Bureau (grant no. QN201102). The authors would like to thank Dr Da Ding and Dr Yan Liu (Genesky Biotechnologies Inc., Shanghai, China) for the technical support.

### References

1. Chen W, Zheng R, Zhang S, *et al.*: Annual report on status of cancer in China, 2010. *Chin J Cancer Res* 26: 48-58, 2014.
2. Berrino F, De Angelis R, Sant M, Rosso S, Bielska-Lasota M, Coebergh JW and Santaquilani M; EUROCARE Working group: Survival for eight major cancers and all cancers combined for European adults diagnosed in 1995-99: Results of the EUROCARE-4 study. *Lancet Oncol* 8: 773-783, 2007.
3. Macfarlane GJ and Boyle P: The epidemiology of oesophageal cancer in the UK and other European countries. *J R Soc Med* 87: 334-337, 1994.
4. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J and Jemal A: Global cancer statistics, 2012. *CA Cancer J Clin* 65: 87-108, 2015.
5. Enzinger PC and Mayer RJ: Esophageal cancer. *N Engl J Med* 349: 2241-2252, 2003.
6. Sun JM, Li Q, Gu HY, Chen YJ, Wei JS, Zhu Q and Chen L: Interleukin 10 rs1800872 T>G polymorphism was associated with an increased risk of esophageal cancer in a Chinese population. *Asian Pac J Cancer Prev* 14: 3443-3447, 2013.
7. Yoshimoto T, Tsutsui H, Tominaga K, Hoshino K, Okamura H, Akira S, Paul WE and Nakanishi K: IL-18, although anti-allergic when administered with IL-12, stimulates IL-4 and histamine release by basophils. *Proc Natl Acad Sci USA* 96: 13962-13966, 1999.
8. Zitvogel L, Kepp O, Galluzzi L and Kroemer G: Inflammasomes in carcinogenesis and anticancer immune responses. *Nat Immunol* 13: 343-351, 2012.
9. Born TL, Thomassen E, Bird TA and Sims JE: Cloning of a novel receptor subunit, AcPL, required for interleukin-18 signaling. *J Biol Chem* 273: 29445-29450, 1998.
10. Torigoe K, Ushio S, Okura T, Kobayashi S, Tani M, Kunikata T, Murakami T, Sanou O, Kojima H, Fujii M, *et al.*: Purification and characterization of the human interleukin-18 receptor. *J Biol Chem* 272: 25737-25742, 1997.
11. Salcedo R, Worschech A, Cardone M, *et al.*: MyD88-mediated signaling prevents development of adenocarcinomas of the colon: Role of interleukin 18. *J Exp Med* 207: 1625-1636, 2010.
12. Cheung H, Chen NJ, Cao Z, Ono N, Ohashi PS and Yeh WC: Accessory protein-like is essential for IL-18-mediated signaling. *J Immunol* 174: 5351-5357, 2005.
13. Sareneva T, Julkunen I and Matikainen S: IFN- $\alpha$  and IL-12 induce IL-18 receptor gene expression in human NK and T cells. *J Immunol* 165: 1933-1938, 2000.
14. Zhernakova A, Festen EM, Franke L, *et al.*: Genetic analysis of innate immunity in Crohn's disease and ulcerative colitis identifies two susceptibility loci harboring CARD9 and IL18RAP. *Am J Hum Genet* 82: 1202-1210, 2008.
15. Thompson SR and Humphries SE: Interleukin-18 genetics and inflammatory disease susceptibility. *Genes Immun* 8: 91-99, 2007.

16. Sánchez E, Palomino-Morales RJ, Ortego-Centeno N, Jiménez-Alonso J, González-Gay MA, López-Nevot MA, Sánchez-Román J, de Ramón E, González-Escribano MF, Pons-Estel BA, *et al*; Italian collaborative group: Identification of a new putative functional IL18 gene variant through an association study in systemic lupus erythematosus. *Hum Mol Genet* 18: 3739-3748, 2009.
17. Koskinen LL, Einarsdottir E, Dukes E, Heap GA, Dubois P, Korponay-Szabo IR, Kaukinen K, Kurppa K, Zibera F, Vatta S, *et al*: Association study of the IL18RAP locus in three European populations with coeliac disease. *Hum Mol Genet* 18: 1148-1155, 2009.
18. Suppiah V, Moldovan M, Ahlenstiel G, Berg T, Weltman M, Abate ML, Bassendine M, Spengler U, Dore GJ, Powell E, *et al*: IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 41: 1100-1104, 2009.
19. Eurich D, Boas-Knoop S, Ruehl M, Schulz M, Carrillo ED, Berg T, Neuhaus R, Neuhaus P, Neumann UP and Bahra M: Relationship between the interleukin-28b gene polymorphism and the histological severity of hepatitis C virus-induced graft inflammation and the response to antiviral therapy after liver transplantation. *Liver Transpl* 17: 289-298, 2011.
20. Chen X, Li S, Yang Y, Yang X, Liu Y, Liu Y, Hu W, Jin L and Wang X: Genome-wide association study validation identifies novel loci for atherosclerotic cardiovascular disease. *J Thromb Haemost* 10: 1508-1514, 2012.
21. Babar M, Ryan AW, Anderson LA, Segurado R, Turner G, Murray LJ, Murphy SJ, Johnston BT, Comber H, Reynolds JV, *et al*: Genes of the interleukin-18 pathway are associated with susceptibility to Barrett's esophagus and esophageal adenocarcinoma. *Am J Gastroenterol* 107: 1331-1341, 2012.
22. Spielman RS, Bastone LA, Burdick JT, Morley M, Ewens WJ and Cheung VG: Common genetic variants account for differences in gene expression among ethnic groups. *Nat Genet* 39: 226-231, 2007.