

Development of a polymerase chain reaction-restriction fragment length polymorphism method for identification of the *Fusarium* genus using the transcription elongation factor-1 α gene

MAJID ZARRIN^{1,2}, FARZANEH GANJ² and SAMA FARAMARZI²

¹Health Research Institute, Infectious and Tropical Diseases Research Center; ²Department of Medical Mycology, Medical School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz 61357-15794, Iran

Received July 14, 2016; Accepted September 28, 2016

DOI: 10.3892/br.2016.783

Abstract. *Fusarium* species are well-known plant pathogens and food contaminants that have also appeared as one of the most important groups of medically significant fungi. The sequences of the translation elongation factor (TEF)-1 α gene have been broadly employed for species detection. A total of 50 strains of *Fusarium* spp., including environmental, clinical and reference isolates were used for the current study. The primer sets, Fu3f and Fu3r, were used to amplify an ~420-bp DNA fragment of the TEF-1 α gene. Double digestion with two restriction enzymes, *Xho*I and *Sdu*I was used for discrimination of the *Fusarium* species in the TEF-1 α gene fragment. Double digestion of the TEF-1 α gene fragment from five clinically important *Fusarium* species were clearly differentiated from each other: The *F. solani* species complex, *F. oxysporum* species complex, *F. verticillioides*, *F. proliferatum* and *F. fujikuroi*. This method facilitates detection and enables verification of the *Fusarium* genus; therefore, it may be applied for disease control.

Introduction

Fusarium species are ubiquitous fungi extensively distributed in soil, plants and various organic substrates. This genus is an important plant pathogen, which causes different diseases and is responsible for important economic losses on crops. In humans, the *Fusarium* species causes a broad range of diseases, including superficial, locally invasive, or disseminated infections. Disseminated infections occur almost exclusively in severely immunocompromised patients and, currently,

disseminated infections are the second most common mold that causes invasive fungal infections in immunosuppressed hosts, and is associated with high morbidity and mortality rates (1,2). Furthermore, the *Fusarium* species causes allergic diseases, such as sinusitis in immunocompetent individuals and mycotoxicosis following ingestion of food that is contaminated with toxin-producing *Fusarium* (3,4). This genus contains >70 species (5); a literature review of 259 cases of fusariosis between 2001 and 2005 demonstrated that 12 species were associated with infection. The *F. solani* species complex was the most common (50% of cases), followed by the *F. oxysporum* species complex (20% of cases) and *F. verticillioides* and *F. moniliforme* (10% of cases for each) (6).

Morphological identification of the *Fusarium* species is the primary, but most difficult, step in the detection procedure. However, for the species that cannot be reliably recognized by morphological characterization, additional analysis, such as DNA sequencing and species-specific polymerase chain reaction (PCR) assays, must be performed.

Translation elongation factor (TEF) 1- α consistently presents as a single-copy gene in the *Fusarium* genus. This gene demonstrates a high level of sequence polymorphism among the closely associated species of *Fusarium*, even compared with the intron-rich portions of protein-coding genes, such as β -tubulin, calmodulin and histone H3. Therefore, TEF has become the choice marker as a single-locus detection tool in *Fusarium* (7,8). The strategy that was developed in the present study consisted of novel PCR-restriction fragment length polymorphism (RFLP) analysis for detecting DNA polymorphisms in the TEF-1 α gene and for discrimination of the *Fusarium* genus.

Materials and methods

Microorganisms. Fifty strains of *Fusarium* spp. (including environmental, clinical and reference isolates) were used in the present study. The following reference strains were used: *F. solani* complex PTCC 5284, *F. solani* complex PTCC 5285, *F. oxysporum* complex IBRC-M 30067, *F. oxysporum* complex PTCC 5115, *F. verticillioides* PFCC 53-131, *F. verticillioides* PFCC 15-89, *F. proliferatum* PFCC 48-125, *F. proliferatum* PFCC 12-86 and *F. fujikuroi* PTCC 5144. The environmental

Correspondence to: Dr Majid Zarrin, Department of Medical Mycology, Medical School, Ahvaz Jundishapur University of Medical Sciences, 1 Golestan Street, Ahvaz 61357-15794, Iran
E-mail: mjjzarrin@yahoo.co.uk

Key words: *Fusarium*, polymerase chain reaction, transcription elongation factor-1 α gene

strains were obtained from soil, and two strains used in the present study were clinical, which included *F. solani* complex PTCC 5284 and B988.

DNA extraction. Thick spore suspension (1 ml) was inoculated in Ehrlenmeyer flasks containing yeast extract peptone dextrose medium and incubated on an incubator shaker at 200 rpm under agitation for 72 h at 25°C for mycelia growth. The harvested mycelia were washed with 0.5 M EDTA and sterile dH₂O. The mycelia were ground into a fine powder using liquid nitrogen and a mortar and pestle.

Approximately 100 mg powdered mycelium was transferred into a 1.5-ml tube containing 400 μ l lysis buffer (100 mM Tris-HCl, pH 8.0, 30 mM EDTA, pH 8.0, 5% SDS w/v). After microtubes were boiled at 100°C for 20 min, 3 M acetate potassium (150 μ l) was added to each tube. The suspension was maintained at -20°C for 10 min and centrifuged at 14,000 x g in 4°C for 10 min. The supernatant was carefully transferred to a fresh 1.5-ml Eppendorf tube and 250 μ l phenol:chloroform:isoamyl alcohol (25:24:1, v/v) was added. The microtube was vortexed briefly and centrifuged at 4°C at 14,000 x g for 10 min. After transferring the supernatant to a 1.5-ml microtube, 250 μ l chloroform:isoamyl alcohol was added. The tubes were briefly vortexed and centrifuged at 4°C at 14,000 x g for 10 min. The supernatant was transferred to a fresh microtube, an equal volume of ice-cold 2-propanol was added, maintained at -20°C for 10 min and centrifuged at 14,000 x g for 10 min. The upper aqueous phase was discarded and the pellet was washed with 70% ethanol (300 μ l). The ethanol was discarded and the DNA pellets were air dried and resuspended in 50 μ l dH₂O.

PCR amplification. The primer sets, Fu3f (5'-GGTATCGA CAAGCGAACCAT-3') and Fu3r (5'-TAGTAGCGGGGA GTCTCGAA-3') was used to amplify an ~420-bp DNA fragment of the TEF-1 α gene (9). PCR reactions were performed with a volume of 50 μ l, comprised of 5 μ l 10X reaction buffer, 2.2 mM MgCl₂, 200 μ M each dNTP, 2.5 units of *Taq* DNA polymerase (CinnaGen, Tehran, Iran), 30 ng template DNA and 50 pmol of each primer.

An initial denaturation step for 5 min at 94°C was followed by 30 cycles of denaturation at 94°C for 1 min, annealing at 58°C for 1 min and extension at 68°C for 2 min. The amplified PCR product (5 μ l) was electrophoresed on 1% agarose gel in TAE buffer at 100 V for 1 h and stained with ethidium bromide. The PCR amplification of TEF-1 α gene resulted in an ~420-bp fragment.

RFLP analysis. Digestion with one restriction enzyme was not sufficient to discriminate the 420-bp DNA fragment of the TEF-1 α gene in the *Fusarium* species. Therefore, double digestion with two restriction enzymes, *Xho*I and *Sdu*I (Thermo Fisher Scientific, Inc., Waltham, MA, USA) was used for discrimination. The restriction digestion reaction was performed in a total volume of 20 μ l containing 5 units of each enzyme, 2 μ l Buffer O (Thermo Fisher Scientific, Inc.), 5 μ l PCR product, and Ultrapure water (CinnaGen, Karaj, Iran) to reach a volume of 20 μ l. Digested PCR products were electrophoresed at 50 V for 3 h on 2% agarose gel in TAE buffer and stained with ethidium bromide.

Table I. Restriction fragment size (bp) of the *Fusarium* species TEF-1 α gene, double digested with two restriction enzymes, *Xho*I and *Sdu*I.

<i>Fusarium</i> species	TEF-1 α fragment prior to digestion (bp)	<i>Xho</i> I and <i>Sdu</i> I (bp)
<i>F. oxysporum</i> species complex	420	45,62,103,170
<i>F. verticillioide</i> s	420	6,30,56,47,55,186
<i>F. proliferatum</i>	420	25,168,187
<i>F. fujikuroi</i>	420	27,62,99,192
<i>F. solani</i> species complex	420	308,110

TEF, transcription elongation factor.

Results

PCR amplification of the TEF-1 α gene. The PCR amplification of TEF-1 α gene with Fu3f and Fu3r primers produced a unique band of ~420 bp for all tested *Fusarium* isolates (Fig. 1). The TEF-1 α gene fragment was sequenced for certain isolates, including the reference strains. The BLAST search in NCBI (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) demonstrated the TEF-1 α gene fragment from five clinically important *Fusarium* reference strains, including *F. solani* species complex, *F. oxysporum* species complex, *F. verticillioide*s, *F. proliferatum* and *F. fujikuroi* exhibited 99% homology with the associated sequences deposited in the GenBank database.

Restriction patterns for the *Fusarium* strains. Double digestion of the fragment with restriction enzymes, *Xho*I and *Sdu*I clearly discriminated the *F. solani* species complex, *F. oxysporum* species complex, *F. verticillioide*s, *F. proliferatum* and *F. fujikuroi* from each other (Table I and Fig. 2).

The restriction patterns of one clinical and six environmental *Fusarium* strains following double digestion using *Xho*I and *Sdu*I are presented in Fig. 3. The digestion of the 420-bp fragment from these strains demonstrated different patterns. Strains E4, E16 and E25 were sequenced. A BLAST search showed that strains E4 and E16 exhibited 100% homology with the *F. equiseti* and *F. solani* species complex, respectively and strain E25 exhibited 99% homology with *F. incarnatum*. Therefore, the restriction pattern strain E16 (Fig. 3) was similar to the *F. solani* complex PTCC 5284 (Fig. 2).

Discussion

Identification of filamentous fungi at the species level using classical techniques, such as morphological methods, is difficult and time-consuming. Novel rapid techniques are required in order to verify the *Fusarium* genus on time, particularly for clinical administration of patients. Rapid molecular approaches, such as PCR, DNA hybridization and DNA microarray have been developed and they may replace the classical methods. The major advantages of molecular




Figure 1. Agarose gel electrophoresis of transcription elongation factor-1 α gene products (420 bp) of the *Fusarium* species. Lane M, 100-bp ladder; lane 1, *F. oxysporum* complex IBRC-M 30067; lane 2, *F. verticillioides* PFCC 15-89; lane 3, *F. proliferatum* PFCC 48-125; lane 4, *F. fujikuroi* PTCC 5144; lane 5, *F. solani* complex PTCC 5284. TEF, transcription elongation factor.




Figure 2. Agarose gel electrophoresis of transcription elongation factor-1 α gene products (420 bp) of the *Fusarium* species following double digestion with *Xho*I and *Sdu*I. Lane M, 100-bp ladder; lane 1, *F. oxysporum* complex IBRC-M 30067; lane 2, *F. verticillioides* PFCC 15-89; lane 3, *F. proliferatum* PFCC 48-125; lane 4, *F. fujikuroi* PTCC 5144; lane 5, *F. solani* complex PTCC 5284.

approaches are their specificity and that they are completely discriminative even for closely associated species (8,9).

The majority of molecular techniques are PCR-based, where the primers are typically directed to conserved regions of the ribosomal DNA gene, particularly towards the internal transcribed spacer (ITS) regions. With regard to *Fusarium* spp., analysis of ITS sequencing is considered unreliable for detection of strains, as they contain two paralogous, discrepant ITS sequence forms, which may cause confusion (10,11). The TEF-1 α gene has shown optimal results for the identification of *Fusarium* spp. (12-14).

Guevara-Suarez *et al* (15) used a TEF-1 α gene fragment and performed a multi-locus sequence analysis of the ITS region with the RNA-dependent polymerase subunit II (Rpb2) genes, and recognized the phylogenetic species and circulating haplotypes for *Fusarium* isolates from onychomycosis. The pathogenic isolates to the pecan tree were identified, based on the TEF-1 α gene, as belonging to the *F. chlamydosporum* species complex, *F. graminearum* species complex, *F. proliferatum*, and *F. oxysporum* (16). A TEF-1 α -RFLP technique was described




Figure 3. Agarose gel electrophoresis of transcription elongation factor-1 α gene products (420 bp) of the *Fusarium* species (lane 1, clinical isolate; lanes 2-7, environmental isolates) following double digestion with *Xho*I and *Sdu*I. Lane M, 100-bp ladder; lane 1, B988; lane 2, E4; lane 3, E16; lane 4, E17; lane 5, E18; lane 6, E20; lane 7, E25.

for the identification of the three clades of *F. oxysporum* (17). The particularly effective TEF-1 α gene of the *Fusarium* spp. encouraged the present development of a PCR-RFLP technique as an advanced, simple and reliable method for determination and discrimination of the clinically important *Fusarium* species.

In the current study, molecular identification was performed using the TEF-1 α gene and RFLP, and it was possible to discriminate between all five clinically important *Fusarium* species. However, further analyses are required for discrimination between other *Fusarium* species.

The Primer set, TEF-Fu3 resulted in an ~420-bp product for five of the *Fusarium* species, including *F. solani* species complex, *F. oxysporum* species complex, *F. verticillioides*, *F. proliferatum* and *F. fujikuroi*. RFLP, using double digestion with two restriction enzymes, *Xho*I and *Sdu*I differentiated between the species. This method may facilitate detection, verify the *Fusarium* genus, and be applied for disease control. This PCR-RFLP method is rapid, economical and efficient for detection and discrimination of the *Fusarium* genus.

Acknowledgements

The present study was supported by the Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences (Ahvaz, Iran) (grant no. 92118).

References

1. Consigny S, Dhedin N, Datry A, Choquet S, Leblond V and Chosidow O: Successful voriconazole treatment of disseminated fusarium infection in an immunocompromised patient. *Clin Infect Dis* 37: 311-313, 2003.
2. Nucci M and Anaissie E: Cutaneous infection by *Fusarium* species in healthy and immunocompromised hosts: Implications for diagnosis and management. *Clin Infect Dis* 35: 909-920, 2002.
3. Twarużek M, Soszczyńska E, Winiarski P, Zwierz A and Grajewski J: The occurrence of molds in patients with chronic sinusitis. *Eur Arch Otorhinolaryngol* 271: 1143-1148, 2014.
4. Beccari G, Caproni L, Tini F, Uhlig S and Covarelli L: Presence of fusarium species and other toxigenic fungi in malting barley and multi-mycotoxin analysis by liquid chromatography-high resolution mass spectrometry. *J Agric Food Chem* 64: 4390-4399, 2016.

5. Al-Hatmi AM, Meis JF and de Hoog GS: *Fusarium*: Molecular Diversity and Intrinsic Drug Resistance. *PLoS Pathog* 12: e1005464, 2016.
6. Nucci M and Anaissie E: *Fusarium* infections in immunocompromised patients. *Clin Microbiol Rev* 20: 695-704, 2007.
7. Geiser DM, Jiménez-Gasco MM, Kang S, Makalowska I, Veeraraghavan N, Ward TJ, Zhang N, Kulda GA and O'Donnell K: FUSARIUM-ID v. 1.0: A DNA sequence database for identifying *Fusarium*. *Eur J Plant Pathol* 110: 473-479, 2004.
8. Hsuan HM, Salleh B and Zakaria L: Molecular identification of *Fusarium* species in *Gibberella fujikuroi* species complex from rice, sugarcane and maize from Peninsular Malaysia. *Int J Mol Sci* 12: 6722-6732, 2011.
9. Arif M, Chawla S, Zaidi NW, Rayar JK, Variar M and Singh US: Development of specific primers for genus *Fusarium* and *F. solani* using rDNA sub-unit and transcription elongation factor (TEF-1 α) gene. *Afr J Biotechnol* 11: 444-447, 2012.
10. Dornbusch HJ, Buzina W, Summerbell RC, Lass-Flörl C, Lackner H, Schwinger W, Sovinz P and Urban C: *Fusarium* verticillioides abscess of the nasal septum in an immunosuppressed child: Case report and identification of the morphologically atypical fungal strain. *J Clin Microbiol* 43: 1998-2001, 2005.
11. O'Donnell K and Cigelnik E: Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Mol Phylogenet Evol* 7: 103-116, 1997.
12. Palacios SA, Susca A, Haidukowski M, Stea G, Cendoya E, Ramírez ML, Chulze SN, Farnochi MC, Moretti A and Torres AM: Genetic variability and fumonisin production by *Fusarium proliferatum* isolated from durum wheat grains in Argentina. *Int J Food Microbiol* 201: 35-41, 2015.
13. Watanabe M, Yonezawa T, Sugita-Konishi Y and Kamata Y: Utility of the phylotoxigenic relationships among trichothecene-producing *Fusarium* species for predicting their mycotoxin-producing potential. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 30: 1370-1381, 2013.
14. Scheel CM, Hurst SF, Barreiros G, Akiti T, Nucci M and Balajee SA: Molecular analyses of *Fusarium* isolates recovered from a cluster of invasive mold infections in a Brazilian hospital. *BMC Infect Dis* 13: 49, 2013.
15. Guevara-Suarez M, Cano-Lira JF, de García MC, Sopo L, De Bedout C, Cano LE, García AM, Motta A, Amézquita A, Cárdenas M, *et al*: Genotyping of *Fusarium* isolates from onychomycoses in Colombia: Detection of two new species within the *Fusarium solani* species complex and in vitro antifungal susceptibility testing. *Mycopathologia* 181: 165-174, 2016.
16. Lazarotto M, Milanesi PM, Muniz MF, Reiniger LR, Beltrame R, Harakava R and Blume E: Morphological and molecular characterization of *Fusarium* spp pathogenic to pecan tree in Brazil. *Genet Mol Res* 13: 9390-9402, 2014.
17. Bogale M, Wingfield BD, Wingfield MJ and Steenkamp ET: Species-specific primers for *Fusarium redolens* and a PCR-RFLP technique to distinguish among three clades of *Fusarium oxysporum*. *FEMS Microbiol Lett* 271: 27-32, 2007.