Interleukin-6 gene variants are associated with reduced risk of chronicity in hepatitis B virus infection in a Malaysian population

BEHNAZ RIAZALHOSSEINI¹, ZAHURIN MOHAMED¹,
YAMUNAH DEVI APALASAMY², NOOR SHAFILA SHAFIE¹ and ROSMAWATI MOHAMED³

¹Department of Pharmacology, Faculty of Medicine; ²Social Wellbeing Research Centre, Faculty of Economics and Administration; ³Department of Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur 50603, Malaysia

Received February 7, 2018; Accepted June 29, 2018

DOI: 10.3892/br.2018.1126

Abstract. Interleukin-6 (IL-6) is a cytokine with a critical role in regulating the immune response to infectious disease. Studies have indicated that polymorphisms in the *IL-6* gene may be linked to hepatitis B virus (HBV) infection. The purpose of the present study was to examine the association among *IL-6* SNPs and haplotypes with HBV infection risk in a Malaysian population. A total of 1,246 Malaysian subjects with and without chronic hepatitis B were recruited for this study. Three *IL-6* polymorphisms (rs2069837, rs1800796 and rs2066992) were genotyped using a Sequenom MassARRAY[®] platform. The results suggested that GC and CC genotypes of rs1800796 as well as GT and TT genotypes of rs2066992 were associated with protection against HBV infection ($P < 0.001$). Furthermore, haplotypes GG and CT exhibited a significant association with protection against HBV ($P = 0.003$ and $= 0.005$, respectively); and haplotypes GG and CT exhibited a significant association with clearance of HBV infection ($P = 0.035$ and $= 0.037$, respectively). The present study indicates that two *IL-6* SNPs (rs1800796 and rs2066992) are associated with clearance of chronic HBV or protection against HBV infection at allelic, genotypic and haplotypic levels.

Introduction

Hepatitis B virus (HBV) infection is a potentially severe disease affecting ~2 billion people worldwide (1,2). It was reported by the World Health Organization in 2015 that an estimated ~240 million individuals are chronically infected with the HBV virus and that 780,000 mortalities occur per year due to the consequences of HBV infection including liver cirrhosis (LC) and hepatocellular carcinoma (HCC) (3). The regions with high endemic rates of HBV infection primarily include Southeast Asia, the Western Pacific and Africa, in which incidences of more than 8% of the population have been reported (4). Thus HBV infection is a major public health issue, particularly since it exhibits variable epidemiology between different populations (4).

Numerous studies have revealed complex interactions between the host (genetic background and immune response), the virus and the environment, which may contribute to the outcomes of HBV infection (5-7). In particular, familial and twin studies have confirmed a significant role of host genetic factors in the persistence of HBV infection (8,9).

Several studies, including genome wide association studies (GWAS), have revealed that single nucleotide polymorphisms (SNPs) in certain genes influence the outcomes of HBV infection (10-16). Nevertheless, there are geographical and ethnic differences between these genetic factors and the outcomes of HBV infection (17,18).

Interleukin-6 (*IL-6*) is among the GWAS-identified genes and has been reported to be linked to susceptibility to HBV infection, clearance rate of HBV and also progression to severe liver disease (19-21). *IL-6* is a cytokine with a vital role in the regulation of immune reactions in numerous target cells including hepatocytes (19). Previous data has indicated that host innate immunity serves a crucial role in controlling HBV infection and that *IL-6* is a main factor involved in regulation of the innate immune response (19). It has been reported that *IL-6* derived from activated monocytes serves a significant role in stimulating the immune response required for viral control (22). However, as a mediator of inflammation, it may also be involved in the progression of HBV-associated cirrhosis and HCC (22). Previous study documented that *IL-6* was capable of suppressing HBV replication and inhibiting

Correspondence to: Dr Behnaz Riazalhosseini, Department of Pharmacology, Faculty of Medicine, University of Malaya, University Street, Kuala Lumpur 50603, Malaysia
E-mail: beh.riazi@gmail.com

Abbreviations: IL-6, interleukin-6; HBV, hepatitis B virus; SNPs, single nucleotide polymorphisms; LC, liver cirrhosis; HCC, hepatocellular carcinoma; GWAS, genome wide association studies; HBsAg, HBV surface antigen; OR, odd ratio; CI, confidence interval; HWE, Hardy-Weinberg equilibrium; LD, linkage disequilibrium

Key words: interleukin-6 gene, hepatitis B virus, protective allele, polymorphism, infection resolution

the accumulation of HBV covalently closed circular DNA in human hepatoma cells (23). By contrast, other studies have reported increased serum levels of *IL-6* in patients with chronic hepatitis B (CHB) infection who developed LC/HCC. Hence, the association between *IL-6* gene polymorphisms with susceptibility to HBV infection, clearance of HBV and progression of HBV infection is inconclusive. Therefore, the present study aimed at investigating the association between *IL-6* gene polymorphisms (rs2069837, rs1800796 and rs2066992) and predisposition to HBV infection, clearance rate of the HBV infection and also progression of HBV to LC/HCC. A hospital-based case-control study was performed to examine the association among *IL-6* SNPs and haplotypes with HBV infection risk in a Malaysian population, comprised of Malay, Chinese and Indian ethnic groups.

Materials and methods

Study participants. A total of 1,246 Malaysian subjects were recruited from the University of Malaya Medical Centre, Kuala Lumpur, Malaysia, from June 2012 to October 2014. The patient groups consisted of 423 patients with chronic HBV [persistence of HBV surface antigen (HBsAg) longer than 6 months] without cirrhosis or HCC and 103 patients with chronic HBV with LC and/or HCC (chronic HBV-related LC/HCC). The study cohort also consisted of 97 HBV-resolved subjects (HBsAg negative and anti-HB core antibody positive) and 623 healthy individuals (HBsAg and anti-HB core antibody negative) as controls.

LC among the chronic HBV-infected patients was confirmed by liver biopsies, or a combination of biochemical, clinical and/or radiological indications of cirrhosis (24). Diagnoses of HCC were based on radiological evidence of liver mass with arterial hypervascularity and washout in the venous-delayed phase on dynamic imaging and/or by liver histology (25). Patients with other possible causes of liver disease including primary biliary cirrhosis, Wilson's disease, hepatitis C infection, autoimmune hepatitis, hemochromatosis and α -1 anti-trypsin deficiency were excluded from the study. The study protocol was approved by the Medical Ethics Committee of the University of Malaya Medical Centre (ref. no. 938.42). Ethnicities of the subjects were confirmed by a self-report. Signed and dated informed consent was obtained from all of the subjects.

Selection of SNPs. To examine whether variants in the *IL-6* gene are capable of affecting HBV infection risk in a Malaysian population, three *IL-6* polymorphisms (rs2069837 A>G, rs1800796 G>C and rs2066992 G>T) were selected based on the criteria of previous HBV studies (19-21) in other populations involving candidate genes with minor allele frequency >0.1 from the NCBI online database (<https://www.ncbi.nlm.nih.gov>).

Genotyping of *IL-6* SNPs. Genomic DNA was extracted from buffy coat (isolated from 4 ml peripheral blood following collection in anticoagulant tubes and centrifugation at 1,600 x g for 10 min at room temperature) using a GeneAll[®] Exgene[™] DNA purification kit (GeneAll Biotechnology, Co., Ltd., Seoul, South Korea) according to the manufacturer's instructions. The 3 selected *IL-6* polymorphisms (rs2069837,

rs1800796 and rs2066992) were genotyped using a Sequenom MassARRAY[®] platform (Sequenom, Inc., San Diego, CA, USA) according to relevant guidelines (26). A volume of 1 μ l genomic DNA with a mean average concentration between 10-20 ng/ μ l was used in each amplification reaction. A blank and five duplicates were presented for quality control, and were excluded if any of the following occurred: assay with >10% call rate in blank checks; >25% call rate within the blank control; <99.5% concordance in duplicate checks and assay with <80% call rate in the same SpectroCHIP.

Statistical analysis. Data analysis was performed using SPSS version 16.0 (SPSS, Inc., Chicago, IL, USA). The allele and genotype distributions of the *IL-6* SNPs between the groups were assessed by χ^2 test. Logistic regression was performed to determine the association of *IL-6* SNPs with (i) chronic HBV infection and disease progression, (ii) between chronic and resolved HBV subjects and (iii) between chronic HBV patients and healthy controls. Odd ratios (ORs) with 95% confidence intervals (CI) were calculated following correction for confounding factors including sex and age. The SNPs were tested for deviation from Hardy-Weinberg equilibrium (HWE). Subgroup analysis based on ethnicity of allele and genotype distribution of the *IL-6* SNPs was not performed due to the small sample size of each ethnic group.

Additionally, construction of linkage disequilibrium (LD) blocks and haplotype analysis were performed using Haploview 4.2 software (Broad Institute, Cambridge, MA, USA). Haplotype frequencies were compared between (i) chronic HBV cases with and without progression to LC/HCC, (ii) chronic and resolved HBV subjects and (iii) between chronic HBV patients and healthy controls, by calculating the permutation P-values created using χ^2 statistics with one degree of freedom. $P < 0.05$ was considered to indicate statistical significance. The data were presented as frequency (percentage) or median (range).

Results

General characteristics of subjects. The demographics of the study participants are listed in Table I. The median ages of chronic HBV patients without LC/HCC, chronic HBV patients with LC/HCC, HBV-resolved subjects and healthy controls were 54 (range, 37-69), 66 (range, 51-79), 31 (range, 20-42) and 33 (range, 19-47), respectively.

***IL-6* allele frequencies.** The allelic distributions and frequencies of the 3 *IL-6* SNPs amongst chronic HBV patients, HBV-resolved subjects and healthy controls are presented in Table II. A significant difference in the allelic frequencies of *IL-6* rs2069837, rs1800796 and rs2066992 was observed between chronic HBV patients and healthy controls (OR=0.60, 95% CI=0.47-0.78, $P < 0.001$; OR=0.40, 95% CI: 0.34-0.48, $P < 0.001$; and OR=0.40, 95% CI: 0.33-0.47, $P < 0.001$, respectively). For rs1800796 and rs2066992, the frequency of allele G (mutate allele) among healthy controls (0.303) was higher than that among chronic HBV patients (0.258 for rs1800796 and 0.265 for rs2066992). There were no significant differences in allelic frequencies for rs2069837, rs1800796 and rs2066992 between chronic HBV cases with and without LC/HCC or between chronic HBV

Table I. Demographic of the study subjects.

Variable	Healthy controls (n=623)	Resolved HBV (n=97)	Chronic HBV (n=423)	Chronic HBV-related LC/HCC (n=103)	P-value chronic HBV vs. healthy controls	P-value chronic HBV vs. resolved HBV	P-value chronic HBV with vs. without LC/HCC
Age, years	33 (19-47)	31 (20-42)	54 (37-69)	66 (51-79)	<0.0001	<0.0001	<0.0001
Ethnicity							
Malay	228 (36.6)	41 (42.3)	117 (27.7)	29 (28.2)	0.849	0.461	0.312
Chinese	316 (50.7)	36 (37.1)	295 (69.7)	72 (69.9)	0.527	0.698	0.448
Indian	79 (12.7)	20 (20.6)	11(2.6)	2 (1.9)	0.760	0.391	0.698
Sex							
Male	441 (70.8)	68 (70.1)	218 (51.5)	66 (64.1)	0.084	0.043	0.046
Female	182 (29.2)	29 (29.9)	205 (48.5)	37 (35.9)	0.061	0.039	0.082

Data are presented as frequency (percentage) or median (range). HBV, hepatitis B virus; LC, liver cirrhosis; HCC, hepatocellular carcinoma.

Table II. Allelic distributions and frequencies for *IL-6* gene polymorphisms in chronic HBV patients, HBV-resolved subjects and healthy controls.

Allele	Healthy controls (n=623)	Resolved HBV (n=97)	Chronic HBV (n=423)	Chronic HBV-related LC/HCC (n=103)	Chronic HBV vs. healthy controls		Chronic HBV vs. resolved HBV		Chronic HBV vs. chronic HBV-related LC/HCC		
					OR (95% CI)	P-value	OR (95% CI)	P-value	OR (95% CI)	P-value	
IL6-rs2069837											
A	1,060 (85.1)	164 (84.5)	716 (84.6)	171 (83.0)	0.60 (0.47-0.78)	<0.001	0.99 (0.63-1.56)	0.972	1.16 (0.75-1.79)	0.506	
G	186 (14.9)	30 (15.5)	130 (15.4)	35 (17.0)							
IL6-rs1800796											
G	378 (30.3)	63 (32.5)	219 (25.8)	64 (31.0)	0.40 (0.34-0.48)	<0.001	1.37 (0.97-1.92)	0.066	0.77 (0.55-1.08)	0.130	
C	868 (69.7)	131 (67.5)	627 (74.2)	142 (69.0)							
IL6-rs2066992											
G	389 (31.4)	65 (33.5)	224 (26.5)	52 (27.6)	0.40 (0.33-0.47)	<0.001	1.39 (0.99-1.94)	0.053	0.80 (0.55-1.04)	0.183	
T	857 (68.6)	129 (66.5)	622 (73.5)	154 (72.4)							

Data are presented as frequency (percentage). OR and 95% CI were calculated using logistic regression, adjusted for sex and age. P-values were determined using χ^2 test. IL-6, interleukin-6; HBV, hepatitis B virus; LC, liver cirrhosis; HCC, hepatocellular carcinoma; OR, odds ratio; CI, confidence interval.

Table III. Genotypic distributions for *IL-6* gene polymorphisms in chronic HBV patients, resolved HBV subjects and healthy controls.

Genotype	Healthy controls (n=623)	Resolved HBV (n= 97)	Chronic HBV (n=423)	Chronic HBV-related LC/HCC (n=103)	Chronic HBV vs. healthy controls		Chronic HBV vs. resolved HBV		Chronic HBV vs. chronic HBV-related LC/HCC		
					OR (95% CI)	P-value	OR (95% CI)	P-value	OR (95% CI)	P-value	
IL6-rs2069837											
AA	436 (70.1)	67 (69.1)	299 (70.7)	67 (65.7)	Ref.		Ref.		Ref.		
AG	186 (29.9)	30 (30.9)	118 (27.9)	35 (34.3)	0.91 (0.64-1.29)	0.600	1.03 (0.59-1.83)	0.901	1.34 (0.82-2.19)	0.248	
GG	0	0	6 (1.4)	0	NA		NA		NA		
HWE P-value	<0.001	0.071	0.136	0.036							
IL6- rs1800796											
GG	61 (9.8)	12 (12.3)	28 (6.6)	8 (7.8)	Ref.		Ref.		Ref.		
GC	256 (41.2)	39 (40.1)	163 (38.5)	48 (46.6)	0.06 (0.04-0.09)	<0.001	1.68 (0.66 - 4.25)	0.275	1.15 (0.47- 2.84)	0.759	
CC	305 (49)	46 (47.6)	232 (54.8)	47 (45.6)	0.07 (0.05-0.11)	<0.001	1.83 (0.73-4.54)	0.195	0.77 (0.32-1.90)	0.570	
HWE P-value	0.498	0.412	0.930	0.371							
IL6- rs2066992											
GG	58 (9.3)	12 (12.4)	31 (7.3)	8 (7.8)	Ref.		Ref.		Ref.		
GT	260 (41.8)	41 (42.3)	162 (38.3)	48 (46.6)	0.06 (0.04-0.09)	<0.001	1.53 (0.62-3.82)	0.363	1.34 (0.55-3.27)	0.520	
TT	304 (48.9)	44 (45.4)	230 (54.4)	47 (45.6)	0.07 (0.05-0.11)	<0.001	1.84 (0.75-4.55)	0.186	0.89 (0.37-2.17)	0.805	
HWE P-value	0.823	0.612	0.736	0.371							

Data are presented as frequency (percentage). OR and 95% CI were calculated using logistic regression, adjusted for sex and age. P-values were determined using χ^2 test. IL-6, interleukin-6; HBV, hepatitis B virus; LC, liver cirrhosis; HCC, hepatocellular carcinoma; OR, odds ratio; CI, confidence interval; HWE, Hardy Weinberg equilibrium; NA, not applicable.

Data are presented as frequency (percentage). OR and 95% CI were calculated using logistic regression, adjusted for sex and age. P-values were determined using χ^2 test. IL-6, interleukin-6; HBV, hepatitis B virus; LC, liver cirrhosis; HCC, hepatocellular carcinoma; OR, odds ratio; CI, confidence interval; HWE, Hardy Weinberg equilibrium; NA, not applicable.

Table IV. Haplotype analysis of *IL-6* rs1800796 and rs2066992 association between chronic HBV patients, healthy controls and HBV-resolved subjects.

Comparison	Haplotype	Frequency					D'
		All	Case ^a	Control ^b	χ^2	P-value	
Chronic HBV vs. healthy controls	Block 1						0.98
	CT	0.709	0.745	0.688	7.727	0.005	
	GG	0.282	0.245	0.304	8.423	0.003	
Chronic HBV vs. resolved HBV	Block 1						0.99
	CT	0.720	0.752	0.694	4.309	0.037	
	GG	0.269	0.237	0.295	4.443	0.035	
Chronic HBV vs. chronic HBV- related LC/HCC	Block 1						0.99
	CT	0.724	0.689	0.733	0.129	2.312	
	GG	0.266	0.311	0.255	0.114	2.108	

^aCase for comparison 1 is chronic HBV patients, for comparison 2 is chronic HBV patients, for comparison 3 is HBV-related LC/HCC; ^bcontrol for comparison 1 is healthy controls, for comparison 2 is resolved HBV subjects and for comparison 3 is chronic HBV. *IL-6*, interleukin-6; HBV, hepatitis B virus; LC, liver cirrhosis; HCC, hepatocellular carcinoma.

patients and HBV-resolved subjects; although association was indicated for *IL-6* rs2066992 between chronic HBV patients and HBV-resolved HBV subjects ($P=0.053$).

***IL-6* genotype frequencies.** The genotypic distributions of the 3 *IL-6* SNPs among chronic HBV patients, HBV-resolved subjects and healthy controls are presented in Table III. A significant difference in the genotype frequencies of *IL-6* rs1800796 was identified between chronic HBV patients and healthy controls (OR=0.06, 95% CI: 0.04-0.09, $P<0.001$ for GC genotype and OR=0.07, 95% CI: 0.05-0.11, $P<0.001$ for CC genotype). Similarly, a significant association was observed in genotype frequencies of *IL-6* rs2066992 between chronic HBV patients and healthy controls (OR=0.06; 95% CI: 0.04-0.09, $P<0.001$ for GT genotype and OR=0.07, 95% CI: 0.05-0.11, $P<0.001$ for TT genotype). There were no significant differences in the genotypic frequencies of rs2069837, rs1800796 and rs2066992 between chronic HBV cases with and without LC/HCC or between chronic HBV patients and HBV-resolved subjects. The genotype distribution of rs2069837, rs1800796 and rs2066992 were in HWE ($P>0.05$ for each) except for rs2069837 in chronic HBV-related LC/HCC cases ($P=0.036$) and the healthy control group ($P<0.001$).

***IL-6* haplotype frequencies.** The results of three haplotype analyses performed between (i) chronic HBV cases with and without LC/HCC, (ii) chronic HBV patients and HBV-resolved subjects and (iii) chronic HBV patients and healthy controls are presented in Table IV. The haplotype combination, CT and GG, was generated between *IL6* rs1800796 and rs2066992. *IL6* rs2069837 was excluded from analysis due to deviation from HWE. Significant associations with strong LD ($D'=0.98$) was observed of haplotypes CT and GG in the comparison of chronic HBV patients vs. healthy controls ($P=0.005$, frequency 0.709 and $P=0.003$, frequency 0.282, respectively). Similarly, the haplotype results on comparing the chronic HBV and HBV-resolved groups indicated significant association with strong LD ($D'=0.99$).

Figure 1. LD analysis of two *IL-6* SNPs (rs1800796 and rs2066992) in three comparisons. (A) Pairwise LD between chronic HBV patients and healthy controls. (B) Pairwise LD between chronic HBV patients and HBV-resolved subjects. (C) Pairwise LD between chronic HBV patients with and without cirrhosis/HCC. LD, linkage disequilibrium; *IL-6*, interleukin-6; SNPs, single nucleotide polymorphisms; HBV, hepatitis B virus; HCC, hepatocellular carcinoma.

Haplotype CT ($P=0.037$, frequency 0.720) and haplotype GG ($P=0.035$, frequency 0.269) were significantly associated with clearance of the HBV virus. Although the *IL6* rs1800796 and rs2066992 SNPs between chronic HBV cases with and without LC/HCC were in strong LD ($D'=0.99$), there were no significant associations with progression of chronic HBV infection. The LD pattern in the three comparisons is depicted in Fig. 1.

Discussion

In the present study, two SNPs (rs1800796 and rs2066992) in the *IL-6* gene were identified to be associated with protection against HBV infection in a Malaysian population. It was observed that the G allele in both SNPs (rs1800796 and rs2066992) may have a significant protective role against HBV infection. The results also indicated that the GG and CT haplotypes of *IL6* rs1800796 and rs2066992 were associated with protection against HBV infection (in comparison with healthy controls) as well as with clearance of HBV (in comparison with HBV-resolved subjects). For *IL-6* rs1800796, the genotypes GG and GC containing the protective allele G were more frequently identified among healthy controls as compared with chronic HBV patients. Saxena *et al* (27) similarly reported that the GG genotype of *IL-6* rs1800796 was more frequent in healthy controls as compared with chronic HBV patients. However, for *IL-6* rs2066992, genotypes GG and GT containing the protective allele G were predominant among healthy controls as compared with chronic HBV patients (27).

There was no significant difference between the ethnicity and the study subjects. Nonetheless, the study had higher number of Chinese participants compared to Malays and Indian. This is probably due to the fact that CHB infection is generally common among Chinese in Malaysia (28).

Host genetic background and its interaction with HBV may affect the outcomes of long-term infection (29). *IL-6* has been demonstrated to be involved in HBV infection *in vitro* and *in vivo* (30). Emerging data has revealed that the *IL-6* polymorphism rs1800796 may be associated with differences in the outcomes of hepatitis B infection including progression to LC/HCC as well as clearance of the virus (19,20). It is speculated that genetic alteration in the *IL-6* gene has an effect on immune dysfunction, which may be involved in the pathogenesis of HBV infection (31). It has been reported that variation in the *IL-6* gene resulting in functional alteration, which may be responsible for changes in the levels of transcription, protein expression and serum *IL-6*, were correlated with HBV disease outcomes (22). Furthermore, it has been indicated that *IL-6* contributes to the interaction between the hepatocyte plasma membrane and the HBV particle (32); HBV could then bind to the preS1 domain on human hepatocytes, mediating HBV infection (31). As a result of activating these signaling pathways, *IL-6* may increase the activity of the HBV enhancer 1 to control the expression of HBV X protein and the replication of HBV (32). A study by Chou *et al* (33) reported that *IL-6* in hepatoma cells promoted HBV transcription by activating signal transducer and activator of transcription 3, which may interact with hepatocyte nuclear factor 3 bound to the HBV enhancer.

In the present study, the rs1800796-G allele (mutant allele) was identified to be the minor allele compared with the C allele (wild-type allele) among healthy controls. A similar result was observed for rs2066992-G allele (mutant allele) as compared with the T allele (wild-type allele) in the same group. In addition, it was identified that rs1800796 and rs2066992 genotypes were significantly associated with protection from HBV infection. However, no association was observed on comparison between chronic HBV and HBV-resolved subjects or between chronic

HBV cases with and without LC/HCC. This is contradictory to a study by Lu *et al* (19), which reported that genetic variation in the SNP *IL-6*-572 C/G may be associated with clearance of HBV infection. Furthermore, the results of another study revealed that genetic variation in the SNP *IL-6*-572 may be associated with HCC incidence in males (34).

In the present study no significant difference was identified in the genotype/allele frequencies of rs2069837, rs1800796 and rs2066992 between chronic HBV patients with and without LC/HCC. Hence, these SNPs are likely not associated with progression to liver disease in HBV-infected patients in the Malaysian population. Furthermore, it was observed that these *IL-6* variants were not associated with clearance of HBV infection. Similarly, Park *et al* (35) reported that *IL-6* variants were not associated with HBV infection or the progression of HBV-associated liver disease.

Considering that haplotypes may be more effective markers of association regarding disease-linked genes than SNPs (36), the association of *IL-6* haplotypes with progression and clearance of HBV infection was examined. All three SNPs are common variants including the functional polymorphism of the *IL-6* promoter (rs1800796) and the SNPs in the intron 2 region (rs2069837 and rs2066992) (37). Significant associations were observed for haplotypes GG and CT on comparison of individuals with chronic HBV with healthy controls and HBV-resolved subjects. Notably, the haplotype results indicated that the wild-type haplotype (GG) had higher frequency in healthy controls as compared with chronic HBV patients. Similarly, the haplotype GG had higher frequency in HBV-resolved patients as compared with chronic HBV patients. However, the haplotype CT exhibited a significant association with chronic HBV as compared with healthy controls and HBV-resolved subjects, with the highest frequency observed in chronic HBV patients.

The present study to the best of our knowledge is the first to investigate the association among *IL-6* SNPs and HBV infection risk in a Malaysian population. Nonetheless, the present study had limitations with regard to functional insights into the role of *IL-6* in HBV infection. Thus, functional studies using mammalian cell lines are required in order to investigate the transcriptional activities of the *IL-6* SNPs.

In conclusion, association was identified between polymorphic variants of the wild-type alleles of rs1800796 and rs2066992 of *IL-6* and protection from HBV infection among a Malaysian population. Similarly, Lu *et al* reported that allele G was associated with the natural elimination of HBV infection (19). Furthermore, study has indicated that the G allele is associated with significantly elevated circulating levels of *IL-6* (27). These findings are consistent with current results and support the apparent protective role of allele G in HBV infection. It was also indicated that haplotypes GG and CT of *IL-6* may have a significant protective role with regard to hepatitis B disease outcomes in Malaysian individuals. However, functional studies and more diverse cohorts are required to provide wider insight into the role of *IL-6* in HBV infection.

Acknowledgements

Not applicable.

Funding

The present study was supported by the High Impact Research Ministry of Higher Education Malaysia (HIR MOHE; grant no. H-20001-00-E000063) and by a University of Malaya postgraduate research grant (grant no. PG146-2014B). The funders had no role in study design, data collection or analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Authors' contributions

BR, ZM and RM conceived and designed the study. BR and NSS performed sample collection and DNA extraction. BR performed the assays, analysed the data and wrote the manuscript. ZM, RM and YDA provided critical revisions on the article. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The present study was approved by the Medical Ethics Committee of the University of Malaya Medical Centre, Kuala Lumpur, Malaysia (ref. no. 938.42). Signed and dated informed consent was obtained from all of the subjects prior to their participation in the study.

Patient consent for publication

Written informed consent was obtained from each participant agreeing to the publication of their data on the terms of data anonymization.

Competing interests

The authors declare that they have no competing interests.

References

- Tang S, Liu Z, Zhang Y, He Y, Pan D, Liu Y, Liu Q, Zhang Z and Yuan Y: Rather than Rs1800796 polymorphism, expression of interleukin-6 is associated with disease progression of chronic HBV infection in a Chinese Han population. *Dis Markers* 35: 799-805, 2013.
- Zhang P, Li N, Zhu Q, Li F, Yang C, Zeng X, Lv Y, Zhou Z, Han Q and Liu Z: Association between TNFAIP3 nonsynonymous single-nucleotide polymorphism rs2230926 and chronic hepatitis B virus infection in a Chinese Han population. *Virol J* 12: 33, 2015.
- World Health Organization (WHO): Hepatitis B. Fact sheet No. 204. WHO, Geneva, 2015. http://www.who.int/mediacentre/factsheets/fs204_Jul2014/en/. Updated July, 2015.
- Papastergiou V, Lombardi R, MacDonald D and Tsochatzis EA: Global Epidemiology of Hepatitis B Virus (HBV) Infection. *Curr Hepatol Rep* 14: 171-178, 2015.
- Zeng Z: Human genes involved in hepatitis B virus infection. *World J Gastroenterol* 20: 7696-7706, 2014.
- Araujo NM, Waizbort R and Kay A: Hepatitis B virus infection from an evolutionary point of view: How viral, host, and environmental factors shape genotypes and subgenotypes. *Infect Genet Evol* 11: 1199-1207, 2011.
- Ali A, Abdel-Hafiz H, Suhail M, Al-Mars A, Zakaria MK, Fatima K, Ahmad S, Azhar E, Chaudhary A and Qadri I: Hepatitis B virus, HBx mutants and their role in hepatocellular carcinoma. *World J Gastroenterol* 20: 10238-10248, 2014.
- He YL, Zhao YR, Zhang SL and Lin SM: Host susceptibility to persistent hepatitis B virus infection. *World J Gastroenterol* 12: 4788-4793, 2006.
- Guo X, Zhang Y, Li J, Ma J, Wei Z, Tan W and O'Brien SJ: Strong influence of human leukocyte antigen (HLA)-DP gene variants on development of persistent chronic hepatitis B virus carriers in the Han Chinese population. *Hepatology* 53: 422-428, 2011.
- Mbarek H, Ochi H, Urabe Y, Kumar V, Kubo M, Hosono N, Takahashi A, Kamatani Y, Miki D, Abe H, *et al*: A genome-wide association study of chronic hepatitis B identified novel risk locus in a Japanese population. *Hum Mol Genet* 20: 3884-3892, 2011.
- Al-Qahtani A, Khalak HG, Alkuraya FS, Al-hamoudi W, Alswat K, Al Balwi MA, Al Abdulkareem I, Sanai FM and Abdo AA: Genome-wide association study of chronic hepatitis B virus infection reveals a novel candidate risk allele on 11q22.3. *J Med Genet* 50: 725-732, 2013.
- Hu Z, Liu Y, Zhai X, Dai J, Jin G, Wang L, Zhu L, Yang Y, Liu J, Chu M, *et al*: New loci associated with chronic hepatitis B virus infection in Han Chinese. *Nat Genet* 45: 1499-1503, 2013.
- Jiang DK, Sun J, Cao G, Liu Y, Lin D, Gao YZ, Ren WH, Long XD, Zhang H, Ma XP, *et al*: Genetic variants in STAT4 and HLA-DQ genes confer risk of hepatitis B virus-related hepatocellular carcinoma. *Nat Genet* 45: 72-75, 2013.
- Li S, Qian J, Yang Y, Zhao W, Dai J, Bei JX, Foo JN, McLaren PJ, Li Z, Yang J, *et al*: GWAS identifies novel susceptibility loci on 6p21.32 and 21q21.3 for hepatocellular carcinoma in chronic hepatitis B virus carriers. *PLoS Genet* 8: e1002791, 2012.
- Kim YJ, Kim HY, Lee JH, Yu SJ, Yoon JH, Lee HS, Kim CY, Cheong JY, Cho SW, Park NH, *et al*: A genome-wide association study identified new variants associated with the risk of chronic hepatitis B. *Hum Mol Genet* 22: 4233-4238, 2013.
- Nishida N, Sawai H, Matsuura K, Sugiyama M, Ahn SH, Park JY, Hige S, Kang JH, Suzuki K, Kurosaki M, *et al*: Genome-wide association study confirming association of HLA-DP with protection against chronic hepatitis B and viral clearance in Japanese and Korean. *PLoS One* 7: e39175, 2012.
- El-Serag HB: Hepatocellular carcinoma. *N Engl J Med* 365: 1118-1127, 2011.
- Karlsen TH, Melum E and Franke A: The utility of genome-wide association studies in hepatology. *Hepatology* 51: 1833-1842, 2010.
- Lu Y, Peng J, Wang C, Zhu Y, Wang F and Sun Z: IL-6 promoter functional polymorphism -572C/G affects spontaneous clearance of hepatitis B virus infection. *Clin Lab* 60: 1903-1907, 2014.
- Tang S, Yuan Y, He Y, Pan D, Zhang Y, Liu Y, Liu Q, Zhang Z and Liu Z: Genetic polymorphism of interleukin-6 influences susceptibility to HBV-related hepatocellular carcinoma in a male Chinese Han population. *Hum Immunol* 75: 297-301, 2014.
- Zhao X-M, Gao Y-F, Zhou Q, Pan F-M and Li X: Relationship between interleukin-6 polymorphism and susceptibility to chronic hepatitis B virus infection. *World J Gastroenterol* 19: 6888-6893, 2013.
- Lan T, Chang L, Wu L and Yuan Y-F: IL-6 plays a crucial role in HBV infection. *J Clin Transl Hepatol* 3: 271-276, 2015.
- Kuo TM, Hu CP, Chen YL, Hong MH, Jeng KS, Liang CC, Chen ML and Chang C: HBV replication is significantly reduced by IL-6. *J Biomed Sci* 16: 41, 2009.
- Chook JB, Ngeow YF, Yap SF, Tan TC and Mohamed R: Combined use of wild-type HBV precore and high serum iron marker as a potential tool for the prediction of cirrhosis in chronic hepatitis B infection. *J Med Virol* 83: 594-601, 2011.
- Bruix J and Sherman M: American Association for the Study of Liver Diseases: Management of hepatocellular carcinoma: An update. *Hepatology* 53: 1020-1022, 2011.
- Bradić M, Costa J and Chelo IM: Genotyping with Sequenom. *Methods Mol Biol* 772: 193-210, 2011.
- Saxena R, Chawla YK, Verma I and Kaur J: IL-6(-572/-597) polymorphism and expression in HBV disease chronicity in an Indian population. *Am J Hum Biol* 26: 549-555, 2014.
- Yap SF: Chronic hepatitis B infection in Malaysians. *Malays J Pathol* 16: 3-6, 1994.
- Tunçbilek S: Relationship between cytokine gene polymorphisms and chronic hepatitis B virus infection. *World J Gastroenterol* 20: 6226-6235, 2014.
- Galun E, Nahor O, Eid A, Jurim O, Rose-John S, Blum HE, Nussbaum O, Ilan E, Daudi N, Shouval D, *et al*: Human interleukin-6 facilitates hepatitis B virus infection in vitro and in vivo. *Virology* 270: 299-309, 2000.

31. De Meyer S, Gong ZJ, Suwandhi W, van Pelt J, Soumillion A and Yap SH: Organ and species specificity of hepatitis B virus (HBV) infection: A review of literature with a special reference to preferential attachment of HBV to human hepatocytes. *J Viral Hepat* 4: 145-153, 1997.
32. Ohno H, Kaneko S, Kobayashi K and Murakami S: Human hepatitis B virus enhancer 1 is responsive to human interleukin-6. *J Med Virol* 52: 413-418, 1997.
33. Chou CH, Chen PJ, Jeng YM, Cheng AL, Huang LR and Cheng JC: Synergistic effect of radiation and interleukin-6 on hepatitis B virus reactivation in liver through STAT3 signaling pathway. *Int J Radiat Oncol Biol Phys* 75: 1545-1552, 2009.
34. Liu S, Qiu XQ, Zeng XY, Bai H, Bei CH and Yang Y: Relationship between IL6 -572G/C polymorphism and hepatocellular carcinoma in men. *Zhonghua Gan Zang Bing Za Zhi* 20: 463-467, 2012 (In Chinese).
35. Park BL, Lee HS, Kim YJ, Kim JY, Jung JH, Kim LH and Shin HD: Association between interleukin 6 promoter variants and chronic hepatitis B progression. *Exp Mol Med* 35: 76-82, 2003.
36. Manolio TA, Brooks LD and Collins FS: A HapMap harvest of insights into the genetics of common disease. *J Clin Invest* 118: 1590-1605, 2008.
37. Chen J, Liu RY, Yang L, Zhao J, Zhao X, Lu D, Yi N, Han B, Chen XF, Zhang K, *et al*: A two-SNP IL-6 promoter haplotype is associated with increased lung cancer risk. *J Cancer Res Clin Oncol* 139: 231-242, 2013.