

Protective functions of *Lycium barbarum* polysaccharides in H₂O₂-injured vascular endothelial cells through anti-oxidation and anti-apoptosis effects

SHUJING XUE^{1*}, XIAOHUI HU^{1*}, LINGIN ZHU², LIHONG NIE¹ and GUANGHUA LI^{1,2}

¹School of Basic Medical Science and ²School of Public Health and Management, Ningxia Medical University, Yinchuan, Ningxia 750004, P.R. China

Received January 15, 2018; Accepted January 18, 2018

DOI: 10.3892/br.2019.1240

Abstract. Cell injury in the cardiovascular endothelia caused by oxidative stress is among the major inducers of endothelium dysfunction and serves an important role in initiating cardiovascular diseases (CVDs). Therefore, protecting and improving the normal function of endothelial cells are considered key measures against CVDs. As a traditional Chinese medicinal component, *Lycium barbarum* is regarded to have high medicinal value. The present study aimed to investigate the potential anti-apoptosis and anti-oxidation effects of *Lycium barbarum* polysaccharides (LBPs) on injured rat artery endothelial cells, to demonstrate the experimental and medicinal values of LBPs. In the present study, the aortic endothelial cells of rats were cultivated and randomly divided into five groups: A control group, H₂O₂-injured group (H₂O₂ group), H₂O₂+LBPs (110 µg/ml) group (low-dose group, LT), H₂O₂+LBPs (220 µg/ml) group (medium-dose group, MT) and H₂O₂+LBPs (440 µg/ml) group (high-dose group, HT). Among these, the activity of superoxide dismutase (SOD), and the levels of malondialdehyde (MDA) and nitric oxide (NO) were detected by colorimetry. Additionally, the expression of B-cell lymphoma-2 (Bcl-2) and Bcl-2-associated X protein (Bax) were detected by western blotting. It was observed that SOD activity and NO content decreased while MDA content increased significantly in the H₂O₂ group (P<0.05 vs. control); that SOD activity in the MT and HT group, and NO content in all three LBP groups were increased, while MDA content in the three LBP groups was decreased, compared with the H₂O₂ group (all P<0.05); that Bcl-2 expression decreased

significantly in the H₂O₂ group while the expression of Bax increased significantly compared with the control group (both P<0.05); and that Bcl-2 expression in all three LBP groups increased, while Bax expression in the MT and HT groups decreased compared with the H₂O₂ group (all P<0.05), with these altered Bax levels being statistically similar to those in the control group (P>0.05). On light microscopy, the cells in the control group exhibited spindle-shaped morphology, consistent sizes, defined boundaries, and distinct nuclei of equivalent sizes with round or oval morphology. Additionally, the chromatin in the nuclei was evenly distributed, and all cells were adhered in a paving-stone arrangement. Notably, only few cells died. Conversely, the cells in the H₂O₂ group exhibited signs of damage and enlarged gaps, and focal cells died. In the HT group, the cells once again appeared adherent and exhibited similar morphological status to the normal cells. Overall, these results indicate that LBPs serve a protective role in oxidative-injured vascular endothelial cells through anti-apoptosis and anti-oxidation effects.

Introduction

Cardiovascular diseases (CVDs), also termed circulatory system diseases, affect the heart and corresponding blood vessels and are established as a worldwide leading cause of mortality (1). In 2014, over 1.7 million fatalities occurred worldwide as a result of CVDs, with a total mortality rate of ~31%, and due to the availability of healthcare, populations in developing countries are more susceptible to CVDs (2). By 2030, it is estimated that the total number of mortalities from CVDs will exceed 2.3 million. To combat this, the World Health Organization plans to reduce the rates of non-communicable diseases (due to living, behavioral and environmental causes) to 25% by 2025, particularly with regard to CVDs (3).

Vascular endothelial cells (VECs) form a continuous cell mass layer over the endangium of vessels in the body. They not only complete the metabolic exchange of blood and interstitial fluid, but also serve as the largest endocrine gland in organisms (4). VECs produce and secrete over ten types of bioactive substances that participate in normal blood flow and blood coagulation, adjust vessel tensions, and control the proliferation and functions of smooth muscles (5). In addition,

Correspondence to: Dr Guanghua Li, School of Basic Medical Science, Ningxia Medical University, 1160 Shengli Road, Yinchuan, Ningxia 750004, P.R. China
E-mail: ghlee0404@163.com

*Contributed equally

Key words: oxidative stress, *Lycium barbarum*, endothelial cells, cell apoptosis

VECs also serve an important role in adjusting the steady state of the cardiovascular system (6). The circulatory system, particularly the blood vessel component, is among the main targets of oxidative stress. Consequently, VECs serve a physiological defense function of 'first line of defense' against stress injuries to organisms (7,8). The function of endothelial cells is associated with the occurrence and development of CVDs (9). Among numerous causes for injury to endothelial cells, oxidative stress is an important inducer of vascular endothelial dysfunction (10). Therefore, the apoptosis of endothelial cells in the cardiovascular system is a major indicator of oxidative stress injury, and a prelude for the occurrence and evolution of various CVDs (11,12).

Lycium barbarum polysaccharides (LBPs), water-soluble protein polysaccharides, are a constituent separated and extracted from the red medlar (also known as wolfberry) fruit of *L. barbarum*, a commonly used medicinal herb in Ningxia, China (13). LBPs have diverse functions, including eye-protective, anti-aging, anti-oxidant, immune adjustment, neuroprotective and anti-tumor effects (14-18). Notably, LBPs may protect the rat reproductive system through an anti-oxygenation mechanism; they may also prolong the apoptosis time of rat epithelial cells, inhibit ultraviolet light-induced peroxidase and free radical-induced cytochrome C expression, and reduce the oxidation of fractured DNA in rat testis cells. Furthermore, LBPs have been reported to markedly promote serum sex hormone levels and superoxide dismutase (SOD) activity, decrease malondialdehyde (MDA) level, decrease injury to spermatogenic cells caused by high temperature and H₂O₂, and promote normal development of germ cells of the testis (19). However, there is limited research on whether LBPs protect against oxidative stress in artery endothelial cells to prevent and cure CVDs. Therefore, the present study aimed to investigate the protective effects of LBPs, particularly regarding anti-apoptosis and anti-oxidation, in injured rat artery endothelial cells (RAECs), to thus provide experimental and theoretical foundations for the medicinal use of LBPs.

Materials and methods

Materials and equipment. The materials used in the present study included RAECs (Jiangsu Chi Scientific Co., Ltd.), complete Dulbecco's modified Eagle's medium (DMEM) containing FBS (both 3-7202; Jiangsu Chi Scientific Co., Ltd.), a Cell Counting kit-8 (CCK-8; Dojindo Molecular Technologies, Inc.), LBPs (Ningxia Qiyuan Pharmaceutical, Co., Ltd.), anti-B-cell lymphoma 2 (Bcl-2) antibody (cat. no. ab59348; Abcam), anti-Bcl-2-associated X protein (Bax) antibody [E63] (cat. no. ab32503; Abcam), and β -actin monoclonal antibody (cat. no. TA811000), horseradish peroxidase (HRP)-goat anti-rat immunoglobulin G (IgG) (cat. no. TA130038) and HRP-goat anti-rabbit IgG (cat. no. TA130023) all from OriGene Technologies, Inc. A total protein extraction kit (KGP250) and bicinchoninic acid assay (BCA) kit (KGP902; Nanjing KeyGen Biotech. Co., Ltd.), a SOD assay kit-WST-1, a Microscale MDA assay kit (thiobarbituric acid method) and a nitric oxide (NO) assay kit (A012-1-2; Colorimetric; Nanjing Jiancheng Bio-Engineering Institute Co., Ltd.) were also used. The major instruments included electrophoresis apparatus, a Trans-Blot SD Semi-Dry Transfer Cell (Bio-Rad Laboratories,

Inc.), a light microscope (Olympus Corporation), an Amersham Imager 600 (GE Healthcare Life Sciences) and a CO₂ incubator (Heraeus).

Cultivation and experimental grouping. RAECs cell line were cultivated in complete Dulbecco's modified Eagle's medium (DMEM) to a confluent state (80-90%) for passage at 37°C in the CO₂ incubator at 5% CO₂. The cells were then divided into five groups: A control group, H₂O₂ injury group (H₂O₂ group), H₂O₂+LBPs (110 μ g/ml) group (low-dose group, LT), H₂O₂+LBPs (220 μ g/ml) group (medium-dose group, MT) and H₂O₂+LBPs (440 μ g/ml) group (high-dose group, HT). The doses of LBPs and H₂O₂ were selected on the basis of preliminary experiments, as follows, in which they were safe and effective without causing toxicity to the RAECs.

Oxidative injury model induced by H₂O₂ analyzed through CCK-8. RAECs, cultured in DMEM supplemented with 10% FBS were seeded in 96-well plates at 2x10⁴ cells/well, treated with 0, 50, 100, 200, 300, 400 and 500 μ mol/l H₂O₂ and incubated at 37°C with 5% CO₂ for 2 h. Subsequently, cell viability was detected using CCK-8 as described previously (20) and cells were detected under a microplate reader at wavelength 450 nm. The optical densities (ODs) were recorded and used to calculate the cell survival rate at each H₂O₂ concentration. The H₂O₂ concentration that retained 50% survival of the cells was selected and used to establish an RAEC oxidative injury model for subsequent experiments.

Influence of LBPs on RAEC survival rate analyzed through CCK-8. Cells were seeded in 96-well plates at 2x10⁴ cells/well and treated with 0.0 (control), 27.5, 55.0, 110.0, 220.0, 440.0, 880.0, 4,400 and 8,800.0 μ g/ml LBPs at 37°C with 5% CO₂ for 2 h. The CCK-8 assay was performed as above. The cells were then detected under a microplate reader at wavelength 450 nm. The ODs were recorded and used to calculate the cell survival rate at each LBP concentration. The concentrations that exerted an inhibitory effect on RAECs were removed.

Determination of LBP effect on oxidative injury in RAECs. Cells were treated with H₂O₂ at the 50% inhibitory concentration and the LBP concentrations that did not result in inhibition of cell viability. Under these established interventions, cells were seeded in 96-well plates at 2x10⁴ cells/well and cultured at 37°C with 5% CO₂ for 2 h. The CCK-8 assay was performed as above. Subsequently, the cells were detected under a microplate reader at wavelength 450 nm. The ODs were recorded and used to calculate the cell survival rates of all groups. The optimum protective clinical concentration of LBPs against oxidative injury in the RAECs was screened out and used in the following experiments.

Morphological observation of all groups. The cells in the established treatment groups (control, H₂O₂, LT, MT and HT groups) were placed under an inverted light microscope. Following removal of the medium, the cell morphologies and quantities were observed at x200 magnification.

Measurement of SOD, MDA and NO. Oxidative stress participates in numerous pathways associated with pathological

changes in endothelial cells (3). To determine whether LBP intervention could remove reactive oxygen species (ROS) and inhibit ROS-mediated oxidative stress, the cell culture was collected and centrifuged at room temperature for 5 min at 3,000 x g, and the commercial kits were used to detect the levels of SOD activity, MDA and NO by colorimetric methods, according to the manufacturer's instructions.

Western blot analysis of Bcl-2 and Bax expression. Oxidative stress ultimately causes cell apoptosis, and thus apoptosis factors were studied to determine whether LBPs could inhibit apoptosis caused by oxidative stress. The typical index of the Bcl class was selected, including the inhibitor of apoptosis protein Bcl-2 and the pro-apoptotic protein Bax, which were used to indirectly assess whether LBPs prevented apoptosis. Total protein was extracted from cells using the protein extraction kit, and quantified using the BCA kit, according to the manufacturer's instructions. Proteins (80 μ g) were separated on 12% SDS-PAGE gels and transferred to polyvinylidene fluoride membranes. Membranes were then blocked (5% non-fat dry milk in PBS plus 0.1% Tween-20; room temperature; 2 h), incubated with primary antibody (4°C; overnight) and second antibody (room temperature; 2 h) and then quantified. The primary antibodies used were anti-Bcl-2, diluted in 1% bovine serum albumin (BSA; Thermo Fisher Scientific, Inc.) to 1:200, anti-Bax, diluted in 1% BSA to 1:3,000, and anti- β -actin, diluted in 1% BSA to 1:500; the secondary antibodies were HRP-goat anti-rat IgG and HRP-goat anti-rabbit IgG, diluted in 1% BSA to 1:5,000. Bands were quantified following detection with the ECL kit using Image Lab 5.0 (Bio-Rad Laboratories, Inc.).

Statistical analysis. Statistical analysis was conducted on SPSS 16.0 (SPSS, Inc.). Data were expressed as the mean \pm standard deviation of at least three independent replicate experiments. The groups were compared by one-way analysis of variance followed by Fisher's least significant difference post hoc analysis, with a significance level of $P < 0.05$.

Results

Optimal concentration of H_2O_2 for the oxidative injury model. H_2O_2 injured cells in a dose-dependent manner. The concentration of H_2O_2 that caused ~50% death rate in RAECs was 100 μ mol/l ($P < 0.05$ vs. control; Fig. 1). Therefore, 100 μ mol/l was selected for the oxidative injury model to be tested in subsequent assays.

Selection of LBP dosage. The RAECs was not affected by different concentrations of LBPs except at the concentration of 8,800 μ g/ml, which significantly reduced cell viability compared with the control ($P < 0.05$; Fig. 2). Thus, 8,800 μ g/ml was removed from further experiments.

Protective effect of LBPs. Following selection of the safe LBP dosages, the oxidative injury model was treated with different concentrations of LBPs to test the effect of the LBPs on cell injury (Fig. 3). LBPs at ≤ 55 or 4,400 μ g/ml did not exert a protective effect in the oxidative-injured RAECs. By contrast, LBP concentrations between 110 and 880 μ g/ml significantly




Figure 1. Effect of H_2O_2 on rat artery endothelial cell viability. * $P < 0.05$ vs. control.




Figure 2. Effect of LBPs on rat artery endothelial cell viability. * $P < 0.05$ vs. control. LBPs, *Lycium barbarum* polysaccharides.




Figure 3. Effect of LBPs on cell viability of rat artery endothelial following H_2O_2 -associated oxidative stress injury. * $P < 0.05$ vs. control. LBPs, *Lycium barbarum* polysaccharides.

protected the RAECs ($P < 0.05$ vs. control), in an apparent dose-dependent manner between 110 and 440 μ g/ml. Thus, the concentration of 440 μ g/ml LBPs exerted the most marked protective effect, and 110, 220 and 440 μ g/ml were used as low, medium and high intervention dosages of LBPs, respectively, to characterize their protective effects against oxidative cell injury.




Figure 4. Morphology of rat artery endothelial cells in the different groups. Arrows indicate cells with typical morphological changes (magnification, x200).




Figure 5. SOD activity and MDA and NO contents. *P<0.05 vs. control; #P<0.05 vs. H₂O₂. SOD, superoxide dismutase; MDA, malondialdehyde; NO, nitric oxide; LT/MT/HT, low/moderate/high-dose *Lycium barbarum* polysaccharide groups.




Figure 6. Western blot analysis of Bcl-2, Bax and Bcl-2/Bax protein expression. (A) Gel bands; (B) graph of quantified expression levels. *P<0.05 vs. control; #P<0.05 vs. H₂O₂. Bcl-2, B-cell lymphoma-2; Bax, Bcl-2-associated X protein; LT/MT/HT, low/moderate/high-dose *Lycium barbarum* polysaccharide groups.

Morphology of the cells. Normal RAECs under the microscope (magnification, x200) exhibited shuttle-like morphology with round or oval, clear nuclei and an ordered, paving stone arrangement (Fig. 4). Furthermore, the cell nuclei were of similar sizes, with evenly distributed chromatin, and were located centrally in the cells. These cells formed a relatively confluent monolayer. Conversely, under the stimulation of 100 μmol/l H₂O₂, the gaps in the RAEC monolayer were

enlarged and the adherence of cells was lost. Regions of the cell nuclei were strongly stained, indicating nuclei shrinkage. When the nuclei shrank, the chromatin became darker, which was matched by cell enlargement in certain cells. Granular, condensed chromatin was rarely identified. In the presence of 440 μg/ml LBPs, the majority of cells exhibited the same morphological characteristics as the normal control group (Fig. 4).

SOD activity, MDA and NO levels in cell culture supernatant. As depicted in Fig. 5, SOD activity, NO and MDA contents were altered between the groups. Notably, SOD activity and NO content significantly decreased while MDA content significantly increased in the H₂O₂ group compared with control group (P<0.05). In turn, SOD activity in the MT and HT group, and NO content in all three LBP groups were increased, while MDA content in the three LBP groups was decreased, compared with the H₂O₂ group (all P<0.05); however, the levels remained significantly altered compared with the control (P<0.05; Fig. 5).

Bcl-2 and Bax protein expression. Under the simulation of H₂O₂, the protein expression of apoptosis factors was also altered (Fig. 6). Notably, Bcl-2 expression decreased and Bax expression increased in the H₂O₂ group compared with control group (both P<0.05). In turn, Bcl-2 expression in all three LBP groups increased, while Bax expression in the MT and HT groups decreased compared with the H₂O₂ group (all P<0.05), with these altered Bax levels being statistically similar to those in the control group (P>0.05; Fig. 6B). The changes observed among the LBP groups appeared dose-dependent (Fig. 6A and B).

Furthermore, the expression ratio of Bcl-2/Bax protein in the H₂O₂ group decreased compared with the control group (P<0.05). The Bcl-2/Bax ratios in the MT and HT groups increased compared with the H₂O₂ group (both P<0.05), with the HT group deemed statistically similar to the control group (P>0.05; Fig. 6B). These changes again appeared dose-dependent (Fig. 6A and B).

Discussion

Due to the environmental, diet and lifestyle factors in modern society, CVDs have become increasingly frequent and tend to occur at younger ages (21-23). Thus, there has been increasing focus on the prevention of these diseases. Red medlar is a traditional medical and food product used in China (24). In traditional Chinese medicine, medlar is considered to nourish the liver, kidney and heart, and balance 'yin' and 'yang' in the body (25). Therefore, the biological effects of medlar should be a focus of clinical research. Despite its history of use, the mechanisms underlying the effects of medlar in body remain unclear. LBPs, as the effective constituents of medlar, may directly block the action of toxins in cells and activate microglial cells in the central nervous system (26). Compared with the potent stimulatory effects of glossy ganoderma and ginseng reported in the immune system, LBPs may moderately stimulate the immunoreactivity of the central nervous system (27). A number of studies have indicated that LBPs can exert marked anti-oxidant and anti-aging effects against oxidation-induced cell injury (28,29). Notably, LBPs could protect lens epithelial cells from H₂O₂ stress by promoting the expression of anti-apoptotic protein Bcl-2 and upregulating Bcl-2/Bax expression (30). Furthermore, LBPs protected seminiferous epithelial cells from thermal-induced injuries, protected DNA from oxidation injury in rat testis cells and markedly decreased the blood glucose of diabetic rats (31).

Oxidative stress occurs under normal conditions in the body, and there are a series of adaptive mechanisms that

protect cells from associated injuries. Normally, the production and elimination of ROS are under dynamic equilibrium, and thus they exert no lasting damage to cells. However, strong harmful stimulation may disrupt this balance and generate abundant levels of ROS that exceed the scavenging activity of the antioxidant system. As a consequence, the body generates an oxidative stress response to promote cell apoptosis, though pathological injury may still result (32). It is established that the ROS-mediated oxidative stress is a key inducer of cell apoptosis (33). ROS may also serve as a messenger in cells that is able to promote the activation of apoptosis to indirectly cause cell injury (34). The cell injury also induces the production of more ROS in cells. Taken together, oxidative stress is an important mechanism of organ injury and participates in the occurrence and development of various diseases (35).

Previous animal experiments by our group demonstrated that under exhaustive exercise and heat stress conditions, LBPs markedly inhibited oxidative stress and lipid peroxidation, improved compliance of the thoracic aorta, vessel diastolic function and exercise tolerance, and adjusted oxidase activities (36); however, the detailed mechanisms were undetermined. Therefore, the aim of the present study was to investigate the effect of LBP on H₂O₂-induced oxidative stress injury in RAECs, to thus determine the mechanism of the LBP protective effects. Initially, various H₂O₂ concentration concentrations were tested to determine the optimum dose for an oxidative stress model, and the H₂O₂ concentration of 100 μmol/l was selected based on its induction of ~50% cell survival.

Subsequently the effect of LBPs on the viability of RAECs was assessed, where it was observed that the cells were not inhibited by concentrations of LBPs <8,800 μg/ml. However, following the application of different concentrations of LBPs in the model cells, it was observed that the LBP concentrations below 55 or higher than 4,400 μg/ml did not exert a protective effect in the oxidative-injured RAECs; the effective concentration of LBPs was between 110-880 μg/ml, with the protective effect was most marked at 440 μg/ml. Therefore, 110-440 μg/ml doses were applied in subsequent assays to evaluate the mechanism of the LBP protective effect.

It is established that SOD is a primary scavenger of free radicals in the body and may also eliminate hazardous metabolic substances (37,38). Ageing has been associated with the production and accumulation of oxygen radicals in the body (39). It has been clinically demonstrated that SOD is associated with physiological pathology and the production/development of various diseases in humans (40,41). SOD is able to adjust blood lipids and, it may prevent atherosclerosis and CVDs caused by hyperlipidemia and reduce the content of lipid peroxide (42,43). The activity level of SOD, as a free-radical scavenger, reflects the ability of cells clear free radicals *in vivo* (44). MDA, as a major product of membrane lipid peroxidation, is also an injury marker that reflects the extent of cell injury. Additionally, MDA content is an important index that reflects membrane lipid peroxidation (45). Thus, the present study detected the change in both SOD activity and MDA content prior to and following LBP treatment in H₂O₂-injured cells.

It was observed that the treatment with LBPs at moderate and high doses (220 and 440 μg/ml) significantly enhanced

SOD activity in the H₂O₂-induced oxidative stress group; notably, SOD activity was improved to a statistically similar level to that of the control group with 440 µg/ml LBPs. This may be associated with lycium polysaccharide reducing the fluidity of the plasma membrane and thus reducing damage by free radicals. The MDS levels in the oxidative injury group were also inhibited in a dose-dependent manner by LBPs, probably due to the LBPs enhancing the activity of anti-oxidase enzymes in free radical clearance.

The common pathological cause of many CVDs is endothelial dysfunction. NO is an important vasodilatation factor produced by VECs. The reduced rate of clearance and excessive production of ROS are important causes of endothelial dysfunction. The present data indicate that LBPs may clear high levels of ROS to thus exert an antioxidant effect, and thereby enhance NO usage rate and adjust vasodilatation function, which would ultimately improve the resistance of cells to oxidative stress. Collectively these results indicate that LBPs may prevent H₂O₂-induced cell apoptosis and modulate, at least in part, the ROS-NO axis.

Cell apoptosis is a structured form of cell death controlled by gene under certain physiological or pathological conditions. This programmed cell death involves the activation, regulation and expression of diverse genes (46). Cell apoptosis is essential in maintaining the body at steady state and the normal physiological functions of organs (47). Cell apoptosis results from stimulation by external factors and pathological events in cells. It is controlled by specific genes, and is induced and inhibited by stimulating factors (including H₂O₂) in the cell microenvironment (30). Exogenous factors may also exert stimulatory effects through certain signaling pathways (48). The Bcl-2 gene family serves a meditative and regulatory role in cell apoptosis (49). Key anti-apoptosis and pro-apoptosis factors in this family are Bcl-2 and Bax (50). As a proto-oncogene, Bcl-2 is a mitochondrial membrane protein and is considered the most important anti-apoptosis gene (51). Conversely, Bax expression may enhance antagonism of Bcl-2 and thereby promote cell apoptosis; H₂O₂ may activate this Bax expression (52). The apoptosis precursor proteins, including Bax, may alter translocation through mitochondrial membranes, but their activation may be resisted by the anti-apoptotic proteins of mitochondrial membranes, which form heterodimers and maintain membrane integrity (53). For instance, the anti-apoptotic protein Bcl-2 protects mitochondria by inhibiting Bax activation and subsequent translocation of cytochrome C through the mitochondrial membrane (54); in this regard, a lack of Bcl-2 would allow the breakdown of mitochondrial outer membrane integrity and the release of cytochrome C (55). Thus, the Bcl-2 family adjusts cell apoptosis mediated by the mitochondrial pathway through anti-apoptosis and pro-apoptosis factors, as well as participating in the crosstalk of the mitochondria and death receptor pathways (56). In cases of apoptosis mediated by oxidative stress, the expression of Bcl-2-related proteins may be altered, affecting cell apoptosis. Notably, the present data indicated that oxidative stress reduced the ratio of Bcl-2/Bax. This was improved following LBP intervention. Thus, LBPs may moderately protect cells from H₂O₂-induced injury and produce an apoptosis-inhibitory effect, ultimately attenuating the oxidative stress condition. Notably, the higher-dose LBP intervention almost recovered cell condition to a normal state.

Overall these results may aid to develop methods for attenuating or preventing oxidative stress in endothelial cells for the clinical treatment of CVDs.

In conclusion, H₂O₂-induced oxidative stress in RAECs, while LBPs upregulated SOD and NO and downregulated MDA through an apparent antioxidant defense mechanism. Furthermore, LBPs increased Bcl-2/Bax expression potentially through an antiapoptosis system. Therefore, LBPs may protect VECs from H₂O₂-induced injury through anti-oxidation and anti-apoptosis effects.

Acknowledgements

Not applicable.

Funding

The present study was supported by the National Natural Science Foundation of China (grant no. 81560052), the University-level scientific research project of Ningxia Medical University (grant no. XM2018002), and the Ningxia High School First-Class Disciplines (West China First-Class Disciplines Basic Medical Sciences at Ningxia Medical University; grant no. NXYLXK2017B07).

Availability of data and materials

All data sets generated or analyzed during the study are included in the published article.

Authors' contributions

GL conceived and designed the experiments. SX performed the experiments and prepared the manuscript. XH performed experiments. LZ and LN analyzed the data. All authors read and approved the final version of the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Mladěnka P, Applová L, Patočka J, Costa VM, Remiao F, Pourová J, Mladěnka A, Karlíčková J, Jahodář L, Vopršalová M, *et al.*: Comprehensive review of cardiovascular toxicity of drugs and related agents. *Med Res Rev* 38: 1332-1403, 2018.
2. Kim KS, Song CG and Kang PM: Targeting oxidative stress using nanoparticles as atheranostic strategy for cardiovascular diseases. *Antioxid Redox Signal* 30: 733-746, 2019.
3. World Health Organization: Comprehensive global monitoring framework, including indicators, and a set of voluntary global targets for the prevention and control of noncommunicable diseases. World Health Organization, 2012.

4. Zeng Z: Internal secretion of vascular endothelial cells. *Chin J Int Med* 37: 77-77, 1998 (In Chinese). <http://www.cqvip.com/Main/Detail.aspx?id=2964468>.
5. Blatter LA: Tissue specificity: SOCE: Implications for Ca²⁺ handling in endothelial cells. *Adv Exp Med Biol* 993: 343-361, 2017.
6. Rajendran P, Rengarajan T, Thangavel T, Nishigaki Y, Sakthisekaran D, Sethi G and Nishigaki I: The vascular endothelium and human diseases. *Int J Biol Sci* 9: 1057-1069, 2013.
7. Yamada T, Egashira N, Bando A, Nishime Y, Tonogai Y, Imuta M, Yano T and Oishi R: Activation of p38 MAPK by oxidative stress underlying epirubicin-induced vascular endothelial cell injury. *Free Radic Boil Med* 52: 1285-1293, 2012.
8. Closhen D, Bender B, Luhmann HJ and Kuhlmann CR: CRP-induced levels of oxidative stress are higher in brain than aortic endothelial cells. *Cytokine* 50: 117-120, 2010.
9. Higashi Y: Mechanisms of impairment of endothelial cell. *Nihon Rinsho* 70: 1519-1523, 2012 (In Japanese).
10. Bei Y, Das S, Rodosthenous RS, Holvoet P, Vanhaverbeke M, Monteiro MC, Monteiro VVS, Radosinska J, Bartekova M, Jansen F, *et al*: Extracellular vesicles in cardiovascular theranostics. *Theranostics* 7: 4168-4182, 2017.
11. Rodrigo R, Libuy M, Feliú F and Hasson D: Oxidative stress-related biomarkers in essential hypertension and ischemia-reperfusion myocardial damage. *Dis Markers* 35: 773-790, 2013.
12. Xu J and Li Y: Effects of salidroside on exhaustive exercise-induced oxidative stress in rats. *Mol Med Rep* 6: 1195-1198, 2012.
13. Luo Q, Li Z, Huang X, Yan J, Zhang S and Cai YZ: *Lycium barbarum* polysaccharides: Protective effects against heat-induced damage of rat testes and H₂O₂-induced DNA damage in mouse testicular cells and beneficial effect on sexual behavior and reproductive function of hemicastrated rats. *Life Sci* 79: 613-621, 2006.
14. Ho YS, Yu MS, Yik SY, So KF, Yuen WH and Chang RC: Polysaccharides from wolfberry antagonizes glutamate excitotoxicity in rat cortical neurons. *Cell Mol Neurobiol* 29: 1233-1244, 2009.
15. Wu HT, He XJ, Hong YK, Ma T, Xu YP and Li HH: Chemical characterization of *Lycium barbarum* polysaccharides and its inhibition against liver oxidative injury of high-fat mice. *Int J Biol Macromol* 46: 540-543, 2010.
16. Tang WM, Chan E, Kwok CY, Lee YK, Wu JH, Wan CW, Chan RY, Yu PH and Chan SW: A review of the anticancer and immunomodulatory effects of *Lyciumbarbarum* fruit. *Inflammopharmacology* 20: 307-314, 2012.
17. Chang HM and But PPH (eds): *Pharmacology and Applications of Chinese Materia Medica, Vol. II*. World Scientific Publishing Company Incorporated, Singapore, 1986. <https://www.worldscientific.com/worldscibooks/10.1142/0377>.
18. Potterat O: Goji (*Lycium barbarum* and *L. chinense*): Phytochemistry, pharmacology and safety in the perspective of traditional uses and recent popularity. *Planta Med* 76: 7-19, 2010.
19. Yu N, Song N, Liu CY and Yang GL: The estrogen-like protective effect of *Lycium barbarum* polysaccharides in reducing oxidative stress on myocardial cells from ovariectomized rats. *Molecular Medicine Reports* 19: 2271-2278, 2019.
20. Li J, Zou Y, Ge J, Zhang D, Guan A, Wu J and Li L: The effects of G-CSF on proliferation of mouse myocardial microvascular endothelial cells. *Int J Mol Sci* 12: 1306-1315, 2011.
21. Younus A, Aneni EC, Spatz ES, Osondu CU, Roberson L, Ogunmoroti O, Malik R, Ali SS, Aziz M, Feldman T, *et al*: A systematic review of the prevalence and outcomes of ideal cardiovascular health in US and Non-US populations. *Mayo Clin Proc* 91: 649-670, 2016.
22. Fang N, Jiang M and Fan Y: Ideal cardiovascular health metrics and risk of cardiovascular disease or mortality: A meta-analysis. *Int J Cardiol* 214: 279-283, 2016.
23. Guo L and Zhang S: Association between ideal cardiovascular health metrics and risk of cardiovascular events or mortality: A meta-analysis of prospective studies. *Clin Cardiol* 40: 1339-1346, 2017.
24. Yang D, Li SY, Yeung CM, Chang RC, So KF, Wong D and Lo AC: *Lycium barbarum* extracts protect the brain from blood-brain barrier disruption and cerebral edema in experimental stroke. *PLoS One* 7: e33596, 2012.
25. Shan X, Zhou J, Ma T and Chai Q: *Lycium barbarum* polysaccharides reduce exercise-induced oxidative stress. *Int J Mol Sci* 12: 1081-1088, 2011.
26. Teng P, Li Y, Cheng W, Zhou L, Shen Y and Wang Y: Neuroprotective effects of lycium barbarum polysaccharides in lipo-polysaccharide-induced BV2 microglial cells. *Mol Med Rep* 7: 1977-1981, 2013.
27. Chang RC and So KF: Use of anti-aging herbal medicine, *Lycium barbarum*, against aging-associated diseases. What do we know so far? *Cell Mol Neurobiol* 28: 643-652, 2008.
28. Liu Y and Zhang Y: *Lycium barbarum* polysaccharides alleviate hydrogen peroxide-induced injury by up-regulation of miR-4295 in human trabecular meshwork cells. *Exp Mol Pathol* 106: 109-115, 2019.
29. Niu T, Jin L, Niu S, Gong C and Wang H: *Lycium barbarum* polysaccharides alleviates oxidative damage induced by H₂O₂ through down-regulating MicroRNA-194 in PC-12 and SH-SY5Y cells. *Cell Physiol Biochem* 50: 460-472, 2018.
30. Qi B, Ji Q, Wen Y, Liu L, Guo X, Hou G, Wang G and Zhong J: *Lycium barbarum* polysaccharides protect human lens epithelial cells against oxidative stress-induced apoptosis and senescence. *PLoS One* 9: e110275, 2014.
31. Schulz C, Farkas L, Wolf K, Kratzel K, Eissner G and Pfeifer M: Differences in LPS-induced activation of bronchial epithelial cells (BEAS-2B) and type II-like pneumocytes (A-549). *Scand J Immunol* 56: 294-302, 2002.
32. Zhao W, Pan X, Li T, Zhang C and Shi N: *Lycium barbarum* polysaccharides protect against trimethyltin chloride-induced apoptosis via sonic hedgehog and PI3K/Akt signaling pathways in mouse neuro-2a cells. *Oxid Med Cell Longev* 2016: 9826726, 2016.
33. Zhang Y, Wang G, Wang T, Cao W, Zhang L and Chen X: Nrf2-Keap1 pathway-mediated effects of resveratrol on oxidative stress and apoptosis in hydrogen peroxide-treated rheumatoid arthritis fibroblast-like synoviocytes. *Ann N Y Acad Sci* 6: 2019. <https://doi.org/10.1111/nyas.14196>.
34. Maritim AC, Sanders RA and Watkins JB III: Diabetes, oxidative stress and antioxidants: A review. *J Biochem Mol Toxicol* 17: 24-38, 2003.
35. Wang Z, Su G, Zhang Z, Dong H, Wang Y, Zhao H, Zhao Y and Sun Q: 25-Hydroxyl-protopanaxatriol protects against H₂O₂-induced H9c2 cardiomyocytes injury via PI3K/Akt pathway and apoptotic protein down-regulation. *Biomed Pharmacother* 99: 33-42, 2018.
36. Zhao Z, Luo Y, Li G, Zhu L, Wang Y and Zhang X: Thoracic aorta vasoreactivity in rats under exhaustive exercise: Effects of *Lycium barbarum* polysaccharides supplementation. *J Int Soc Sports Nutr* 10: 47, 2013.
37. Yang CH, Xu JH, Ren QC, Duan T, Mo F and Zhang W: Melatonin promotes secondary hair follicle development of early post-natal cashmere goat and improves cashmere quantity and quality by enhancing antioxidant capacity and suppressing apoptosis. *J Pineal Res* 67: e12569, 2019.
38. Majumdar A and Kar RK: Orchestration of Cu-Zn SOD and class III peroxidase with upstream interplay between NADPH oxidase and PM H⁺-ATPase mediates root growth in *Vigna radiata* (L.) Wilczek. *J Plant Physiol* 232: 248-256, 2019.
39. Liochev SI: Reactive oxygen species and the free radical theory of aging. *Free Radic Biol Med* 60: 1-4, 2013.
40. Kumar M, Chand R and Shah K: Evidences for growth-promoting and fungicidal effects of low doses of tricyclazole in barley. *Plant Physiol Biochem* 103: 176-182, 2016.
41. Lu Z, Xu X, Hu X, Zhu G, Zhang P, van Deel ED, French JP, Fassett JT, Oury TD, Bache RJ and Chen Y: Extracellular superoxide dismutase deficiency exacerbates pressure overload-induced left ventricular hypertrophy and dysfunction. *Hypertension* 51: 19-25, 2008.
42. Song QQ, Niu JP, Zhang SY, Liang TT, Zhou J and Feng SS: Effects of simulated heat wave and ozone on high fat diet ApoE deficient mice. *Biomed Environ Sci* 31: 757-768, 2018.
43. Shatoor AS, Al Humayed S, Alkhateeb MA, Shatoor KA, Aldera H, Alassiri M and Shati AA: *Crataegus Aronia* protects and reverses vascular inflammation in a high fat diet rat model by an antioxidant mechanism and modulating serum levels of oxidized low-density lipoprotein. *Pharm Biol* 57: 38-48, 2019.
44. Saran M, Michel C and Bors W: Radical functions in vivo: A critical review of current concepts and hypotheses. *Z Naturforsch C* 53: 210-227, 1998.
45. Bai YX, Fang F, Jiang JL and Xu F: Extrinsic calcitonin gene-related peptide inhibits hyperoxia-induced alveolar epithelial type II cells apoptosis, oxidative stress, and reactive oxygen species (ROS) production by enhancing notch 1 and homocysteine-induced endoplasmic reticulum protein (HERP) expression. *Med Sci Monit* 23: 5774-5782, 2017.

46. Omer FAA, Hashim NBM, Ibrahim MY, Dehghan F, Yahayu M, Karimian H, Salim LZA and Mohan S: Beta-mangostin from *Cratoxylum arborescens* activates the intrinsic apoptosis pathway through reactive oxygen species with downregulation of the HSP70 gene in the HL60 cells associated with a G₀/G₁ cell-cycle arrest. *Tumour Biol* 39: 1010428317731451, 2017.
47. Radović N, Cucić S and Altarac S: Molecular aspects of apoptosis. *Acta Med Croatica* 62: 249-256, 2008 (In Croatian).
48. Qi B, Ji Q, Wen Y, Liu L, Guo X, Hou G, Wang G and Zhong J: *Lycium barbarum* polysaccharides protect human lens epithelial cells against oxidative stress-induced apoptosis and senescence. *PLoS One* 9 (10): e110275, 2014.
49. Edlich F: BCL-2 proteins and apoptosis: Recent insights and unknowns. *Biochim Biophys Res Commun* 500: 26-34, 2018.
50. Xi H, Fan X, Zhang Z, Liang Y, Li Q and He J: Bax and Bcl-2 are involved in the apoptosis induced by local testicular heating in the boar testis. *Reprod Domest Anim* 52: 359-365, 2017.
51. Gross A: BCL-2 family proteins as regulators of mitochondria metabolism. *Biochim Biophys Acta* 1857: 1243-1246, 2016.
52. Wang X, Li Z, Bai J, Song W and Zhang F: miR-17-5p regulates the proliferation and apoptosis of human trabecular meshwork cells by targeting phosphatase and tensin homolog. *Mol Med Rep* 19: 3132-3138, 2019.
53. Goping IS, Gross A, Lavoie JN, Nguyen M, Jemmerson R, Roth K, Korsmeyer SJ and Shore GC: Regulated targeting of BAX to mitochondria. *J Cell Biol* 143: 207-215, 1998.
54. Cheng EH, Wei MC, Weiler S, Flavell RA, Mak TW, Lindsten T and Korsmeyer SJ: Bcl-2, Bcl-X_L sequester BH3 domain-only molecules preventing Bax-and Bak-mediated mitochondrial apoptosis. *Mol Cell* 8: 705-711, 2001.
55. Yang J, Liu X, Bhalla K, Kim CN, Ibrado AM, Cai J, Peng TI, Jones DP and Wang X: Prevention of apoptosis by Bcl-2: Release of cytochrome C from mitochondria blocked. *Science* 275: 1129-1132, 1997.
56. Hsu HC, Liu YS, Tseng KC, Tan BC, Chen SJ and Chen HC: Lgr5 regulates survival through mitochondria-mediated apoptosis and by targeting the Wnt/β-catenin signaling pathway in colorectal cancer cells. *Cellular Signalling* 26: 2333-2342, 2014.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.