

# Urinary expression of long non-coding RNA TUG1 in non-diabetic patients with glomerulonephritides

FERNANDO JAVIER SALAZAR-TORRES<sup>1,2</sup>, MIGUEL MEDINA-PEREZ<sup>1</sup>, ZESERGIO MELO<sup>3</sup>,  
CLAUDIA MENDOZA-CERPA<sup>4</sup> and RAQUEL ECHAVARRIA<sup>3</sup>

<sup>1</sup>Departamento de Nefrología, UMAE-Hospital de Especialidades,  
CMNO, Instituto Mexicano del Seguro Social, Guadalajara, Jalisco 44340;

<sup>2</sup>Unidad de Medicina Familiar con Unidad Médica de Atención Ambulatoria UMF/UMAA 39,  
Instituto Mexicano del Seguro Social, Matamoros, Tamaulipas 87344;

<sup>3</sup>CONACyT-Centro de Investigación Biomédica de Occidente, Instituto Mexicano del Seguro Social;

<sup>4</sup>Departamento de Patología, UMAE-Hospital de Especialidades,  
CMNO, Instituto Mexicano del Seguro Social,  
Guadalajara, Jalisco 44340, México

Received March 4, 2020; Accepted September 29, 2020

DOI: 10.3892/br.2020.1393

**Abstract.** Metabolic alterations serve a significant role in the pathogenesis of kidney disease. Long non-coding RNA (lncRNA) taurine upregulated gene 1 (TUG1) is a known regulator of podocyte health and mitochondrial biogenesis. Although TUG1 protects against podocyte loss in models of diabetic nephropathy, it is unknown if urinary TUG1 expression is associated with clinical and histopathological findings in non-diabetic patients diagnosed with glomerulonephritides. In the present study, the expression of TUG1, podocyte-specific markers (nephrin and podocin) and mitochondrial biogenesis-associated mRNAs (transcription factor A mitochondrial, cytochrome C oxidase subunit 5A and peroxisome proliferator-activated receptor  $\gamma$  coactivator 1 $\alpha$ ) were examined in urinary sediment of non-diabetic patients

with biopsy-confirmed glomerulonephritides and healthy controls. Urinary expression of TUG1 was significantly lower in patients with glomerulonephritides, particularly those diagnosed with Focal Segmental Glomerulosclerosis (FSGS). Furthermore, TUG1 levels were associated with urinary expression of podocyte-specific markers and mRNAs associated with mitochondrial biogenesis. Loss of TUG1 expression in urinary sediment was strongly associated with FSGS, highlighting the potential of this lncRNA and its mitochondrial biogenesis-associated targets as non-invasive biomarkers of assessing podocytopathy.

## Introduction

Glomerulonephritides are a group of rare diseases that often affects younger individuals and can lead to end-stage renal disease (ESRD) (1). Although the clinical presentation of glomerulonephritides and their outcomes are variable, well-established factors such as persistent proteinuria, hypertension, diabetes and cardiovascular disease increase the risk of ESRD progression (2). Podocyte injury is a hallmark of renal diseases, presenting clinically with proteinuria, glomerulosclerosis and kidney failure (3). Terminally differentiated podocytes have a limited capacity to self-replicate; hence, direct cellular damage contributes to the onset and progression of glomerular diseases such as Focal Segmental Glomerulosclerosis (FSGS), Minimal Change Disease and Diabetic Nephropathy (DN) (4). Additionally, immune-mediated damage is responsible for the establishment and progression of secondary glomerulonephritides, including lupus nephritis (LN) and ANCA-associated vasculitis (AAV) (5).

The gold standard for confirming a diagnosis of glomerulonephritides diagnosis is renal biopsy (6). However, biopsies are invasive, may have complications and are hard to interpret due to the presence of highly complex pathogenic mechanisms that translate into limited histological responses of kidney injury (7). Thus, identifying non-invasive biomarkers

---

*Correspondence to:* Dr Raquel Echavarría, CONACyT-Centro de Investigación Biomédica de Occidente, Instituto Mexicano del Seguro Social, Sierra Mojada 800 Col. Independencia, Guadalajara, Jalisco 44340, México  
E-mail: rechavarría@conacyt.mx

**Abbreviations:** lncRNA, long non-coding RNA; TUG1, taurine upregulated gene 1; TFAM, transcription factor A mitochondrial; COX5A, cytochrome C oxidase subunit 5A; and PPARGC1A, peroxisome proliferator-activated receptor  $\gamma$  coactivator 1 $\alpha$ ; ESRD, end-stage renal disease; FSGS, focal segmental glomerulosclerosis; DN, diabetic nephropathy; AGS, advanced glomerular sclerosis; NSPH1, nephrin; NSPH2, podocin; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; LN, lupus nephritis; AAV, ANCA-associated vasculitis; GFR, glomerular filtration rate

**Key words:** TUG1, lncRNA, glomerulonephritides, FSGS, urinary sediment

of glomerular disease may result in improved diagnosis and guide therapeutic choices in the field of nephrology (8).

Long non-coding RNAs (lncRNAs) are RNA molecules comprised of >200 nucleotides with no protein-coding capacity. They are a class of RNAs comprised of heterogeneous intergenic transcripts, sense or antisense transcripts that overlap with other genes, or enhancer RNAs with a range of functions that includes the regulation of gene expression, chromatin remodelling, microRNA (miRNA/miR)-sponging and protein scaffolding (9,10). lncRNAs modulate several biological processes, including homeostasis, cellular metabolism, proliferation, apoptosis and differentiation (10,11). Dysregulated expression of lncRNAs such as metastasis-associated lung adenocarcinoma transcript 1, LOC105374325, LOC105375913, X-inactive specific transcript and RP11-2B6.2 contribute to the pathogenesis of various kidney diseases (12-17). Moreover, circulating lncRNAs are stable and easily detectable in plasma, serum and urine, characteristics that make them convenient for use diagnostically (18,19).

The lncRNA taurine upregulated gene 1 (TUG1), located on chromosome 22q12, regulates podocyte health and glomerulosclerosis by altering the expression of peroxisome proliferator-activated receptor  $\gamma$  Coactivator 1 $\alpha$  (PGC1A) (20-23). PGC1A is a transcriptional coactivator that controls mitochondrial biogenesis and is encoded by the *PPARGC1A* gene in humans (24,25). A decrease in PGC1A expression contributes to the onset of metabolic diseases such as DN, and transgenic expression of *PPARGC1A* in tubular cells protects mice from developing acute and chronic forms of kidney disease (26,27). Long *et al* (28) found that podocyte-specific transgenic expression of lncRNA TUG1 protects against DN in a mouse model, and demonstrated that lower glomerular filtration rates (GFR) were correlated with a decrease in TUG1 intrarenal expression in human subjects. However, it is unknown if TUG1 can be detected in urine and if its expression levels are correlated with histopathological findings in kidney biopsies from patients diagnosed with glomerulonephritides other than DN. In the present study, the urinary expression of TUG1 in non-diabetic patients with glomerulonephritides was characterized and it was shown that a decrease in TUG1 expression was significantly associated with FSGS.

## Materials and methods

**Ethical considerations.** The present study complied with the ethical principles for medical research specified in the Declaration of Helsinki (29) and was approved by the Local Ethics and Research Committee at the Hospital de Especialidades NUM. 1, Bajío, Leon, Guanajuato, Instituto Mexicano del Seguro Social (approval no. CLIEIS R-2018-1001-114). Each participant provided written informed consent prior to enrolment in the study.

**Patient enrolment.** A total of 11 patients with biopsy-confirmed glomerulonephritides (7 females and 4 males; median age, 31 years; range 19-59 years), and 10 healthy controls (6 females and 4 males; median age 34 years; range 22-54) were enrolled in the present study at UMAE-Hospital de Especialidades, CMNO in Guadalajara, Mexico. Inclusion criteria were non-diabetic patients aged  $\geq 18$  years. Exclusion criteria were intake of non-steroid anti-inflammatory drugs or platelet anti-aggregation

medication a week before sampling, administration of anticoagulant medication 96 h before the biopsy, blood pressure >160/90 mmHg, active urinary infection or anaemia. Urine sample collection was taken shortly before biopsy collection. Only 1 patient enrolled in the study was unable to provide a urine sample and, therefore, was excluded from further analysis.

**Data collection.** Biochemical parameters corresponding to each glomerulonephritides patient at the time of biopsy were assayed by routine laboratory techniques at UMAE-Hospital de Especialidades and retrieved from patients' electronic medical records.

**Reverse transcription-quantitative (RT-q)PCR.** Urine specimens from patients and healthy controls were collected and centrifuged at 4,500  $\times$  g for 30 min at 4°C. The urinary sediments were recovered by discarding the supernatant, and total RNA was extracted using RiboZol reagent (Amresco LLC) according to the manufacturer's protocol. cDNA was synthesized by reverse transcription using 1  $\mu$ g total RNA and a QuantiTect Reverse Transcription kit according to the manufacturer's protocol (Qiagen, Inc.). The cDNA was used to perform quantitative PCR using EvaGreen 5x qPCR mix (qARTA Bio Inc.) and 10 pmol/ $\mu$ l each of the reverse and forward primers in a Lightcycler 96 (Roche Diagnostics). The sequences of the primers used to amplify human lncRNA TUG1 (30), nephrin (NSPH1), podocin (NSPH2), transcription factor A, mitochondrial (TFAM), cytochrome C oxidase subunit 5A (COX5A), PPARGC1A and GAPDH are shown in Table I. The amplification conditions consisted of 1 cycle of initial denaturation at 95°C for 15 min; followed by 40 cycles of denaturation at 95°C for 15 sec, annealing at 56°C for 20 sec and extension at 72°C for 20 sec. GAPDH was used as the internal control to normalize gene expression data. The relative mRNA expression was calculated using the  $2^{-\Delta\Delta C_q}$  method (31).

**In silico identification of TUG1/miR-204-5p axis mRNA targets.** The VarElect phenotype program from GeneAnalytics (geneanalytics.genecards.org), a pathway enrichment analysis tool, was used to identify predicted mRNA targets of the TUG1/miR-204-5p axis involved in mitochondrial biogenesis based on the genomic information stored in GeneCards (32). In total, 453 predicted mRNA targets of hsa-miR-204-5p downloaded from TargetScan version 7.2 (targetscan.org) (33) were profiled with GeneAnalytics. The entries were scored and ranked using an algorithm that enables matching of genomic expression, sequencing and microarray datasets to tissues and cells.

**Statistical analysis.** Data are expressed as the mean  $\pm$  standard error of the mean. Non-parametric variables were compared using a Mann-Whitney U-test or a Kruskal-Wallis test with a Dunn's post-hoc test, as appropriate. A Spearman's rank correlation analysis was used to analyse correlations. Statistical analysis was performed in GraphPad Prism Version 6.01 (GraphPad Software, Inc.).  $P < 0.05$  was considered to indicate a statistically significant difference.

## Results

**Clinical and biochemical characteristics of glomerulonephritides patients and healthy controls.** The present study included

Table I. Sequences of the primers.

Gene	Sequence, 5'-3'	Size, bp
TUG1		150
Forward	TAGCAGTTCCCCAATCCTTG	
Reverse	CACAAATTCCTCATTCCTCC	
NPHS1		124
Forward	GTGCACTATGCTCCCACCAT	
Reverse	TCTCCCAGTTGAACATGCCC	
NPHS2		160
Forward	GAGGAAGGTACCAATCCTCCG	
Reverse	GCAGATGTCCCAGTCGGAATA	
TFAM		105
Forward	ATGGAGGCAGGAGTTTCGTT	
Reverse	CCTAGATGAGTTCTGCCTGCT	
COX5A		160
Forward	AGATGCCTGGGAATTGCGTA	
Reverse	AGGTCCTGCTTTGTCCTTAACA	
PPARGC1A		158
Forward	TGTGGAGTCCCTGGAATGGA	
Reverse	AAGATCGTGTTGGGCGAGAG	
GAPDH		153
Forward	CCCACTCCTCCACCTTTGAC	
Reverse	TGGTCCAGGGTCTTACTCC	

10 patients with biopsy-confirmed diagnosis, 4 males and 6 females, and 10 sex-matched healthy controls. The mean age at renal biopsy was  $34.4 \pm 11.1$  years old, the mean body mass index was  $30.1 \pm 5.1$ , and the diagnoses were as follows: 5 cases of FSGS, 3 cases of LN, 1 case of AAV and 1 case of advanced glomerular sclerosis (AGS). Table II shows the biochemical parameters of the glomerulonephritides patients. A total of 80% of the patients had haemoglobin values within the normal range. All glomerulonephritides patients had proteinuria, with the highest levels present in the patient diagnosed with AGS. However, CKD staging, classified according to the level of GFR (34) varied greatly. Only 2 patients had a normal GFR of  $>90$  ml/min/1.73 m<sup>2</sup>; 3 FSGS patients were Stage 2; 1 AAV patient was Stage 3B; 3 patients were Stage 4 and only 1 LN patient was Stage 5 with a GFR  $<15$  ml/min/1.73m<sup>2</sup>. The presence of glomerular damage in the absence of renal failure highlights the importance of identifying novel circulating markers able to detect glomerulonephritides during the early stages of CKD.

**Glomerulonephritides patients have low levels of urinary TUG1 expression.** Cell-free lncRNAs are stable in urine and may potentially be used as non-invasive biomarkers to identify diseases, such as lupus nephritis and membranous nephropathy (18,19). To assess alterations in TUG1 expression in the urinary sediment of patients with glomerulonephritides, RT-qPCR was used. Urinary expression of TUG1 was reduced in glomerulonephritides patients compared with the healthy controls (Fig. 1A). Furthermore, significantly lower TUG1 expression levels in urinary sediment was observed in patients with FSGS when compared with all




Figure 1. Urinary expression of TUG1 is decreased in FSGS patients. (A) RT-qPCR results showing the log<sub>2</sub>-fold change in expression of TUG1/GAPDH in the urinary sediment of healthy controls (n=10) and glomerulonephritides patients (n=10). (B) Reverse transcription-quantitative PCR results showing the log<sub>2</sub>-fold change in expression of TUG1/GAPDH in the urinary sediment of healthy controls (n=10), FSGS patients (n=5) and other diagnoses (lupus nephritis, n=3; ANCA-associated vasculitis, n=1; and advanced glomerular sclerosis, n=1). Data are presented as the mean  $\pm$  standard error of the mean. \*P<0.05, \*\*P<0.01. TUG1, taurine upregulated gene 1; FSGS, focal segmental glomerulosclerosis.

other diagnoses (Fig. 1B). These results suggest that the urinary expression of TUG1 decreases with podocytopathy.

**Urinary expression of TUG1 correlates with podocyte marker expression.** Glomerular injury is associated with podocyte loss, a phenomenon that can be monitored by detecting podocyte-specific mRNAs in the urine (35). In the present study, two podocyte-specific mRNAs, NPHS1 and NPHS2, were detected using RT-qPCR. Although the expression levels of urinary NPHS1 was not significantly different between glomerulonephritides patients and healthy controls, NPHS1 urinary expression was significantly lower in the FSGS patients when compared with other diagnoses (Fig. 2A). In contrast, NPHS2 mRNA did not differ significantly amongst the groups (Fig. 2B). Interestingly, a positive correlation was observed between TUG1 urinary expression and both podocyte-specific mRNAs (NPHS1,  $r=0.6486$ , \*\*P<0.01; NPHS2,  $r=0.4782$ , \*P<0.05; Fig. 2C and D). These results further suggest an association between TUG1 urinary expression and podocyte damage.

**Urinary expression of mitochondrial biogenesis markers is correlated with TUG1 expression levels.** Several studies have validated the functional role of TUG1 as a lncRNA in directly binding to specific miRNAs to regulate post-transcriptional processing via a competing endogenous RNA mechanism (21,23,28,30). One such target is miR-204-5p (30). Thus, the VarElect phenotype program from Gene Analytics was used to predict mRNA targets of the TUG1/miR-204-5p

Table II. Biochemical parameters of glomerulonephritides patients.

Factor	Patient no.									
	1	2	3	4	5	6	7	8	9	10
Diagnosis	FSGS	FSGS	AAV	LN	AGS	FSGS	LN	LN	FSGS	FSGS
Hemoglobin, g/dl	14.7	13.2	12.8	11.6	8.7	10.5	15.6	13.8	16.7	16.3
sCr, mg/dl	0.84	0.57	1.5	3.7	3.9	2.1	0.86	8.4	1.4	1.5
GFR, ml/min/1.73m <sup>2</sup>	90	126	38	17	15	28	107	8	66	63
Proteinuria, g/24 h	1.28	2.68	1.9	2.24	5.8	3.68	1.59	2.9	2.72	1.79
Albumin, g/dl	3.7	2.1	4.1	3.6	2.5	2.3	2.9	3	3.5	4.3
pANCA	+	-	+	+++	-	+	+++	+++	-	-
cANCA	-	-	-	-	-	-	-	-	-	-
Antinuclear antibodies	-	0.09722	0.26389	0.48611	0.09722	0.09722	0.15278	0.15278	0.15278	-
Anti-DNA antibodies	-	-	-	+++	-	-	++	++	-	-
C3 fraction, mg/dl	124	112	93.9	55	104	127	59.7	92.2	143	125
C4 fraction, mg/dl	28.6	38.4	24.4	7	31.8	28.9	9.29	19.4	33.5	33.5
CRP, mg/l	3.75	<3.23	5.53	<3.23	15.8	<3.23	16	11.1	<3.23	<3.23
ESR, mm/h	37	42	16	12	18	-	28	26	-	10

FSGS, focal segmental glomerulosclerosis; LN, lupus nephritis; AAV, ANCA-associated vasculitis; AGS, advanced global sclerosis; sCr, serum creatinine; pANCA, perinuclear/nuclear antineutrophil cytoplasmic antibody; cANCA, cytoplasmic antineutrophil cytoplasmic antibody; C3, complement component 3; C4, complement component 4; CRP, C reactive protein; ESR, erythrocyte sedimentation rate; -, negative; +, low; ++, moderate; +++, high.




Figure 2. TUG1 urinary expression is correlated with podocyte marker mRNA expression in glomerulonephritides patients and healthy controls. Reverse transcription-quantitative PCR results showing the log<sub>2</sub>-fold change in expression of (A) NPHS1 and (B) NPHS2 in the urinary sediment of healthy controls (n=10), FSGS patients (n=5) and other diagnoses (lupus nephritis, n=3; ANCA-associated vasculitis, n=1; and advanced glomerular sclerosis, n=1). Data are presented as the mean  $\pm$  standard error of the mean. \*P<0.05. Correlation between lncRNA TUG1 transcript levels and (C) NPHS1 or (D) NPHS2 in the urinary sediment of healthy controls (n=10) and glomerulonephritides patients (n=10). \*P<0.05, \*\*P<0.01. TUG1, taurine upregulated gene 1; NPHS1, nephrin; NPHS2, podocin; ns, not significant.



Table III. Predicted targets of TUG1/miR-204-5p axis involved in mitochondrial biogenesis.

Symbol	Description	Global Rank <sup>a</sup>	Score <sup>a,b</sup>
PPARGC1A	PPARG Coactivator 1 $\alpha$	1	18.76
TFAM	Transcription Factor A, Mitochondrial	4	11.67
CREB1	CAMP Responsive Element Binding Protein 1	25	4.98
SIRT1	Sirtuin 1	26	4.80
COX5A	Cytochrome C Oxidase Subunit 5A	39	3.13
ATF2	Activating Transcription Factor 2	52	2.42
MAPK1	Mitogen-Activated Protein Kinase 1	56	2.32
ESR1	Estrogen Receptor 1	61	1.96
CAMK2D	Calcium/Calmodulin Dependent Protein Kinase II $\delta$	68	1.68
NFATC3	Nuclear Factor of Activated T Cells 3	68	1.68
OGT	O-Linked N-Acetylglucosamine (GlcNAc) Transferase	69	1.59
MXI1	MAX Interactor 1, Dimerization Protein	69	1.59
ESRRG	Estrogen Related Receptor $\gamma$	75	0.68
CPOX	Coproporphyrinogen Oxidase	75	0.68
ESR2	Estrogen Receptor 2	80	0.36
TFAP2A	Transcription Factor AP-2 $\alpha$	80	0.36
ZBTB20	Zinc Finger and BTB Domain Containing 20	81	0.26
NPAS3	Neuronal PAS Domain Protein 3	81	0.26
NR4A2	Nuclear Receptor Subfamily 4 Group A Member 2	81	0.26
RICTOR	RPTOR Independent Companion of mTOR Complex 2	81	0.26
CAMK1	Calcium/Calmodulin Dependent Protein Kinase I	81	0.26
TET2	Tet Methylcytosine Dioxygenase 2	81	0.26
ATXN1	Ataxin 1	81	0.26
VHL	Von Hippel-Lindau Tumor Suppressor	81	0.26
HMGCR	3-Hydroxy-3-Methylglutaryl-CoA Reductase	81	0.26
ZNF521	Zinc Finger Protein 521	81	0.26

FSGS, focal segmental glomerulosclerosis; LN, lupus nephritis; AVV, ANCA-associated vasculitis; AGS, advanced global sclerosis; sCr, serum creatinine; CRP, C reactive protein; ESR, erythrocyte sedimentation rate. <sup>a</sup>Obtained from a total of 453 genes. <sup>b</sup>Based on analysis in VarElect of miR-204-5p predicted targets from TargetScan.

axis involved in mitochondrial biogenesis. Through gene data profiling, a total of 27 genes significantly associated with mitochondrial biogenesis were identified (Table III).

Next, the urinary expression of two predicted TUG1/miR-204 mRNA targets significantly associated with mitochondrial biogenesis, TFAM and COX5A, as well as one validated target, PPARGC1A (36), were quantified (Fig. 3). The relative expression levels of COX5A and PPARGC1A were lower in the urinary sediment of glomerulonephritides patients when compared with healthy controls (Fig. 3C and E). Additionally, there was a significant correlation between TUG1 urinary expression and all three mitochondrial biogenesis mRNAs quantified in the present study (TFAM,  $r=0.6328$ ,  $^{**}P<0.01$ ; COX5A,  $r=0.4872$ ,  $^{*}P<0.05$ ; PPARGC1A,  $r=0.7206$ ,  $^{**}P<0.01$ ; Fig. 3B-F). These results suggest that urinary RNAs may reflect molecular processes involved in the pathophysiology of renal dysfunction.

## Discussion

The diagnosis of glomerulonephritides in current clinical practice is primarily dependent on the histological analysis of

renal biopsies (6). However, this traditional approach is being challenged by molecular techniques that go beyond description and examine disease mechanisms (37,38). Multiple studies have shown the association between dysregulated expression of certain lncRNAs and the development and progression of pathological disease states that affect the kidney (12-17). Urinary non-coding RNAs are a promising non-invasive tool able to reflect renal disease, aid in its appropriate diagnosis, and guide therapeutic choices (12,39,40). In the present study, it was shown that lncRNA TUG1 was present in the urinary sediment, and its expression was significantly reduced in patients with biopsy-confirmed glomerulonephritides, particularly those diagnosed with FSGS. Moreover, there was a positive correlation between urinary TUG1 expression and mRNAs known to be involved in pathogenic mechanisms associated with podocyte loss.

Mutations in NPHS1 or NPHS2 lead to proteinuria and rapid ESRD progression, highlighting the critical role of these podocyte-specific structural proteins in glomerular filtration barrier function (41,42). Although NPHS1 and NPHS2 are detectable in the urine of glomerulonephritides patients and




Figure 3. Levels of urinary lncRNA TUG1 correlate with mitochondrial biogenesis mRNAs predicted to be modulated by the TUG1/miR-2014-5p axis. Reverse transcription-quantitative PCR results showing the log<sub>2</sub>-fold change in expression of (A) TFAM, (C) COX5A and (E) PPARGC1A in the urinary sediment of healthy controls (n=10) and glomerulonephritides patients (n=10). Data are presented as the mean  $\pm$  standard error of the mean. Correlation between lncRNA TUG1 transcript levels and (B) TFAM, (D) COX5A and (F) PARGC1A in the urinary sediment of healthy controls (n=10) and glomerulonephritides patients (n=10). \* $P<0.05$ , \*\* $P<0.01$ . TUG1, taurine upregulated gene 1; miR, microRNA; TFAM, transcription factor A mitochondrial, COX5A, cytochrome C oxidase subunit 5A; and PPARGC1A, peroxisome proliferator-activated receptor  $\gamma$  coactivator 1 $\alpha$ .

have been described as clinically valuable biomarkers of podocyte injury, the results between studies are inconsistent and raise questions regarding the reasons behind the contrasting findings (43). In the present study, a decrease in urinary NPHS1 expression in FSGS was observed when compared to other diagnoses, but no differences in NPHS2 levels within the groups was found. These results partially agree with previous studies that showed urinary NPHS1 and NPHS2 mRNA levels were reduced in patients with MCN and FSGS, a feature that correlates with the degree of proteinuria (44,45). However, TUG1 expression levels showed no correlation with GFR, proteinuria or albuminuria in the present study

(data not shown). Although it has been proposed that these podocyte-specific biomarkers can detect nephropathy before the development of albuminuria (46-48), a larger patient cohort is required to assess the relationship between TUG1 urinary expression and clinical markers of renal dysfunction traditionally used in the diagnosis of glomerulonephritides. In the present study, a positive correlation between NPHS1 and NPHS2 mRNA levels and TUG1 expression was observed, which suggests that the molecular mechanisms described *in vitro* and in animal models of DN regarding the relationship between TUG1 and podocyte loss are reflected in the urine of non-diabetic patients (28).

A decrease in PGC1A, a transcriptional coactivator that regulates energy homeostasis and mitochondrial biogenesis, has been implicated in the development of acute kidney injury, DN and renal fibrosis (49). The expression levels and activity of PGC1A depend on multiple transcriptional and post-transcriptional mechanisms, some mediated by non-coding RNAs (22,28). Interactions between TUG1 and the promoter of PGC1A enhances its transcription, and increases mitochondrial content and cellular ATP levels whilst reducing mitochondrial reactive oxygen species levels. In addition, TUG1 acts as a competitive endogenous RNA for miRNAs such as miR-145, miR-144 and miR-204-5p (30,50,51). miRNA miR-204-5p is involved in multiple cellular processes, including angiogenesis, vascular disease, metabolism and glucose homeostasis (30,52). Through *in silico* analysis, targets of the TUG1/miR-204-5p axis involved in mitochondrial biogenesis were predicted and it was subsequently shown that PGC1A, COX5A and TFAM were also detectable in the urine and were correlated with TUG1 expression. PGC1A is a known posttranscriptional target of miR-204-5p (53,54); however, COX5A and TFAM still require experimental validation. Nonetheless, the results suggest that the TUG1/miR-204 axis and their mitochondrial biogenesis targets are relevant in FSGS pathogenesis and may serve as potential biomarkers.

The present study has some limitations, including the small number of glomerulonephritides patients enrolled and the variability of diagnosis, which included primary and secondary glomerulonephritides. Nonetheless, our findings support a role for lncRNA TUG1 in the development of glomerulonephritides, and justify further exploration of urinary TUG1 as a potential biomarker of FSGS. More extensive studies, with carefully selected subjects are required to confirm the relationship between urinary TUG1 expression, clinical markers of renal dysfunction and glomerular damage observed in kidney biopsies; and to validate lncRNA TUG1 as a biomarker of FSGS. Additionally, more *in vitro* and animal experiments focused on the TUG1/miR-204-5p axis should be performed to increase our understanding of these non-coding RNAs in mitochondrial bioenergetics and podocyte loss.

In conclusion, the results of the present study highlight the potential of urinary non-coding RNAs, such as TUG1, in the diagnosis of renal disease, and suggests that this lncRNA and its mitochondrial-associated pathways are relevant in glomerulonephritides other than DN. Further studies are required to evaluate urinary TUG1 as a potential biomarker of podocytopathy, and to determine its association with kidney dysfunction and patient prognosis.

## Acknowledgements

We would like to thank Claudia Susana Meza Calvillo for her technical assistance as part of the Scientific Research Summer Program of the Mexican Academy of Science 2019.

## Funding

This study was supported by Consejo Nacional de Ciencia y Tecnología (CONACyT), Mexico (grant no. SALUD-2018-02-B-S-42687).

## Availability of data and materials

The datasets used and/or analysed during the present study are available from the corresponding author on reasonable request.

## Authors' contributions

FJS-T, MM-P and RE conceived the study. FJS-T and MM-P collected and analysed the data. ZM performed the experiments. CM-C performed the histopathological analysis of renal biopsies. RE drafted the manuscript. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

The present study complied with the ethical principles for medical research specified in the Declaration of Helsinki and was approved by the Local Ethics and Research Committee at the Hospital de Especialidades NUM. 1, Bajío, León, Guanajuato, Instituto Mexicano del Seguro Social (approval no. CLIEIS R-2018-1001-114). Each participant provided written informed consent prior to enrolment in the study.

## Patient consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

1. Floege J and Amann K: Primary glomerulonephritides. *Lancet* 387: 2036-2048, 2016.
2. Caliskan Y and Kiryluk K: Novel biomarkers in glomerular disease. *Adv Chronic Kidney Dis* 21: 205-216, 2014.
3. Matovinović MS: Podocyte injury in glomerular diseases. *EJIFCC* 20: 21-27, 2009.
4. Mallipattu SK and He JC: The podocyte as a direct target for treatment of glomerular disease? *Am J Physiol Renal Physiol* 311: F46-F51, 2016.
5. Bagavant H and Fu SM: Pathogenesis of kidney disease in systemic lupus erythematosus. *Curr Opin Rheumatol* 21: 489-494, 2009.
6. Hogan J, Mohan P and Appel GB: Diagnostic tests and treatment options in glomerular disease: 2014 update. *Am J Kidney Dis* 63: 656-666, 2014.
7. Parikh SV, Ayoub I and Rovin BH: The kidney biopsy in lupus nephritis: Time to move beyond histology. *Nephrol Dial Transplant* 30: 3-6, 2015.
8. Rovin BH, Almaani S and Malvar A: Reimagining the kidney biopsy in the era of diagnostic biomarkers of glomerular disease. *Kidney Int* 95: 265-267, 2019.
9. Kopp F and Mendell JT: Functional classification and experimental dissection of long noncoding RNAs. *Cell* 172: 393-407, 2018.
10. Mongelli A, Martelli F, Farsetti A and Gaetano C: The dark that matters: Long Non-coding RNAs as master regulators of cellular metabolism in Non-communicable diseases. *Front Physiol* 10: 369, 2019.
11. Lorenzen JM and Thum T: Long noncoding RNAs in kidney and cardiovascular diseases. *Nat Rev Nephrol* 12: 360-373, 2016.
12. Ignarski M, Islam R and Müller RU: Long Non-coding RNAs in kidney disease. *Int J Mol Sci* 20: 3276, 2019.
13. Tian H, Wu M, Zhou P, Huang C, Ye C and Wang L: The long non-coding RNA MALAT1 is increased in renal ischemia-reperfusion injury and inhibits hypoxia-induced inflammation. *Ren Fail* 40: 527-533, 2018.

14. Hu S, Han R, Shi J, Zhu X, Qin W, Zeng C, Bao H and Liu Z: The long noncoding RNA LOC105374325 causes podocyte injury in individuals with focal segmental glomerulosclerosis. *J Biol Chem* 293: 20227-20239, 2018.
15. Han R, Hu S, Qin W, Shi J, Zeng C, Bao H and Liu Z: Upregulated long noncoding RNA LOC105375913 induces tubulointerstitial fibrosis in focal segmental glomerulosclerosis. *Sci Rep* 9: 716, 2019.
16. Jin LW, Pan M, Ye HY, Zheng Y, Chen Y, Huang WW, Xu XY and Zheng SB: Down-regulation of the long non-coding RNA XIST ameliorates podocyte apoptosis in membranous nephropathy via the miR-217-TLR4 pathway. *Exp Physiol* 104: 220-230, 2019.
17. Liao Z, Ye Z, Xue Z, Wu L, Ouyang Y, Yao C, Cui C, Xu N, Ma J, Hou G, *et al*: Identification of renal long Non-coding RNA RP11-2B6.2 as a positive regulator of type I interferon signaling pathway in lupus nephritis. *Front Immunol* 10: 975, 2019.
18. Santer L, López B, Ravassa S, Baer C, Riedel I, Chatterjee S, Moreno MU, González A, Querejeta R, Pinet F, *et al*: Circulating long noncoding RNA LIPCAR predicts heart failure outcomes in patients without chronic kidney disease. *Hypertension* 73: 820-828, 2019.
19. Huang YS, Hsieh HY, Shih HM, Sytwu HK and Wu CC: Urinary Xist is a potential biomarker for membranous nephropathy. *Biochem Biophys Res Commun* 452: 415-421, 2014.
20. Forbes JM and Thorburn DR: Mitochondrial dysfunction in diabetic kidney disease. *Nat Rev Nephrol* 14: 291-312, 2018.
21. Young TL, Matsuda T and Cepko CL: The noncoding RNA taurine upregulated gene 1 is required for differentiation of the murine retina. *Curr Biol* 15: 501-512, 2005.
22. Li SY, Park J, Qiu C, Han SH, Palmer MB, Arany Z and Susztak K: Increasing the level of peroxisome proliferator-activated receptor  $\gamma$  coactivator-1 $\alpha$  in podocytes results in collapsing glomerulopathy. *JCI Insight* 2: e92930, 2017.
23. Li SY and Susztak K: The long noncoding RNA Tug1 connects metabolic changes with kidney disease in podocytes. *J Clin Invest* 126: 4072-4075, 2016.
24. LeBleu VS, O'Connell JT, Gonzalez Herrera KN, Wikman H, Pantel K, Haigis MC, de Carvalho FM, Damascena A, Domingos Chinen LT, Rocha RM, *et al*: PGC-1 $\alpha$  mediates mitochondrial biogenesis and oxidative phosphorylation in cancer cells to promote metastasis. *Nat Cell Biol* 16: 992-1003, 1-15, 2014.
25. Uldry M, Yang W, St-Pierre J, Lin J, Seale P and Spiegelman BM: Complementary action of the PGC-1 coactivators in mitochondrial biogenesis and brown fat differentiation. *Cell Metab* 3: 333-341, 2006.
26. Finck BN and Kelly DP: PGC-1 coactivators: Inducible regulators of energy metabolism in health and disease. *J Clin Invest* 116: 615-622, 2006.
27. Kang HM, Ahn SH, Choi P, Ko YA, Han SH, Chinga F, Park AS, Tao J, Sharma K, Pullman J, *et al*: Defective fatty acid oxidation in renal tubular epithelial cells has a key role in kidney fibrosis development. *Nat Med* 21: 37-46, 2015.
28. Long J, Badal SS, Ye Z, Wang Y, Ayanga BA, Galvan DL, Green NH, Chang BH, Overbeek PA and Danesh FR: Long noncoding RNA Tug1 regulates mitochondrial bioenergetics in diabetic nephropathy. *J Clin Invest* 126: 4205-4218, 2016.
29. Association WM: World medical association declaration of Helsinki: Ethical principles for medical research involving human subjects. *JAMA* 310: 2191-2194, 2013.
30. Yu C, Li L, Xie F, Guo S, Liu F, Dong N and Wang Y: LncRNA TUG1 sponges miR-204-5p to promote osteoblast differentiation through upregulating Runx2 in aortic valve calcification. *Cardiovasc Res* 114: 168-179, 2018.
31. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
32. Ben-Ari Fuchs S, Lieder I, Stelzer G, Mazor Y, Buzhor E, Kaplan S, Bogoch Y, Plaschkes I, Shitrit A, Rappaport N, *et al*: GeneAnalytics: An integrative gene set analysis tool for next generation sequencing, RNAseq and microarray data. *OMICS* 20: 139-151, 2016.
33. Agarwal V, Bell GW, Nam JW and Bartel DP: Predicting effective microRNA target sites in mammalian mRNAs. *Elife* 4: e05005, 2015.
34. Levey AS, Eckardt KU, Tsukamoto Y, Levin A, Coresh J, Rossert J, De Zeeuw D, Hostetter TH, Lameire N and Eknoyan G: Definition and classification of chronic kidney disease: A position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 67: 2089-2100, 2005.
35. Fukuda A, Wickman LT, Venkatarreddy MP, Wang SQ, Chowdhury MA, Wiggins JE, Shedden KA and Wiggins RC: Urine podocin:nephrin mRNA ratio (PNR) as a podocyte stress biomarker. *Nephrol Dial Transplant* 27: 4079-4087, 2012.
36. Civelek M, Hagopian R, Pan C, Che N, Yang WP, Kayne PS, Saleem NK, Cederberg H, Kuusisto J, Gargalovic PS, *et al*: Genetic regulation of human adipose microRNA expression and its consequences for metabolic traits. *Hum Mol Genet* 22: 3023-3037, 2013.
37. Saez-Rodriguez J, Rinschen MM, Floege J and Kramann R: Big science and big data in nephrology. *Kidney Int* 95: 1326-1337, 2019.
38. Kretzler M, Cohen CD, Doran P, Henger A, Madden S, Gröne EF, Nelson PJ, Schlöndorff D and Gröne HJ: Repuncturing the renal biopsy: Strategies for molecular diagnosis in nephrology. *J Am Soc Nephrol* 13: 1961-1972, 2002.
39. Sun IO and Lerman LO: Urinary microRNA in kidney disease: Utility and roles. *Am J Physiol Renal Physiol* 316: F785-F793, 2019.
40. Brandenburger T, Salgado Somoza A, Devaux Y and Lorenzen JM: Noncoding RNAs in acute kidney injury. *Kidney Int* 94: 870-881, 2018.
41. Patrakka J, Kestilä M, Wartiovaara J, Ruotsalainen V, Tissari P, Lenkkeri U, Männikkö M, Visapää I, Holmberg C, Rapola J, *et al*: Congenital nephrotic syndrome (NPHS1): Features resulting from different mutations in Finnish patients. *Kidney Int* 58: 972-980, 2000.
42. Boute N, Gribouval O, Roselli S, Benessy F, Lee H, Fuchshuber A, Dahan K, Gubler MC, Niaudet P and Antignac C: NPHS2, encoding the glomerular protein podocin, is mutated in autosomal recessive steroid-resistant nephrotic syndrome. *Nat Genet* 24: 349-354, 2000.
43. Sekulic M and Pichler Sekulic S: A compendium of urinary biomarkers indicative of glomerular podocytopathy. *Patholog Res Int* 2013: 782395, 2013.
44. Hara M, Yanagihara T and Kihara I: Urinary podocytes in primary focal segmental glomerulosclerosis. *Nephron* 89: 342-347, 2001.
45. Szeto CC, Wang G, Chow KM, Lai FM, Ma TK, Kwan BC, Luk CC and Li PK: Podocyte mRNA in the urinary sediment of minimal change nephropathy and focal segmental glomerulosclerosis. *Clin Nephrol* 84: 198-205, 2015.
46. Pätäri A, Forsblom C, Havana M, Taipale H, Groop PH and Holthöfer H: Nephropathy in diabetic nephropathy of type 1 diabetes. *Diabetes* 52: 2969-2974, 2003.
47. Ng DP, Tai BC, Tan E, Leong H, Nurbaya S, Lim XL, Chia KS, Wong CS, Lim WY and Holthöfer H: Nephropathy associates with multiple renal traits in type 2 diabetes. *Nephrol Dial Transplant* 26: 2508-2514, 2011.
48. Chang JH, Paik SY, Mao L, Eisner W, Flannery PJ, Wang L, Tang Y, Mattocks N, Hadjadj S, Goujon JM, *et al*: Diabetic kidney disease in FVB/NJ Akita mice: Temporal pattern of kidney injury and urinary nephrin excretion. *PLoS One* 7: e33942, 2012.
49. Lynch MR, Tran MT and Parikh SM: PGC1 $\alpha$  in the kidney. *Am J Physiol Renal Physiol* 314: F1-F8, 2018.
50. Tan J, Qiu K, Li M and Liang Y: Double-negative feedback loop between long non-coding RNA TUG1 and miR-145 promotes epithelial to mesenchymal transition and radioresistance in human bladder cancer cells. *FEBS Lett* 589: 3175-3181, 2015.
51. Cai H, Xue Y, Wang P, Wang Z, Li Z, Hu Y, Li Z, Shang X and Liu Y: The long noncoding RNA TUG1 regulates blood-tumor barrier permeability by targeting miR-144. *Oncotarget* 6: 19759-19779, 2015.
52. Flores-Pérez A, Marchat LA, Rodríguez-Cuevas S, Bautista-Piña V, Hidalgo-Miranda A, Ocampo EA, Martínez M, Palma-Flores C, Fonseca-Sánchez MA, Astudillo-de la Vega H, *et al*: Dual targeting of ANGPT1 and TGFBR2 genes by miR-204 controls angiogenesis in breast cancer. *Sci Rep* 6: 34504, 2016.
53. Maniyadath B, Chattopadhyay T, Verma S, Kumari S, Kulkarni P, Banerjee K, Lazarus A, Kokane SS, Shetty T, Anamika K and Kolthur-Seetharam U: Loss of hepatic oscillatory fed microRNAs Abrogates Refed transition and causes liver dysfunctions. *Cell Rep* 26: 2212-2226, 2019.
54. Houzelle A, Dahlmans D, Nascimento EBM, Schaart G, Jörgensen JA, Moonen-Kornips E, Kersten S, Wang X and Hoeks J: MicroRNA-204-5p modulates mitochondrial biogenesis in C2C12 myotubes and associates with oxidative capacity in humans. *J Cell Physiol* 235: 9851-9863, 2020.