

Whole rat DNA array survey for candidate genes related to hypertension in kidneys from three spontaneously hypertensive rat substrains at two stages of age and with hypotensive induction caused by hydralazine hydrochloride

KOSHO KINOSHITA, MOHAMMAD SAID ASHENAGAR, MASAKI TABUCHI and HIDEAKI HIGASHINO

Department of Pharmacology, Kinki University School of Medicine, Osaka 589-8511, Japan

Received October 1, 2010; Accepted December 28, 2010

DOI: 10.3892/etm.2011.193

Abstract. Clarification of the genetic nature and more effective care for hypertension are required, given the high incidences of cardiovascular and cerebrovascular mortality. Thus, we surveyed candidate genes for hypertension with rat whole gene DNA microarrays using three novel methods. Gene expression analyses were conducted as follows: Method 1, three types of spontaneously hypertensive rat (SHR) substrains, SHR, stroke-prone SHR (SHRSP) and malignant type of SHRSP (M-SHRSP) were used and compared to normotensive Wistar Kyoto rats; Method 2, the expressed genes between rats of different ages were compared for different blood pressures; and Method 3, genes that were expressed in rats treated with or without an acute hypotensive stimulus, the antihypertensive hydralazine hydrochloride, were compared. This approach identified dozens of genes, including *Dusp15*, *Cyp8b1*, *Armc 3*, *Gtpbp4*, *Mettl2*, *Mapk14*, *Prkar2b*, *frame 12*, *Anxa13*, *Ephx2*, *Myr8* and *Pcdh9* by Method 1; *Cyp2C* and *Atp12a* by Method 2; and *Kcnc3*, *Vnn1*, *TC560558* and *Gabrq* and a number of unknown genes by Methods 2 and 3, as probable candidate genes for hypertension in SHR substrains. *Ephx2* was previously reported as a candidate gene in SHRs; however other genes were identified for the first time in this study. Since it was not always possible to completely demonstrate that these genes are responsible for hypertension in SHRs, further research into true candidate genes that participate in the genesis of hypertension in SHR substrains is warranted.

Introduction

Hypertension is one of the most prevalent diseases in humans who live a modern life style. It is a silent disease and does

not markedly affect quality of life when it is moderate and not serious. However, when left untreated it leads to life-threatening diseases associated with atherosclerosis, including myocardial infarction, renal failure and strokes (1-3). In Japan, where general medical care should be sufficient to treat hypertension, more than 300,000 patients per year die from diseases that are related to hypertension (4,5). Clarification of the genetic nature and more effective care for hypertension are required.

There are a number of methods for investigating the genetic nature of hypertension. The polygenic nature of human essential hypertension has made it challenging to isolate the genes involved in the genesis of the disease. DNA microarrays are a potentially powerful tool for studying the genetics of hypertension, as they facilitate the measurement of the expression of thousands of genes simultaneously (6-8). Inbred homozygous rodent models of human essential hypertension are ideal for microarray research, and animal models of essential hypertension have been studied using microarrays (9).

In this study, we present a comparison of kidney gene expression in three hypertensive rat models: spontaneously hypertensive rats (SHRs) (10) and two substrains derived from SHR – stroke-prone SHRs (SHRSPs) (11) and malignant-type SHRSPs (M-SHRSPs) (12). SHR, the current model for essential hypertension research, were developed in a breeding program based solely on the selection of elevated blood pressure (BP) in Wistar Kyoto (WKY) rats (10). Normotensive WKY rats, from which the SHR strain was derived, were used as controls. SHRSPs were established from SHRs by selective inbreeding for stroke proneness (11), and M-SHRSPs were selected and established through the brother-sister mating of selected SHRSPs that showed higher BP following treatment with hydralazine hydrochloride to prevent the development of hypertension and stroke (12). An inbred strain of M-SHRSPs shows BPs of 250 mmHg or higher before 14 weeks of age and, compared to SHRSPs, M-SHRSPs showed more rapid and severe increases in BP and stroke in almost all animals. Our facility is one of the original places that bred the three types of SHR substrains and families. The former director, Professor Kozo Okamoto, introduced these animals to our school when relocating from the Kyoto University School of Medicine, the original site of SHR breeding. Using these three

Correspondence to: Dr Hideaki Higashino, Department of Pharmacology, Kinki University School of Medicine, Ohno-Higashi, Osaka-Sayama, Osaka 589-8511, Japan
E-mail: higashino@med.kindai.ac.jp

Key words: kidney, DNA array, hypertension, spontaneously hypertensive rat, stroke-prone spontaneously hypertensive rat, malignant type of stroke-prone spontaneously hypertensive rat, hydralazine

SHR substrains, we previously reported (13) that a number of BP-regulating genes, including *sparc/osteonectin (Spock2)*, *kynureninase (Kynu)*, *regulator of G-protein signaling 2 (Rgs2)* and *gap junction protein $\alpha 1$ (Gjal)* were identified as up-regulated, and *urotensin 2 (Uts2)*, *cytoplasmic epoxide hydrolase 2 (Ephx2)*, *apelin (Apln)*, *insulin-like growth factor 1 receptor (Igflr)* and *angiotensin II receptor-associated protein (Agtrap)* were identified as down-regulated in the adrenal glands at 6 weeks of age.

Kidneys are logical candidate organs for studying hypertension due to their direct influence on body fluids and endocrine, cardiovascular and sympathetic functions (14,15). The relationship between kidney function and blood pressure is known to be influenced by numerous intrinsic and extrinsic factors, such as the renin-angiotensin system and catecholamine and aldosterone hormones (16,17). Previously, *Styl* (18), *Edg1*, *Vcam1* (19), *C1q*, *CD24* (20), *SPON1* (21), *Gstm1* (22), *ACE-2* (23), *AMPK*, *APLP2* (24), *Ephx2*, *Elal* (25) and *Egln1* (26) were shown to be hypertension-related genes in SHR or SHRSP kidneys using a DNA microarray. In contrast to the reports, this study is the first attempt to compare gene expression profiles in the kidneys of three SHR substrains, SHR, SHRSP and M-SHRSP, employing WKY rats as controls and using DNA microarrays to survey for genes related to hypertension in these SHR substrains. In addition to analyzing gene expression in these three types of SHR substrains (Method 1), young rats whose BP levels were not yet elevated (6 weeks of age) and slightly older rats that developed hypertension (9 weeks of age) were used to survey candidate blood pressure elevating genes and to examine the relationship between blood pressure elevation and gene expression (Method 2). Next, a hypotensive drug with no known receptor, hydralazine hydrochloride, was administered to each group of rats to induce acute hypotension to detect hypertension-associated genes. Thus, genes induced by acute hypotension were identified (Method 3). This study aimed to use three analytical methods to comprehensively identify candidate genes involved in the genesis of hypertension in the kidneys of SHR substrains.

Materials and methods

Animals. The experiments were performed on rats at 6 and 9 weeks of age. A total of 3 rats were used in each experimental group. WKY/Izm was used as a wild-type control strain, and SHR/Kpo (10), SHRSP/Kpo (11) and M-SHRSP/Kpo (12) were hypertensive model rats. WKY/Izm were purchased from SLC Co. (Shizuoka, Japan) and the other three substrains were purchased from the Animal Center of Kinki University School of Medicine. The animals used in this experiment were handled with due care according to the guidelines established by the Japanese Association for Laboratory Animal Science, which complies with international rules and policies. This study was performed under approval (KAME-19-078 on April 1, 2007) of the Animal Care and Use Committee of Kinki University. Measures were taken to minimize the pain and discomfort of the experimental animals. Hydralazine hydrochloride (30 mg/kg/day) mixed with SP-2 chow (Funahashi Farm, Chiba, Japan) was administered to half of each rat group (n=3) to induce an acute decline in blood pressure for 2 days prior to euthanasia.

Systolic blood pressure measurements. Systolic BP (SBP) was measured using the tail-cuff method with a UR-5000 instrument (Ueda, Tokyo, Japan). Briefly, three consecutive SBP readings were taken between 09:00-11:00 after warming the body at 35°C for 5 min in a heater box. The SBP values were expressed as the mean \pm SEM.

Tissue processing and RNA isolation. After the kidneys were harvested under sodium pentobarbital anesthesia (50 mg/kg i.p.), the organs were homogenized at a pitch speed of 22 strokes/sec for 2 min (twice) in a 2-ml plastic tube with 5-mm diameter glass beads with a Qiagen Tissue Lyser (Retsch GmbH & Co., Haan, Germany). Total RNA was extracted with an RNeasy Mini kit (Qiagen GmbH, Hilden, Germany) according to the manufacturer's protocol. The quality of RNA was checked with RNA Nano Chips (Agilent Technologies, Waldbornn, Germany) with an Agilent 2100 Bioanalyzer, and the RNA was then used in the microarray experiments. For the microarray analysis, tissues from 3 rats per group of all hypertensive and normotensive strains were used in each experiment.

Analysis of gene expression profiling with oligonucleotide arrays. To examine the gene expression profiles of rat kidneys, cRNA labeled with cyanine 3-CTP (PerkinElmer, Boston, MA, USA) was synthesized from 1 μ g of DNase I-treated total RNA with a Low RNA Input Amplification kit (Agilent Technologies), and this was hybridized by incubating with a Whole Rat Genome Microarray (4x44K formatted) (Agilent Technologies) in a rotor oven (Agilent Technologies) for 17 h at 65°C, followed by washing. The hybridized slides were scanned with an Agilent GenPix Scanner 4000 (Agilent Technologies). The data were extracted, and the overall raw signal intensities on each array were normalized to the median value of all rat probes with BRB-Array Tool software ver. 3.7.0. (Biometric Research Branch) (27). A significance level ($P < 0.01$) for each probe was set using the univariate Student's t-test.

Annotation of differentially expressed genes. A BLASTN search of the NCBI RefSeq database was performed, employing corresponding 60-nucleotide probes (NCBI, GEO accession: GPL7294) to identify homologous genes with functional annotations (28). After running the BLASTN search, clones showing either a score higher than 50 or an E-value below 5e-05 were defined as annotated clones, and the remaining clones were defined as non-annotated clones or unknown genes. The annotated gene and protein symbols are shown in italics and regular font, respectively.

Strategies to survey candidate genes related to hypertension. The following three different analyses were adopted to identify candidate genes related to or causing hypertension as described above. Method 1: data from the comparison among the SHR substrains were used to survey candidate genes among the SHRs, SHRSPs and M-SHRSPs according to the ascending order of hypertension. Method 2: data from the comparison between the 6- and 9-week-old rats of each SHR substrain were used to survey the candidate genes in ascending or descending order of expression between age groups in each substrain. Method 3: data obtained from the comparison of

Table I. SBP levels (mmHg) of the WKYs, SHRs, SHRSPs and M-SHRSPs with or without hydralazine administration for 2 days.

Rats	6 weeks of age (n=3)		9 weeks of age (n=3)	
	No treatment	30 mg/kg hydralazine	No treatment	30 mg/kg hydralazine
WKY	130±4.1	118±4.9 ^d	137±5.2	122±1.2 ^d
SHR	139±2.5 ^a	127±6.4 ^d	158±7.1 ^{b,c}	146±4.3 ^d
SHRSP	140±7.6	126±1.2 ^d	180±10 ^{b,c}	161±5.8 ^d
M-SHRSP	153±2.1 ^a	142±3.4 ^d	217±10 ^{b,c}	198±6.0 ^d

Significant differences at P<0.05; ^avs. 6-week-old WKY; ^bvs. 9-week-old WKY; ^cvs. the same subgroup of rats aged 6 weeks, ^dvs. without hydralazine (no treatment);

Table II. Genes (16) commonly expressed more than four times higher in the kidneys of the SHRs, SHRSPs and M-SHRSPs compared to WKY rats at 6 weeks of age.

Fold increases (/WKY)			Clones	GB account	Description
SHR	SHRSP	M-SHRSP			
1. 91.961	27.387	49.827	A_43_P13995	Gc	Group-specific component
2. 26.341	29.378	20.778	A_44_P863709	AW143870	RGICB87 5' end, mRNA sequence
3. 18.587	17.952	17.263	A_44_P868694	TC538548	Sugt 1, kinetochore function
4. 18.822	17.034	17.395	A_44_P562830	Dusp15_predicted	Dual specificity phosphatase-like 15 (predicted)
5. 14.245	15.181	9.883	A_44_P653949	TC558814	Unknown
6. 13.108	6.051	9.843	A_44_P387780	Cyp8b1	Cytochrome P450, family 8, subfamily b, polypeptide 1
7. 8.900	7.453	9.406	A_44_P371125	Sult1b1	Sulfotransferase family 1B, member 1
8. 6.604	7.894	7.490	A_44_P670594	TC528756	O87809 EprE protein
9. 9.898	5.456	6.217	A_42_P655890	XM_225610	Armc3, multiple functions in signal transduction
10. 6.068	7.768	7.310	A_44_P470661	Serpina3m	Serine (or cysteine) proteinase inhibitor, member 3M
11. 7.768	6.944	5.705	A_42_P732437	Bri3bp	Bri3 binding protein
12. 7.222	6.495	6.503	A_44_P227132	RGD1564324_predicted	cDNA clone UI-R-CV1-bvu-e-07-0-UI 3'
13. 6.473	6.865	5.205	A_44_P928190	RGD1561961_predicted	Similar to IQ motif and WD repeats 1 (predicted)
14. 5.301	5.194	5.294	A_42_P578953	Pthr1_predicted	Peptidyl-tRNA hydrolase 1 homolog (predicted)
15. 5.733	5.427	4.624	A_44_P176831	BG378920	BG378920 UI-R-CV1-bvu-e-07-0-UI.s1 UI-R-CV1
16. 4.841	4.507	4.438	A_44_P1033521	Trps1_predicted	Trichorhinophalangeal syndrome I (predicted)

Gene names in bold print represent genes commonly expressed at 6 and 9 weeks of age.

hypotensive effects, with or without hydralazine hydrochloride treatment, in the SHR substrains compared to the WKY rats were used to survey the genes induced by this treatment in each SHR substrain.

Analyses of genes expressed in biochemical pathways. Analyses of the role of genes expressed in biochemical pathways were performed using Skypainter of REACTOME, a free and open-source database (<http://www.reactome.org/>), offered on the website of the Cold Spring Harbor Laboratory, The European Bioinformatics Institute and The Gene Ontology Consortium.

Statistical analyses. Comparisons between the means of the data in each group were performed using one-way analysis of variance (ANOVA) and Scheffe's multiple comparisons test. Differences were considered significant at P<0.05 and P<0.01 for BP measurements and DNA array measurements, respectively.

Results

Blood pressure. SBP was measured in the WKY rats and the three SHR substrains at 6 and 9 weeks of age before and 2 days after receiving 30 mg/kg/day p.o. hydralazine hydrochloride

Table III. Genes (37) commonly expressed more than four times higher in the kidneys of the SHR_s, SHRSP_s and M-SHRSP_s compared to the WKY rats at 9 weeks of age.

Fold increases (/WKY)			Clones	GB account	Description
SHR	SHRSP	M-SHRSP			
1. 29.913	28.674	39.915	A_44_P863709	AW143870	cDNA clone RGICB87 5' end
2. 22.900	29.660	36.970	A_44_P562830	Dusp15_predicted	Dual specificity phosphatase-like 15 (predicted)
3. 40.920	14.342	25.341	A_44_P457153	RGD1564999_predicted	Similar to isopentenyl-diphosphate δ isomerase 2
4. 26.305	15.898	14.860	A_42_P655890	XM_225610	Armc 3, multiple functions in signal transduction
5. 25.413	12.309	17.761	A_44_P387780	Cyp8b1	Cytochrome P450, family 8, subfamily b, polypeptide 1
6. 15.568	14.383	23.048	A_44_P653949	TC558814	Unknown
7. 13.884	9.754	15.786	A_44_P330188	Acox2	Acyl-Coenzyme A oxidase 2
8. 9.098	10.313	16.107	A_44_P868694	TC538548	Sugt 1, kinetochore function
9. 7.847	9.047	16.460	A_44_P970369	BG664685	cDNA clone DRABHF03 5'
10. 10.039	8.030	13.229	A_44_P132729	Rdh2	Retinol dehydrogenase 2
11. 9.048	8.655	11.654	A_42_P826202	Zfp597	Zinc finger protein 597
12. 11.879	4.934	10.385	A_43_P12900	Gtpbp4	GTP binding protein 4
13. 6.778	7.495	10.912	A_44_P470661	Serpina3m	Serine (or cysteine) proteinase inhibitor, member 3M
14. 6.829	7.566	11.821	A_44_P746348	RGD1562658_predicted	Similar to RIKEN cDNA 1700009P17 (predicted)
15. 8.605	5.376	11.989	A_43_P13995	Gc	Group-specific component
16. 11.364	4.291	9.047	A_44_P1036339	XR_006738	Similar to nucleolar GTP-binding protein 1
17. 6.639	6.330	11.704	A_44_P1029892	Tmem14a_predicted	Transmembrane protein 14A (predicted)
18. 4.650	7.172	12.044	A_43_P23115	XM_347233	Similar to indolethylamine N-methyltransferase
19. 10.275	5.217	8.145	A_44_P792207	TC539990	O63614 (O63614) ATP synthase subunit 8
20. 6.619	8.338	8.469	A_44_P777328	TC540923	Phosphatidylinositol 3 kinase regulator
21. 5.459	5.549	9.791	A_44_P670594	TC528756	O87809 (O87809) EprE protein
22. 6.291	5.570	8.778	A_44_P464942	BF545795	cDNA clone UI-R-BT0-qc-d-07-0-UI 5'
23. 6.257	5.488	8.339	A_44_P117119	Gloxd1	Glyoxalase domain containing 1
24. 6.115	5.583	7.945	A_42_P645467	Fbxo36_predicted	F-box only protein 36 (predicted)
25. 8.799	6.587	4.183	A_43_P16225	J01879	Rat brain-specific identifier sequence RNA, clone p2a120
26. 5.900	4.490	7.362	A_44_P556895	Ddit4	DNA-damage-inducible transcript 4
27. 5.505	5.534	6.463	A_44_P928190	RGD1561961_predicted	Similar to IQ motif and WD repeats 1 (predicted)
28. 4.469	4.354	8.532	A_42_P649672	Sv2a	Synaptic vesicle glycoprotein 2a
29. 6.838	5.859	4.258	A_44_P218896	Cyr61	Cysteine rich protein 61
30. 5.950	4.612	6.371	A_44_P333374	RGD1560736_predicted	Similar to solute carrier family 9 (predicted)
31. 4.734	4.573	6.306	A_44_P180259	Dpt_predicted	Dermatopontin (predicted)
32. 4.471	4.216	6.773	A_44_P534791	Mettl2_predicted	Methyltransferase like 2 (predicted)
33. 4.846	4.425	5.821	A_44_P360409	Mapk14	Mitogen activated protein kinase 14
34. 5.594	4.498	4.891	A_44_P497253	LOC689240	Similar to amyotrophic lateral sclerosis 2 chromosome region
35. 4.151	4.143	5.636	A_42_P732437	Bri3bp	Bri3 binding protein
36. 4.203	4.772	4.697	A_44_P466118	Slc11a1	Solute carrier family 11, member 1
37. 4.512	4.038	4.971	A_44_P316122	Prkar2b	Protein kinase, cAMP dependent regulatory, type II β

Gene names in bold print represent genes commonly expressed either at 6 or 9 weeks of age.

(Table I). SBP levels in the SHR_s and M-SHRSP_s were significantly higher than levels in the WKY rats at 6 weeks of age. At 9 weeks of age, SBP levels increased in the order

WKY rats, SHR_s, SHRSP_s and M-SHRSP_s as compared to those at 6 weeks of age, and every value was higher than that noted in the WKY rats. Moreover, hydralazine hydrochloride

Table IV. Genes (8) commonly expressed more than four times in the kidneys of the SHRs, SHRSPs and M-SHRSPs compared to the WKY rats at 6 and 9 weeks of age.

Fold increases (/WKY): 6 or 9 weeks						
	SHR	SHRSP	M-SHRSP	Clones	GB account	Description
1.	91.961 8.605	27.387 5.376	49.827: 6W 11.989: 9W	A_43_P13995	<i>Gc</i>	Group-specific component
2.	18.822 22.900	17.034 29.660	17.395: 6W 36.970: 9W	A_44_P562830	<i>Dusp15</i> _predicted	Dual specificity phosphatase-like 15 (predicted)
3.	14.245 15.568	15.181 14.383	9.883: 6W 23.048: 9W	A_44_P653949	TC558814	Unknown
4.	18.587 9.098	17.952 10.313	17.263: 6W 16.107: 9W	A_44_P868694	TC538548	<i>Sugt 1</i> , kinetochore function
5.	13.108 25.413	6.051 12.309	9.843: 6W 17.761: 9W	A_44_P387780	<i>Cyp8b1</i>	Cytochrome P450, family 8, subfamily b, polypeptide 1
6.	9.898 26.305	5.456 15.898	6.217: 6W 14.860: 9W	A_42_P655890	XM_225610	<i>Armc 3</i> , multiple functions in signal transduction
7.	6.068 6.778	7.768 7.495	7.310: 6W 10.912: 9W	A_44_P470661	<i>Serpina3m</i>	Serine (or cysteine) proteinase inhibitor, member 3M
8.	7.768 4.151	6.944 4.143	5.705: 6W 5.636: 9W	A_42_P732437	<i>Bri3bp</i>	Bri3 binding protein

6W and 9W represent 6 and 9 weeks of age, respectively.

(30 mg/kg/day p.o.) significantly decreased SBP levels in every rat at 6 and 9 weeks of age 2 days after drug administration compared to the values prior to drug administration

Microarray findings. Since this study aimed to identify the candidate genes involved in the genesis of hypertension in three SHR substrains, gene expression profiles were compared to DNA microchips that contained 41,012 probes using the mRNAs extracted from the kidneys of the three SHR substrains at 6 and 9 weeks of age, with or without the hypotensive agent, hydralazine hydrochloride, which dilates resistant arteries distributed in the whole body through an unknown receptor. The numbers of up- and down-regulated genes were less than the number of probes due to redundancy in the probe sets (i.e., in some cases, two or three probes represent one gene).

Method 1: Comparison of the spontaneously hypertensive rat substrains. Significantly increased or decreased DNA array data from each SHR substrain were obtained and compared to age-matched WKY rats. In 6-week-old rats, the numbers of significantly up-regulated genes compared to the WKY rats at the level of a probability ratio <0.01 were 217, 405 and 224 in SHRs, SHRSPs and M-SHRSPs, respectively. The numbers of commonly expressed genes between the SHRs and SHRSPs, SHRSPs and M-SHRSPs, and M-SHRSPs and SHRs were 135, 123 and 79, respectively. A total of 63 genes were commonly expressed in the SHRs, SHRSPs and M-SHRSPs. Among these genes, 16 were expressed more than four times higher than in the WKY rats (Table II). Of these 16 genes, *Gc*, *Sugt 1*, *Dusp15*, *Cyp8b1*, *Sult1b1*, *EprE*,

Armc 3, *Serpina3m*, *Bri3bp*, *Ptrhl* and *Trps1* were identified as known functional genes in addition to five previously unidentified genes. In 9-week-old rats, the numbers of significantly up-regulated genes compared to the WKY rats at the level of a probability ratio <0.01 were 5,447, 4,172 and 9,549 in the SHRs, SHRSPs and M-SHRSPs, respectively. The numbers of commonly expressed genes between the SHRs and SHRSPs, SHRs and M-SHRSPs, and SHRSPs and M-SHRSPs were 2,764, 3,647 and 3,018, respectively. In the SHRs, SHRSPs and M-SHRSPs, 2,103 genes were commonly expressed. Of these genes, 37 genes were expressed more than four times higher compared to the WKY rats (Table III). Of these 37 genes, *Dusp15*, *Armc 3*, *Cyp8b1*, *Acox2*, *Sugt 1*, *Rdh2*, *Zfp597*, *Gtpbp4*, *Serpina3m*, *Gc*, *XR_006738* (similar to nucleolar GTP-binding protein 1), *Tmem14a*, *XM_347233* (similar to indolethylamine N-methyltransferase), *TC539990* (ATP synthase subunit 8), *TC540923* (phosphatidylinositol 3 kinase regulator), *TC528756* (EprE protein), *Gloxd1*, *Fbxo36*, *Ddit4*, *Sv2a*, *Cyr61*, *RGD1560736* (similar to solute carrier family 9), *Dpt*, *Mett12*, *Mapk14*, *LOC689240* (similar to amyotrophic lateral sclerosis 2 chromosome region), *Bri3bp*, *Slc11a1* and *Prkar2b* were identified as known functional genes in addition to ten previously unidentified genes. A total of 8 genes were identified that were commonly expressed at a higher level in the SHRs, SHRSPs and M-SHRSPs compared to the WKY rats at 6 and 9 weeks of age (Table IV). A total of 7 known genes, *Gc*, *Dusp15*, *Sugt 1*, *Cyp8b1*, *Armc 3*, *Serpina3m* and *Bri3bp*, and one previously unknown gene were detected.

In 6-week-old rats, the numbers of significantly down-regulated genes compared to the WKY rats at the level of a

Table V. Genes (6) whose expression was commonly lower in the kidneys of the SHR_s, SHRSP_s and M-SHRSP_s compared to the WKY rats at 6 weeks of age.

	Fold changes (/WKY)			Clones	GB account	Description
	SHR	SHRSP	M-SHRSP			
1.	0.088	0.088	0.091	A_44_P808578	TC525845	<i>Sc1B</i>, collagen-like surface protein of <i>Streptococcus</i>
2.	0.104	0.094	0.088	A_44_P362486	BQ780215	cDNA clone UI-R-FF0-cpb-k-24-0-UI 3'
3.	0.044	0.179	0.114	A_44_P292314	<i>Hmmr</i>	Hyaluronan mediated motility receptor
4.	0.127	0.141	0.102	A_44_P402948	TC540893	Unknown
5.	0.156	0.161	0.113	A_44_P306008	AI179380	<i>Frame 12</i>, RSPCG42 3' end
6.	0.174	0.174	0.220	A_42_P719350	RGD1562974_predicted	Similar to DKFZp434P0316 (predicted)

Gene names in bold print represent genes commonly expressed either at 6 or 9 weeks of age.

Table VI. Genes (18) whose expression was commonly lower in the kidneys of the SHR_s, SHRSP_s and M-SHRSP_s compared to WKY at 9 weeks of age.

	Fold changes (/WKY)			Clones	GB account	Description
	SHR	SHRSP	M-SHRSP			
1.	0.033	0.039	0.032	A_44_P157134	<i>Anxa13</i> _predicted	Annexin A13 (predicted)
2.	0.071	0.073	0.055	A_44_P396474	RGD1563398_predicted	RGD1563398 (predicted)
3.	0.062	0.054	0.093	A_44_P362486	BQ780215	cDNA clone UI-R-FF0-cpb-k-24-0-UI 3'
4.	0.077	0.080	0.077	A_44_P808578	TC525845	<i>Sc1B</i>, collagen-like surface protein of <i>Streptococcus</i>
5.	0.044	0.045	0.158	A_44_P540755	<i>Olr1455</i> _predicted	Olfactory receptor 1455 (predicted)
6.	0.067	0.095	0.085	A_44_P402948	TC540893	Unknown
7.	0.086	0.056	0.157	A_44_P306008	AI179380	<i>Frame 12</i>, RSPCG42 3' end
8.	0.100	0.083	0.121	A_44_P531870	<i>Ephx2</i>	Epoxide hydrolase 2
9.	0.073	0.112	0.146	A_44_P138838	<i>Kb9</i>	Type II keratin Kb9
10.	0.112	0.125	0.193	A_44_P850331	BF554752	cDNA clone UI-R-C2p-ol-e-12-0-UI 5'
11.	0.150	0.108	0.179	A_44_P142391	LOC360919	Similar to α -fetoprotein
12.	0.158	0.163	0.127	A_44_P510436	<i>Myr8</i>	Myosin heavy chain
13.	0.122	0.142	0.223	A_43_P10290	<i>Tspan1</i>	Tetraspanin 1
14.	0.130	0.144	0.247	A_42_P719350	RGD1562974_predicted	Similar to DKFZp434P0316 (predicted)
15.	0.217	0.128	0.206	A_44_P900530	<i>Pcdh9</i> _predicted	Protocadherin 9 (predicted)
16.	0.198	0.135	0.249	A_44_P324806	AI103119	cDNA clone REMBX47 3' end
17.	0.223	0.206	0.212	A_44_P223009	CA506853	HIV-I <i>Nef</i> : negative effector of Fas and TNF
18.	0.205	0.223	0.221	A_44_P780781	TC564995	Unknown

Gene names in bold print were expressed at lower levels at either 6 or 9 weeks of age.

probability ratio <0.01 were 1,611, 1,260 and 1,361 in the SHR_s, SHRSP_s and M-SHRSP_s, respectively. The numbers of commonly down-regulated genes between the SHR_s and SHRSP_s, SHR_s and M-SHRSP_s, and SHRSP_s and M-SHRSP_s were 1,094, 1,021 and 851, respectively. In the SHR_s, SHRSP_s and M-SHRSP_s, 767 genes were commonly down-regulated. Of these genes, 6 were expressed less than 1/4 the levels noted in the WKY rats (Table V). *Sc1B*, *Hmmr* and *frame 12* were identified as known genes in addition to

three previously unidentified genes. In 9-week-old rats, the numbers of significantly down-regulated genes compared to those in the WKY rats at the level of a probability ratio <0.01 were 1,330, 1,465 and 176 in the SHR_s, SHRSP_s and M-SHRSP_s, respectively. The numbers of commonly down-regulated genes between the SHR_s and SHRSP_s, SHR_s and M-SHRSP_s, and SHRSP_s and M-SHRSP_s were 844, 121 and 125, respectively. In the SHR_s, SHRSP_s and M-SHRSP_s, 121 commonly down-regulated genes were identified. Of




Figure 1. Sample analysis using the Reactome database in Method 1 [more up-regulated genes in the spontaneously hypertensive rats (SHRs) substrains than in the Wistar Kyoto (WKY) rats]. Sample analysis of genes that were up-regulated >2 times in the malignant-type stroke-prone SHRs than in the WKY rats at (A) 6 weeks of age, n=193, and (B) 9 weeks of age, n=228.

these genes, 18 were expressed less than 1/4 the levels noted in the WKY rats (Table VI) and included *Anxa13*, *Sc1B*, *Olr1455*, *frame 12*, *Ephx2*, *Kb9*, *Myr8*, *Tspan1*, *Pcdh9* and *CA506853* (HIV-I *Nef* negative effector of Fas and TNF) as known genes in addition to 8 previously unidentified genes. A total of 5 genes were found to be commonly expressed at lower levels in SHR, SHRSP and M-SHRSP compared to WKY at 6 and 9 weeks of age (genes shown in bold in Table V) and included *Sc1B*, *Hmmr* and *frame 12*.

Up-regulated genes in the SHR substrains compared to the WKY rats were separately analyzed with the Reactome database

to determine the functional relationship in hypertension. An example of analysis of genes up-regulated more than two times in the M-SHRSPs compared to the WKY rats at 6 weeks of age (n=193) is shown in Fig. 1A. Analysis of genes up-regulated more than four times in the M-SHRSPs compared to the WKY rats at 9 weeks of age (n=228) is shown in Fig. 1B. A total of 4 genes, including *Yc2*, *Cyp2c*, *Gsta3* and *Cyp8b1*, participate in biological oxidation with a relatively strong relationship at the $P=1.4 \times 10^{-2}$ level at 6 weeks of age. A total of 7 genes, including *RGD1564999*, *Hmgcs2*, *Apob*, *Aptlc1*, *Acox2*, *Angptl4* and *Cyp8b1*, participate in the metabolism of lipids and lipoproteins with a relatively stronger relationship at the $P=4.1 \times 10^{-2}$ level at 9 weeks of age.

Method 2: Comparison of two different ages of each spontaneously hypertensive rat substrain. SBP was elevated in each SHR strain during the period from 6 to 9 weeks of age as shown in Table I. Therefore, candidate genes that were up- or down-regulated during this time may be related to hypertension. Multiple methods were used to survey candidate genes related to increased blood pressure during the ages of 6 to 9 weeks in each substrain. The numbers of genes that were up-regulated to a greater extent in rats at 9 compared to 6 weeks of age were 302, 680, 881 and 1,352 in the WKY rats, SHRs, SHRSPs and M-SHRSPs, respectively. Of these genes, 79, 124 and 7 genes were more highly up-regulated in the SHRs, SHRSPs and M-SHRSPs compared to the WKY rats, respectively. A total of 8 genes were up-regulated >1.5 times between rats 6 to 9 weeks of age in two or more substrains or in the M-SHRSPs (Table VII). The known genes were *Nef3*, *Slc26a4*, *Cyp2C*, *Gfral* and *Resp18*, and three previously unidentified genes were noted as well. The numbers of genes showing reduced expression in rats at 9 weeks of age compared to 6 weeks of age were 988, 285, 1,285 and 38 in the WKY rats, SHRs, SHRSPs and M-SHRSPs, respectively. Of these genes, 5, 51 and 0 genes showed reduced expression in the SHRs, SHRSPs and M-SHRSPs compared to the WKY rats, respectively. A total of 2 genes, *Atp12a* and *Hbb*, were

Table VII. Genes (8) whose expression was higher in the kidneys of the SHRs, SHRSPs and M-SHRSPs at 9 compared to 6 weeks of age and whose expression was higher in the SHR substrains than in the WKY rats.

	Clones	Fold increases (9/6 weeks of age)			GB account	Description
		SHR	SHRSP	M-SHRSP		
1.	A_42_P752336	5.281	8.974	10.278	<i>Nef3</i>	Neurofilament 3, medium
2.	A_44_P669819	2.513	2.886	5.548	TC543467	Unknown
3.	A_42_P800771		8.739	4.865	RGD1563825_predicted	Similar to ENSANGP00000020885, SOD
4.	A_44_P302221		5.804	5.057	LOC689753	Similar to K06A9.1b
5.	A_44_P1017035	1.475		2.020	<i>Slc26a4</i>	Solute carrier family 26, member 4, controls the balance of charged ions
6.	A_44_P280786	3.749	7.594		<i>Cyp2C</i>	Cytochrome P450, subfamily lic
7.	A_43_P11634			2.117	<i>Gfral</i>	Glial cell line derived neurotrophic factor family receptor α 1
8.	A_43_P12005			2.165	<i>Resp18</i>	Regulated endocrine-specific protein 18

Table VIII. Genes (2) whose expression was lower in the kidney of the SHR_s and SHRSP_s at 9 compared to 6 weeks of age and was lower compared to the WKY_s.

	Clones	Fold changes (9/6 weeks of age)		GB account	Description
		SHR	SHRSP		
1.	A_42_P684885	0.229	0.118	<i>Atp12a</i>	ATPase, H ⁺ /K ⁺ transporting, non-gastric, α polypeptide
2.	A_44_P306307	0.162	0.247	<i>Hbb</i>	Hemoglobin β chain complex




Figure 2. Sample analysis using the Reactome database in Method 2. Genes up-regulated >2 times (n=391) at 9 compared to 6 weeks of age in malignant-type stroke-prone spontaneously hypertensive rats.

expressed at less than 1/4 the levels at 6 compared to 9 weeks of age in more than two substrains (Table VIII).

Using the Reactome database as a representative sample, genes that were up-regulated more than two times (n=391) at 9 compared to 6 weeks of age in M-SHRSP were separately analyzed to determine the functional relationship in hypertension. Although three genes (*Slc28a1*, *Xdh* and *Gda*) were related to the metabolism of nucleotides at a $P=9.9 \times 10^{-2}$ level, no significant relationship with other processes was observed (Fig. 2).

Method 3: Comparison between genes expressed with and without hydralazine hydrochloride-induced hypotensive effects in the Wistar Kyoto rats and the spontaneously hypertensive rat substrains. BP was decreased in all rats after 30 mg/kg/day hydralazine hydrochloride treatment for 2 days. In this paradigm, candidate genes related to hydralazine hydrochloride-induced hypotension may be up- or down-regulated, particularly in the SHR substrains. Following administration of hydralazine hydrochloride, 22, 35, 26 and 66 genes in the WKY_s, SHR_s, SHRSP_s and M-SHRSP_s were up-regulated 4, 2, 2 and 8 times at 6 weeks of age, respectively. The numbers of genes up-regulated >1.0-fold in the WKY rats and M-SHRSP_s, SHR_s and SHRSP_s, SHR_s and M-SHRSP_s, and SHRSP_s and M-SHRSP_s at 6 weeks of age were 67, 5, 9 and 23, respectively. Following administration

of hydralazine hydrochloride, 9, 34, 60 and 4 genes in the WKY rats, SHR_s, SHRSP_s and M-SHRSP_s were up-regulated 1.2, 1.2, 1.5 and 1.2 times at 9 weeks of age, respectively. The numbers of genes up-regulated >1.0-fold in the WKY rats and M-SHRSP_s, SHR_s and SHRSP_s, SHR_s and M-SHRSP_s, and SHRSP_s and M-SHRSP_s at 9 weeks of age were 67, 5, 9 and 23, respectively. On the other hand, following administration of hydralazine hydrochloride, 13, 20, 11 and 18 genes showed expression that was reduced 0.6, 0.2, 0.6 and 0.25 times in the WKY rats, SHR_s, SHRSP_s and M-SHRSP_s at 6 weeks of age, respectively. The numbers of genes showing expression that was reduced by <1.0-fold in the WKY rats and M-SHRSP_s, SHR_s and SHRSP_s, SHR_s and M-SHRSP_s, and SHRSP_s and M-SHRSP_s at 6 weeks of age were 3, 0, 21 and 3, respectively. In addition, the numbers of genes showing expression that was reduced <0.8 times following administration of hydralazine hydrochloride were 11, 7, 7 and 41 in the WKY rats, SHR_s, SHRSP_s and M-SHRSP_s, respectively, at 9 weeks of age. The numbers of genes showing expression that was reduced <1.0-fold in the WKY rats and M-SHRSP_s, SHR_s and SHRSP_s, SHR_s and M-SHRSP_s, and SHRSP_s and M-SHRSP_s at 9 weeks of age were 3, 0, 21 and 3, respectively.

Using this method, few genes were identified that were both up- and down-regulated and commonly expressed between substrains. Thus, the analyses were modified as follows. Genes identified with Method 3 and those identified with Method 2 were combined and considered candidate hypertension-related genes. Using this method, genes that were up-regulated >1.2 times or down-regulated <0.8 times in two or more substrains were focused on. After identifying the genes that were expressed in the WKY rats and satisfying the criteria of Method 2, ten genes satisfied both conditions at once. These 10 genes are strongly suggested to be candidate genes (Table IXA), and they included *TC550463* (farnesyl pyrophosphate synthetase), *Kcnc3*, *Vnn1* and *RGD1561143* (similar to cell surface receptor FDFACT) in addition to 6 previously unidentified genes. In addition, 4 genes that were up-regulated >4 times following administration of hydralazine hydrochloride for 2 days in 6-week-old M-SHRSP and that satisfied the criteria of Method 2 were identified as possible candidate genes (Table IXB). These genes included *TC560558* (FK506-binding protein 1B), *TC564079* (*Drosophila melanogaster*), *XM_343516* (similar to sulfotransferase K2) and one previously unidentified gene.

When genes that were up-regulated 1.5 times (n=505) following hydralazine hydrochloride administration in 6-week-old SHRSP_s were analyzed further with the Reactome

Table IX. Genes (14) that were up- or down-regulated following administration of hydralazine, and/or were commonly more highly expressed in the kidneys of the SHR, SHRSP or M-SHRSPs compared to the WKY rats at 9 compared to 6 weeks of age.

A, Genes strongly suggested to be associated with hypertension

	Fold changes (hydralazine/none)	Strains	Clones	GB account	Description
1. ^a	9.646	6W M-SHRSP	A_44_P566390	TC550463	Farnesyl pyrophosphate synthetase
2. ^a	9.551	6W M-SHRSP	A_44_P623610	TC525804	Unknown
3. ^a	8.615	6W M-SHRSP	A_43_P10474	TC527985	Unknown
4. ^a	8.419	6W M-SHRSP	A_44_P808679	TC541828	Unknown
5. ^a	8.387	6W M-SHRSP	A_44_P104687	<i>Kcnc3</i>	Potassium voltage-gate channel protein
6. ^a	2.044	6W SHRSP	A_44_P871211	TC566645	Unknown
7. ^a	0.664	9W SHR	A_44_P732488	TC567669	Unknown
8.	2.367 2.529	6W SHR 6W SHRSP	A_42_P811256	<i>Vnn1</i>	Participate in an oxidative-stress response
9.	2.205 8.872	6W SHRSP 6W M-SHRSP	A_44_P854454	TC543180	Unknown
10.	1.260 2.115	9W SHR 6W SHRSP	A_44_P435955	RGD1561143_predicted	Similar to cell surface receptor FDFACT

B, Genes possibly suggested to be associated with hypertension

1. ^b	9.245	6M M-SHRSP	A_44_P610417	TC560558	FK506-binding protein 1B
2. ^c	8.997	6W M-SHRSP	A_44_P841829	TC564079	CG4187-PA, <i>Drosophila melanogaster</i>
3. ^b	8.969	6W M-SHRSP	A_44_P917243	TC533180	Unknown
4. ^b	8.533	6W M-SHRSP	A_44_P408114	XM_343516	Similar to sulfotransferase K2

^aMore highly expressed at 9 than at 6 weeks of age in either SHR or SHRSP; ^bMore highly expressed at 9 than at 6 weeks of age in SHRSP;

^cMore highly expressed at 9 than at 6 weeks of age in SHR.

Table X. Genes (3) down-regulated following administration of hydralazine in the SHR and M-SHRSPs at 6 weeks of age.

	Fold changes (hydralazine/none)	Strains	Clones	GB account	Description
1.	0.090 0.168	6W SHR 6W M-SHRSP	A_44_P541438	RGD1308699	Similar to 1700060H10Rik protein
2.	0.132 0.129	6W SHR 6W M-SHRSP	A_43_P15862	<i>Gabrq</i>	γ -aminobutyric acid A receptor
3.	0.113 0.204	6W SHR 6W M-SHRSP	A_44_P281788	BQ196033	cDNA clone UI-R-CN1-cmn-f-01-0-UI 3'

database, 5 genes ($P=2.7 \times 10^{-2}$) involved in DNA replication, 14 genes ($P=9.7 \times 10^{-2}$) involved in diabetes pathways, 5 genes ($P=5.9 \times 10^{-3}$) involved in APC/C:Cdh1-mediated degradation of Skp2, 5 genes ($P=8.3 \times 10^{-3}$) involved in Cdc20: phospho-APC/C-mediated degradation of cyclin A, 2 genes ($P=9.2 \times 10^{-3}$) involved in the inhibition of hormone sensitive

lipase (HSL), 3 genes ($P=6.3 \times 10^{-2}$) involved in the metabolism of vitamins and cofactors, and 4 genes ($P=2.1 \times 10^{-2}$) involved in Wnt signaling were detected with high significance (Fig. 3). Thus, expression of numerous genes related to DNA replication and cell proliferation, including *Psmc6*, *Psm2*, *Psm6* and *LOC311078* [proteasome (prosome, macropain) subunits],




Figure 3. Sample analysis using the Reactome database in Method 3. Genes up-regulated >1.5 times (n=505) following treatment with 30 mg/kg/day hydralazine hydrochloride for 2 days in 6-week-old stroke-prone spontaneously hypertensive rats.

was modified by hydralazine hydrochloride treatment of the SHRSPs with high significance.

As shown in Table X, the expression of three possibly hypertension-related genes was found to be down-regulated 0.25 times by hydralazine hydrochloride administration for 2 days in either 6-week-old SHR or M-SHRSP, including *Gabrq* (γ -aminobutyric acid A receptor) and two previously unidentified genes.

Discussion

SHRs (10) were created at the Kyoto University School of Medicine, Japan, around 1963 through continuous brother-sister mating of six generations of normotensive WKY rats, and they have a slightly higher BP. Using this strain with high blood pressure, numerous studies concerning essential hypertension, including pathophysiology and the effects of food, have been carried out (29,30). Descendants of SHRs, SHRSPs (11) were created through continuous brother-sister mating in a closed colony of SHRs after approximately one decade of mating. M-SHRSPs (12) were established in 1985 through the brother-sister mating of the SHRSP strain under hypotensive treatment with hydralazine hydrochloride to avoid stroke during mating and litter care. An inbred strain of M-SHRSP shows SBPs of 250 mmHg or higher before 14 weeks of age, and shows more rapid and severe increases in BP and stroke in almost all animals (31). These three types of hypertensive rat substrains, SHR, SHRSP and M-SHRSP, have been used in multiple investigations of pathophysiology, preventive medicine, pharmacology and drug development worldwide (29-32). Our facility is one of the original places that bred the three types of SHR substrains and families. For our present study, we decided to use DNA array methodology and these three hypertensive substrains of rats and the normotensive WKY to identify genes related to hypertension and, when possible, to identify candidate genes that cause hypertension.

For theoretical reasons, we should have investigated hereditary subjects using congenic or consomic animal models of disease (33,34). We considered the three substrains

of hypertensive rats, SHR, SHRSP and M-SHRSP, to belong to one family, although several decades have passed since the inbred strains were established. SHRs are derived from WKY rats, SHRSPs are derived from SHRs and M-SHRSPs are derived from SHRSPs on the basis of higher blood pressure and/or high incidences of stroke as a selection criterion. M-SHRSPs show a higher and earlier elevation of blood pressure that is accompanied by a higher and earlier incidence of stroke compared to SHRSPs. This is also true for SHRSPs compared to SHRs. Therefore, candidate genes for hypertension must be condensed, increased or expressed at higher/lower levels to a greater extent in M-SHRSP than in SHRSP and SHR substrains. On this basis, we searched for candidate genes for hypertension with these three hypertensive SHR substrains compared to normotensive WKY rats.

The kidneys were thought to be the most appropriate tissue for studying hypertension due to their direct influence on body fluids and endocrine, cardiovascular and sympathetic functions (14). There are numerous intrinsic and extrinsic factors, including the renin-angiotensin system and catecholamine and aldosterone hormones, that control the relationship between kidney function and blood pressure (14,15). This study is the first attempt to use DNA microarrays to compare the gene expression profiles of the kidneys of SHRs, SHRSPs and M-SHRSPs employing WKY rats as a control. In addition to analyzing the genes expressed in these three types of SHR substrains (Method 1), young rats (6 weeks old), whose blood pressure was not yet elevated, and slightly older rats (9 weeks old) that had developed hypertension were employed to detect candidate genes related to blood pressure elevation (Method 2). Furthermore, a hypotensive drug, hydralazine hydrochloride, which acts through an unknown receptor, was administered to each group of rats to induce acute hypotension to detect hypertension-associated genes (Method 3). This study aimed to identify candidate genes for hypertension in the kidneys of the SHR substrains using these three analytical methods.

BP was elevated in 9-week-old SHR substrains compared to 6-week-old rats, and tended to increase from WKY to SHR, SHRSP and M-SHRSP in this order at 6 and 9 weeks of age. Hydralazine hydrochloride administration for 2 days decreased blood pressure in all the rats (Table I). Therefore, Methods 1, 2 and 3 may be appropriate for surveying candidate genes related to BP.

In the Method 1 analysis at 6 weeks of age, 16 genes were significantly more highly expressed in SHR, SHRSP and M-SHRSP (Table II). Of these genes, *Sugt 1* (kinetochore function), *Dusp15* (dual specificity phosphatase-like 15), *Armc 3* (multiple functions in signal transduction) and *Serpina3m* (serine or cysteine protease inhibitor, member 3M) were related to cell proliferation, protein modification or signal transduction. At 9 weeks of age, *Cyp8b1* (cytochrome P450, family 8, subfamily b, polypeptide 1), *Zfp597* (zinc finger protein 597), *Gtphp4* (GTP binding protein 4), *Tmem14a* (transmembrane protein 14A), *TC540923* (phosphatidylinositol 3 kinase regulator), *Sv2a* (synaptic vesicle glycoprotein 2a), *Mapk14* (mitogen activated protein kinase 14) and *Prkar2b* (protein kinase, cAMP dependent regulatory type II β) were involved in cell proliferation, protein modification or signal transduction (Table III). At 6 and 9 weeks of age, *Gc* (group specific

component), *Dusp15*, *TC558814* (unknown), *Sugt 1*, *Cyp8b1*, *Armc 3*, *Serpina3m* and *Bri3bp* (Bri3 binding protein) were significantly more highly expressed (Table IV). A number of genes were significantly expressed at lower levels and included *Sc1B* (collagen-like surface protein of *Streptococcus*), *Hmmr* (hyaluronan mediated motility receptor) and *frame 12* (RSPCG42 3' end) at 6 weeks of age (Table V), and *Anxa13* (annexin A13), *Ephx2* (epoxide hydrolase 2), *Kb9* (type II keratin Kb9), *Myr8* (myosin heavy chain), *Tspan1* (tetraspanin 1) and *Pcdh9* (protocadherin 9) at 9 weeks of age (Table VI). A number of previously unidentified genes were also found.

The genes that were identified by Method 1 as being more highly expressed were investigated using the Reactome database to determine how and where they work in biochemical processes. As shown in Fig. 1A and B, these genes were highly related to biological oxidation at 6 weeks of age and metabolism of lipids and lipoproteins at 9 weeks of age. However, these genes were not involved in signal transduction or muscle function in M-SHRSPs compared to WKY rats. Therefore, the data derived from the Method 1 analysis may show differences in metabolic characteristics mainly between SHR substrains and WKY rats, although further study is required.

In the Method 2 analysis, which compared two different ages of each SHR substrain, *Nef3* (neurofilament 3), *Slc26a4* (solute carrier family 26, member 4, controls the balance of ions), *Cyp2C* (cytochrome P450, subfamily 1C), *Gfral* (glial cell line-derived neurotrophic factor family receptor α 1) and *Resp18* (regulated endocrine-specific protein 18) were identified as known genes that were up-regulated in addition to three previously unidentified genes (Table VII). In addition, *Atp12a* (ATPase, H⁺/K⁺ transporting, nongastric, α polypeptide) and *Hbb* (hemoglobin β chain complex) were identified as genes that were down-regulated with age in the SHR substrains (Table VIII). Functional relationship analysis with the Reactome database of genes from 6-week-old M-SHRSP showed three genes involved in the metabolism of nucleotides, including *Slc28a1*, *Xdh* and *Gda*. Therefore, the genes identified with this method, including *Slc26a4*, *Cyp2C*, *Gfral*, *Resp18* and *Atp12a*, may be closely related to hypertension as they are related to energy production and consumption or ion exchange. Further investigation is required to determine the importance of these identified genes.

From analysis with Method 3 which compared genes expressed with or without hydralazine hydrochloride-induced hypotensive effects in the SHR substrains to WKY rats, *Vnn1* (participates in the oxidative-stress response) was identified as a gene that was more highly expressed, and *Gabrq* (γ -aminobutyric acid A receptor) was identified as a suppressed gene. These two genes were commonly expressed in more than two SHR substrains. Numerous other genes were uniquely expressed in one SHR substrain, but not expressed commonly among the three substrains. Therefore, other candidate genes were identified by combining Methods 2 and 3. From this process, *TC550463* (farnesyl pyrophosphate synthetase), *Kcnc3* (potassium voltage-gate channel protein), *TC560558* (FK506-binding protein 1B) and *XM_343516* (sulfotransferase K2) were selected in addition to a number of previously unidentified genes. When genes that were

up-regulated more than 1.5 times (n=505) by hydralazine hydrochloride administration in the 6-week-old SHRSP were analyzed with the Reactome database software, five genes involved in DNA replication, five genes involved in the APC/C:Cdh1-mediated degradation of Skp2, five genes involved in the Cdc20:phospho-APC/C-mediated degradation of cyclin A, two genes involved in the inhibition of HSL and four genes involved in Wnt signaling were detected with high significances (Fig. 3). A number of genes related to DNA replication and cell proliferation, including *Psmc6*, *Psmc2*, *Psmc6* and *LOC311078* [proteasome (prosome, macropain) subunits], were identified in 6-week-old SHRSP. Therefore, genes identified with Methods 2 and 3 may be the candidate genes closely related to hypertension for which we are searching.

Although current gene expression arrays permit the simultaneous analysis of thousands of rat genes, this method is not yet capable of addressing all functional genes in the genome. However, as rat genome annotation progresses and arrays continue to improve in their extent of genomic coverage, a more complete analysis will be possible. Our present approach identified dozens of genes, including *Dusp15*, *Cyp8b1*, *Armc 3*, *Gtpbp4*, *Mettl2*, *Mapk14*, *Prkar2b*, *frame 12*, *Anxa13*, *Ephx2*, *Myr8* and *Pcdh9* from Method 1; *Cyp2C* and *Atp12a* from Method 2; and *Kcnc3*, *Vnn1*, *TC560558* and *Gabrq* from Methods 2 and 3, in addition to a number of previously unidentified genes, as probable candidate genes that cause hypertension in SHR substrains, as determined by common biochemical knowledge. Of these genes, only *Ephx2* has been previously reported as a strongly related gene in SHRs (24). A key question has arisen regarding *Ephx2*. *Ephx2* was reported to be a significantly up-regulated gene in SHRs by 3.39- and 4.30-fold at 3 and 9 weeks of age, respectively, compared to WKY rats (24). However, we identified *Ephx2* as a significantly down-regulated gene at 9 weeks of age (but not at 6) in the SHRs, SHRSPs and M-SHRSPs by 0.100-, 0.083- and 0.121-fold, respectively (Table VI). Fornage *et al* reported lower expression of this gene in the kidney of 4- to 5-week-old SHRs compared to WKY rats (35), and Corenblum *et al* observed lower expression of *Ephx2* in the brain of SHRSPs compared to stroke-resistant SHRs (36). Although a number of studies have reported that soluble epoxide hydrolase elevates blood pressure by degrading vasodilative epoxyeicosatrienoic acids by means of an inhibitor (37-39), the role of *Ephx2* in hypertension remains controversial. Considering our data and several other reports, *Ephx2* may not be a candidate gene for hypertension as the expression of *Ephx2* was elevated or decreased in some SHR substrains at some ages and, thus, results were not consistent. *Ephx2* possibly controls the adaptation of BP changes.

Why the surveyed genes related to hypertension vary among reports has yet to be elucidated. In addition, we believe that discovering the roles of unknown genes is crucial, as genes that are strongly related to hypertension may exist among these candidates, and these genes may be related to the cause of stroke in SHRSPs and M-SHRSPs. Since the majority of these genes have not yet been demonstrated to be responsible for hypertension in SHRs, we must continue to search for true candidate genes that participate in the genesis of hypertension in SHR substrains using current technology.

Acknowledgements

This study was supported by a Japanese Grant-in-Aid for Scientific Research (no. 19500699), an Aid Grant to private universities to cover current expenses from the MEXT in Japan, the Fund of Mishima-Kaiun Memorial Foundation, a donation from Ueshima Coffee Co., Ltd., and the Kinki University School of Medicine. We thank the National Center for Biotechnology Information, USA, and the DNA Data Bank of Japan for access to the network servers.

References

1. Acelajado MC and Oparil S: Hypertension in the elderly. *Clin Geriatr Med* 25: 391-412, 2009.
2. Ghiadoni L, Bruno RM, Stea F, Virdis A and Taddei S: Central blood pressure, arterial stiffness, and wave reflection: new targets of treatment in essential hypertension. *Curr Hypertens Rep* 11: 190-196, 2009.
3. Kannel WB: Table of contents. Hypertension: reflections on risks and prognostication. *Med Clin North Am* 93: 541-558, 2009.
4. Imaizumi Y: Mortality in the elderly population aged over 40 in Japan, 1947-1988. *Jinko Mondai Kenkyu* (in Japanese) 47: 40-57, 1991.
5. Kesteloot H, Yuan XY and Joossens JV: Changing mortality patterns in men. *Acta Cardiol* 43: 133-139, 1988.
6. Drmanac R, Drmanac S, Strezoska Z, *et al*: DNA sequence determination by hybridization: a strategy for efficient large-scale sequencing. *Science* 260: 1649-1652, 1993.
7. Schena M, Heller RA, Thieriault TP, Konrad K, Lachenmeier E and Davis RW: Microarrays: biotechnology's discovery platform for functional genomics. *Trends Biotechnol* 16: 301-306, 1998.
8. Hacia JG, Brody LC and Collins FS: Applications of DNA chips for genomic analysis. *Mol Psychiatry* 3: 483-492, 1998.
9. McBride MW, Charchar FJ, Graham D, *et al*: Functional genomics in rodent models of hypertension. *J Physiol* 554: 56-63, 2004.
10. Okamoto K and Aoki K: Development of a strain of spontaneously hypertensive rat. *Jpn Circ J* 27: 282-293, 1963.
11. Okamoto K, Yamori Y and Nagaoka A: Establishment of the stroke-prone spontaneously hypertensive rats (SHR). *Circ Res* 34 (Suppl 1): 143-153, 1974.
12. Okamoto K, Yamamoto K, Morita N and Ohta Y: Establishment and characteristics of rat with precocious and severe hypertension (M-SHRSP). *Acta Med Kinki Univ* 10: 73-95, 1985.
13. Ashenagar MS, Tabuchi M, Kinoshita K, *et al*: Gene expression in the adrenal glands of three spontaneously hypertensive rat substrains. *Mol Med Rep* 3: 213-222, 2010.
14. Guyton AC: Dominant role of the kidneys and accessory role of whole-body autoregulation in the pathogenesis of hypertension. *Am J Hypertens* 2: 575-585, 1989.
15. Guyton AC: Abnormal renal function and autoregulation in essential hypertension. *Hypertension* 18 (Suppl 5): 49-53, 1991.
16. Gavras H, Brunner HR and Laragh JH: Renin and aldosterone and the pathogenesis of hypertensive vascular damage. *Prog Cardiovasc Dis* 17: 39-49, 1974.
17. Satoh T, Owada S and Ishida M: Recent aspects in the genetic renal mechanisms involved in hypertension. *Intern Med* 38: 919-926, 1999.
18. Ely D, Milsted A, Dunphy G, *et al*: Delivery of sry1, but not sry2, to the kidney increases blood pressure and SNS indices in normotensive WKY rats. *BMC Physiol* 9: 10, 2009.
19. Graham D, McBride MW, Gaasenbeek M, *et al*: Candidate genes that determine response to salt in the stroke-prone spontaneously hypertensive rat: congenic analysis. *Hypertension* 50: 1134-1141, 2007.
20. Kreutz R, Schulz A, Sietmann A, Stoll M, Daha MR, de Heer E and Wehland M: Induction of C1q expression in glomerular endothelium in a rat model with arterial hypertension and albuminuria. *J Hypertens* 25: 2308-2316, 2007.
21. Clemitson JR, Dixon RJ, Haines S, *et al*: Genetic dissection of a blood pressure quantitative trait locus on rat chromosome 1 and gene expression analysis identifies SPON1 as a novel candidate hypertension gene. *Circ Res* 100: 992-999, 2007.
22. McBride MW, Brosnan MJ, Mathers J, *et al*: Reduction of Gstm1 expression in the stroke-prone spontaneously hypertensive rat contributes to increased oxidative stress. *Hypertension* 45: 786-792, 2005.
23. Zhong JC, Huang DY, Yang YM, Li YF, Liu GF, Song XH and Du K: Upregulation of angiotensin-converting enzyme 2 by all-trans retinoic acid in spontaneously hypertensive rats. *Hypertension* 44: 907-912, 2004.
24. Kurdi M, Cerutti C, Randon J, McGregor L and Bricca G: Macroarray analysis in the hypertrophic left ventricle of renin-dependent hypertensive rats: identification of target genes for renin. *Renin Angiotensin Aldosterone Syst* 5: 72-78, 2004.
25. Seubert JM, Xu F, Graves JP, *et al*: Differential renal gene expression in prehypertensive and hypertensive spontaneously hypertensive rats. *Am J Renal Physiol* 289: F552-F561, 2005.
26. Charchar FJ, Kaiser M, Bingham AJ, Fotinatos N, Ahmady F, Tomaszewski M and Samani NJ: Whole genome survey of copy number variation in the spontaneously hypertensive rat: relationship to quantitative trait loci, gene expression, and blood pressure. *Hypertension* 55: 1231-1238, 2010.
27. Simon R, Lam A, Li M-C, *et al*: Analysis of gene expression data using BRB-array tools. *Cancer Informatics* 3: 11-17, 2007.
28. Altschul SF, Thomas LM, Alejandro AS, *et al*: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389-3402, 1997.
29. Toal C and Leenen FH: Dietary sodium restriction and development of hypertension in spontaneously hypertensive rats. *Am J Physiol* 245: H1081-H1084, 1983.
30. Wexler BC: Inhibition of the pathogenesis of spontaneous hypertension in spontaneously hypertensive rats by feeding a high fat diet. *Endocrinology* 108: 981-989, 1981.
31. Okamoto K, Ohta Y, Chikugo T, Shiokawa H and Morita N: Chronic treatment with captopril, SQ29,852, hydralazine and a 33% fish meal diet in malignant stroke-prone spontaneously hypertensive rats. *J Hypertens* 9: 1105-1117, 1991.
32. Yamori Y: Implication of hypertensive rat models for primordial nutritional prevention of cardiovascular diseases. *Clin Exp Pharmacol Physiol* 26: 568-572, 1999.
33. Pravenec M and Kurtz TW: Recent advances in genetics of the spontaneously hypertensive rat. *Curr Hypertens Rep* 12: 5-9, 2010.
34. Cowley AW Jr, Roman RJ and Jacob HJ: Application of chromosomal substitution techniques in gene-function discovery. *J Physiol* 554: 46-55, 2004.
35. Fornage M, Hinojos CA, Nurowska BW, Boerwinkle E, Hammock BD, Morisseau CHP and Doris PA: Polymorphism in soluble epoxide hydrolase and blood pressure in spontaneously hypertensive rats. *Hypertension* 40: 485-490, 2002.
36. Corenblum MJ, Wise VE, Georgi K, Hammock BD, Doris PA and Fornage M: Altered soluble epoxide hydrolase gene expression and function and vascular disease risk in the stroke-prone spontaneously hypertensive rat. *Hypertension* 51: 567-573, 2008.
37. Chiamvimonvat N, Ho CM, Tsai HJ and Hammock BD: The soluble epoxide hydrolase as a pharmaceutical target for hypertension. *J Cardiovasc Pharmacol* 50: 225-237, 2007.
38. Li J, Carroll MA, Chander PN, Falck JR, Sangras B and Stier CT: Soluble epoxide hydrolase inhibitor, AUDA, prevents early salt-sensitive hypertension. *Front Biosci* 13: 3480-3487, 2008.
39. Yousif MH and Benter IF: Role of 20-hydroxyeicosatetraenoic and epoxyeicosatrienoic acids in the regulation of vascular function in a model of hypertension and endothelial dysfunction. *Pharmacology* 86: 149-156, 2010.