

# Polymorphisms of *IGF1* contribute to the development of ischemic stroke

HAK JAE KIM<sup>1\*</sup>, SU KANG KIM<sup>2\*</sup>, HAE JEONG PARK<sup>2</sup>, JOO-HO CHUNG<sup>2</sup>,  
JINMAN CHUN<sup>3</sup>, DONG HWAN YUN<sup>3</sup> and YOUNG OCK KIM<sup>4</sup>

<sup>1</sup>Soonchunhyang Medical Research Institute, College of Medicine, Soonchunhyang University, Chunan;  
Departments of <sup>2</sup>Pharmacology and Kohwang Medical Research Institute, and <sup>3</sup>Physical Medicine  
and Rehabilitation, School of Medicine, Kyung Hee University, Seoul; <sup>4</sup>Development of Ginseng and  
Medical Plants Research Institute, Rural Administration, Eumseong, Republic of Korea

Received August 1, 2011; Accepted September 1, 2011

DOI: 10.3892/etm.2011.372

**Abstract.** Insulin-like growth factor 1 (IGF1) is neuroprotective in animal models of focal brain ischemia and correlates with ischemic stroke (IS) outcome in the elderly. In this study, we investigated whether single nucleotide polymorphisms (SNPs) of the *IGF1* gene are associated with the development and clinical features of IS in a Korean population. A total of 119 patients with IS and 289 control subjects were recruited. Stroke patients were classified into subgroups according to the scores of the National Institutes of Health Stroke Survey (NIHSS; <6 and ≥6) and the Modified Barthel Index (MBI; <60 and ≥60). Among the SNPs of the *IGF1* gene, five SNPs were selected and analyzed by direct sequencing: rs2162679 (intron), rs2195239 (intron), rs978458 (intron), rs1520220 (intron) and rs6214 (3' untranslated region; 3'UTR). Multiple logistic regression models were conducted to analyze genetic data. SNPStats, SNPAnalyzer Pro and HelixTree programs were used to calculate odds ratios (ORs), 95% confidence intervals (CIs) and p-values. Two SNPs, rs2162679 and rs6214, were associated with the development of IS. After Bonferroni correction (p<sup>c</sup>), the A and G alleles of rs2162679 and rs6214 had significant differences between patients with IS and the controls [rs2162679, OR (95% CI) = 1.64 (1.17-2.31), p=0.004, p<sup>c</sup>=0.02; rs6214, OR (95% CI) = 1.52 (1.12-2.07), p=0.007, p<sup>c</sup>=0.035], respectively. However, the five selected SNPs were not related to the NIHSS and MBI scores. These results suggest that *IGF1* may be associated with the development of IS.

## Introduction

Stroke is a neurological disease which causes long-term disability and is the third leading cause of death in the United States. Ischemic stroke (IS) accounts for approximately 85% of all strokes. The other 9% are caused by intracerebral hemorrhage and 4-5% by subarachnoid hemorrhage (1,2). Environmental and genetic factors are related to the pathogenesis of IS. Several lines of genetic studies have reported the relationship between IS and single nucleotide polymorphisms (SNPs) of candidate genes, such as the transforming growth factor β1 (TGFB1), transforming growth factor β receptor II (70/80 kDa) (TGFB2) and tumor necrosis factor (TNF) (3-5).

Insulin-like growth factor 1 (somatomedin C) (IGF1) is similar to insulin in function and plays a crucial role in mammalian growth and development. The IGF1 level declines during the normal aging process, and a low IGF1 level correlates with decreased cognitive abilities (6). *In vitro* studies have shown that IGF1 reduces neuronal cell death in various injury insults (7,8) and IGF1 has a protective effect in ischemic animal models (9,10). Several reports have reviewed the roles of IGF1 in stroke severity and outcome (11,12). They have suggested that IGF levels may be associated with neurological recovery and functional outcome, and have also proposed IGF1 as a predictor of stroke outcome.

In this study, we investigated the association between *IGF1* SNPs and IS in a Korean population. We also assessed the relationship between *IGF1* SNPs and the clinical phenotypes according to the scores of National Institutes of Health Stroke Survey (NIHSS) and the Modified Barthel Index (MBI).

## Materials and methods

**Study population and clinical phenotypes.** IS patients were enrolled among participants visiting the Departments of Neurosurgery and Physical Medicine and Rehabilitation, Kyung Hee Medical Center (Seoul, Republic of Korea). Patients with transient ischemic attack, cerebrovascular malformation, congenital brain disorders and accidental or iatrogenic stroke, were excluded. Stroke patients were diagnosed by computed tomography, magnetic resonance imaging, angiography and

---

**Correspondence to:** Dr Young-Ock Kim, Ginseng and Medical Plants Research Institute Rural Development Administration, Chunbuk 369-873, Republic of Korea  
E-mail: kyo9128@korea.kr

\*Contributed equally

**Key words:** insulin-like growth factor 1, ischemic stroke, Modified Barthel Index, National Institutes of Health Stroke Survey, single nucleotide polymorphism



Figure 1. Gene map and single nucleotide polymorphisms in the *IGF1* gene on chromosome 12q23.2. Exons are marked with a box. The coding regions are represented by black boxes and untranslated regions by white boxes.

duplex sonography. The control subjects were recruited among healthy volunteers to examine the general health check-up program. Patients with neurological diseases, ischemic heart diseases and other severe diseases, were excluded. All stroke patients were classified into clinical phenotypes according to the NIHSS and MBI scores. For the neurological functional level of IS patient, the severity of 13 neurological symptoms was assessed by the NIHSS score. For the daily living activity of IS patients, the quality of 10 general life activities was evaluated by the MBI score. This study was approved by the Ethics Review Committee of the Medical Research Institute, School of Medicine, Kyung Hee University. Written informed consent was obtained from all patients. If patients were incommunicative, it was obtained from a guardian or close relatives.

**SNP selection and genotyping.** We searched the coding SNPs (cSNPs) of the *IGF1* gene in the SNP database of the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/SNP>, BUILD 132). The cSNPs with a heterozygosity of <0.05 and/or a minor allele frequency (MAF) of <0.05, were excluded. Out of five missense and two synonymous SNPs, there were no cSNPs with a heterozygosity of >0.05 or a MAF of >0.05. Therefore, we searched the untranslated and intron SNPs of the *IGF1* gene and previous studies (13-15). Finally, five SNPs [rs2162679 (intron), rs2195239 (intron), rs978458 (intron), rs1520220 (intron) and rs6214 (3' untranslated region; 3'UTR)] were selected (Fig. 1). Genotypes of the five selected SNPs were analyzed by direct sequencing (Macrogen, Seoul, Republic of Korea). Polymerase chain reactions (PCRs) were conducted using the forward and reverse primers of each SNP (Table II). PCRs were performed under the following conditions: 35 cycles at 95°C for 30 sec, 58°C for 30 sec, 72°C for 30 sec, and 1 cycle at 72°C for 5 min for the final reaction. The PCR products were sequenced by an ABI PRISM 3730XL analyzer (PE Applied Biosystems, Foster City, CA, USA) and genotypes of each SNP were analyzed by SeqManII software (DNASTAR, Madison, WI, USA).

**Statistical analysis.** SNPStats (<http://bioinfo.iconcologia.net/index.php?module=Snpstats>), SNPAnalyzer Pro (ISTECH, Goyang, Republic of Korea), and Helixtree (Golden Helix, Bozeman, MT, USA) were used to obtain odds ratios (ORs), 95% confidence intervals (CIs) and p-values. Hardy-Weinberg equilibrium (HWE) was calculated using the Chi-square test. Multiple logistic regression models were conducted using the following models: codominant1 (major allele homozygotes vs. heterozygotes), codominant2 (major allele homozygotes

Table I. Clinical characteristics in stroke patients and control subjects.

	IS	Control
Total no.	119	289
Male/female (n)	66/53	150/139
Age (mean $\pm$ SD, years)	65.8 $\pm$ 12.1	62.9 $\pm$ 9.3
NIHSS (score)		
<6	55	
$\geq$ 6	57	
MBI (score)		
<60	71	
$\geq$ 60	24	

IS, ischemic stroke; SD, standard deviation; NIHSS, National Institutes of Health Stroke Survey; MBI, Modified Barthel Index. Stroke patients with inappropriate clinical data were excluded.

vs. minor allele homozygotes), dominant (major allele homozygotes vs. heterozygotes and minor allele homozygotes), recessive (major allele homozygotes and heterozygotes vs. minor allele homozygotes) and log-additive (major allele homozygotes vs. heterozygotes vs. minor allele homozygotes) (16,17). Bonferroni correction was performed to obtain further statistical significance. Linkage disequilibrium (LD) blocks were estimated using Haploview version 4.2 (Daly Lab, Cambridge, MA, USA). The required case size in each SNP was estimated using the genetic power calculator (<http://pngu.mgh.harvard.edu/~purcell/gpc/cc2.html>) to obtain the statistical power. The statistical significant level was set at a value of  $p < 0.05$ .

## Results

**Demographic and clinical features of study subjects.** Table I shows the demographic and clinical features of the IS patients and the control subjects. The age of the IS patients and control subjects was 65.8 $\pm$ 12.1 (mean  $\pm$  SD) and 62.9 $\pm$ 9.3 years, respectively. The number of IS patients and control subjects was 119 (66 male/53 female) and 289 (150 male/139 female), respectively. All IS patients were divided into two clinical subgroups according to the NIHSS (<6 and  $\geq$ 6) and MBI scores (<60 and  $\geq$ 60). The numbers of IS patients with a NIHSS score of <6 and  $\geq$ 6 were 55 (49.1%) and 57 (50.9%), respectively. The

Table II. Primer sequences for each single nucleotide polymorphism (SNP).

SNP	Forward (5'-3')	Reverse (5'-3')	Size (bp)
rs2162679	TCTGCAAGATCAATCACAGGTT	AAAAACCAAAACCCCTGTCTCT	386
rs2195239	CAAATTTTCAGCTGCGACTTTATC	GAGCCAAAACCATCTCTACACC	306
rs978458	TCCACTAGAGCCAAAGAAGAGC	GTGAAATGGTGGAGGATGATTT	381
rs1520220	AAAGGATCTAGAGGCCAGAAGG	AGTTCTTGTTTCCTGCACTCCC	390
rs6214	CCAGACATACAGGTTCTGTGGA	TTGGAGAGGATTATGTGTTGGA	304

Table III. Genotype and allele frequencies of *IGF* single nucleotide polymorphisms in controls and IS patients.

SNP	Type	Control		IS		Model	OR (95% CI)	p-value	p <sup>c</sup>
		n	%	n	%				
rs2162679 Intron	A/A	120	41.5	63	53.9	Codominant1	0.68 (0.43-1.07)	0.0900	0.4500
	A/G	129	44.6	48	41.0	Codominant2	0.27 (0.11-0.67)	<b>0.0050</b>	<b>0.0250</b>
	G/G	40	13.8	6	5.1	Dominant	0.58 (0.37-0.90)	<b>0.0150</b>	0.0800
						Recessive	0.32 (0.13-0.79)	<b>0.0059</b>	<b>0.0295</b>
						Log-additive	0.59 (0.41-0.83)	<b>0.0020</b>	<b>0.0100</b>
						1			
	G	209	36.2	60	25.6				
	A	369	63.8	174	74.4		1.64 (1.17-2.31)	<b>0.0040</b>	<b>0.0200</b>
	G/G	93	32.2	43	36.1	Codominant1	0.84 (0.53-1.36)	0.4800	1.0000
						Codominant2	0.78 (0.39-1.54)	0.4700	1.0000
						Dominant	0.83 (0.53-1.30)	0.4200	1.0000
						Recessive	0.86 (0.46-1.61)	0.6400	1.0000
						Log-additive	0.87 (0.63-1.21)	0.4100	1.0000
rs2195239 Intron	G	338	58.5	146	61.3		1		
	C	240	41.5	92	38.7		0.89 (0.65-1.21)	0.4500	1.0000
	G/G	90	31.1	41	34.5	Codominant1	0.92 (0.57-1.48)	0.7200	1.0000
						Codominant2	0.70 (0.37-1.36)	0.3000	1.0000
						Dominant	0.86 (0.54-1.35)	0.5100	1.0000
						Recessive	0.74 (0.41-1.34)	0.3200	1.0000
						Log-additive	0.85 (0.62-1.17)	0.3200	1.0000
						1			
	G	324	56.1	142	59.7				
	A	254	43.9	96	40.3		0.86 (0.64-1.17)	0.3400	1.0000
	C/C	90	31.1	41	34.8	Codominant1	0.93 (0.57-1.50)	0.7500	1.0000
						Codominant2	0.64 (0.33-1.25)	0.2000	1.0000
						Dominant	0.85 (0.53-1.34)	0.4700	1.0000
						Recessive	0.68 (0.37-1.23)	0.1900	0.9500
						Log-additive	0.83 (0.60-1.13)	0.2300	1.0000
rs1520220 Intron	C	323	55.9	142	60.2		1		
	G	255	44.1	94	39.8		0.84 (0.62-1.14)	0.2600	1.0000
	G/G	73	25.3	44	37.0	Codominant1	0.66 (0.41-1.08)	0.1000	0.5000
						Codominant2	0.44 (0.23-0.83)	<b>0.0110</b>	0.0600
						Dominant	0.59 (0.37-0.93)	<b>0.0250</b>	0.1300
						Recessive	0.57 (0.32-1.00)	<b>0.0420</b>	0.2100
						Log-additive	0.66 (0.49-0.90)	<b>0.0088</b>	<b>0.0440</b>
						1			
	A	288	49.8	94	39.5				
	G	290	50.2	144	60.5		1.52 (1.12-2.07)	<b>0.0070</b>	<b>0.0350</b>

The p-values were calculated from logistic regression analysis adjusting age and gender. Bold numbers indicate significant associations. IS, ischemic stroke; SNP, single nucleotide polymorphism; OR, odds ratio; CI, confidence interval.

Table IV. Genotype frequencies of *IGF* single nucleotide polymorphisms in stroke subgroups according to the NIHSS and MBI scores.

SNP	Type	Subgroups				Model	OR (95% CI)	p-value
		n	%	n	%			
NHISS		( $<6$ )		( $\geq 6$ )				
rs2162679	A/A	27	49.1	32	58.2	Codominant1	0.63 (0.29-1.38)	0.25
Intron	A/G	26	47.3	20	36.4	Codominant2	1.26 (0.19-8.23)	0.81
	G/G	2	3.6	3	5.5	Dominant	0.67 (0.31-1.45)	0.31
						Recessive	1.56 (0.25-9.79)	0.63
						Log-additive	0.79 (0.41-1.53)	0.49
rs2195239	G/G	23	41.8	16	28.1	Codominant1	2.00 (0.87-4.59)	0.10
Intron	C/G	24	43.6	33	57.9	Codominant2	1.46 (0.45-4.73)	0.53
	C/C	8	14.6	8	14.0	Dominant	1.87 (0.84-4.12)	0.12
						Recessive	0.97 (0.34-2.80)	0.95
						Log-additive	1.36 (0.77-2.38)	0.28
rs978458	G/G	22	40.0	15	26.3	Codominant1	1.89 (0.82-4.39)	0.14
Intron	A/G	25	45.5	32	56.1	Codominant2	1.87 (0.60-5.87)	0.28
	A/A	8	14.6	10	17.5	Dominant	1.89 (0.85-4.21)	0.12
						Recessive	1.27 (0.46-3.51)	0.65
						Log-additive	1.46 (0.83-2.54)	0.18
rs1520220	C/C	22	40.0	15	26.8	Codominant1	1.77 (0.76-4.10)	0.18
Intron	C/G	26	47.3	31	55.4	Codominant2	2.12 (0.66-6.86)	0.21
	G/G	7	12.7	10	17.9	Dominant	1.84 (0.83-4.12)	0.13
						Recessive	1.50 (0.52-4.28)	0.45
						Log-additive	1.52 (0.86-2.67)	0.14
rs6214	G/G	20	36.4	21	36.8	Codominant1	0.96 (0.42-2.20)	0.92
3'UTR	A/G	27	49.1	26	45.6	Codominant2	1.22 (0.40-3.73)	0.73
	A/A	8	14.6	10	17.5	Dominant	1.02 (0.47-2.23)	0.96
						Recessive	1.25 (0.45-3.45)	0.67
						Log-additive	1.07 (0.63-1.84)	0.79
MBI		( $<60$ )		( $\geq 60$ )				
rs2162679	A/A	37	53.6	15	62.5	Codominant1	0.96 (0.35-2.65)	0.94
Intron	A/G	29	42.0	9	37.5	Codominant2	0.00 (0.00-NA)	
	G/G	3	4.3	0	0.0	Dominant	0.91 (0.33-2.50)	0.85
						Recessive	0.00 (0.00-NA)	0.32
						Log-additive	0.83 (0.32-2.15)	0.70
rs2195239	G/G	25	35.2	8	33.3	Codominant1	1.06 (0.36-3.12)	0.91
Intron	C/G	36	50.7	13	54.2	Codominant2	0.86 (0.17-4.33)	0.86
	C/C	10	14.1	3	12.5	Dominant	1.02 (0.36-2.88)	0.97
						Recessive	0.83 (0.19-3.61)	0.80
						Log-additive	0.96 (0.46-2.03)	0.92
rs978458	G/G	24	33.8	7	29.2	Codominant1	1.40 (0.46-4.25)	0.55
Intron	A/G	35	49.3	14	58.3	Codominant2	0.83 (0.17-4.15)	0.82
	A/A	12	16.9	3	12.5	Dominant	1.26 (0.43-3.66)	0.67
						Recessive	0.67 (0.16-2.82)	0.58
						Log-additive	1.00 (0.48-2.07)	1.00
rs1520220	C/C	24	34.3	7	29.2	Codominant1	1.44 (0.47-4.36)	0.52
Intron	C/G	35	50.0	14	58.3	Codominant2	0.90 (0.18-4.57)	0.90
	G/G	11	15.7	3	12.5	Dominant	1.31 (0.45-3.81)	0.62
						Recessive	0.72 (0.17-3.06)	0.65
						Log-additive	1.04 (0.50-2.17)	0.92

Table IV. Continued.

SNP	Type	Subgroups				Model	OR (95% CI)	p-value
		n	%	n	%			
rs6214	G/G	26	36.6	10	41.7	Codominant1	0.41 (0.13-1.32)	0.14
3'UTR	A/G	37	52.1	8	33.3	Codominant2	1.56 (0.40-6.13)	0.53
	A/A	8	11.3	6	25.0	Dominant	0.62 (0.22-1.75)	0.37
						Recessive	2.50 (0.73-8.61)	0.15
						Log-additive	1.06 (0.52-2.17)	0.86

The p-values were evaluated from logistic regression analysis adjusting age and gender. Subjects with an undetected genotype were excluded. SNP, single nucleotide polymorphism; NIHSS, National Institutes of Health Stroke Survey; MBI, Modified Barthel Index; OR, odds ratio; CI, confidence interval.

numbers of IS patients with a MBI score of <60 and ≥60 were 71 (74.7%) and 24 (25.3%), respectively. In two clinical subgroups, patients with inappropriate or insufficient clinical data, were excluded (Table I).

**Genetic analysis of IGF1 SNPs.** Table III represents the genotype and allele frequencies of the five examined SNPs (rs2162679, rs2195239, rs978458, rs1520220 and rs6214) in the IS patients and the control subjects. The HWE of the five SNPs showed no difference in the control group (rs2162679,  $p=0.61$ ; rs2195239,  $p=0.18$ ; rs978458,  $p=0.91$ ; rs1520220,  $p=1.00$ ; rs6214,  $p=1.00$ ). Multiple logistic regression analysis adjusting for age and gender was performed. An intron SNP (rs2162679) was associated with the development of IS [ $p=0.0050$ , OR = 0.27, 95% CI = 0.11-0.67 in the codominant2 model (A/A vs. G/G);  $p=0.0150$ , OR = 0.58, 95% CI = 0.37-0.90 in the dominant model (A/A vs. A/G and G/G);  $p=0.0059$ , OR = 0.32, 95% CI = 0.13-0.79 in the recessive model (A/A and A/G vs. G/G); and  $p=0.0020$ , OR = 0.59, 95% CI = 0.41-0.83 in the log-additive model (A/A vs. A/G vs. G/G)]. A 3'UTR SNP (rs6214) was also associated with the development of IS [ $p=0.0110$ , OR = 0.44, 95% CI = 0.23-0.83 in the codominant2 model (G/G vs. A/A);  $p=0.0250$ , OR = 0.59, 95% CI = 0.37-0.93 in the dominant model (G/G vs. A/G and A/A);  $p=0.0420$ , OR = 0.57, 95% CI = 0.32-1.00 in the recessive model (G/G and A/G vs. A/A); and  $p=0.0088$ , OR = 0.66, 95% CI = 0.49-0.90 in the log-additive model (G/G vs. A/G vs. A/A)]. The A allele frequency of rs2162679 was higher in the IS group (74.4%) than in the control group (63.8%) ( $p=0.004$ , OR = 1.64, 95% CI = 1.17-2.31). The G allele frequency of rs6214 was higher in the IS group (60.5%) than in the control group (50.2%) ( $p=0.007$ , OR = 1.52, 95% CI = 1.12-2.07). After Bonferroni correction ( $p^c$ ), the allele frequencies of rs2162679 and rs6214 showed significant differences between IS and the controls (rs2162679,  $p^c=0.004$ ; rs6214,  $p^c=0.007$ ). The other SNPs (rs2195239, rs978458 and rs1520220) had no differences between IS and the controls (Table III). The LD block was estimated using Haploview version 4.2. One LD block was strongly made between rs1520220 and rs978458 ( $D'=1.0$  and  $r^2=0.993$  in the control group). However, the haplotypes of these two SNPs were not associated with the development of IS (data not shown). IS patients were classified into two clinical subgroups according to the NIHSS (<6 and ≥6) and

MBI (<60 and ≥60) scores. As shown in Table IV, the five tested SNPs were not associated with the NIHSS and MBI scores.

**Sample power.** We estimated the sample power using a genetic power calculator to obtain the required sample size. The sample powers ( $\alpha=0.05$ , genotype relative risk 2-fold, number of case for 70% power) of each SNP in the IS group were 0.757 for rs2162679 ( $n=104$ ), 0.800 for rs2195239 ( $n=93$ ), 0.801 for rs978458 ( $n=93$ ), 0.801 for rs1520220 ( $n=93$ ) and 0.801 for rs6214 ( $n=93$ ). Therefore, the results of the five examined SNPs in the *IGF1* gene had statistical confidence.

## Discussion

IGF1 is an endogenous survival factor for neurons, glial cells and endothelial cells. IGF1 plays an important role in tissue repair and cell proliferation. IGF1 induces the synthesis of elastin and prevents apoptosis of vascular smooth muscle cells. Therefore, low levels of IGF1 may be a risk factor of stroke. As shown in previous studies, the expression of IGF1 increased after hypoxic injury in regions with neuronal loss (18) and IGF1 reduced infarct volume and improved neurological function after ischemia in an animal study (19).

There are several reports that *IGF1* polymorphisms are associated with certain diseases, such as adenocarcinoma (EAC) and colorectal cancer (15,20). McElholm *et al* observed that *IGF1* SNP rs6214 was associated with Barrett's esophagus (BE) in EAC (15). Using GG genotype as reference, OR for BE in AA (wild-type) was 0.43 (95% CI 0.24-0.75). Feik *et al* also suggested that rs6214 could have an impact on developing colorectal cancer and colorectal polyps with villous elements (20). Based on previous studies, we believed that the SNPs of *IGF1* were associated with the development of IS and clinical features according to the NIHSS and MBI scores in Korean stroke patients. In the present study, we found a significant association between *IGF1* SNPs and IS. The G allele frequency of rs6214 in the *IGF1* gene was higher in the IS group (60.5%) than in the control group (50.2%). The A allele of rs2162679 was also higher in the IS group (74.4%) than in the control group (74.4%). Thus, our results suggest that *IGF1* may be a risk factor of IS development. However, all five SNPs (rs2162679, rs2195239, rs978458, rs1520220 and rs6214) of *IGF1* did not contribute to the NIHSS and MBI scores of



IS. To our knowledge, this is the first study on whether *IGF1* SNPs are associated with the development of IS in a Korean population. Additional studies with a larger number of cases or different populations are required in order to confirm our results.

In conclusion, we suggest that an intron SNP (rs2162679) and 3'UTR SNP (rs6214) of the *IGF1* gene may be associated with the development of IS.

## Acknowledgments

This work was supported by the Biogreen 21 Program (Code PJ007187), Rural Development Administration

## References

- Grysiewicz RA, Thomas K and Pandey DK: Epidemiology of ischemic and hemorrhagic stroke: incidence, prevalence, mortality, and risk factors. *Neurol Clin* 26: 871-895, 2008.
- Van Asch CJ, Luitse MJ, Rinkel GJ, *et al*: Incidence, case fatality, and functional outcome of intracerebral haemorrhage over time, according to age, sex, and ethnic origin: a systematic review and meta-analysis. *Lancet Neurol* 9: 167-176, 2010.
- Tao HM, Chen GZ, Lu XD, Chen GP and Shao B: TGF-beta1 869T/C polymorphism and ischemic stroke: sex difference in Chinese. *Can J Neurol Sci* 37: 803-807, 2010.
- Tong Y, Geng Y, Xu J, *et al*: The role of functional polymorphisms of the TNF-alpha gene promoter in the risk of ischemic stroke in Chinese Han and Uyghur populations: two case-control studies. *Clin Chim Acta* 411: 1291-1295, 2010.
- Lim YH, Jeong YS, Kim SK, *et al*: Association between TGFBR2 gene polymorphism (rs2228048, Asn389Asn) and intracerebral hemorrhage in Korean Population. *Immunol Invest* 40: 569-580, 2011.
- Kalmijn S, Janssen JA, Pols HA, Lamberts SW and Breteler MM: A prospective study on circulating insulin-like growth factor I (IGF-I), IGF-binding proteins, and cognitive function in the elderly. *J Clin Endocrinol Metab* 85: 4551-4555, 2000.
- Tagami M, Yamagata K, Nara Y, *et al*: Insulin-like growth factors prevent apoptosis in cortical neurons isolated from stroke-prone spontaneously hypertensive rats. *Lab Invest* 76: 603-612, 1997.
- Galli C, Meucci O, Scorziello A, *et al*: Apoptosis in cerebellar granule cells is blocked by high KCl, forskolin, and IGF-1 through distinct mechanisms of action: the involvement of intracellular calcium and RNA synthesis. *J Neurosci* 15: 1172-1179, 1995.
- Kooijman R, Sarre S, Michotte Y and De Keyser J: Insulin-like growth factor I: a potential neuroprotective compound for the treatment of acute ischemic stroke? *Stroke* 40: e83-e88, 2009.
- Guan J, Gunn AJ, Sirimanne ES, *et al*: The window of opportunity for neuronal rescue with insulin-like growth factor-1 after hypoxia-ischemia in rats is critically modulated by cerebral temperature during recovery. *J Cereb Blood Flow Metab* 20: 513-519, 2000.
- De Smedt A, Brouns R, Uyttenboogaart M, *et al*: Insulin-like growth factor I serum levels influence ischemic stroke outcome. *Stroke* 42: 2180-2185, 2011.
- Bondanelli M, Ambrosio MR, Onofri A, *et al*: Predictive value of circulating insulin-like growth factor I levels in ischemic stroke outcome. *J Clin Endocrinol Metab* 91: 3928-3934, 2006.
- Henningson M, Hietala M, Torngren T, Olsson H and Jernstrom H: IGF1 htSNPs in relation to IGF-1 levels in young women from high-risk breast cancer families: implications for early-onset breast cancer. *Fam Cancer* 10: 173-185, 2010.
- Hahn WH, Suh JS and Cho BS: Polymorphisms of insulin-like growth factor-1 (IGF-1) and IGF-1 receptor (IGF-1R) contribute to pathologic progression in childhood IgA nephropathy. *Growth Factors* 29: 8-13, 2010.
- McElholm AR, McKnight AJ, Patterson CC, *et al*: A population-based study of IGF axis polymorphisms and the esophageal inflammation, metaplasia, adenocarcinoma sequence. *Gastroenterology* 139: 204-212, 2010.
- Kim SK, Park HJ, Lee JS, *et al*: Association of niemann-pick disease, type c2 (NPC2) polymorphisms with obesity in Korean population. *Mol Cell Toxicol* 6: 395-400, 2010.
- Lim TW, Kim SK, Ban JY, *et al*: Polymorphisms of transmembrane channel-like 1 gene are associated with Kawasaki disease in Korean population. *Mol Cell Toxicol* 5: 291-297, 2009.
- Gluckman P, Klempt N, Guan J, *et al*: A role for IGF-1 in the rescue of CNS neurons following hypoxic-ischemic injury. *Biochem Biophys Res Commun* 182: 593-599, 1992.
- Liu XF, Fawcett JR, Thorne RG and Frey WH II: Non-invasive intranasal insulin-like growth factor-I reduces infarct volume and improves neurologic function in rats following middle cerebral artery occlusion. *Neurosci Lett* 308: 91-94, 2001.
- Feik E, Baierl A, Hieger B, *et al*: Association of IGF1 and IGFBP3 polymorphisms with colorectal polyps and colorectal cancer risk. *Cancer Causes Control* 21: 91-97, 2010.