

Neuroprotective effect of paeoniflorin on H₂O₂-induced apoptosis in PC12 cells by modulation of reactive oxygen species and the inflammatory response

PENG LI¹ and ZHAOHUI LI²

¹Medical College of Nankai University, Tianjin 300071; ²Department of Ophthalmology, General Hospital of PLA, Beijing 100853, P.R. China

Received July 25, 2014; Accepted February 9, 2015

DOI: 10.3892/etm.2015.2360

Abstract. Paeoniflorin (PF) is a product derived from *Paeoniae Radix* and is commonly prescribed in traditional Chinese medicine. PF has been reported to exhibit neuroprotective, anti-ischemic, antioxidant, anti-inflammatory and anticancer effects. The neuroprotective properties of PF have been demonstrated in animal models of various neuropathologies. The present study investigated the effects of PF on hydrogen peroxide (H₂O₂)-induced apoptosis in PC12 cells, to improve the understanding of the mechanisms underlying its neuroprotective properties. The H₂O₂-induced apoptosis of PC12 cells resulted in a reduction in the B-cell lymphoma 2 (Bcl-2)/Bcl-2-associated X protein ratio and the activation of caspase-3. PF treatment was observed to reverse the apoptotic process and to modulate the expression levels of a number of apoptosis-associated proteins. Furthermore, PF significantly mitigated the H₂O₂-induced reduction in cell viability, in addition to scavenging reactive oxygen species and preventing the release of lactate dehydrogenase from the PC12 cells. In addition, the apoptosis-associated activation of nuclear factor (NF)-κB was inhibited in the PF-treated cells, and the expression levels of tumor necrosis factor α and interleukin (IL)-1β were reduced. In conclusion, the present study demonstrated that PF was able to reduce H₂O₂-induced toxicity by blocking the activation of the neuroinflammatory factor NF-κB. These results suggest that PF may be a valuable neuroprotective agent for the treatment of neurological disease and injury.

Introduction

Paeoniae Radix is a well-known herb and is used widely as a component of traditional Chinese prescriptions to treat certain types of dementia, traumatic injury and inflammation. Paeoniflorin (PF), a product derived from *Paeoniae Radix*, has been reported to exhibit neuroprotective, anti-ischemic, antioxidative, anti-inflammatory and anticancer effects. The neuroprotective potential of PF has been demonstrated in animal models of various neuropathologies (1-4).

Reactive oxygen species (ROS) are produced by various enzymatic reactions and chemical processes, which are essential for numerous physiological functions, in addition to serving as secondary messengers in the human body (5). A number of neurodegenerative diseases, including Alzheimer's, Parkinson's and Huntington's, are characterized by severe and/or prolonged oxidative stress (6). The primary outcome of oxidative stress is the irreversible damage of macromolecules by ROS (7). The association between oxidative stress and inflammation is due to the activation of nuclear factor (NF)-κB and activator protein-1, and the inhibition of nuclear factor (erythroid-derived 2)-like 2, peroxynitrite-mediated endothelial dysfunction, altered nitric oxide levels and macrophage migration (8). Previous studies have indicated that PF protects neurons against ischemia-reperfusion injury by reducing the expression levels of intracellular adhesion molecule 1 and tumor necrosis factor α (TNF-α), resulting in reduced inflammation in infarcted brain regions, and PF prevents chronic cognitive damage by downregulating the expression of NF-κB in hippocampal astrocytes (4,9). The present study investigated the neuroprotective effect of PF following H₂O₂-induced injury in PC12 cells and the possible signaling pathways involved.

Materials and methods

Reagents and cell line. PF (purity, 98.5%) was purchased from Nanjing Zelang Medical Technology Co., Ltd. (Nanjing, China). The PC12 cell line was obtained from the American Type Culture Collection (Manassas, VA, USA).

MTT cell proliferation assay. Cell viability was measured using an MTT assay as described in a previous study (9).

Correspondence to: Dr Zhaohui Li, Department of Ophthalmology, General Hospital of PLA, 28 Fuxing Road, Beijing 100853, P.R. China
E-mail: li_zhao_hui@163.com

Key words: paeoniflorin, neuroprotective effects, PC12 cells, inflammatory response

The PC12 cells received different treatments, including no treatment (control), 200 μM H_2O_2 alone or 200 μM H_2O_2 in combination with 20, 40 or 80 μM PF. Briefly, the cells were seeded into 96-well plates (3.0×10^3 /well) and cultured for 6 h. MTT solution (5 mg/ml; Sigma-Aldrich, St. Louis, MO, USA) was added to each well and incubated for 4 h. Next, 150 μl dimethyl sulfoxide (DMSO; Sigma-Aldrich) was added to dissolve the formazan precipitate. Absorbance was then measured at 570 nm using a ThermoMax microplate reader (Molecular Devices LLC, Sunnyvale, CA, USA). Cell viability is expressed as a percentage relative to the untreated control.

Lactate dehydrogenase (LDH) release assay. The rate of cell death was further assessed by measuring the leakage of LDH into the surrounding medium, as described in a previous study (6). Briefly, following treatment of the PC12 cells, the supernatants of each group were collected. The quantity of LDH released was determined using a Neutral Red LDH Cytotoxicity Assay Kit according to the manufacturer's instructions (Beyotime Institute of Biotechnology, Wuhan, China). Optical absorbance was measured at 440 nm using the ThermoMax microplate reader.

Measurement of intracellular ROS levels. Intracellular H_2O_2 and low-molecular weight peroxides are able to oxidize 2',7'-dichlorofluorescein diacetate (DCFH-DA) to dichlorofluorescein (DCF), which is highly fluorescent under absorption analysis. A DCFH-DA fluorescent probe from a Reactive Oxygen Species Assay kit (Beyotime Institute of Biotechnology) was used to measure ROS generation, as previously reported (6). Following treatment, cells were incubated with 10 mM DCFH-DA for 30 min at 37°C and washed twice with phosphate-buffered saline. Subsequently, the DCF fluorescence was measured using the ThermoMax microplate spectrofluorometer at excitation and emission wavelengths of 485 and 530 nm, respectively.

Hoechst 33258 staining. PC12 cells at the logarithmic-growth phase were seeded into 96-well plates (1×10^4 /well). The cells were cultured in H_2O_2 alone or with 80 μM PF. A third group of cells received no treatment and was used as a control group. Next, the cells were fixed with 3.7% paraformaldehyde for 30 min at room temperature, then washed and stained with Hoechst 33258 (Sigma-Aldrich) for 30 min at 37°C. PC12 cells were observed under a Nikon 80i fluorescence microscope equipped with a UV filter (Nikon Corporation, Tokyo, Japan).

Western blot analysis. PC12 cells were seeded in 6-well plates (3.0×10^5 /well) and pretreated with 200 μM H_2O_2 alone or 200 μM H_2O_2 + 80 μM PF for 6 h. A third group of cells received no treatment and was used as a control group. After incubation the culture medium was collected for detection of the levels of TNF- α and interleukin (IL)-1 β . Cells were collected and lysed in a buffer containing 50 mM HEPES (pH 7.4), 150 mM NaCl, 0.1% Triton X-100, 1.5 mM MgCl_2 , 1 mM EDTA, 2 mM sodium orthovanadate, 4 mM sodium pyrophosphate, 100 mM NaF and protease inhibitor mixture (1:500; Sigma-Aldrich) for cell lysates. Cell lysates were subjected to 10% SDS-polyacrylamide gel




Figure 1. Protective effect of PF on the H_2O_2 -induced reduction of cell viability in PC12 cells. * $P < 0.01$ vs. control group, # $P < 0.05$ and ## $P < 0.01$ vs. the H_2O_2 group. H_2O_2 , hydrogen peroxide; PF, paeoniflorin.




Figure 2. Effects of PF on H_2O_2 -induced increase in ROS accumulation and LDH release in PC12 cells. * $P < 0.01$ vs. the control group, # $P < 0.05$ and ## $P < 0.01$ vs. the H_2O_2 group. H_2O_2 , hydrogen peroxide; PF, paeoniflorin; ROS, reactive oxygen species; LDH, lactate dehydrogenase.

(Invitrogen, Thermo Fisher Scientific, USA) electrophoresis, then transferred onto polyvinylidene fluoride membranes (EMD Millipore, Billerica, MA, USA). The membranes were subsequently probed with antibodies, including rabbit polyclonal caspase-3 (#9662), cleaved poly(ADP-ribose) polymerase (PARP; #9541), B-cell lymphoma 2 (Bcl-2; #2872) and Bcl-2-associated X (Bax; #2772) antibodies purchased from Cell Signaling Technology, Inc. (1:1,000; Danvers, MA, USA). Mouse monoclonal NF- κB -p65RelA (1:800; sc-8008) and rabbit polyclonal p-NF- κB Ser536 (1:500; sc-33020) antibodies were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA) and mouse monoclonal β -actin antibody (1:10,000; #ab6276) from Abcam (Cambridge, MA, USA). Immunoblots were developed using horseradish peroxidase (HRP)-conjugated secondary antibodies. Bound antibodies were visualized using Immobilon Western Chemiluminescent HRP substrate (EMD Millipore) and quantified by densitometry using a ChemiDoc XRS system (Bio-Rad Laboratories Inc., Berkeley, CA, USA). Densitometric analyses of bands were adjusted against β -actin, which functioned as a loading control. The percentage increase or reduction in protein




Figure 3. PF protected PC12 cells against H₂O₂-induced apoptosis. (A) Morphological analysis of nuclear chromatin in a) control, b) H₂O₂ and c) 80 μM PF treated cells with Hoechst 33258 staining. Cells displayed condensed chromatin and apoptotic nuclei following H₂O₂ treatment, while PF treatment markedly reversed these effects. (B) Bax:Bcl-2 ratio in the control, H₂O₂ and 80 μM PF groups. (C) PF reversed the H₂O₂-induced increase in the expression levels caspase-3 and cleaved PARP in PC12 cells. *P<0.01 vs. the control group and ##P<0.01 vs. the H₂O₂ group. H₂O₂, hydrogen peroxide; PF, paeoniflorin; PARP, poly(ADP-ribose) polymerase; Bcl-2, B-cell lymphoma 2; Bax, Bcl-2-associated X.

expression levels was estimated by comparison to a vehicle control. Experiments were performed in triplicate, separately.

TNF-α and IL-1β assays. The culture medium was collected in microcentrifuge tubes and subjected to centrifugation for 10 min. The supernatants were separated out and the expression levels of TNF-α and IL-1β were detected using Human TNF-alpha Quantikine (DTA00C) and Human IL-1 beta/IL-1F2 Quantikine ELISA kits (R&D Systems Inc., Minneapolis, MN, USA) according to the manufacturer's instructions.

Statistical analysis. Statistical analysis was performed using SPSS software for Windows, version 18.0 (SPSS, Inc., Chicago, IL, USA). Data are presented as the mean ± standard deviation and were analyzed by one-way analysis of variance. Multiple comparisons between groups were performed using the Student-Newman-Keuls method and P<0.05 was considered to indicate a statistically significant difference.

Results

Effect of PF on cell viability. The viability of cells incubated with 200 μM H₂O₂ was 58.6±2.4% of the control value (P<0.01; Fig. 1). The viabilities of cells treated with 200 μM H₂O₂ + 20, 40 or 80 μM PF were increased in a dose-dependent manner to 66.3±1.6 (P<0.05 vs. H₂O₂), 75.9±1.1 and 83.4±1.7% (P<0.01 vs. H₂O₂) of the control values, respectively (n=3; Fig. 1). These results clearly indicate that PF attenuated the H₂O₂-induced cytotoxicity in the PC12 cells.

Effect of PF on LDH and ROS levels. The neuroprotective effect of PF was further investigated by measuring ROS accumulation and levels of LDH release following treatment.

Pretreatment with PF attenuated the H₂O₂-induced increase in levels of ROS and LDH release (Fig. 2).

PF protects PC12 cells against H₂O₂-induced apoptosis. Alterations of cellular morphology were assessed using Hoechst 33258 staining in order to characterize the degree of H₂O₂-induced PC12 cell death (Fig. 3A). The nuclei of the PC12 cells treated with H₂O₂ appeared fragmented, indicating that apoptosis affected the morphology of the cells. However, treatment with 80 μM PF for 6 h clearly reduced the percentage of necrotic and apoptotic cells.

Expression levels of the anti-apoptotic protein Bcl-2 and the pro-apoptotic protein Bax were measured using western blot analysis. H₂O₂ was observed to increase the Bax:Bcl-2 ratio in the PC12 cells, while the PF treatment produced an opposite effect (Fig. 3B). Furthermore, the H₂O₂-induced elevation of the expression levels of caspase-3 and cleaved PARP appeared significantly reduced in cells treated with PF (Fig. 3).

PF suppresses the expression levels of NF-κB and its associated inflammatory factors. Western blot analysis indicated that the expression levels and activity of NF-κB were elevated in cells treated with H₂O₂ alone. Treatment with PF appeared to significantly reduce this H₂O₂-induced NF-κB activity (Fig. 4A). The levels of total NF-κB protein displayed a marked reduction in cells treated with PF (P<0.01). Further analysis indicated that the levels of p-NF-κB (Ser536), the active form of NF-κB, were significantly reduced by the PF treatment (P<0.01).

Inhibitory effect of PF on the expression levels of TNF-α and IL-1β. Western blot analysis indicated that treatment with PF reversed the H₂O₂-induced elevation of the expression levels




Figure 4. PF suppressed the expression of NF- κ B and its associated inflammatory factors. (A) PF reduced the H₂O₂-induced increase in the expression levels of (A) NF- κ B and its active form p-NF- κ B Ser536, (B) IL-1 β and (C) TNF- α . * P <0.01 vs. the control group and ## P <0.01 vs. the H₂O₂ group. H₂O₂, hydrogen peroxide; PF, paeoniflorin; NF- κ B, nuclear factor- κ B; IL-1 β , interleukin 1 β ; TNF- α , tumor necrosis factor α .

of IL-1 β (P <0.01; Fig. 4C) and TNF- α (P <0.01; Fig. 4B) in the PC12 cells.

Discussion

PF is the main component of *Paeoniae Radix* used in traditional Chinese medicine, and has been reported to exhibit numerous pharmacological effects. The results of the present study indicated that PF may protect PC12 cells from H₂O₂-induced oxidative injury. PF was observed to regulate H₂O₂-induced oxidative stress, indicated by the modulation of LDH and ROS levels in PC12 cells. PF was also demonstrated to reduce H₂O₂-induced apoptosis and promote overall cell survival. Furthermore, the results indicated that PF down-regulated H₂O₂-induced neuroinflammation by regulating NF- κ B-associated inflammatory signals.

Oxidative stress has been extensively implicated in the pathophysiology of cerebral ischemia and stroke (10). Hypoxia is a crucial initiator of the loss of neurocytes and apoptosis is considered to be a pivotal source of damage to neurocytes during this process (11). The accumulation of ROS may lead to various forms of oxidative modification of proteins, lipids and DNA, resulting in cellular damage (12). The present study demonstrated that 200 μ M H₂O₂ was able to signifi-

cantly stimulate the accumulation of ROS and the release of LDH (203.1 and 270.1% of control values, respectively) in PC12 cells. Furthermore, the cell survival rate in the H₂O₂ group was 58.6 \pm 2.4% of the control value.

PF treatment appeared to markedly improve these oxidative conditions. The ROS levels in the 20, 40 and 80 μ M PF treatment groups were 171.8, 141.6 and 117.4% of the control group, respectively. The LDH expression levels of the 20, 40 and 80 μ M PF treatment groups were 217.0, 175.1 and 133.8% of the control group, respectively. These results indicated that the PF treatment produced a significant reduction in H₂O₂-induced toxicity and oxidative stress in the PC12 cells.

ROS are widely recognized to be key mediators of cell survival, proliferation, differentiation and apoptosis (5,13,14). Previous studies have demonstrated that proteins of the Bcl-2 family, including Bax and Bcl-2, are associated with apoptosis induced by ROS-generating agents (Ji BS, Renaud, Pan). In addition, ROS may activate caspase-3, which results in the cleavage of PARP, a 116-kDa nuclear poly (ADP-ribose) polymerase, which appears to be involved in DNA repair in response to environmental stress. PARP may be cleaved by numerous caspase-1-like caspases *in vitro* and is one of the primary cleavage targets of caspase-3 *in vivo*. Furthermore, ROS may activate caspase-3, which results in the cleavage of PARP into an 89-kDa fragment (6,15-17,20). In the present study, a Hoechst 33258 staining assay indicated that treatment with 200 μ M H₂O₂ alone induced notable cell apoptosis in PC12 cells, while 80 μ M PF produced a reduction in the extent of apoptosis-associated nuclear fragmentation (Fig. 3A). Furthermore, H₂O₂-induced apoptosis was associated with an increase in the Bax:Bcl-2 ratio and with the activation of caspase-3. Treatment with PF was observed to downregulate the expression of the pro-apoptotic protein Bax, and to upregulate the anti-apoptotic protein Bcl-2. The results of the present study also demonstrated that caspase-3 and cleaved PARP were modulated by PF treatment.

Oxidative stress-induced neuroinflammation has been reported to be a vital factor in nerve injury and associated diseases (5,8,21). Numerous studies have suggested that chronic inflammation is implicated in neurodegenerative disease and injury (1,4,21-23). A number of well-established inflammatory target proteins, including matrix metalloproteinase-9, cyclooxygenase-2, inducible nitric oxide synthase and certain adhesion molecules have been associated with ROS generation, which is also induced by proinflammatory cytokines, peptides, peroxidants and infection (5,13,24,25). Increasing inflammatory stress has been reported to correlate with oxidative stress during the progression of neurodegenerative disease (5,19,26). NF- κ B, a proinflammatory transcription factor, functions as the 'first responder' to various generators of cellular stress, including free radicals and pro-inflammatory cytokines (e.g. TNF- α) and bacterial biomolecules (27). In the present study, H₂O₂ was observed to induce an inflammatory response involving NF- κ B and its associated signals. Following H₂O₂ treatment, the levels of NF- κ B and its active form, p-NF- κ B (Ser536), were elevated, as were the levels of TNF- α and IL-1 β . However, cells cocultured with 80 μ M PF exhibited reduced levels of these inflammatory factors, indicating that PF modified the apoptotic process, in addition to correcting the abnormal inflammatory signals induced by H₂O₂.

In conclusion, PF treatment significantly reduced H₂O₂-induced apoptosis and ROS accumulation, promoted cell survival and downregulated neuroinflammation in PC12 cells. Thus, PF may serve as a protective agent against oxidative stress and scavenger of intracellular ROS, and may offer a novel pharmacological preventative or palliative treatment for ischemic cerebral injury.

References

- Liu HQ, Zhang WY, Luo XT, Ye Y and Zhu XZ: Paeoniflorin attenuates neuroinflammation and dopaminergic neurodegeneration in the MPTP model of Parkinson's disease by activation of adenosine A1 receptor. *Br J Pharmacol* 148: 314-325, 2006.
- Liu DZ, Xie KQ, Ji XQ, Ye Y, Jiang CL and Zhu XZ: Neuroprotective effect of paeoniflorin on cerebral ischemic rat by activating adenosine A1 receptor in a manner different from its classical agonists. *Br J Pharmacol* 146: 604-611, 2005.
- Tang LM, Liu IM and Cheng JT: Stimulatory effect of paeoniflorin on adenosine release to increase the glucose uptake into white adipocytes of Wistar rat. *Planta Med* 69: 332-336, 2003.
- Kapoor S: Neuroprotective effects of paeoniflorin: An emerging concept in neurology. *Folia Neuropathol* 51: 92, 2013.
- Hsieh HL and Yang CM: Role of redox signaling in neuroinflammation and neurodegenerative diseases. *Biomed Res Int* 2013: 484613, 2013.
- Jia BS and Gao Y: Protective effect of trihexyphenidyl on hydrogen peroxide-induced oxidative damage in PC12 cells. *Neurosci Lett* 437: 50-54, 2008.
- Dasgupta A, Zheng J and Bizzozero OA: Protein carbonylation and aggregation precede neuronal apoptosis induced by partial glutathione depletion. *ASN Neuro* 4: e00084, 2012.
- Sandireddy R, Yerra VG, Areti A, Komirishetty P and Kumar A: Neuroinflammation and oxidative stress in diabetic neuropathy: Futuristic strategies based on these targets. *Int J Endocrinol* 2014: 674987, 2014.
- Liu J, Jin DZ, Xiao L and Zhu XZ: Paeoniflorin attenuates chronic cerebral hypoperfusion-induced learning dysfunction and brain damage in rats. *Brain Res* 1089: 162-170, 2006.
- Li RC, Morris MW, Lee SK, Pouranfar F, Wang Y and Gozal D: Neuroglobin protects PC12 cells against oxidative stress. *Brain Res* 1190: 159-166, 2008.
- Vilalta A and Brown GC: Deoxyglucose prevents neurodegeneration in culture by eliminating microglia. *J Neuroinflammation* 11: 58, 2014.
- Giorgio M, Trinei M, Migliaccio E and Pelicci PG: Hydrogen peroxide: A metabolic by-product or a common mediator of ageing signals? *Nat Rev Mol Cell Biol* 8: 722-728, 2007.
- Dröge W: Free radicals in the physiological control of cell function. *Physiol Rev* 82: 47-95, 2002.
- von Bernhardt R and Eugén J: Alzheimer's disease: Redox dysregulation as a common denominator for diverse pathogenic mechanisms. *Antioxid Redox Signal* 16: 974-1031, 2012.
- Wang R, Zhang HY and Tang XC: Huperzine A attenuates cognitive dysfunction and neuronal degeneration caused by beta-amyloid protein-(1-40) in rat. *Eur J Pharmacol* 421: 149-156, 2001.
- Lee SY, Ha TY, Son DJ, Kim SR and Hong JT: Effect of sesaminol glucosides on β -amyloid-induced PC12 cell death through anti-oxidant mechanisms. *Neurosci Res* 52: 330-341, 2005.
- Shelton SN, Shawgo ME, Matthews SB, Lu Y, Donnelly AC, Szabla K, Tanol M, *et al*: KU135, a novel novobiocin-derived C-terminal inhibitor of the 90-kDa heat shock protein, exerts potent antiproliferative effects in human leukemic cells. *Mol Pharmacol* 76: 1314-1322, 2009.
- Renaud J, Bournival J, Zottig X and Martinoli MG: Resveratrol protects DAergic PC12 cells from high glucose-induced oxidative stress and apoptosis: Effect on p53 and GRP75 localization. *Neurotox Res* 25: 110-123, 2014.
- Pan C, Giraldo GS, Prentice H and Wu JY: Taurine protection of PC12 cells against endoplasmic reticulum stress induced by oxidative stress. *J Biomed Sci* 17 (Suppl 1): S17, 2010.
- Cheung HH, Lynn Kelly N, Liston P and Korneluk RG: Involvement of caspase-2 and caspase-9 in endoplasmic reticulum stress-induced apoptosis: A role for the IAPs. *Exp Cell Res* 312: 2347-2357, 2006.
- Di Napoli M and Shah IM: Neuroinflammation and cerebrovascular disease in old age: A translational medicine perspective. *J Aging Res* 2011: 857484, 2011.
- Guo RB, Wang GF, Zhao AP, Gu J, Sun XL and Hu G: Paeoniflorin protects against ischemia-induced brain damages in rats via inhibiting MAPKs/NF- κ B-mediated inflammatory responses. *PLoS One* 7: e49701, 2012.
- Bettcher BM and Kramer JH: Inflammation and clinical presentation in neurodegenerative disease: A volatile relationship. *Neurocase* 19: 182-200, 2013.
- Lee IT and Yang CM: Role of NADPH oxidase/ROS in pro-inflammatory mediators-induced airway and pulmonary diseases. *Biochem Pharmacol* 84: 581-90, 2012.
- Chiurchiù V and Maccarrone M: Chronic inflammatory disorders and their redox control: From molecular mechanisms to therapeutic opportunities. *Antioxid Redox Signal* 15: 2605-2641, 2011.
- Urrutia PJ, Mena NP and Núñez MT: The interplay between iron accumulation, mitochondrial dysfunction, and inflammation during the execution step of neurodegenerative disorders. *Front Pharmacol* 5: 38, 2014.
- Liu HP, Gao ZH, Cui SX, Sun DF, Wang Y, Zhao CR, Lou HX and Qu XJ: Inhibition of intestinal adenoma formation in APC(Min/+) mice by Riccardin D, a natural product derived from liverwort plant *Dumortiera hirsuta*. *PLoS One* 7: e33243, 2012.