

Effects of DLX2 overexpression on the osteogenic differentiation of MC3T3-E1 cells

HAO SUN*, ZHIXU LIU*, BIAO LI, JIEWEN DAI and XUDONG WANG

Department of Oral and Cranio-maxillofacial Surgery, Shanghai Ninth People's Hospital,
Shanghai Jiao Tong University School of Medicine, Shanghai 200011, P.R. China

Received June 5, 2014; Accepted January 30, 2015

DOI: 10.3892/etm.2015.2378

Abstract. Distal-less genes (DLX) play important roles in regulating organism development. DLX2 is crucial for the differentiation and development of the primordium, which determines the subsequent development and phenotype of the maxillofacial skeletal patterns, and is the primary candidate gene that regulates the development of the first branchial arch. The aim of the present study was to investigate the effects of DLX2 overexpression on the osteogenic differentiation of MC3T3-E1 cells *in vitro*. A DLX2-expression retrovirus vector was constructed by subcloning with a murine stem cell virus (MSCV) and verified by sequencing. MC3T3-E1 cells were transfected with pMSCV-DLX2 and stable clones were selected with puromycin. The mRNA and protein expression levels of DLX2 were determined using quantitative polymerase chain reaction (PCR) and western blot analysis, respectively. In addition, the expression levels of the osteogenic biomarkers, alkaline phosphatase (ALP), osteocalcin (OCN), runt-related transcription factor (RUNX)2 and Msh homeobox (MSX)2, were assessed by quantitative PCR. ALP detection and Alizarin red staining were conducted to evaluate the effect of DLX2 overexpression on osteogenic differentiation. The data were analyzed by analysis of variance using the Student-Newman-Keuls method. Successful pMSCV-DLX2 construction, as verified by direct sequencing, enabled DLX2

overexpression *in vitro*. Enhanced ALP activity and Alizarin red staining were observed in the MC3T3-E1-DLX2 cells when compared with the control group. During osteogenic induction, DLX2 overexpression was demonstrated to upregulate ALP and MSX2 expression at the early stage and OCN expression at the late stage, while no statistically significant difference was observed in RUNX2 expression when compared with the control group. Therefore, DLX2 overexpression *in vitro* induced the osteogenic differentiation of MC3T3-E1 cells via upregulating bone formation-associated genes, such as ALP and MSX2.

Introduction

Distal-less genes (DLX) are divergent homeobox genes of the Hox gene family, including DLX1-6, which play important roles in regulating organism development (1,2). DLX2 overexpression occurs in the middle area of the first branchial arch, namely the maxillary process, between days 9.5 and 10.5 of mouse embryo development. This overexpression is crucial to the differentiation and development of the primordium, as demonstrated by experimental embryology, since it determines the subsequent development and phenotype of the maxillofacial skeletal patterns (3). Thereafter, DLX2 is the primary candidate gene that regulates the development of the first branchial arch (4). Craniofacial bone defects, cranial parietal bone ossification delay and cortical bone dysplasia in long bones have been observed in DLX2 knockout mouse models (5,6), indicating that the main function of DLX2 is to regulate the proper migration of ectomesenchymal cells and the normal osteogenic differentiation of preosteoblast cells. However, the function of DLX2 in regulating osteoblast differentiation and the underlying molecular mechanism remain controversial (7-9).

Therefore, the aim of the present study was to investigate the effect of DLX2 on osteogenic differentiation, and the possible underlying molecular mechanism, through *in vitro* transfection of a murine stem cell virus containing DLX2 (pMSCV-DLX2) into the preosteoblast cell line, MC3T3-E1.

Materials and methods

Materials. An MC3T3-E1 cell line was purchased from the Cell Bank of the Chinese Academy of Sciences (Shanghai,

Correspondence to: Professor Xudong Wang, Department of Oral and Cranio-maxillofacial Surgery, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, 639 Zhizaoiu Road, Shanghai 200011, P.R. China
E-mail: xudongwang70@hotmail.com

*Contributed equally

Abbreviations: DLX, distal-less gene; ALP, alkaline phosphatase; OCN, osteocalcin; RUNX2, runt-related transcription factor 2; MSX2, Msh homeobox 2; BMP, bone morphogenetic protein; MSCV, murine stem cell virus; PCR, polymerase chain reaction; FCS, fetal calf serum; PNPP, p-nitrophenylphosphate; MEM, Minimal Essential Medium; PBS, phosphate-buffered saline; OD, optical density

Key words: distal-less gene 2, osteogenic differentiation, MC3T3-E1 cells

China). Fetal calf serum (FCS) was obtained from the Beyotime Institute of Biotechnology (Haimen, China) and α -Minimal Essential Medium (MEM) was purchased from Gibco Life Technologies (Carlsbad, CA, USA). α -MEM osteogenesis induction culture medium (α -MEM plus 1.0×10^{-7} mol/l dexamethasone, 2.5×10^{-4} mol/l vitamin C and 1.0×10^{-2} mol/l β -sodium glycerophosphate) and α -nitrophenylphosphate (PNPP) were purchased from Sigma-Aldrich (St. Louis, MO, USA). A 3180P INC Heraeus Hera Cell CO₂ incubator was obtained from Heraeus Holding GmbH (Hanau, Germany) and a Leica DM IRB inverted phase contrast microscope was purchased from Leica Microsystems GmbH (Wetzlar, Germany). The Z2 Cell Counter was purchased from Beckman Coulter (Brea, CA, USA). A Prime Script™ RT Reagent Kit and SYBR Premix Ex Taq™ were purchased from Takara Biotechnology Co., Ltd. (Dalian, China). CAGGS/DLX2 was obtained from Dr Rubenstein (University of California, San Francisco, CA, USA).

Establishment of pMSCV-puro-DLX2. Restriction enzyme identification with *Xho*I and *Eco*RI (Takara Biotechnology Co., Ltd.) was applied to pCAGGS/DLX2 and pMSCV-puro (GeneCopoeia, Rockville, MD, USA), respectively. The target fragments were collected through gel extraction according to the manufacturer's instructions (QIAquick Gel extraction kit, Qiagen, Hilden, Germany) and stored at 4°C overnight. pMSCV-puro and DLX2 were subsequently combined by T4 ligase (Takara Biotechnology Co. Ltd.) at 16°C overnight and sequenced.

293FT viral packaging. Transfection was performed when the cell density of the 293FT cell line reached 80–90%. The 293FT cell line was used to construct the retrovirus. Vesicular stomatitis virus glycoprotein (10 μ g), GAG-pol (15 μ g) and pMSCV (15 μ g) were prepared for the retroviral packaging system. At 4–6 h after transfection, the culture medium in the T75 flask was replaced with 10 ml α -MEM containing 10% FCS, after which the virus was collected. This was done by collecting the pseudovirus-containing medium from the 293FT cell culture into sterile capped tubes for 48 h post-transfection. The tubes were then centrifuged at 550 x g for 10 min to remove cell debris and the supernatant was filtered through 45 μ m polyethersulfone low protein-binding filters. The virus pellet was resuspended in phosphate-buffered saline (PBS) and stored at 4°C.

MC3T3-E1 transfection. The virus used for transfection was determined through dosage titration at 1:1, 10:1, 100:1 and 1,000:1. Polybrene (1/1,000; Sigma-Aldrich) was added to the medium, after which the plate was centrifuged for 3 min at 300 x g at 37°C. The culture medium was replaced 24 h after transfection. Following incubation in the culture medium for 48 h, 3 μ g/ml puromycin was applied for growth selection. The cells that survived the selection were collected and passaged. Controls included pMSCV (dosage titration at 1:1, 10:1 and 100:1).

DLX2 expression detection with quantitative polymerase chain reaction (PCR) and western blot analysis. At 72 h after puromycin selection, the total RNA was extracted for quantita-

tive PCR. This was performed using the SYBR Green system (Takara Bio, Inc, Shiga, Japan) with a total volume of 20 μ l in a 96-well microwell plate. The reaction consisted of 10 μ l SYBR Premix Ex Taq (Takara Biotechnology Co. Ltd.), 0.8 μ l mixed primers, 1 μ l cDNA template and 8.2 μ l RNase-free distilled water. The PCR program was set at 95°C for 4 min followed by 40 cycles at 95°C for 20 sec, 60 °C for 15 sec and 72 °C for 20 sec. The threshold cycle values were calculated using the iQ5 software (Bio-Rad, Hercules, CA, USA). All the primers used in the experiments are shown in Table I. The analysis of the data was based on calculating the relative expression levels of these genes compared to the expression of the control (glyceraldehyde 3-phosphate dehydrogenase, a housekeeping gene).

Cells were collected, washed 3 times with pre ice-cold PBS and lysed in SDS-lysis buffer containing protease inhibitors (Roche Diagnostics GmbH, Mannheim, Germany) for 20 min on ice. The lysates were centrifuged at 12,000 x g at 4°C for 30 min and the supernatants were boiled in sodium dodecyl sulfate sample buffer containing 0.5 M of mercaptoethanol. The samples were then separated on a 10% sodium dodecyl sulfate polyacrylamide gel and transferred to a polyvinylidene difluoride membrane using a semi-dry transfer apparatus (Bio-Rad, Hercules, CA, USA). The membrane was blocked with 5% milk for 1 h and then incubated with primary antibodies in modified D-PBS tween-20 buffer overnight. The primary antibodies were anti-DLX2 (1:800; Abcam, Cambridge, UK) and β -actin (1:1,000; Sigma-Aldrich). Following the primary antibody incubation, they were incubated with horse-radish peroxidase-conjugated secondary antibodies, including goat anti-rabbit and goat anti-mouse IgG (1:5,000; Beyotime Institute of Biotechnology, Haimen, China). The protein bands were visualised using an enhanced chemiluminescent western blot analysis kit (Pierce Biotechnology, Inc., Rockford, IL, USA).

Detection of DLX2 expression in the stable cell line following transfection. Following transfection and selection, the stable cell line was seeded into a six-well plate at a cell density of 1.5×10^4 /cm². α -MEM osteogenesis induction culture medium was added to the wells and the cells were cultured in conditions of 5% CO₂ at 37°C. Cells were collected and protein samples were obtained on days 1, 4, 7 and 14 for western blot analysis.

Analysis of effect of DLX2 overexpression on osteogenesis-associated genes via quantitative PCR. Stable cell lines, following transfection and selection, were seeded into a six-well plate at a cell density of 1.5×10^4 /cm². The wells contained α -MEM osteogenesis induction culture medium and the cells were cultured in 5% CO₂ at 37°C. Cells were collected and the total RNA was extracted on days 1, 4, 7 and 14 for quantitative PCR. The primers used for the PCR assays of ALP, osteocalcin (OCN), runt-related transcription factor (RUNX)2 and MSX2 are included in Table I.

ALP activity detection. Stable cell lines, following transfection and selection, were seeded into a 24-well plate at a cell density of 1.5×10^4 /cm². α -MEM osteogenesis induction culture medium was added to the wells and the cells were cultured in 5% CO₂ at 37°C. ALP activity detection was performed on days 4,

Table I. Primers for used for quantitative polymerase chain reaction.

Gene	Forward sequence	Reverse sequence
DLX2	5'-CATGGGCTCCTACCAGTACCAC-3'	5'-TCGGATTTCAGGCTCAAGGTC-3'
GAPDH	5'-GGTGAAGGTGGTGTGAACG-3'	5'-CTCGTCTCCTGGAAGATGGTG-3'
ALP	5'-TGGGCATTGTGACTACCACTCGG-3'	5'-CCTCTGGTGGCATCTCGTTATCC-3'
OCN	5'-GGACCATCTTTCTGCTCACTCTG-3'	5'-GTTCACTACCTTATTGCCCTCCTG-3'
RUNX2	5'-AACTTCCTGTGCTCCGTGCTG-3'	5'-TCGTTGAACCTGGCTACTTGG-3'
MSX2	5'-GGAGCACCGTGGATACAGGA-3'	5'-AGGCTAGAAGCTGGGATGTGG-3'

DLX, distal-less gene; ALP, alkaline phosphatase; OCN, osteocalcin; RUNX2, runt-related transcription factor; MSX, Msh homeobox.




Figure 1. Analysis of DLX2 mRNA expression levels in the MC3T3-E1 cell line using quantitative polymerase chain reaction. DLX, distal-less gene; MSCV, murine stem cell virus. GAPDH, glyceraldehyde 3-phosphate dehydrogenase.




Figure 2. Analysis of DLX2 protein expression levels in the MC3T3-E1 cell line using western blot analysis. DLX, distal-less gene; MSCV, murine stem cell virus.

7 and 14 using PNPP. The cells were washed with PBS and lysed with ALP buffer containing 0.2% Triton X-100, 42.74 mg $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 20 μl HCl (10 mol/l), 19.4 ml diethylamine and 40 mg sodium azide dissolved in 200 ml distilled water. After 4 h in the incubator at 37°C, 190 μl supernatant was collected and added to a 96-well plate containing 190 μl substrate buffer per well. The optical density (OD) value was measured at 405 nm following incubation for 30 min at 37°C. A 10- μl sample of the supernatant was added to 200 μl protein analysis buffer and the OD value at 630 nm was detected after 5 min. The intracellular protein content was determined using a Micro BCA protein assay kit (Thermo Fisher Scientific, Waltham, MA, USA) and applied for the standard curve. The alkaline phosphatase activity was determined by measuring the optical density values for absorbance at 405 nm after incubation with *p*-nitrophenyl phosphate for 30 min at 37°C (Sigma-Aldrich).

Alizarin red staining. Alizarin red staining was used in a biochemical assay to determine, quantitatively by colorimetry,

the presence of calcific deposition by cells of an osteogenetic lineage. Following transfection and selection, the stable cell lines were seeded into a six-well plate at a cell density of $1.5 \times 10^4/\text{cm}^2$ with α -MEM osteogenesis induction culture medium. The cells were cultured in conditions of 5% CO_2 at 37°C. Alizarin red staining (Sigma-Aldrich) was performed on days 3, 21 and 27, however, the stain was not that notable amongst the blank, pMSCV and pMSCV-Dlx2 groups. Each group was scanned for detection by a Canon LiDE 110 scanner (Canon, Beijing, China) following the staining.

Statistical analysis. Data were analyzed by analysis of variance using the Student-Newman-Keuls method. Statistical analysis was performed using SAS 6.04 software package (SAS Institute, Cary, NC, USA), where $P < 0.05$ was considered to indicate a statistically significant difference.

Results

DLX2 expression in the stably transfected cell line. At 72 h after puromycin selection, mRNA expression of DLX2 was observed in the pMSCV-DLX2 group, but not in the pMSCV or non-transfection groups (Fig. 1). As shown in the western blot analysis, a 33-kDa band presented in the pMSCV-DLX2 group; however, this band was not observed in the pMSCV or non-transfection groups (Fig. 2). These observations indicated that the transfection was successful and expression of DLX2 had been induced.

DLX2 expression in the stable clones selected with osteogenesis induction. On day 1 following osteogenesis induction, DLX2 expression was detected in the pMSCV-DLX2 group, which was shown to remain stable until day 14 post-transfection. Increasing endogenous DLX2 expression was observed in the pMSCV group (Fig. 3).

Effect of DLX2 overexpression on the expression levels of osteogenesis-associated genes. Expression levels of osteogenesis-associated genes are shown in Fig. 4. The relative expression levels were calculated, as compared with the control group on day 1, and are outlined in Table II. Statistically significant differences between the control and experimental cells were observed in the expression of ALP ($P < 0.01$), OCN ($P < 0.05$, on day 14) and MSX2 ($P < 0.01$), but not in the expression of RUNX2 ($P > 0.05$).




Figure 3. Western blot analysis showing DLX2 protein expression levels following osteogenic induction in pMSCV-DLX2 stably transfected cells. DLX, distal-less gene; MSCV, murine stem cell virus; D, day.




Figure 4. Quantitative polymerase chain reaction analysis showing the relative mRNA expression levels of the bone formation-associated genes, ALP, OCN, RUNX2 and MSX2. Values were normalized against the mRNA expression level of the control group on day 1. * $P < 0.05$, pMSCV-DLX2 group vs. pMSCV group ($n = 3$). DLX, distal-less gene; ALP, alkaline phosphatase; OCN, osteocalcin; RUNX2, runt-related transcription factor; MSX, Msh homeobox; MSCV, murine stem cell virus.

ALP activity. Higher ALP activity levels were observed on days 4, 7 and 14 after osteogenesis induction in the transfection group (pMSCV-DLX2), and statistically significant differences were observed when compared with the control group ($P < 0.01$). The highest activity was observed on day 14, as shown in Fig. 5. The ALP activity levels are listed in Table III.

Alizarin red staining. Positive results from the Alizarin red staining were observed on day 21 following osteogenesis induction, as shown in Fig. 6. The results for staining at day 27 were almost the same to those on day 21 and are therefore not discussed. In the DLX2 transfection group, the alizarin red staining was used to show the calcium nodule which indi-

cated that DLX2 promoted the osteogenic differentiation of MC3T3-E1 cells.

Discussion

DLX are divergent homeobox genes that are part of the Hox gene family, which are involved in the regulation of maxillofacial bone development. However, the function of DLX remains controversial and the regulatory mechanisms are yet to be elucidated. In two previous studies, Qiu *et al* (5,6) created a DLX2^{-/-} mouse model through gene knockout technology. Newborn DLX2^{-/-} mice were found to die after birth with structural abnormalities observed in the bones originating from the first branchial arch maxillary process,

Table II. Quantitative polymerase chain reaction analysis of the relative expression levels of bone formation-associated genes.

Gene	Day 1	Day 4	Day 7	Day 14
ALP				
pMSCV	1	1.18±0.045	3.719±0.23	7.971±0.32
pMSCV-DLX2	1.193±0.076	3.387±0.172	7.93±0.223	9.117±0.305
OCN				
pMSCV	1	1.084±0.058	3.79±0.271	5.56±0.217
pMSCV-DLX2	1.013±0.075	1.057±0.087	3.873±0.215	6.482±0.035
RUNX2				
pMSCV	1	3.104±0.083	2.86±0.395	3.626±0.217
pMSCV-DLX2	1.081±0.104	3.209±0.149	3.02±0.308	3.481±0.03
MSX2				
pMSCV	1	1.104±0.083	2.882±0.313	3.54±0.046
pMSCV-DLX2	8.901±0.223	12.676±0.641	13.02±0.308	13.615±0.236

Statistically significant differences between the control and experimental groups were observed in the expression of ALP ($P<0.01$, on days 4, 7, 14), OCN ($P<0.05$, on day 14) and MSX2 ($P<0.01$, on day 1, 4, 7, 14), but not in the expression of RUNX2 ($P>0.05$). DLX, distal-less gene; ALP, alkaline phosphatase; OCN, osteocalcin; RUNX2, runt-related transcription factor; MSX, Msh homeobox; MSCV, murine stem cell virus.

Table III. Semi-quantitative analysis of ALP activity.

Group	Day 4	Day 7	Day 14
pMSCV	0.521±0.089	0.668±0.066	2.041±0.094
pMSCV-DLX2	1.673±0.073	2.561±0.087	2.881±0.097

A statistically significant difference was observed when compared with the control group ($P<0.01$). DLX, distal-less gene; ALP, alkaline phosphatase; MSCV, murine stem cell virus.





Figure 6. Alizarin red staining of MC3T3-E1 cells on day 21. DLX, distal-less gene; MSCV, murine stem cell virus.

Figure 5. Semi-quantitative analysis of ALP activity. * $P<0.05$, pMSCV-DLX2 group vs. pMSCV group (n=3). ALP, alkaline phosphatase; MSCV, murine stem cell virus; OD, optical density; DLX, distal-less gene.

such as the basisphenoid, greater wing of the sphenoid bone and pterygoid lamina, which was accompanied by a cleft palate, cranial parietal bone ossification delay and cortical

bone dysplasia in long bones. These observations indicated that DLX2 was crucial in regulating craniofacial bone development and differentiation. In addition, gene expression microarray analysis revealed that DLX2 is an early response gene in the regulation of bone morphogenetic protein (BMP)2-mediated osteogenic differentiation (10). Therefore, the DLX knockout mouse models and DLX overexpression experiments indicated that DLX promotes skeleton formation; however, the effect of DLX on osteogenic differentiation remains unknown. Lee *et al* (11) found that DLX5, which is closely associated to DLX2, suppressed the expression of OCN, in contrast to DLX2. In addition, overexpression of DLX5 was reported to result in bone formation disorder in immunodeficient mice and mineralized matrix deposition of cells cultured *in vitro*. *In vitro*, overexpression of DLX5 was shown to have no effect on chondroblast differentiation (12). In the present study, MC3T3-E1 cells were transfected with pMSCV-DLX2 and stable clones were selected with puromycin. Subsequently, the mRNA and protein expression levels of DLX2 were determined by quantitative PCR and western

blot analysis, respectively. This stable transfection cell line established the basis for investigating the effect of DLX2 overexpression on osteogenic differentiation and the possible underlying mechanisms.

ALP is hypothesized to be an indicator of early osteogenesis (13), and is known to hydrolyze numerous types of phosphates under alkaline conditions to promote cell maturation and calcification (14). ALP has been shown to be highly expressed during early osteogenic differentiation (day 7) through induction by morphogenetic proteins (15,16). In the present study, ALP activity in the MC3T3-E1 cell line was upregulated in the early stage of osteogenesis between days 7 and 14, as induced by the osteogenesis culture medium. As indicated in the quantitative PCR and ALP activity experiments, DLX2 upregulated ALP activity as early as day 4. On day 14, no statistically significant difference was observed between the groups, indicating that DLX2 primarily functions in the early stages of osteogenic differentiation, which is different from that of BMPs (15). OCN is the most abundant non-procollagen protein in the bone tissue, and can promote osteogenic differentiation by combining with minerals (17). In osteoblasts cultured *in vitro*, OCN upregulation has been observed during the mineralization stage; however, OCN expression has subsequently decreased until the end of osteogenic differentiation (13,18,19). The results of the present study demonstrated that low expression levels of OCN were observed in each group during the early stage of osteogenesis on days 1, 4 and 7, and no statistically significant differences were observed among the groups. However, on day 14, higher expression levels of OCN were observed in the transfection group ($P < 0.05$). In addition, positive Alizarin red staining indicated that DLX2 promoted the expression of OCN and osteogenesis at the later stage.

RUNX2 belongs to the Runt domain gene family and has been demonstrated to be the one of the most important transcription factors involved in osteogenic differentiation (20). However, no statistically significant difference in RUNX2 expression was observed at any of the four time points when comparing the transfection and control groups ($P > 0.05$), which indicated that the RUNX2 pathway may not be involved in the mechanism underlying DLX2-induced promotion of MC3T3-E1 osteogenesis. MSX2 belongs to the Hox gene family and is involved in the regulation of osteogenesis. Downregulation of MSX2 has been shown to result in preosteoblast dysmaturity (21). DLX genes have been reported to function in transforming growth factor- β (TGF- β)/BMP-mediated osteogenesis regulation through combining with and antagonizing MSX2 (22,23). In the present study, overexpression of DLX2 was shown to upregulate MSX2 expression. Thus, it was hypothesized that DLX2 and MSX2 regulate the transcription of osteogenesis-associated genes in a synergistic manner by forming a transcription factor complex. However, future investigation is required to further support this hypothesis.

In conclusion, through DLX2 transfection, examination of osteogenesis-associated gene expression, ALP activity and Alizarin red staining, the present study demonstrated that DLX2 overexpression induces the osteogenic differentiation of MC3T3-E1 cells via upregulating bone formation-associated genes, such as ALP and MSX2. Future studies will focus on

osseointeractions between DLX2 and MSX2. Further studies will aim to investigate the role of the TGF- β signalling pathway on the induction of osteogenesis by DLX2. Such a study may contribute to potential clinical application of bone mesenchymal stem cells transduce with Dlx2 for bone augmentation in patients suffering from bone deficiency.

Acknowledgements

This study was supported by grants from the Combined Engineering and Medicine Project of Shanghai Jiao Tong University (no. YG2014QN02) and the National Nature Science Foundation of China (no. 81271122).

References

1. Depew MJ, Simpson CA, Morasso M and Rubenstein JL: Reassessing the Dlx code: the genetic regulation of branchial arch skeletal pattern and development. *J Anat* 207: 501-561, 2005.
2. Jin Y: *Mouse Development Biology and Embryo Research Method*. Volume 2, 1st edition. People's Medical Publishing House, Beijing, pp70-73, 2005.
3. Ferguson CA, Tucker AS and Sharpe PT: Temporospatial cell interactions regulating mandibular and maxillary arch patterning. *Development* 127: 403-412, 2000.
4. Depew MJ, Lufkin T and Rubenstein JL: Specification of jaw subdivisions by Dlx genes. *Science* 298: 381-385, 2002.
5. Qiu M, Bulfone A, Ghattas I, *et al*: Role of the Dlx homeobox genes in proximodistal patterning of the branchial arches: mutations of Dlx-1, Dlx-2 and Dlx-1 and -2 alter morphogenesis of proximal skeletal and soft tissue structures derived from the first and second arches. *Dev Biol* 185: 165-184, 1997.
6. Qiu M, Bulfone A, Martinez S, *et al*: Null mutation of Dlx-2 results in abnormal morphogenesis of proximal first and second branchial arch derivatives and abnormal differentiation in the forebrain. *Genes Dev* 9: 2523-2538, 1995.
7. Merlo GR, Zerega B, Paleari L, *et al*: Multiple functions of Dlx genes. *Int J Dev Biol* 44: 619-626, 2000.
8. Panganiban G and Rubenstein JL: Developmental functions of the Distal-less/Dlx homeobox genes. *Development* 129: 4371-4386, 2002.
9. Li H, Marijanovic I, Kronenberg MS, *et al*: Expression and function of Dlx genes in the osteoblast lineage. *Dev Biol* 316: 458-470, 2008.
10. Harris SE, Guo D, Harris MA, *et al*: Transcriptional regulation of BMP-2 activated genes in osteoblasts using gene expression microarray analysis: role of Dlx2 and Dlx5 transcription factors. *Front Biosci* 8: s1249-s1265, 2003.
11. Lee MH, Kwon TG, Park HS, *et al*: BMP-2-induced Osterix expression is mediated by Dlx5 but is independent of Runx2. *Biochem Biophys Res Commun* 309: 689-694, 2003.
12. Muraglia A, Perera M, Verardo S, *et al*: DLX5 overexpression impairs osteogenic differentiation of human bone marrow stromal cells. *Eur J Cell Biol* 87: 751-761, 2008.
13. Karsenty G and Wagner EF: Reaching a genetic and molecular understanding of skeletal development. *Dev Cell* 2: 389-406, 2002.
14. Beck GR Jr, Zerler B and Moran E: Phosphate is a specific signal for induction of osteopontin gene expression. *Proc Natl Acad Sci USA* 97: 8352-8357, 2000.
15. Hu Z, Peel SA, Ho SK, *et al*: Role of bovine bone morphogenetic proteins in bone matrix protein and osteoblast-related gene expression during rat bone marrow stromal cell differentiation. *J Craniofac Surg* 16: 1006-1014, 2005.
16. Vaes BL, Decherin KJ, Feijen A, *et al*: Comprehensive microarray analysis of bone morphogenetic protein 2-induced osteoblast differentiation resulting in the identification of novel markers for bone development. *J Bone Miner Res* 17: 2106-2118, 2002.
17. Glowacki J, Rey C, Glimcher MJ, *et al*: A role for osteocalcin in osteoclast differentiation. *J Cell Biochem* 45: 292-302, 1991.
18. Hu JZ, Jiang XQ, Zhang ZY, *et al*: Effects of Nell-1 gene on bone formation-related gene expression during osteoblastic differentiation of rat bone marrow stromal cells by real-time PCR. *Zhongguo Kou Qiang He Mian Wai Ke Za Zhi* 6: 48-53, 2008 (In Chinese).

19. Hu JZ, Jiang XQ, Zhang ZY and Zhang XL: Effect of Nell-1 gene on osteogenic differentiation of rat bMSCs: An in vitro study. *Zhongguo Kou Qiang He Mian Wai Ke Za Zhi* 7: 38-43, 2009 (In Chinese).
20. Li P, Yu SH, Chen D, *et al*: Overexpression of Runx2 induces osteogenic differentiation in C2C12 cells. *Zhongguo Sheng Wu Hua Xue Yu Fen Zi Sheng Wu Xue Bao* 26: 236-242, 2010 (In Chinese).
21. Cheng SL, Shao JS, Charlton-Kachigian N, *et al*: MSX2 promotes osteogenesis and suppresses adipogenic differentiation of multipotent mesenchymal progenitors. *J Biol Chem* 278: 45969-45977, 2003.
22. Bendall AJ and Abate-Shen C: Roles for Msx and Dlx homeoproteins in vertebrate development. *Gene* 247: 17-31, 2000.
23. Ryoo HM, Lee MH and Kim YJ: Critical molecular switches involved in BMP-2-induced osteogenic differentiation of mesenchymal cells. *Gene* 366: 51-57, 2006.