

In vitro gene expression profile of bovine peripheral blood mononuclear cells in early *Mycobacterium bovis* infection

YAFEN CHENG^{1,2}, CHUNG-HSI CHOU¹ and HSIANG-JUNG TSAI^{1,3}

¹School of Veterinary Medicine, National Taiwan University, Taipei 10617; ²Centers for Disease Control, Ministry of Health and Welfare, Taipei 10050; ³Animal Health Research Institute, Council of Agriculture, Executive Yuan, New Taipei 25118, Taiwan, R.O.C.

Received September 12, 2014; Accepted July 29, 2015

DOI: 10.3892/etm.2015.2814

Abstract. The intracellular parasite *Mycobacterium bovis* (*M. bovis*) causes tuberculosis in cattle and humans. Understanding the interactions between *M. bovis* and host cells is essential in developing tools for the prevention, detection, and treatment of *M. bovis* infection. Gene expression profiles provide a large amount of information regarding the molecular mechanisms underlying these interactions. The present study analyzed changes in gene expression in bovine peripheral blood mononuclear cells (PBMCs) at 0, 4 and 24 h following exposure to *M. bovis*. Using bovine whole-genome microarrays, a total of 420 genes were identified that exhibited significant alterations in expression (≥ 2 -fold). Significantly enriched genes were identified using the Kyoto Encyclopedia of Genes and Genomes database, of which the highest differentially expressed genes were associated with the immune system, signal transduction, endocytosis, cellular transport, inflammation, and apoptosis. Of the genes associated with the immune system, 84.85% displayed downregulation. These findings support the view that *M. bovis* inhibits signaling pathways of antimycobacterial host defense in bovine PBMCs. These *in vitro* data demonstrated that molecular alterations underlying the pathogenesis of tuberculosis begin early, during the initial 24 h following *M. bovis* infection.

Introduction

Bovine tuberculosis (BTB) is caused by the intracellular pathogen *Mycobacterium bovis* (*M. bovis*), which is a facultative intracellular parasite of macrophages. BTB has a significant economic impact and serious implications for human health, particularly in developing countries (1). *M. bovis* can be trans-

mitted to humans by infectious bacilli via respiratory contact with infected cattle, or consumption of unpasteurized dairy products (2). The host immune response to *M. bovis* infection is complex: Following initial exposure, T-helper cell-1 (T_H1) innate immunity is induced. The bacilli are phagocytosed by host macrophages via pathogen-recognition receptors (PRRs), such as Toll-like receptors (TLRs) and C-type lectin receptors (3,4). Signals transduced through these receptors result in the release of endogenous cytokines, which initiate the T-cell secretion of proinflammatory cytokines, including tumor necrosis factor- α (TNF- α) and interferon- γ (IFN- γ). The action of IFN- γ on infected macrophages promotes granuloma formation, which prevents the spread of infection (5). However, the pathogen often persists within granulomas, and this latent infection can recur as active tuberculosis. The mechanisms underlying evasion of the host immune response by *M. bovis* are not entirely understood; however they are known to involve the prevention of host phagosome maturation, the inhibition of apoptosis in infected macrophages, and the suppression of cell signaling pathways and cytokine production (6-8).

Genomic technologies can be used to elucidate the molecular mechanisms underlying immune responses to pathogens. With the availability of the complete *Bos taurus* genome (9), numerous studies have used bovine genome microarrays to analyze transcriptional changes induced by infection of various types of bovine cells with *M. bovis* (10,11). Killick *et al* (12) reported that *M. bovis* infection of peripheral blood leukocytes was associated with decreased expression levels of numerous host genes. Using an Affymetrix bovine genome array to investigate the effects of *M. bovis* challenge on bovine monocyte-derived macrophages *in vitro*, Magee *et al* (6) observed significant alterations in expression of genes associated with the inflammatory response, and cell signaling pathways, including TLRs, PRRs and apoptosis. Furthermore, the suppression of immune-associated genes has been detected *in vivo* in *M. bovis*-infected cattle (10).

These observations strongly suggest that *M. bovis* evades immune surveillance by altering the expression of genes essential to host immunity. The timing and potency of the cellular and immunological events that occur immediately after infection are suggested to be crucial determinants governing the outcome of an infection (13). Therefore, further elucidation of these early events in numerous types of host

Correspondence to: Dr Hsiang-Jung Tsai, School of Veterinary Medicine, National Taiwan University, 1 Roosevelt Road, Taipei 10617, Taiwan, R.O.C.
E-mail: tsaihj@ntu.edu.tw

Key words: *Mycobacterium bovis*, global gene expression, immune response, macrophage

cells is essential for the prevention, detection, and treatment of *M. bovis* infections. The aim of the present study was to evaluate early changes in global gene expression in bovine peripheral blood mononuclear cells (PBMCs) in response to *M. bovis* exposure. Microarray analyses were used to compare PBMC gene expression over a time course of 0, 4, and 24 h following exposure to *M. bovis*. Systems analysis was then used to determine the pathways and networks associated with the affected genes.

Materials and methods

PBMC preparation. The three 3-year-old female Holstein cattle with no recent history of BTB, which were used in the present study, were obtained from the National Taiwan University Experimental Farm (Taipei, Taiwan, R.O.C). The cattle were maintained under uniform housing conditions (temperature, 25–28°C; humidity, 50–70%) and nutritional regimens; the cattle were fed twice a day with alfalfa and pangola grass hay and fresh, farm-grown grass, and all tested negative on tuberculin skin tests. All procedures described in the present study were reviewed and approved by the National Taiwan University Institutional Animal Care and Use Committee (Taipei, Taiwan, R.O.C). From each animal, 50 ml blood from the vein was collected in a sterile heparinized bottle and layered onto ACCUSPIN™ tubes containing Histopaque® 1077 (Sigma-Aldrich, St. Louis, MO, USA). Following density gradient centrifugation (300 x g for 20 min at room temperature), the PBMCs were collected and cultured as previously described by Magee *et al* (6), with numerous modifications, including the use of an antibiotic-free culture media supplemented with NaHCO₃ and 10% fetal bovine serum. Cell cultures in 60 mm dishes (~5x10⁶ cells/dish), were grown in RPMI (Gibco Life Technologies, Grand Island, NY, USA) supplemented with 10% fetal bovine serum (GE Healthcare Life Sciences, Logan, UT, USA) and NaHCO₃ to a final concentration of 26 mM at 37°C for 24 h in a culture incubator containing 5% CO₂. The medium was then replaced with 1 ml fresh medium in order to remove any non-adherent cells. To ensure that the same number of PBMCs were subjected to *M. bovis*-challenge, 80–100% confluent monolayers of PBMCs were generated and counted on day 3, yielding ~5x10⁶ cells per dishes, providing enough total RNA for microarray analysis. The cells were counted in Marienfeld® Thoma counting chamber (Celeromics Technologies, Grenoble, France), and trypan blue solution (Sigma-Aldrich, St. Luis, MO, USA) was used for the exclusion of dead cells.

Bacterial preparations. *M. bovis* strain 331-A1 (Animal Health Research Institute, New Taipei City, Taiwan, R.O.C), which was isolated in 2008 from cattle with tuberculosis, was used. The strain was confirmed by acid-fast staining using the Ziehl-Neelsen method, the gold standard procedure for the diagnosis of tuberculosis, and identified in mycobacterium culture by PCR and spoligotyping (14). The strain used was spoligotype SB0265, which is frequently isolated from cattle in Taiwan. The strain was cultured in Middlebrook 7H9 broth (BD Biosciences, Franklin Lakes, NJ, USA) containing 10% (v/v) Middlebrook albumin-dextrose-catalase (BD Biosciences), 0.05% Tween 80 (Sigma-Aldrich) and

0.40% (w/v) sodium pyruvate (Sigma-Aldrich) at 37°C. Bacterial suspensions were then centrifuged at 3,000 x g for 15 min at 25°C, and the pellets were washed twice in phosphate-buffered saline (PBS, pH 7.0), and resuspended in PBS prior to the determination of bacterial concentrations. A NanoDrop 1000 spectrophotometer (Thermo Fisher Scientific, Inc., Wilmington, DE, USA) was used to determine the optical density of the bacterial culture and calculate bacterial concentration in colony-forming u/ml. The concentration of the bacterial inoculum was 10⁷ colony-forming units/ml. Fresh stocks of bacteria were prepared for each experiment.

In vitro challenge of PBMCs with *M. bovis*. When the PBMCs reached 80% confluence, three dishes for each *M. bovis* infection trial were randomly selected from the 20 dishes grown from cells collected from each cow. All *in vitro* challenge experiments included a non-challenge PBMC control for each time point. The cells were inoculated at a multiplicity of infection (MOI) of ~2:1 (6), and control cultures were treated in the same manner using PBS instead of bacterial suspensions. Inoculated and control cultures were incubated for either 4 h (4-hours post-infection (hpi) group) or 24 h (24-hpi group). Total RNA was then extracted using the RNeasy Mini kit (Qiagen, Inc., Valencia, CA, USA), according to the manufacturer's instructions. The concentration and purity of RNA extracts were verified optically using a spectrophotometer (ND-1000; Nanodrop Technologies, Thermo Fisher Scientific, Inc.) and the Bioanalyzer 2100 (Agilent Technologies, Inc., Santa Clara, CA, USA), respectively.

Microarray analysis. Bovine V2 Oligo 4x44 K microarrays (Agilent Technologies, Inc.) were used to determine the differential gene expression between infected and control cells. For reverse transcription, second-strand cDNA was synthesized from 0.5 µg total RNA using the Fluorescent Linear Amplification kit containing T7 RNA polymerase (Agilent Technologies, Inc.). The cDNA served as template for *in vitro* transcription to produce target cRNA labeled with Cy3-CTP (to label infected cells) and Cy5-CTP (to label control cells) (PerkinElmer, Inc., Waltham, MA, USA). Labeled cRNA (0.825 µg) was fragmented (mean size, ~50–100 nucleotides) in fragmentation buffer (Agilent Technologies, Inc.) at 60°C for 30 min. The prepared cRNA was subsequently hybridized to the microarray at 60°C for 17 h. Two replicates of the microarray assays (M1 and M2) were performed. Hybridized microarray chips were scanned using the Agilent Microarray Scanner with Feature Extraction software 9.5.3 (Agilent Technologies, Inc.). The locally weighted linear regression method was applied to normalize the results by rank consistency filtering.

Statistical analysis of microarray data. Microarray data were analyzed using GeneSpring GX 7.3.1 software (Agilent Technologies, Inc.). With a false discovery rate <0.05, data acquisition was conducted using the following criteria: i) P<0.01 for gene expression difference (GeneSpring); ii) a distinct signal from the microarray image that was flagged by the software; and iii) l-log 2-fold changel ≥2.5. Significantly enriched genes were identified using the Kyoto Encyclopedia of Genes and Genomes (KEGG; <http://www.genome.jp/kegg/>),

Table I. Genes with upregulated expression (≥ 2.5 -fold) following exposure of peripheral blood mononuclear cells to *Mycobacterium bovis*.

Processes	Symbol	Fold change			P-value	Gene name
		0-4 h	4-24 h	0-24 h		
Signal transduction	CLEC4E	1.15	7.70	8.85	0.001	C-type lectin domain family 4 member E
	STAT1	0.71	2.98	2.12	0.002	Signal transducer and activator of transcription 1
	DDIT3	0.78	2.91	2.27	0.004	DNA-damage-inducible transcript 3
	LDHA	0.77	2.81	2.16	<0.001	Lactate dehydrogenase A (LDHA)
Immune system	C3AR1	0.97	6.68	6.49	0.002	Complement component 3a receptor 1
	PDK1	0.86	6.14	5.26	0.001	Pyruvate dehydrogenase kinase, isozyme 1
	DAPPI	0.58	4.16	2.43	<0.001	Dual adaptor of phosphotyrosine and 3-phosphoinositides
	RASGRP1	0.90	3.43	3.09	<0.001	RAS guanyl releasing protein 1 (calcium and DAG-regulated)
	CD244	0.94	3.29	3.08	<0.001	CD244 molecule, natural killer cell receptor 2B4
	IL7	0.61	2.63	1.59	0.001	Interleukin-7 precursor
Endocytosis and transport	COLEC11	1.14	4.78	5.44	<0.001	Collectin sub-family member 11
	LAMP3	0.97	2.81	2.72	0.001	Lysosomal-associated membrane protein 3
	CTSL2	0.39	2.78	1.08	0.005	Cathepsin L2
	EHD1	0.90	2.58	2.33	0.001	EH-domain containing 1
Inflammation and apoptosis	CDKN2D	1.21	03.15	3.82	<0.001	Cyclin-dependent kinase inhibitor 2D (p19, inhibits CDK4)
	CCNG2	0.59	2.60	1.53	0.004	Cyclin G2
Others	ME1	0.95	6.79	6.48	0.002	Malic enzyme 1
	FBXL3	0.90	4.21	3.80	<0.001	F-box and leucine-rich repeat protein 3

Table I. Continued.

Processes	Symbol	Fold change			P-value	Gene name
		0-4 h	4-24 h	0-24 h		
	SQLE	0.97	3.52	3.41	0.003	Squalene epoxidase
	POLD4	1.01	3.34	3.37	0.003	Polymerase (DNA-directed), delta 4
	PTGES	1.12	3.13	3.52	<0.001	Prostaglandin E synthase (PTGES)
	PHOSPHO2	1.03	3.06	3.15	<0.001	Phosphatase, orphan 2
	DCK	1.03	2.99	3.07	<0.001	Deoxycytidine kinase
	PLA2G16	0.84	2.91	2.44	<0.001	Phospholipase A2, group XVI
	CSGALNACT1	1.07	2.87	3.07	0.001	Chondroitinsulfate N-acetylgalactosaminyltransferase 1
	NAPB	0.85	2.83	2.40	0.005	N-ethylmaleimide-sensitive factor attachment protein, beta
	PGAP1	1.00	2.77	2.76	<0.001	Similar to GPI deacylase
	HSD17B7	0.93	2.65	2.47	0.002	Hydroxysteroid (17-beta) dehydrogenase 7
	TRIP10	1.05	2.63	2.75	<0.001	Thyroid hormone receptor interactor 10
	FDFT1	0.92	2.62	2.41	<0.001	Farnesyl-diphosphate farnesyltransferase 1

Fold change was calculated as the mean of triplicate experiments.

Table II. Genes with downregulated expression (≥ 2.5 -fold) following exposure of peripheral blood mononuclear cells to *Mycobacterium bovis*.

Processes	Symbol	Fold change			P-value	Gene name
		0-4 h	4-24 h	0-24 h		
Signal transduction	THBS1	1.33	-27.82	-20.91	<0.001	Thrombospondin 1
	HMOX1	1.29	-10.11	-7.86	<0.001	Heme oxygenase (decycling) 1
	FST	1.19	-7.82	-6.59	<0.001	Follistatin
	FOSL1	-1.08	-5.65	-6.11	0.004	Fos-related antigen 1 (FRA-1)
	CD38	1.05	-5.64	-5.38	0.002	CD38 molecule
	GNG4	1.06	-5.21	-4.93	<0.001	Guanine nucleotide binding protein (G protein), gamma 4
	FZD4	1.20	-4.98	-4.16	<0.001	Frizzled homolog 4
	EDN1	-1.14	-4.51	-5.13	0.002	Endothelin 1
	PTAFR	1.14	-4.01	-3.51	0.005	Platelet-activating factor receptor
	MRAS	1.29	-3.64	-2.82	0.002	Muscle RAS oncogene homolog
	FOS	-1.02	-3.56	-3.64	0.002	FBJ murine osteosarcoma viral oncogene homolog
	ICOS	-1.30	-3.06	-3.99	<0.001	Inducible T-cell co-stimulator
	FN1	1.20	-2.87	-2.40	0.004	Fibronectin 1
	NUMBL	1.24	-2.83	-2.28	0.002	Numb homolog (Drosophila)-like
	CACNG4	1.25	-2.79	-2.24	0.001	Calcium channel, voltage-dependent, gamma subunit 4
	PTGS2	-1.13	-2.71	-3.06	<0.001	Prostaglandin-endoperoxide synthase 2
	TCF7L2	1.21	-2.51	-2.08	<0.001	Transcription factor 7-like 2 (T-cell specific, HMG-box)
Immune system	C1QB	1.21	-26.31	-21.76	0.002	Complement component 1, q subcomponent, B chain
	PPBP	-1.07	-17.11	-18.37	<0.001	Pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)
	CFB	-1.07	-16.33	-17.52	<0.001	Complement factor B (CFB)
	IFNG	1.02	-14.28	-14.06	<0.001	Interferon, gamma
	THBD	-1.00	-14.27	-14.32	<0.001	Thrombomodulin
	GZMB	1.13	-9.25	-8.21	<0.001	Granzyme B (granzyme 2, cytotoxic T-lymphocyte-associated serine esterase 1)
	C1QA	1.08	-7.58	-7.03	<0.001	Complement component 1, q subcomponent, A chain
	SPP1	-1.07	-6.62	-7.05	0.003	Secreted phosphoprotein 1
	PLK3	-1.10	-5.99	-6.60	<0.001	Polo-like kinase 3 (Drosophila)
	F13A1	1.07	-5.12	-4.79	0.002	Coagulation factor XIII, A1 polypeptide
	PLA2G4A	1.07	-4.58	-4.28	0.002	Phospholipase A2, group IVA (cytosolic, calcium-dependent)
	CD55	-1.09	-4.54	-4.97	<0.001	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
	CD14	1.49	-4.41	-2.96	<0.001	CD14 molecule

Table II. Continued.

Processes	Symbol	Fold change			P-value	Gene name
		0-4 h	4-24 h	0-24 h		
Endocytosis and transport	IL10	-1.29	-4.11	5.30	0.002	Interleukin 10
	MAP3K8	-1.17	-4.02	-4.70	<0.001	Mitogen-activated protein kinase kinase kinase 8
	CCR3	1.01	-3.88	-3.84	0.003	Chemokine (C-C motif) receptor 3
	CCR4	-1.06	-3.63	-3.85	0.002	Chemokine (C-C motif) receptor 4
	KLRK1	-1.16	-3.55	-4.12	<0.001	Killer cell lectin-like receptor subfamily K, member 1
	IL2RA	-1.00	-3.46	-3.48	<0.001	Interleukin 2 receptor, alpha (IL2RA)
	CSF1R	1.32	-3.41	-2.58	<0.001	Colony stimulating factor 1 receptor precursor
	CCR1	1.08	-3.18	-2.95	0.003	Chemokine (C-C motif) receptor 1
	CCL4	-1.06	-3.13	-3.31	0.006	Chemokine (C-C motif) ligand 4 (CCL4),
	INHBA	1.04	-3.13	-3.02	0.002	Inhibin, beta A
	PECAM1	1.04	-3.59	-3.47	0.001	Platelet/endothelial cell adhesion molecule
	TNFRSF25	-1.10	-3.08	-3.40	<0.001	Tumor necrosis factor receptor superfamily, member 25
	CCL3	1.35	-3.01	-2.22	0.002	Chemokine (C-C motif) ligand 3
	TLR8	-1.26	-2.85	-3.60	0.006	Toll-like receptor 8
	IL12B	-1.20	-2.83	-3.40	0.004	Interleukin 12B
	CTSB	-1.06	-2.79	-2.97	0.003	Cathepsin B
	JAM3	1.05	-2.67	-2.54	<0.001	Junctional adhesion molecule 3
	SIPA1	1.07	-2.61	-2.43	0.003	Signal-induced proliferation-associated 1
	PGD	-1.16	-2.54	-2.95	0.004	Phosphogluconate dehydrogenase
	RAB7B	1.13	-4.29	-3.79	<0.001	RAB7B, member RAS oncogene family
	CD36	1.02	-3.58	-3.50	0.001	CD36 molecule (thrombospondin receptor)
	ACP2	1.00	-3.56	-3.54	<0.001	Acid phosphatase 2, lysosomal
	SORT1	-1.02	-3.45	-3.51	<0.001	Sortilin 1
	ACTB	1.17	-2.85	-2.44	<0.001	Actin, beta
	APIB1	1.03	-2.78	-2.71	<0.001	Adaptor-related protein complex 1, beta 1 subunit
Inflammation and apoptosis	TNF	-1.02	-6.07	-6.18	<0.001	Tumor necrosis factor (TNF superfamily, member 2)
	IL1RAP	1.17	-4.85	-5.67	0.005	Interleukin 1 receptor accessory protein
	IGFBP3	-1.06	-3.06	-3.23	0.002	Insulin-like growth factor binding protein 3
	CDKN1C	1.33	-2.95	-2.22	<0.001	Cyclin-dependent kinase inhibitor 1C (p57, Kip2)
	AMOTL1	-1.10	-2.71	-2.99	0.001	Angiomotin like 1


Table II. Continued.

Processes	Symbol	Fold change			P-value	Gene name
		0-4 h	4-24 h	0-24 h		
Others	FASLG	-1.19	-2.62	3.12	<0.001	Fas ligand
	CYP3A4	-1.04	-4.65	-4.83	0.002	Cytochrome P450, subfamily IIIA, polypeptide 4
	B4GALT6	-1.05	-4.09	-4.31	0.003	B4GALT6 protein, transcript variant 2
	TBXAS1	1.07	-4.03	-3.78	0.002	Thromboxane A synthase 1
	ST8SIA1	1.07	-3.78	3.52	0.002	Alpha-N-acetylneuraminide alpha-2,8-sialyltransferase
	GAB1	1.29	-3.61	-4.65	<0.001	GRB2-associated binding protein 1
	HSD17B8	1.01	-3.60	-3.57	0.005	Hydroxysteroid (17-beta) dehydrogenase 8
	HS3ST1	1.55	-3.27	-2.11	0.001	Heparan sulfate (glucosamine) 3-O-sulfotransferase 1
	MMP14	1.04	-3.09	-2.96	<0.001	Matrix metalloproteinase 14
	ITGAD	1.03	-2.83	-2.75	<0.001	Integrin, alpha D
	TXNDC5	-1.08	-3.34	-3.60	<0.001	Thioredoxin domain-containing protein 5 precursor
	VIM	1.06	-2.59	-2.44	0.004	Vimentin
	SDS	-1.23	-3.93	-4.85	<0.001	Serine dehydratase
	EME2	1.09	-2.65	2.44	0.003	Endonuclease
	SFRS7	-1.02	-2.82	-2.87	<0.001	Splicing factor, arginine/serine-rich 7
	NUP160	1.15	-3.01	-2.62	0.001	Nucleoporin 160kDa
	PWP2	1.13	-2.92	-2.59	<0.001	Periodic tryptophan protein 2 homolog
	DSE	-1.09	-2.92	-3.17	<0.001	Dermatan sulfate epimerase
	PNPLA4	1.10	-2.83	-2.57	<0.001	Patatin-like phospholipase domain containing 4
	BLVRB	1.00	-2.75	-2.75	<0.001	Biliverdin reductase B
	CHST2	1.04	-2.75	-2.64	<0.001	Carbohydrate (N-acetyl)glucosamine-6-O) sulfotransferase 2
	AGPAT3	1.08	-2.63	-2.43	<0.001	1-acylglycerol-3-phosphate O-acyltransferase 3
	EXT1	-1.19	-2.61	-3.12	0.001	Exostosins (multiple) 1
	GSTK1	1.12	-2.60	-2.90	0.002	Glutathione S-transferase kappa 1

Fold change was calculated as the mean of triplicate experiments.

Table III. Ingenuity pathway analysis profile of genes differentially expressed between 0 and 4 hours post-infection with *Mycobacterium bovis*.

Symbol	Entrez gene name	Fold change	Network	Location	Type(s)
CLDN3	Claudin 3	3.07	3	Plasma membrane	Transmembrane receptor
MMP2	Matrix metalloproteinase 2 (gelatinase A, 72 kDa gelatinase, 72 kDa type IV collagenase)	0.11	3	Extracellular space	Peptidase
MAPK14	Mitogen-activated protein kinase 14	0.05	1	Cytoplasm	Kinase
GEMIN6	Gem (nuclear organelle) associated protein 6	-0.09	4	Nucleus	Other
TNF	Tumor necrosis factor	-0.09	1	Extracellular space	Cytokine
TGFB1	Transforming growth factor, beta 1	-0.13	1	Extracellular space	Growth factor
CREB1	cAMP responsive element binding protein 1	-0.51		Nucleus	Transcription regulator
IL1B	Interleukin 1, beta	-0.52	1	Extracellular space	Cytokine
STRAP	Serine/threonine kinase receptor associated protein	-0.60	4	Plasma membrane	Other
REM1	RAS (RAD and GEM)-like GTP-binding 1	-3.13		Other	Enzyme
MMP13	Matrix metalloproteinase 13 (collagenase 3)	-6.25	1	Extracellular space	Peptidase
ADCY6	Adenylate cyclase 6	-6.67	2	Plasma membrane	Enzyme
GEMIN7	Gem (nuclear organelle) associated protein 7	-11.11	4	Nucleus	Other
SMURF2	SMAD specific E3 ubiquitin protein ligase 2	-16.67	1	Cytoplasm	Enzyme
FMO4	Flavin containing monooxygenase 4	-32.98		Cytoplasm	Enzyme

Figure 1. Significantly differentially expressed genes at each time point following *Mycobacterium bovis* challenge of peripheral blood mononuclear cells. The number of upregulated and downregulated genes relative to uninfected controls are shown for each time point sampled. *Adjusted $P \leq 0.05$.

and the pathways and networks involving these genes were identified using Ingenuity Pathway Analysis (IPA; <http://norris.usc.libguides.com/IPA>), a web-based functional analysis tool. The criteria for gene selection for IPA analysis were as follows: i) A fold-change in expression >2 for comparison between 0 and 4 hpi and >4 for comparison between 0 and 24 hpi, and ii) $P < 0.05$ for changes in gene expression in cells from all three cows.

Results

Kinetics of gene expression during *M. bovis* infection. Comprehensive gene expression profiles of the three PBMC samples with or without *M. bovis* challenge were generated using oligonucleotide bovine microarrays containing 43,803 probe sets. These probe sets interrogated the expression levels of ~29,356 transcripts, some of which mapped to known genes. A total of 3,937 probe sets passed the filtering step, consisting of a t-test with an adjusted P-value threshold ≤ 0.05 . At 4 and 24 hpi, 207 and 3,186 unique probe sets, respectively, were significantly differentially expressed (Fig. 1). To investigate the kinetics of gene expression, a total of 420 genes (including genes of unknown function) were found to be differentially expressed. Genes with an upregulated expression (30 out of 135 genes of known function) following exposure of PBMC to *M. bovis* are listed in Table I, and those with a downregulated expression (84 out of 285 genes of known function) are listed in Table II. As shown in Tables I and II, the genes with a fold change ≥ 2.5 between 0-4, 4-24 and 0-24 h were listed and divided by functions. Inspection of KEGG pathway annotations for these genes detected their association with the immune system (28%), signal transduction (23%), metabolism (21%), transport and catabolism (8%), genetic information processing (6%), cell growth and death (6%), and other organismal systems (8%). Of the affected genes associated with the immune system, 84.85% were downregulated in PBMCs following *M. bovis* challenge (Fig. 2). These results indicate a decreased T_H1 response [down-regulated TNF- α , IFN- γ , and interleukin (IL)-12 β], suggesting *M. bovis* infection may suppress the PBMC immune response.

Table IV. Pathways involving genes differentially expressed between 0 and 4 hours post-infection with *Mycobacterium bovis*.

Ingenuity canonical pathway	$-\log(p)$	Gene ratio ^a	Genes
Granulocyte adhesion and diapedesis	2.57	1.2E-02	MMP13, CLDN3
Agranulocyte adhesion and diapedesis	2.52	1.14E-02	MMP13, CLDN3
Leukocyte extravasation signaling	2.43	1.02E-02	MMP13, CLDN3
Oncostatin M signaling	1.78	2.94E-02	MMP13
Inhibition of matrix metalloproteases	1.73	2.63E-02	MMP13
CDK5 signaling	1.38	1.12E-02	ADCY6
TGF- β signaling	1.38	1.08E-02	SMURF2
IL-1 signaling	1.36	9.8E-03	ADCY6
HIF1 α signaling	1.32	9.8E-03	MMP13
G α i signaling	1.24	7.81E-03	ADCY6
eNOS signaling	1.23	7.75E-03	ADCY6
CXCR4 signaling	1.15	6.25E-03	ADCY6
Gap junction signaling	1.15	6.41E-03	ADCY6
Tight junction signaling	1.14	6.37E-03	CLDN3
PPAR α /RXR α activation	1.11	5.75E-03	ADCY6
RAR activation	1.09	5.59E-03	ADCY6
LPS/IL-1 mediated inhibition of RXR function	1.01	4.5E-03	FMO4

^aNumber of pathway genes differentially expressed/number of genes in pathway.




Figure 2. Profile of differentially expressed genes organized by functional subcategories of Kyoto Encyclopedia of Genes and Genomes pathways. Based on the number of genes with altered expression, *Mycobacterium bovis* exposure had the greatest effect on genes associated with the immune response.

Identification of pathways and networks associated with genes affected by M. bovis infection. Gene set enrichment analysis was performed on the microarray data to identify the specific biological pathways associated with genes differentially expressed upon *M. bovis* infection. IPA of the microarray data identified the functional profiles of 15 genes that were differentially expressed between 0 and 4 hpi (Table III), and their associated pathways (Table IV). A total of 91 genes with

differential expression between 0 and 24 hpi were selected in the present study for IPA; and the results of the top 28 (greatest change in expression) are shown in Table V. The pathways associated with these genes are shown in Table VI. A map of the network of pathways involving genes differentially expressed between 0 and 4 hpi is shown in Figs. 3-6.

The key genes in network 1 (Fig. 3) were transforming growth factor- β and matrix metalloproteinase (MMP)13;

Table V. Ingenuity pathway analysis profile of genes differentially expressed between 0 and 24 hours post-infection with *Mycobacterium bovis*.

Symbol	Entrez gene name	Fold change	Network	Location	Type(s)
GPNMB	Glycoprotein (transmembrane) nmb	22.93	2	Plasma membrane	Enzyme
AK4	Adenylate kinase 4	19.13	4	Cytoplasm	Kinase
KCNH2	Potassium voltage-gated channel, subfamily H (eag-related), member 2	18.27	4	Plasma membrane	Ion channel
CLEC4E	C-type lectin domain family 4, member E	13.36	2	Plasma membrane	Other
CLU	Clusterin	12.13	4	Cytoplasm	Other
C3AR1	Complement component 3a receptor 1	7.93	1	Plasma membrane	G-protein coupled receptor
Akt	Bos taurus v-akt murine thymoma viral oncogene homolog 2	2.06	4	Cytoplasm	Group
BCL2	B-cell CLL/lymphoma 2	-0.29	3	Cytoplasm	Transporter
CASP8	Caspase 8, apoptosis-related cysteine peptidase	-0.60		Nucleus	Peptidase
NFKB1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1	-0.80		Nucleus	Transcription regulator
IL1B	Interleukin 1, beta	-1.04		Extracellular space	Cytokine
FCER1G	Fc fragment of IgE, high affinity I, receptor for; gamma polypeptide	-1.98		Plasma membrane	Transmembrane receptor
CD55	CD55 molecule, decay accelerating factor for complement (Cromer blood group)	-4.76	2	Plasma membrane	Other
TNF	Tumor necrosis factor	-6.45	1	Extracellular space	Cytokine
C1QA	Complement component 1, q subcomponent, A chain	-7.69	1	Extracellular space	Other
PLK3	Polo-like kinase 3	-8.33	3	Nucleus	Kinase
CEBPD	CCAAT/enhancer binding protein (C/EBP), delta	-9.09	4	Nucleus	Transcription regulator
IL17F	Interleukin 17F	-9.52	1	Extracellular space	Cytokine
GZMB	Granzyme B (granzyme 2, cytotoxic T-lymphocyte-associated serine esterase 1)	-11.11	1	Cytoplasm	Peptidase
MMP13	Matrix metalloproteinase 13 (collagenase 3)	-11.11	1	Extracellular space	Peptidase
SMURF2	SMAD specific E3 ubiquitin protein ligase 2	-12.50	3	Cytoplasm	Enzyme
CFB	Complement factor B	-13.33	1	Extracellular space	Peptidase
THBS1	Thrombospondin 1	-13.33	1,2	Extracellular space	Other
FMO4	Flavin containing monooxygenase 4	-14.29		Cytoplasm	Enzyme
IFNG	Interferon, gamma	-20.00	1,3	Extracellular space	Cytokine
THBD	Thrombomodulin	-20.00	1	Plasma membrane	Transmembrane receptor
IL17RB	Interleukin 17 receptor B	-25.00	1	Plasma membrane	Transmembrane receptor
C1QB	Complement component 1, q subcomponent, B chain	-28.57	1	Extracellular space	Other

Table VI. Pathways involving genes differentially expressed between 0 and 24 hours post-infection with *Mycobacterium bovis*.

Ingenuity canonical pathways	-log (P)	Gene ratio ^a	Molecules
Pattern recognition receptors in recognition of bacteria and viruses	3.23E00	4.21E-02	C1QA, C1QB, C3AR1, TNF
Production of nitric oxide and reactive oxygen species in macrophages	3.02E00	2.69E-02	IFNG, MAP3K8, TNF, CLU, RBP4
Cytokines mediating communication between immune cells	2.86E00	5.77E-02	IFNG, IL17F, TNF
T helper cell differentiation	2.55E00	4.35E-02	IFNG, IL17F, TNF
VDR/RXR activation	2.37E00	3.85E-02	IFNG, SPP1, THBD
Crosstalk between dendritic cells and natural killer cells	2.2E00	3.3E-02	IFNG, TNF, ACTG1
Acute phase response signaling	1.47E00	1.73E-02	CFB, TNF, RBP4
Communication between innate and adaptive immune cells	1.33E00	2.15E-02	IFNG, TNF
PI3K/AKT signaling	1.03E00	1.48E-02	GAB1, MAP3K8
TNFR2 signaling	9.51E-01	3.12E-02	TNF
Interferon signaling	8.72E-01	2.94E-02	IFNG
Antigen presentation pathway	8.38E-01	2.5E-02	IFNG
Inhibition of matrix metalloproteases	8.28E-01	2.63E-02	MMP13
NF-κB signaling	7.96E-01	1.14E-02	MAP3K8, TNF
iNOS signaling	7.78E-01	2.13E-02	IFNG
TNFR1 signaling	7.43E-01	1.96E-02	TNF
Cytotoxic T lymphocyte-mediated apoptosis of target cells	7.27E-01	1.92E-02	GZMB
Leukocyte extravasation signaling	7.08E-01	1.02E-02	MMP13, ACTG1
Death receptor signaling	6.62E-01	1.61E-02	TNF
Activation of IRF by cytosolic pattern recognition receptors	6.49E-01	1.59E-02	TNF
Role of PI3K/AKT signaling in the pathogenesis of influenza	6.42E-01	1.49E-02	IFNG
IL-10 signaling	6.01E-01	1.39E-02	TNF
Apoptosis signaling	5.06E-01	1.09E-02	TNF
Fcγ receptor-mediated phagocytosis in macrophages and monocytes	4.87E-01	1.05E-02	ACTG1

^aNumber of pathway genes differentially expressed/number of genes in pathway.

both of which are major genes associated with inflammatory responses. There were only two genes in network 2 (Fig. 4): Retinol-binding protein 4 and adenylyl cyclase 6; both of which have known roles in development, particularly embryonic, skeletal and muscular, and so were unlikely to be associated with infection. The key gene in network 3 (Fig. 5) was claudin 3 (CLDN3); as in the case of network 2, this network is predominantly associated with skeletal and muscular development, and therefore has only a weak association with the infection. The key gene in network 4 (Fig. 6) was gem-associated protein 7, which is associated with cell death and survival, and therefore may be associated with the late response to *M. bovis* infection. The four networks of the pathways comprising genes differentially expressed between 0 and 24 hpi are shown in Figs. 7-10. The key gene in network 1 was TNF-α, which interacts with IFN-γ, MMP13, and thrombospondin 1 (THBS1) (Fig. 7). Expression of these four genes was downregulated. TNF-α also interacts with IL-17 receptor B and thrombomodulin, both of which demonstrated downregulated expression. In addition, TNF-α interacts with activating protein-1, which upregulates complement component 3a receptor 1. The key genes in network 2 were luteinizing hormone (LH) and IL-13. IL-13 upregulates C-type lectin domain family 4 member E, and glycoprotein

(transmembrane) nmb, and downregulates THBS1 and androgen-induced 1 (Fig. 8). The key gene in network 3 was IFN-γ, which regulates the expression of B-cell lymphoma-2 and the nuclear factor (NF)-κB p65 subunit (Fig. 9). The key gene in network 4 was Akt, which regulates the expression of NF-κB, followed by the downregulation of CCAAT/enhancer binding protein, delta (Fig. 10).

Discussion

The present study demonstrated that bovine PBMCs responded to *in vitro* *M. bovis* infection by undergoing large-scale alterations in gene expression. The expression of 420 genes was shown to significantly differ between 4 and 24 hpi, with 135 upregulated and 285 downregulated genes. Inspection of KEGG pathway annotations for these genes demonstrated that the majority was associated with the immune system, signal transduction, and metabolism. Of the affected genes with immune system functions, 84.85% were downregulated. System pathway analysis of differentially expressed genes revealed the key genes in four different networks to be TNF-α, IFN-γ, LH, IL-13, and NF-κB. These results suggested that *M. bovis* may suppress the PBMC immune response soon after infection.




Figure 3. Network 1 of pathways involving genes differentially expressed between 0 and 4 h post-infection with *Mycobacterium bovis*.

The number of differentially expressed PBMC genes increased during the first 24 h following exposure to *M. bovis*. Changes were observed in 207 unique probe sets at 4 hpi and in 3,186 unique probe sets at 24 hpi. Of these, 420 genes displayed significantly altered expression from 4 to 24 hpi, with expression decreasing for 285 genes and increasing for 135 genes. In addition, the fold-change in expression of the downregulated genes was much greater, as compared with that of upregulated genes. Previous studies of transcriptional responses to *M. bovis* infection have reported downregulation of the majority of differentially expressed genes (10,12,15,16). In the present study, more genes were suppressed later post-infection (24 h), as compared with earlier (4 h), suggesting the cellular activities involving these genes progressively declined during *M. bovis* infection.

Inspection of KEGG pathway annotations for the 420 differentially expressed genes revealed their involvement in the immune system (28%), signal transduction (23%), metabolism (21%), transport and catabolism (8%), genetic information processing (6%), cell growth and death (6%), and other organismal systems (8%). Of these genes, >67% (280) exhibited time-dependent decreases in expression associated with signal transduction, immune response, pro-inflammatory cytokines, metabolism, or cell death processes. In addition, 84.85% of the differentially expressed genes associated with immune responses displayed downregulated expression in infected PBMCs. Suppression of host immune response genes is a common finding among studies of gene expression

following *M. bovis* infection (6,12,16). Transcriptome analysis of peripheral blood leukocytes from cattle infected with *M. bovis* previously detected over-representation of differentially expressed genes associated with the immune response; of these genes, 64.5% showed decreased expression, indicating *M. bovis* infection may be associated with the suppression of host immune genes (12). Meade *et al* (16) reported a decrease in the *in vivo* PBMC expression of key innate immune genes in *M. bovis*-infected cattle. Subsequent *in vivo* transcriptional studies of *M. bovis*-infected cattle demonstrated that PBMC genes associated with immunity, inflammatory responses, and apoptosis were among those with the highest differential expression (10). These *in vivo* experiments were conducted in animals following the establishment of an infection, ranging from 2-12 months following inoculation. The *in vitro* data of the present study supports these findings and further reveals that changes in gene expression begin very early in the course of infection.

Notably, *M. bovis* infection resulted in a decrease in expression of the most important components of the T_H1 response: IFN- γ , TNF- α , and IL-12. The significance of the altered expression of IFN- γ and TNF- α is demonstrated by their mapping to key locations in the pathway networks. These genes are crucial to the host immune response against mycobacteria, including granuloma formation and apoptosis. Signaling via the IFN- γ pathway is required for macrophage activation and granuloma formation (5). Such signaling is dependent on the production of IFN- γ by T-cells; and IFN- γ synthesis requires




Figure 4. Network 2 of pathways involving genes differentially expressed between 0 and 4 h post-infection with *Mycobacterium bovis*.






Figure 5. Network 3 of pathways involving genes differentially expressed between 0 and 4 h post-infection with *Mycobacterium bovis*.






Figure 8. Network 2 of pathways involving genes differentially expressed between 0 and 24 h post-infection with *Mycobacterium bovis*.




Figure 9. Network 3 of pathways involving genes differentially expressed between 0 and 24 h post-infection with *Mycobacterium bovis*.




Figure 10. Network 4 of pathways involving genes differentially expressed between 0 and 24 h post-infection with *Mycobacterium bovis*.

the cytokine IL-12. Conversely, the observation in the present study of decreased IFN- γ expression differs from a previous study, which reported that the vaccine *Mycobacterium bovis* bacillus Calmette-Guérin triggers a T_H1 -type response (17). In addition, another study reported the increased expression of IFN- γ in PBMC from cattle infected 4 months previously (10). However, these studies observed that genes downstream of IFN- γ were significantly downregulated, suggesting suppression of IFN- γ signaling despite its increased expression. Strain virulence, MOI, cell type, post-infection harvest time, and specific assays used may also underlie these different results.

Apoptosis of infected macrophages is an innate host defense mechanism against intracellular *M. bovis* and *M. tuberculosis*. The extrinsic cell death pathway involved in apoptosis is induced by the binding of TNF- α to its receptor on the macrophage surface. Macrophages infected with attenuated strains of pathogenic mycobacteria undergo TNF- α -mediated apoptosis, reducing the viability of intracellular bacilli. Virulent *M. tuberculosis* strains have been found to suppress macrophage apoptosis (18,19). Previous studies have detected upregulation of programmed cell death signaling genes, including TNF- α , following live *M. bovis* challenge of bovine macrophages *in vitro* (6,20). Therefore, it is intriguing that TNF- α was the key gene affected in network 1. Through this network of pathways, TNF- α was shown to interact with IFN- γ , MMP13, and THBS1.

The present study has numerous limitations. The methods did not distinguish between *M. bovis*-infected cells and

M. bovis-exposed cells; therefore, no correlations can be made between gene expression levels and infection rates. The *M. bovis* cells were not from a standard strain; therefore, some comparisons to other studies may be less reliable. In addition, the present study does not provide further analysis of specific genes indicated by the microarray data. However, the data from this preliminary screening provide a solid foundation for future investigations.

To the best of our knowledge, this is the first study providing a time-course analysis of global gene expression in bovine PBMCs following *in vitro* exposure to *M. bovis*. Our data indicate that extensive alterations in PBMC gene expression may begin early in infection. The majority of the differentially expressed genes were related to immune responses and cell survival. Changes observed in the expression of genes associated with immune responses suggest that *M. bovis* infection may be associated with the suppression of immune response-related gene expression. In addition, *M. bovis* infection in PBMCs may suppress apoptosis by interfering with TNF- α signaling. The present study provides valuable information for the further characterization of host responses to *M. bovis* infection.

Acknowledgements

The present study was supported by a grant from the Centers for Disease Control, Taiwan, the Republic of China.

References

1. Etter E, Donado P, Jori F, Caron A, Goutard F and Roger F: Risk analysis and bovine tuberculosis, a re-emerging zoonosis. *Ann NY Acad Sci* 1081: 61-73, 2006.
2. Neill SD, Pollock JM, Bryson DB and Hanna J: Pathogenesis of *Mycobacterium bovis* infection in cattle. *Vet Microbiol* 40: 41-52, 1994.
3. Kabara E, Kloss CC, Wilson M, Tempelman RJ, Sreevatsan S, Janagama H and Coussens PM: A large-scale study of differential gene expression in monocyte-derived macrophages infected with several strains of *Mycobacterium avium* subspecies paratuberculosis. *Brief Funct Genomics* 9: 220-237, 2010.
4. Kerrigan AM and Brown GD: Syk-coupled C-type lectins in immunity. *Trends Immunol* 32: 151-156, 2011.
5. Saunders BM and Britton WJ: Life and death in the granuloma: Immunopathology of tuberculosis. *Immunol Cell Biol* 85: 103-111, 2007.
6. Magee DA, Taraktoglou M, Killick KE, Nalpas NC, Browne JA, Park SD, Conlon KM, Lynn DJ, Hokamp K, Gordon SV, *et al*: Global gene expression and systems biology analysis of bovine monocyte-derived macrophages in response to in vitro challenge with *Mycobacterium bovis*. *PLoS One* 7: e32034, 2012.
7. Netea MG, Van der Meer JW and Kullberg BJ: Toll-like receptors as an escape mechanism from the host defense. *Trends Microbiol* 12: 484-488, 2004.
8. Behar SM, Divangahi M and Remold HG: Evasion of innate immunity by *Mycobacterium tuberculosis*: Is death an exit strategy? *Nat Rev Microbiol* 8: 668-674, 2010.
9. Bovine Genome Sequencing and Analysis Consortium; Elsik CG, Tellam RL, Worley KC, Gibbs RA, Muzny DM, Weinstock GM, Adelson DL, Eichler EE, Elnitski L, *et al*: The genome sequence of taurine cattle: A window to ruminant biology and evolution. *Science* 324: 522-528, 2009.
10. Blanco FC, Soria M, Bianco MV and Bigi F: Transcriptional response of peripheral blood mononuclear cells from cattle infected with *Mycobacterium bovis*. *PLoS One* 7: e41066, 2012.
11. Meade KG, Gormley E, O'Farrelly C, Park SD, Costello E, Keane J, Zhao Y and MacHugh DE: Antigen stimulation of peripheral blood mononuclear cells from *Mycobacterium bovis* infected cattle yields evidence for a novel gene expression program. *BMC Genomics* 9: 447, 2008.
12. Killick KE, Browne JA, Park SD, Magee DA, Martin I, Meade KG, Gordon SV, Gormley E, O'Farrelly C, Hokamp K and MacHugh DE: Genome-wide transcriptional profiling of peripheral blood leukocytes from cattle infected with *Mycobacterium bovis* reveals suppression of host immune genes. *BMC Genomics* 12: 611, 2011.
13. Pollock JM and Neill SD: *Mycobacterium bovis* infection and tuberculosis in cattle. *Vet J* 163: 115-127, 2002.
14. Aranaz A, Liébana E, Mateos A, Dominguez L, Vidal D, Domingo M, Gonzolez O, Rodriguez-Ferri EF, Bunschoten AE, Van Embden JD and Cousins D: Spacer oligonucleotide typing of *Mycobacterium bovis* strains from cattle and other animals: A tool for studying epidemiology of tuberculosis. *J Clin Microbiol* 34: 2734-2740, 1996.
15. Meade KG, Gormley E, Park SD, Fitzsimons T, Rosa GJ, Costello E, Keane J, Coussens PM and MacHugh DE: Gene expression profiling of peripheral blood mononuclear cells (PBMC) from *Mycobacterium bovis* infected cattle after in vitro antigenic stimulation with purified protein derivative of tuberculin (PPD). *Vet Immunol Immunopathol* 113: 73-89, 2006.
16. Meade KG, Gormley E, Doyle MB, Fitzsimons T, O'Farrelly C, Costello E, Keane J, Zhao Y and MacHugh DE: Innate gene repression associated with *Mycobacterium bovis* infection in cattle: Toward a gene signature of disease. *BMC Genomics* 8: 400, 2007.
17. Marchant A, Goetghebuer T, Ota MO, Wolfe I, Ceesay SJ, De Groote D, Corrah T, Bennett S, Wheeler J, Huygen K, *et al*: Newborns develop a Th1-type immune response to *Mycobacterium bovis* bacillus Calmette-Guérin vaccination. *J Immunol* 163: 2249-2255, 1999.
18. Divangahi M, Chen M, Gan H, Desjardins D, Hickman TT, Lee DM, Fortune S, Behar SM and Remold HG: *Mycobacterium tuberculosis* evades macrophage defenses by inhibiting plasma membrane repair. *Nat Immunol* 10: 899-906, 2009.
19. Lee J, Remold HG, Jeong MH and Kornfeld H: Macrophage apoptosis in response to high intracellular burden of *Mycobacterium tuberculosis* is mediated by a novel caspase-independent pathway. *J Immunol* 176: 4267-4274, 2006.
20. Widdison S, Watson M and Coffey TJ: Early response of bovine alveolar macrophages to infection with live and heat-killed *Mycobacterium bovis*. *Dev Comp Immunol* 35: 580-591, 2011.