

NSAIDs modulate GABA-activated currents via Ca^{2+} -activated Cl^- channels in rat dorsal root ganglion neurons

LEI ZHAO^{1,2*}, LI LI^{1,2*}, KE-TAO MA^{1,2}, YANG WANG¹⁻³, JING LI¹, WEN-YAN SHI^{1,2},
HE ZHU^{1,2}, ZHONG-SHUANG ZHANG^{1,2} and JUN-QIANG SI¹⁻⁴

¹Department of Physiology; ²Electrophysiological Laboratory, Laboratory of Xinjiang Endemic and Ethnic Diseases, Shihezi University Medical College, Shihezi, Xinjiang 832002; ³Department of Physiology, School of Basic Medical Sciences, Wuhan University, Wuhan, Hubei 430071; ⁴Department of Physiology, School of Basic Medical Sciences, Huazhong University of Science and Technology, Wuhan, Hubei 430030, P.R. China

Received June 23, 2015; Accepted February 11, 2016

DOI: 10.3892/etm.2016.3158

Abstract. The ability of non-steroidal anti-inflammatory drugs (NSAIDs) to modulate γ -aminobutyrate (GABA)-activated currents via Ca^{2+} -activated Cl^- channels in rat dorsal root ganglion neurons (DRG), was examined in the present study. During the preparation of DRG neurons harvested from Sprague-Dawley rats, the whole-cell recording technique was used to record the effect of NSAIDs on GABA-activated inward currents, and the expression levels of the TMEM16A and TMEM16B subunits were revealed. In the event that DRG neurons were pre-incubated for 20 sec with niflumic acid (NFA) and 5-nitro-2-(3-phenylpropylamino) benzoic acid (NPPB) prior to the administration of GABA, the GABA-induced inward currents were diminished markedly in the majority of neurons examined (96.3%). The inward currents induced by 100 $\mu\text{mol/l}$ GABA were attenuated by (0 \pm 0.09%; neurons = 4), (5.32 \pm 3.51%; neurons = 6), (21.3 \pm 4.00%; neurons = 5), (33.8 \pm 5.20%; neurons = 17), (52.2 \pm 5.10%; neurons = 4) and (61.1 \pm 4.12%; neurons = 12) by 0.1, 1, 3, 10, 30 and 100 $\mu\text{mol/l}$ NFA, respectively. The inward currents induced by 100 $\mu\text{mol/l}$ GABA were attenuated by (13.8 \pm 6%;

neurons = 6), (23.2 \pm 14.7%; neurons = 6) and (29.7 \pm 9.1%; neurons = 9) by 3, 10 and 30 $\mu\text{mol/l}$ NPPB, respectively. NFA and NPPB dose-dependently inhibited GABA-activated currents with half maximal inhibitory concentration (IC_{50}) values of 6.7 and 11 $\mu\text{mol/l}$, respectively. The inhibitory effect of 100 $\mu\text{mol/l}$ NFA on the GABA-evoked inward current were also strongly inhibited by nitrendipine (NTDP; an L-type calcium channel blocker), 1,2-bis(2-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid tetrakis (a highly selective calcium chelating reagent), caffeine (a widely available Ca^{2+} consuming drug) and calcium-free extracellular fluid, in a concentration-dependent manner. Immunofluorescent staining indicated that TMEM16A and TMEM16B expression was widely distributed in DRG neurons. The results suggest that NSAIDs may be able to regulate Ca^{2+} -activated chloride channels to reduce GABA_A receptor-mediated inward currents in DRGs.

Introduction

γ -aminobutyrate (GABA) is a crucial inhibitory neurotransmitter in the mammalian peripheral nervous system (PNS) and central nervous system (CNS) (1). GABA_A receptor is a pentamer comprised of multiple subunits (α_6 , β_{1-3} , γ_{1-3} , π , ϵ , δ and θ) with an absolute chloride ion channel and diversiform allosteric binding sites through which rapid inhibitory synaptic neurotransmission may be modulated (1). The GABA_A receptor is a favorable target for therapeutic agents including steroids, barbiturates, benzodiazepines, anesthetics and convulsants (2). Recently, it has also been proposed that the β -subunit has an important role in determining the chloride ion selectivity of GABA_A receptors (3,4).

GABA_A receptor antagonists niflumic acid (NFA) and 5-nitro-2-(3-phenylpropylamino) benzoic acid (NPPB), are the only chloride ion channel blockers able to protect cells from excitotoxicity (5). Accumulating evidence suggests that non-steroidal anti-inflammatory drugs (NSAIDs) modulate GABA_A receptor function in heterologous expression systems (6). NSAIDs at clinically relevant concentrations (low micromolar) are sufficient to potentiate $\beta_{2/3}$ -containing GABA_A receptors (7). The NSAID-sensitive $\alpha_1\beta_2\gamma_2$ receptor

Correspondence to: Professor Jun-Qiang Si, Department of Physiology, Shihezi University Medical College, 106 Beier Road, Shihezi, Xinjiang 832002, P.R. China
E-mail: sijunqiang11@hotmail.com

Abbreviations: GABA, γ -aminobutyrate; PNS, periphery nervous system; CNS, central nervous system; NFA, niflumic acid; NPPB, 5-nitro-2-(3-phenylpropylamino) benzoic acid; NSAIDs, non-steroidal anti-inflammatory drugs; CaCCs, Ca^{2+} -activated Cl^- channels; DRG, dorsal root ganglion; SDRs, Sprague-Dawley rats; NTDP, nitrendipine

*Contributed equally

Key words: non-steroidal anti-inflammatory drugs, niflumic acid, γ -aminobutyrate-induced inward current, modulatory, Ca^{2+} -activated Cl^- channels, dorsal root ganglion

subtype is the predominant and the largest GABA_A receptor population in mammalian PNS and CNS (7). In addition to their effect on GABA_A receptors, fenamate NSAIDs also affect a variety of other ion channels (8,9). Several drugs that have an effect on GABA_A receptor function have been revealed to depend on subtypes (subunits combinations), and on specific amino acids sites of specific subunits (10).

Halliwell *et al* (11) revealed that the regulation of GABA_A receptors by one particular anti-inflammatory agent, mefenamic acid, was dependent on the β -subunit. Conversely, Sinkkonen *et al* (12) reported that the potentiation of $\alpha_1\beta_2\gamma_2$ receptors by NFA was dependent on the presence of a γ_2 subunit, which also effects mefenamic acid modulation (11). Antagonism of the $\alpha_6\beta_2\gamma_2$ receptor subtype by NFA has also been reported (12), and the substitution of an α_4 subunit reduced the mefenamic potentiation of $\alpha_1\beta_2\gamma_2$ receptors by 50% (13). Similar observations have been observed in electrophysiological studies regarding the actions of mefenamic acid, pentobarbital and etomidate (11,14,15).

The aim of the present study is to use conventional whole-cell patch-clamp recordings, immunofluorescence and NSAIDs, including NFA and NPPB, to investigate the effect of Ca²⁺-activated Cl⁻ channels (CaCCs) on GABA-induced currents in the dorsal root ganglion (DRG) of rats. The present study intended to elucidate the diversity of the modulatory effect route of NSAIDs on GABA-activated currents via CaCCs.

Materials and methods

Isolation of DRG neurons. A total of 120 Sprague-Dawley rats (SDRs) were provided by the Experimental Animal Center of Xinjiang Medical University, Urumqi, China (certificate no. SCXK 2003-0001; age, 8-10 weeks; weight, 250-280 g) irrespective of gender. All protocols were approved by the Institutional Animal Care and Use Committee at the Medical College of Shihezi University (Shihezi, China) and were consistent with the Guidelines for the Care and Use of Laboratory Animals published by the US National Institutes of Health (16). SDRs were bred in separate specific pathogen-free cages at a relative humidity of 40-70%, (24±3°C), 100-120 lx/12-h dark:light illumination and free access to food and water. The DRG neuron selection and the separation process are described in our previous studies (17,18). Rats were sacrificed by decapitation.

Electrophysiological recordings. A gap-free recording with a sampling interval of 50 msec (17,18) was performed in the present study. Briefly, with the aid of a whole-cell patch clamp amplifier, perforated patch-clamp recordings in the whole-cell mode were performed. Using an Axon 700B amplifier (Axon, San Jose, CA, USA) and pCLAMP version 0.2 hardware and software (Axon), currents were recorded from the DRG neurons *in vitro*. The room temperature was set at 22-24°C. The resistance of the recording pipette ranged from 3 to 5 M Ω . The experimental procedures were performed according to the Regulations for the Administration of Affairs. Concerning Experimental Animals, formulated by the Ministry of Science and Technology of the People's Republic of China (The Ministry of Science and Technology of the People's Republic

of China. Guidance Suggestions for the Care and Use of Laboratory Animals 2011).

Immunofluorescent staining of TMEM16A and TMEM16B to determine expression in the DRG. Rats were anesthetized with 0.3% (w/v) sodium pentobarbital [Sangon Biotech (Shanghai,) Co., Ltd., Shanghai, China], and perfused through the aorta with 0.9% (w/v) normal saline, followed by fresh 4% (w/v) paraformaldehyde in phosphate-buffered saline [both purchased from Sangon Biotech (Shanghai,) Co., Ltd.] for 10 min for tissue fixation. The lumbar DRG at level L₄₋₆ to the nerve injury was removed rapidly and placed in 4% (w/v) paraformaldehyde in PBS for 24 h. The L₄₋₆ DRG were cut into 5- μ m sections with a freezing microtome (CM1510S; Leica Biosystems, Wetzlar, Germany). Immunofluorescent staining was performed using rabbit anti-TMEM16A polyclonal antibody (1:20; sc-135235) and goat anti-TMEM16B polyclonal antibody (1:20; sc-169622) (both purchased from Cell Signaling Technology, Inc., Danvers, MA, USA) overnight at 4°C. The sections were incubated for 1 h in a solution containing donkey anti-rabbit IgG-fluorescein isothiocyanate (FITC; 1:50; 711-095-152) and donkey anti-goat IgG-tetramethylrhodamine (TRITC; 1:50; 705-025-003) (both purchased from Jackson ImmunoResearch, West Grove, PA, USA) at 37°C. Several tissue sections were selected for double-labeling of TMEM16A and TMEM16B and were incubated in a mixture of primary antibodies against TMEM16A and TMEM16B, followed by donkey anti-rabbit IgG conjugated with FITC and donkey anti-goat IgG conjugated with TRITC. Slides were then examined by confocal microscopy (LSM710; Carl Zeiss AG, Oberkochen, Germany). Quantitative analysis of TMEM16A and TMEM16B expression in the DRG was performed by measuring the mean absorbance at 488 and 550 nm (Zeiss LSM 510 System; Carl Zeiss, Jena, Germany) following laser confocal microscopy and using analysis software (ZEN 2009 Light Edition; Carl Zeiss).

Drug application. GABA, muscimol, bicuculline, NFA, NPPB, caffeine and NTDP were purchased from Sigma-Aldrich, (St. Louis, MO, USA). 1,2-Bis(2-aminophenoxy)ethane-N, N',N'-tetraacetic acid tetrakis (acetoxymethyl ester; BAPTA-AM) was from Merck Millipore (Darmstadt, Germany). Rabbit anti-TMEM16A polyclonal antibody and donkey anti-rabbit IgG-FITC were purchased from Santa Cruz Biotechnology, Inc., (Dallas, TX, USA). All drugs used in electrophysiological recordings were dissolved in extracellular fluid and applied by gravity flow from a home-made perfusion system consisting a row of tubules connected with a series of individual reservoirs (17). This rapid solution exchange system was manipulated by shifting the tubules horizontally with a micromanipulator (17,18). The time of pre-perfusion of antagonists was 0.5-5 min, and the time of pre-perfusion of GABA was 5-10 sec.

Statistical analysis. Statistical analysis of the data was performed using SPSS (version 13.0; SPSS, Inc., Chicago, IL, USA) and the values of GABA-activated currents are presented as mean \pm standard error of the mean and a Student's t test was used to assess the significance. P<0.05 was considered to indicate a statistically significant difference.




Figure 1. Inhibition of GABA_A receptor antagonist on muscimol-induced depolarization. (A) Blockade of 100 $\mu\text{mol/l}$ muscimol-induced inward current by GABA_A receptor antagonist bicuculline (100 $\mu\text{mol/l}$). (B) Statistical results indicating the ability of bicuculline to block muscimol-activated currents. * $P<0.05$ vs. control. Numbers on bars=number of neurons.

Results

GABA-induced inward currents. Treatment with different concentrations of GABA (1-1,000 $\mu\text{mol/l}$) activated an inward current in the majority of cells (94.32%, 150/159) examined. The GABA-induced response was concentration-dependent, and displayed evidence of desensitization at high concentrations (Fig. 1). The activation threshold was ~ 1 $\mu\text{mol/l}$ and the maximal response was achieved at 300 $\mu\text{mol/l}$ GABA. The value of the dissociation constant was ~ 30 $\mu\text{mol/l}$, derived from a concentration-response curve (17,18). The averaged amplitude of 100 $\mu\text{mol/l}$ GABA-evoked inward current was $(1.29 \pm 0.72$ nA; neurons = 52). The selective GABA_A receptor agonist, muscimol (100 $\mu\text{mol/l}$) mimicked the GABA-evoked response (neurons = 8). A selective GABA_A receptor antagonist, bicuculline (100 $\mu\text{mol/l}$), suppressed GABA (neurons = 9) and muscimol-evoked (neurons = 8) inward currents (Fig. 1).

Inhibition of GABA-induced inward currents by NFA and NPPB. NFA and NPPB were pre-incubated for 20 sec prior to application of GABA, resulting in the marked attenuation of the GABA-induced inward current in the majority of the neurons examined (96.3%, 52/54). Inhibition of GABA-induced responses by NFA and NPPB were concentration-dependent. The inward currents induced by 100 $\mu\text{mol/l}$ GABA were suppressed by $(0 \pm 0.09\%$; neurons=4), $(5.32 \pm 3.51\%$; neurons=6), $(21.3 \pm 4.00\%$; neurons=5), $(33.8 \pm 5.20\%$; neurons=17), $(52.2 \pm 5.10\%$; neurons=4), $(61.1 \pm 4.12\%$; neurons=12) and $(57.6 \pm 4.20\%$; neurons=4) by 0.1, 1, 3, 10, 30, 100 and 300 $\mu\text{mol/l}$ NFA, respectively (Fig. 2A). The




Figure 2. Effects of different concentrations of NFA on GABA-activated inward currents. (A) Results of 100 $\mu\text{mol/l}$ GABA-activated inward currents in the different-concentration of NFA (0.1-100 $\mu\text{mol/l}$). (B) Inward current induced by 100 $\mu\text{mol/l}$ GABA was suppressed by $0 \pm 0.09\%$ (neurons=4), $5.32 \pm 3.51\%$ (neurons=6), $21.3 \pm 4.00\%$ (neurons=5), $33.8 \pm 5.20\%$ (neurons=17), $52.2 \pm 5.10\%$ (neurons=4), $61.1 \pm 4.12\%$ (neurons=12) and $57.6 \pm 4.20\%$ (neurons=4) by 0.1, 1, 3, 10, 30, 100 and 300 $\mu\text{mol/l}$ NFA (neurons=3-17), respectively. * $P<0.05$ and ** $P<0.01$ vs. control. Data are results of a paired t test.

inhibition threshold was ~ 0.1 $\mu\text{mol/l}$ and the maximal inhibition was achieved at 300 $\mu\text{mol/l}$ NFA. The IC_{50} value was ~ 6.7 $\mu\text{mol/l}$ derived from the concentration-inhibition curve (Fig. 2B). NFA did not alter the half maximal effective concentration EC_{50} value for GABA (~ 30 $\mu\text{mol/l}$) (17,19), but reduced the maximal GABA currents by $\sim 60\%$. The inward current induced by 100 $\mu\text{mol/l}$ GABA was suppressed by $13.8 \pm 6.7\%$, neurons=6, $P<0.05$; $23.2 \pm 14.7\%$, neurons=6, $P<0.01$ and $(29.7 \pm 9.1\%$, neurons=9, $P<0.01$, by 3, 10 and 30 $\mu\text{mol/l}$ NFA, respectively. The inhibition threshold was ~ 1 $\mu\text{mol/l}$ and the maximal inhibition was achieved by 30 $\mu\text{mol/l}$ NFA. The IC_{50} value was ~ 11 $\mu\text{mol/l}$ (Fig. 3).

Effects of NTDP and extracellular calcium on GABA-induced inward currents. The L-type calcium channel blocker, NTDP (0.1-30 $\mu\text{mol/l}$), inverted the inhibitory effect of 100 $\mu\text{mol/l}$ NFA on 100 $\mu\text{mol/l}$ GABA-induced inward current (Fig. 4). The inhibitory ratio of NFA on inward current induced by GABA were $(59.6 \pm 8.70\%$, neurons=4, $P>0.05$; $43.6 \pm 5.10\%$, neurons=3, $P<0.05$; $32.3 \pm 6.62\%$, neurons=8, $P<0.01$; $8.7 \pm 7.6\%$, neurons=6, $P<0.01$ and $8.6 \pm 7.4\%$, neurons=4, $P<0.01$, in the presence of 0.1, 1, 3, 10 and 30 $\mu\text{mol/l}$ NTDP, respectively. To investigate the effect of extracellular free calcium on NFA, calcium-free extracellular fluid was utilized. The inhibitory effect of NFA on GABA-evoked inward currents was strongly suppressed by calcium-free extracellular fluid ($P<0.01$; Fig. 4). The 100 $\mu\text{mol/l}$ GABA-activated current




Figure 3. Effects of different-concentration of NPPB on GABA-activated inward currents. (A) Results of 100 $\mu\text{mol/l}$ GABA-activated inward currents with different concentrations of NPPB (1–30 $\mu\text{mol/l}$). (B) Inward current induced by GABA (100 $\mu\text{mol/l}$) was suppressed by 13.8 ± 6.7 , 23.2 ± 14.7 and $29.7 \pm 9.1\%$ by 3, 10 and 30 $\mu\text{mol/l}$ NPPB, respectively. * $P < 0.05$, ** $P < 0.01$ vs. control. Results as determined by paired t test. Numbers on bars = number of neurons. NPPB, 5-nitro-2-(3-phenylpropylamino) benzoic acid; GABA, γ -aminobutyrate.

with 100 $\mu\text{mol/l}$ NFA were $1,298.8 \pm 124.4$ pA; neurons = 22 and 775.9 ± 104.9 pA; neurons = 6; $P < 0.01$, in the presence and absence of calcium-free extracellular solution, respectively. The inhibition ratio of NFA on the GABA-evoked inward current was $8.4 \pm 7.2\%$ (neurons = 6, $P < 0.01$) in the calcium-free extracellular solution.

Effect of intracellular calcium on the GABA-induced inward current by intracellular calcium. To investigate the effect of intracellular free calcium on the GABA-induced inward current, BAPTA-AM was utilized and caffeine. The inhibitory effect of NFA on GABA-evoked inward current was strongly suppressed by BAPTA-AM and caffeine. The inhibition ratio of NFA on the GABA-evoked inward current was decreased by $48.2 \pm 15.7\%$ (neurons = 9, $P < 0.01$) and $38.7 \pm 13.2\%$ (neurons = 7, $P < 0.05$) with BAPTA-AM (100 $\mu\text{mol/l}$) and caffeine (30 $\mu\text{mol/l}$), respectively (Fig. 5).

Distribution of TMEM16A and TMEM16B subunits expressed in DRG neurons. Immunofluorescence staining revealed that TMEM16A and TMEM16B expression was widely distributed in DRG neurons, and was predominately located in the cell membranes of various diameters (Fig. 6A and B). Thus, TMEM16A and TMEM16B were co-expressed in the membranes of DRG neurons (Fig. 6C).

Discussion

GABA_A receptor mediates GABA-evoked membrane depolarization responses, or inward current, since the selective GABA_A receptor agonist muscimol mimicked GABA-evoked responses, and the selective GABA_A receptor antagonist




Figure 4. Effects of nitrendipine (NTDP) and extracellular Ca^{2+} on GABA-activated inward currents. (A) Results of 100 $\mu\text{mol/l}$ GABA-activated inward currents in the presence of 100 $\mu\text{mol/l}$ NFA, 100 $\mu\text{mol/l}$ NFA + calcium-free extracellular solution and 100 $\mu\text{mol/l}$ NFA + 10 $\mu\text{mol/l}$ NTDP. (B) Inward current induced by 100 $\mu\text{mol/l}$ GABA was suppressed by 62.2 ± 12.6 , 8.7 ± 7.6 and $8.4 \pm 7.2\%$ by NFA, NFA + NTDP and NFA + calcium-free extracellular solution, respectively. ** $P < 0.01$. Paired t test. Numbers on bars = number of neurons. NFA, niflumic acid; GABA, γ -aminobutyrate.

bicuculline blocked GABA-activated membrane responses in rat DRG neurons (17,19). The results of the present study indicated that NFA and NPPB, non-steroid anti-inflammatory agents, reduced GABA-activated inward currents. The present study also revealed that NTDP, an L-type calcium channel blocker, a calcium-free extracellular solution, BAPTA-AM, which is a membrane permeable Ca^{2+} chelator and caffeine, a Ca^{2+} consuming drug, also reduced the inhibitory effect of NFA on the GABA-activated inward current. Furthermore, the current study demonstrated that the TMEM16A and TMEM16B subunits were expressed in rat DRG neurons.

A number of cells express CaCCs, which have several physiological functions, including their being developmentally adjusted with maximum peak expression in the period of peripheral synaptogenesis in DRG neurons (20). An association was observed between the expression of CaCCs and the growth competence of sensory neurons (21). CaCCs activation augments after depolarization following spike firing of action potential in neonatal rat DRG neurons (22,23). Axotomy upregulates the expression of CaCCs in adult sensory, nodose and sympathetic ganglion neurons (21,24,25). Additionally, with regard to electrical activity, CaCCs have an important role in other basic cellular functions such as cell adhesion, apoptosis and potentially in volume regulation (26,27). In 2008, the




Figure 5. Effects of intracellular Ca^{2+} on GABA-activated inward currents. (A) Results of 100 $\mu\text{mol/l}$ GABA-activated inward currents in the presence of 100 $\mu\text{mol/l}$ NFA, 100 $\mu\text{mol/l}$ NFA + 100 $\mu\text{mol/l}$ BAPTA-AM and 100 $\mu\text{mol/l}$ NFA + 30 $\mu\text{mol/l}$ caffeine. (B) Inward current induced by GABA was suppressed by 61.8 ± 13.4 , 18.9 ± 17.6 and $28.4 \pm 13.2\%$ NFA, NFA + BAPTA-AM and NFA + caffeine, respectively. * $P < 0.05$, ** $P < 0.01$. Results as determined by paired t test. Numbers on bars = number of neurons. NFA, ni-flumic acid; GABA, γ -aminobutyrate; BAPTA, 1,2-bis(2-aminophenoxy) ethane- N,N,N',N' -tetraacetic acid.

announcements that three labs had cloned genes that encoded classical CaCCs generated considerable interest (28-30); the 2 genes that have been definitively shown to encode CaCCs termed TMEM16A and TMEM16B. CaCCs are activated by an increase in intracellular free calcium concentration following either internal calcium release from Ca^{2+} stores, or external calcium entry through Ca^{2+} channel (31). NFA and NPPB have been indicated to suppress the activity of CaCCs, and NPPB inhibits chloride ion flux through other anionic channels (32). The results of the present study show that NFA and NPPB are able to significantly inhibit the GABA-activated inward current, and NTDP, calcium-free extracellular fluid and BAPTA-AM may significantly reduce the inhibitory effect of NFA on GABA-activated inward current. The present results also suggest that NSAIDs have an important role in the GABA-activated inward current via CaCCs in DRG neurons in rats, and support the hypothesis proposed by the present study. GABA activates the GABA_A receptor to open chloride ion channels, the chloride ion efflux induces the depolarization response of the membrane of DRG neurons (17). Conversely, voltage dependent L-type Ca^{2+} channels may be activated by depolarization, and lead to increased intracellular Ca^{2+} . Furthermore, the NFA-induced increase in intracellular Ca^{2+} is likely due to Ca^{2+} release from an intracellular store (33-35). CaCCs are activated by an increase in intracellular calcium concentration, which in turn increases the driving force for




Figure 6. Confocal images of TMEM16A and TMEM16B subunits expressed in the L_{4-6} dorsal root ganglion (DRG). Immunofluorescent staining indicated that (A) TMEM16A and (B) TMEM16B expression were widely distributed in DRG neurons, and were mainly located in cell membranes of various diameters. (C) A and B overlaid (scale bars, 100 μm).

chloride ion efflux (28). Finally, the synergistic action of chloride ion efflux via GABA_A receptors and NFA-sensitive CaCCs results in GABA-activated currents or depolarization responses in rat DRG neurons. The depolarization arising from Cl^- efflux through CaCCs indicates a mechanism of electrical amplification of the GABA-activated currents (Fig. 7).

The binding of intracellular signaling molecules or extracellular ligands activates a conformational change that opens or closes the pores of ligand-gated ion channels such as GABA, glutamate, serotonin and acetylcholine, as well as the cyclic nucleotide-gated ion channels that have important roles in sensory biology (17). However, the structure and function of ligand-gated ion channels and their integral receptors have yet to be elucidated. Therefore, it is important to develop novel tools to investigate the interactions between various receptor subunits, which may benefit the future designing of receptor subtype-selective therapies.




Figure 7. Schematic displaying the effects of CaCCs on GABA-activated inward currents and depolarization. GABA activates the GABA_A receptor to open the Cl⁻ channel and the Cl⁻ efflux induces the depolarization response (inward current) of the membrane of dorsal root ganglion (DRG) neurons. Then, voltage dependent L-type Ca²⁺ channels are activated by the depolarization, and give rise to an increase in intracellular Ca²⁺. CaCCs are activated by an increase in intracellular Ca²⁺ concentration which, in turn, increases the driving force for Cl⁻ efflux. Finally, the synergistic action of the chloride ion efflux through GABA_A receptors and NFA-sensitive CaCCs causes GABA-activated currents or depolarization response in rat DRG neurons. '-' expressed function was inhibitory, '?' expressed function was not clear. GABA, γ -aminobutyrate; NPPB, 5-nitro-2-(3-phenylpropylamino) benzoic acid; NFA, niflumic acid; CaCC, Ca²⁺-activated Cl⁻ channels; BAPTA-AM, 1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid; NTDP, nifentridine.

Ion channel gating is influenced by several factors, including the binding of another ligand and intracellular Ca²⁺ (36). GABA_A integrates the actions of a wide range of therapeutic agents, including steroids, benzodiazepines, barbiturates, convulsants and anesthetics (2). Previous studies have indicated that NFA is able to directly act on GABA_A and NMDA receptors (5,12,37). Notably, another chloride blocker, mefenamic acid, is also able to directly activate GABA_A receptors (38). NFA functions as a positive allosteric modulator of $\alpha_1\beta_2\gamma_2$, and a negative modulator of $\alpha_6\beta_2$ and $\alpha_6\beta_2\gamma_2$ (and $\alpha_1\beta_2$) GABA_A receptors. Despite the knowledge that NFA shares the same site as furosemide to mediate its inhibitory effect, the site for the positive regulation remains elusive, and is dependant on the presence of the γ_2 subunit, yet separable from the benzodiazepine binding site (11,12).

It has been suggested that the niflumate potentiation of GABA_A function is through a pure direct allosteric mechanism (12). Conversely, it has been reported that the activation of GABA_A opens NFA-sensitive anion channels (39,40). Furthermore, GABA_A-mediated chloride ion influx lowers the magnitudes of NFA- and NPPB-sensitive chloride currents in motoneurons. NFA and other fenamate blockers of CaCCs (41,42), and recombinant GABA_A (43). The expression levels of rat homopentamer GABA_A receptors display GABA-independent activation (44). In addition, NFA activates single chloride ion channels, likely to be an isoform of the GABA_A receptors in mouse sperm (45). Therefore, it is

possible that alternative mechanisms exists for NFA action on the GABA_A receptor.

In the present study, NFA did not alter the EC₅₀ value but reduced the maximal response of GABA currents, which is consistent with noncompetitive antagonism. Similar results have been reported for furosemide (46) and NFA (12) previously. Changes in the expression and function of α_2 , but not α_6 subunits of GABA_A, were observed in L₄-L₆ DRG neurons by whole-cell patch-clamp and immunofluorescence.

Further experiments are needed to determine whether NFA and NPPB may directly act upon the GABA_A α_2 subunit in DRG neurons, and the physiological activator of CaCCs may also provide further elucidation regarding the contribution of CaCCs to electrical activity. As NSAIDs are highly subtype-selective, further studies to investigate its behavioral and cognitive effects are warranted (11). We propose that expression of CaCCs in DRG should increase in response to peripheral nerve injury, and enhance the responses of GABA_A receptors. Therefore, the upregulation of CaCCs may prominently enhance 'the presynaptic inhibition' of GABA in the primary afferent endings, and have involvement in pain modulation.

Acknowledgements

The authors thank Dr Hong-Zhen Hu for reading the manuscript and for valuable suggestions. The present study was supported by the National Natural Science Foundation of China (grant no. 30160026 to Dr Jun-Qiang Si) and the Youth Science and Technology Innovation Special Foundation of Xinjiang Production and Construction Corps (China; grant no. 2010JC33 to Dr Li Li).

References

- Whiting PJ, Bonnert TP, McKernan RM, Farrar S, Le Bourdellès B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghji DJ, *et al*: Molecular and functional diversity of the expanding GABA-A receptor gene family. *Ann N Y Acad Sci* 868: 645-653, 1999.
- Mehta AK and Ticku MK: An update on GABAA receptors. *Brain Res Brain Res Rev* 29: 196-217, 1999.
- Smith AJ, Oxley B, Malpas S, Pillai GV and Simpson PB: Compounds exhibiting selective efficacy for different beta subunits of human recombinant gamma-aminobutyric acid A receptors. *J Pharmacol Exp Ther* 311: 601-609, 2004.
- Jensen ML, Timmermann DB, Johansen TH, Schousboe A, Varming T and Ahring PK: The beta subunit determines the ion selectivity of the GABAA receptor. *J Biol Chem* 277: 41438-41447, 2002.
- Babot Z, Cristófol R and Suñol C: Excitotoxic death induced by released glutamate in depolarized primary cultures of mouse cerebellar granule cells is dependent on GABAA receptors and niflumic acid-sensitive chloride channels. *Eur J Neurosci* 21: 103-112, 2005.
- Wallenstein MC: Attenuation of epileptogenesis by nonsteroidal anti-inflammatory drugs in the rat. *Neuropharmacology* 30: 657-663, 1991.
- McKernan RM and Whiting PJ: Which GABAA-receptor subtypes really occur in the brain? *Trends Neurosci* 19: 139-143, 1996.
- Greenwood IA and Large WA: Comparison of the effects of fenamates on Ca-activated chloride and potassium currents in rabbit portal vein smooth muscle cells. *Br J Pharmacol* 116: 2939-2948, 1995.
- Malykhina AP, Shoeb F and Akbarali HI: Fenamate-induced enhancement of heterologously expressed HERG currents in *Xenopus oocytes*. *Eur J Pharmacol* 452: 269-277, 2002.

10. Korpi ER, Gründer G and Lüddens H: Drug interactions at GABA (A) receptors. *Prog Neurobiol* 67: 113-159, 2002.
11. Halliwell RF, Thomas P, Patten D, James CH, Martinez-Torres A, Miledi R and Smart TG: Subunit-selective modulation of GABAA receptors by the non-steroidal anti-inflammatory agent, mefenamic acid. *Eur J Neurosci* 11: 2897-2905, 1999.
12. Sinkkonen ST, Mansikkamäki S, Möykkynen T, Lüddens H, Uusi-Oukari M and Korpi ER: Receptor subtype-dependent positive and negative modulation of GABA (A) receptor function by niflumic acid, a nonsteroidal anti-inflammatory drug. *Mol Pharmacol* 64: 753-763, 2003.
13. Whittemore ER, Yang W, Drewe JA and Woodward RM: Pharmacology of the human gamma-aminobutyric acid A receptor alpha 4 subunit expressed in *Xenopus laevis* oocytes. *Mol Pharmacol* 50: 1364-1375, 1996.
14. Thompson SA, Whiting PJ and Wafford KA: Barbiturate interactions at the human GABAA receptor: Dependence on receptor subunit combination. *Br J Pharmacol* 117: 521-527, 1996.
15. Belelli D, Lambert JJ, Peters JA, Wafford K and Whiting PJ: The interaction of the general anesthetic etomidate with the gamma-aminobutyric acid type A receptor is influenced by a single amino acid. *Proc Natl Acad Sci USA* 94: 11031-11036, 1997.
16. Institute of Laboratory Animal Resources (US). Committee on Care, Use of Laboratory Animals, and National Institutes of Health (US). Division of Research Resources: Guide for the care and use of laboratory animals. 8th edition. National Academies Press, Washington, DC, 2011.
17. Si JQ, Zhang ZQ, Li CX, Wang LF, Yang YL and Li ZW: Modulatory effect of substance P on GABA-activated currents from rat dorsal root ganglion. *Acta Pharmacol Sin* 25: 623-629, 2004.
18. Cheng HJ, Ma KT, Li L, Zhao L, Wang Y and Si JQ: Differential expression of alpha-adrenoceptor subtypes in rat dorsal root ganglion after chronic constriction injury. *Hua Zhong Ke Ji Da Xue Xue Bao* 34: 322-329, 2014 (In Chinese).
19. Ma KT, Si JQ, Zhang ZQ, Zhao L, Fan P, Jin JL, Li XZ and Zhu L: Modulatory effect of CCK-8S on GABA-induced depolarization from rat dorsal root ganglion. *Brain Res* 1121: 66-75, 2006.
20. Bernheim L, Bader CR, Bertrand D and Schlichter R: Transient expression of a Ca²⁺-activated Cl⁻ current during development of quail sensory neurons. *Dev Biol* 136: 129-39, 1989.
21. André S, Boukhaddaoui H, Campo B, Al-Jumaily M, Mayeux V, Greuet D, Valmier J and Scamps F: Axotomy-induced expression of calcium-activated chloride current in subpopulations of mouse dorsal root ganglion neurons. *J Neurophysiol* 90: 3764-3773, 2003.
22. De Castro F, Geijo-Barrientos E and Gallego R: Calcium-activated chloride current in normal mouse sympathetic ganglion cells. *J Physiol* 498: 397-408, 1997.
23. Mayer ML: A calcium-activated chloride current generates the after-depolarization of rat sensory neurones in culture. *J Physiol* 364: 217-239, 1985.
24. Sánchez-Vives MV and Gallego R: Calcium-dependent chloride current induced by axotomy in rat sympathetic neurons. *J Physiol* 475: 391-400, 1994.
25. Lancaster E, Oh EJ, Gover T and Weinreich D: Calcium and calcium-activated currents in vagotomized rat primary vagal afferent neurons. *J Physiol* 540: 543-556, 2002.
26. Abdel-Ghany M, Cheng HC, Elble RC and Pauli BU: Focal adhesion kinase activated by beta (4) integrin ligation to mCLCA1 mediates early metastatic growth. *J Biol Chem* 277: 34391-34400, 2002.
27. Elble RC and Pauli BU: Tumor suppression by a proapoptotic calcium-activated chloride channel in mammary epithelium. *J Biol Chem* 276: 40510-40517, 2001.
28. Yang YD, Cho H, Koo JY, Tak MH, Cho Y, Shim WS, Park SP, Lee J, Lee B, Kim BM, *et al*: TMEM16A confers receptor-activated calcium-dependent chloride conductance. *Nature* 455: 1210-1215, 2008.
29. Caputo A, Caci E, Ferrera L, Pedemonte N, Barsanti C, Sondo E, Pfeffer U, Ravazzolo R, Zegarra-Moran O and Galletta LJ: TMEM16A, a membrane protein associated with calcium-dependent chloride channel activity. *Science* 322: 590-594, 2008.
30. Schroeder BC, Cheng T, Jan YN and Jan LY: Expression cloning of TMEM16A as a calcium-activated chloride channel subunit. *Cell* 134: 1019-1029, 2008.
31. Scott RH, Sutton KG, Griffin A, Stapleton SR and Currie KP: Aspects of calcium-activated chloride currents: A neuronal perspective. *Pharmacol Ther* 66: 535-565, 1995.
32. Frings S, Reuter D and Kleene SJ: Neuronal Ca²⁺-activated Cl⁻channels-homing in on an elusive channel species. *Prog Neurobiol* 60: 247-289, 2000.
33. Cruickshank SF, Baxter LM and Drummond RM: The Cl⁻ channel blocker niflumic acid releases Ca(2+) from an intracellular store in rat pulmonary artery smooth muscle cells. *Br J Pharmacol* 140: 1442-1450, 2003.
34. Li L, Ma KT, Zhao L and Si JQ: Niflumic acid hyperpolarizes the smooth muscle cells by opening BK(Ca) channels through ryanodine-sensitive Ca(2+) release in spiral modiolary artery. *Sheng Li Xue Bao* 60: 743-750, 2008.
35. Li L, Ma KT, Zhao L, Si JQ, Zhang ZS, Zhu H and Li J: Niflumic acid hyperpolarizes smooth muscle cells via calcium-activated potassium channel in spiral modiolary artery of guinea pigs. *Acta Pharmacol Sin* 29: 789-799, 2008.
36. Levitan IB: Signaling protein complexes associated with neuronal ion channels. *Nat Neurosci* 9: 305-310, 2006.
37. Lerma J and Martin del Rio R: Chloride transport blockers prevent N-methyl-D-aspartate receptor-channel complex activation. *Mol Pharmacol* 41: 217-222, 1992.
38. Coyne L, Su J, Patten D and Halliwell RF: Characterization of the interaction between fenamates and hippocampal neuron GABA(A) receptors. *Neurochem Int* 51: 440-446, 2007.
39. Raiteri L, Schmid G, Prestipino S, Raiteri M and Bonanno G: Activation of alpha 6 GABAA receptors on depolarized cerebellar parallel fibers elicits glutamate release through anion channels. *Neuropharmacology* 41: 943-951, 2001.
40. Van Damme P, Callewaert G, Eggermont J, Robberecht W and Van Den Bosch L: Chloride influx aggravates Ca²⁺-dependent AMPA receptor-mediated motoneuron death. *J Neurosci* 23: 4942-4950, 2003.
41. White MM and Aylwin M: Niflumic and flufenamic acids are potent reversible blockers of Ca²⁺ (+)-activated Cl⁻channels in *Xenopus* oocytes. *Mol Pharmacol* 37: 720-724, 1990.
42. Korn SJ, Bolden A and Horn R: Control of action potentials and Ca²⁺ influx by the Ca (2+)-dependent chloride current in mouse pituitary cells. *J Physiol* 439: 423-437, 1991.
43. Woodward RM, Polenzani L and Miledi R: Effects of fenamates and other nonsteroidal anti-inflammatory drugs on rat brain GABAA receptors expressed in *Xenopus* oocytes. *J Pharmacol Exp Ther* 268: 806-817, 1994.
44. Sigel E, Baur R, Malherbe P and Möhler H: The rat beta 1-subunit of the GABAA receptor forms a picrotoxin-sensitive anion channel open in the absence of GABA. *FEBS Lett* 257: 377-379, 1989.
45. Espinosa F, de la Vega-Beltrán JL, López-González I, Delgado R, Labarca P and Darszon A: Mouse sperm patch-clamp recordings reveal single Cl⁻channels sensitive to niflumic acid, a blocker of the sperm acrosome reaction. *FEBS Lett* 426: 47-51, 1998.
46. Korpi ER, Kuner T, Seeburg PH and Lüddens H: Selective antagonist for the cerebellar granule cell-specific gamma-aminobutyric acid type A receptor. *Mol Pharmacol* 47: 283-289, 1995.