

Association between alcohol and Alzheimer's disease (Review)

WEN-JUAN HUANG, XIA ZHANG and WEI-WEI CHEN

Department of Neurology, Xuzhou Central Hospital, Xuzhou, Jiangsu 221009, P.R. China

Received February 18, 2016; Accepted June 14, 2016

DOI: 10.3892/etm.2016.3455

Abstract. Alzheimer's disease (AD) is a neurodegenerative disease characterized by dense deposition of amyloid- β (A β) protein in the brain, failure of the memory and dementia. At present, there is no cure for AD and current treatments only provide a temporary reduction of symptoms. Thus, there is a need for effective preventive/curative strategic approaches. Accordingly, epidemiological studies have reported a reduction in the prevalence of AD in individuals ingesting low amounts of alcohol, while a moderate consumption of ethanol may protect against A β . These data are conflicting with other observations that assigned detrimental effects of heavy alcohol use on brain function, which are apparently similar to those observed in AD. These discrepancies questioned whether or not alcohol is a protective agent against the development of AD, whether the probable protective effects are influenced by the quantity and/or frequency of drinking. These issues are addressed in this review with the aim to suggest the real risk of alcohol for developing or preventing AD.

Contents

1. Introduction
2. Alcohol consumption as a risk factor for developing AD
3. Potential benefits of alcohol on AD
4. Conclusion

1. Introduction

Alzheimer's disease (AD), the most common form of dementia affecting populations aged over 65 years worldwide, is sporadic, genetically non-obvious, and rarely inherited (1). The main well-known pathological features of the disease

include the abnormal extracellular deposition of misfolded amyloid- β (A β) senile plaques in brain parenchyma and cerebral vessels, the intracellular accumulation of hyperphosphorylated tau (τ) in neurofibrillary tangles (NFTs), chronic neuroinflammation, neuronal loss and severe brain atrophy as well as progressive loss of memory (2,3). Consequently, in the last 30 years, the deregulation of A β metabolism (oligomerization, aggregation and plaque formation) has been the major target for therapeutic intervention (4), and only marginal effects have been registered (5-7), suggesting the need for preventive/curative treatments or alternative solutions.

Therefore, although the causes of AD remain unknown and cures or universally effective treatments, are not available, most experts have highlighted a broad constellation of contributing risk factors. Among these risk factors, alcohol consumption, associated with extensive cognitive problems (8), including alcoholic dementia (9), has been targeted. Similar features have been denoted between the effects of alcohol on cognition, brain disorder and brain biochemistry with the biological effects of AD, suggesting that the use of alcohol may constitute a risk for aggravating or developing AD (10). In line with this view, AD patients with a habitual drinking history have shown cognitive improvement during the clinical course of abstinence (11). By contrast, this effect was not observed in patients consuming high amounts of alcohol prior to diagnosis of AD (11). In line with these data, there is another emerging body of literature that contends alcohol consumption, particularly red wine, may rather serve as a protective factor for cognitive decline (12,13). Several epidemiological studies have shown that low or moderate wine consumption can be effective in retarding age-related cognitive decline (14), possibly linked to polyphenols present in beverages.

However, the studies promoting the benefits of alcohol on AD exclusively focus on moderate alcohol consumption (12), the restriction of alcohol to only an elderly population (13), and broad classification of cognitive decline (15). Relevant issues remained regarding the protection, the aggravating or detrimental effects of alcohol consumption or whether protective effects are simply influenced by the quantity and/or frequency of drinking. Despite the limiting factors identified regarding the beneficial effects of alcohol consumption (16), a single-target therapeutic strategy appears to produce only suboptimal results and a broader neuroprotective approach, at least theoretically, appears more appealing (17). Thus, the aim of the present review was to discuss the association between alcohol consumption and AD.

Correspondence to: Dr Wen-Juan Huang, Department of Neurology, Xuzhou Central Hospital, 199 Jiefang South Road, Xuzhou, Jiangsu 221009, P.R. China
E-mail: ftmlb8556215@163.com

Key words: Alzheimer's disease, risk factor, alcohol consumption

2. Alcohol consumption as a risk factor for developing AD

Heavy alcohol consumption impairs cognitive performance with immediate and long-term effects on the brain anatomy and neuropsychological functioning (18-21). Cognitive impairment is related to clinical dementia as it accelerates shrinkage and atrophy of the brain, leading to critical determinant of neurodegenerative changes and cognitive decline in aging (22). However, these morphological changes induced by alcohol consumption may be reversible unlike AD or aging (23), as atrophy decreases, with cognitive improvement after abstinence from alcohol (24,25). Other data showed that morphological changes in the brain are associated with the loss of a number of nerve cells occurring in the white matter, which largely comprises nerve fibers that connect neurons (26) and/or cortical cholinergic neurons (24), known to be affected in AD. This link renders plausible that alcohol use may be linked to AD as the cholinergic system plays an important role in memory. This role is confirmed and its deficits are well established in AD (27). Chronic alcohol use causes degeneration of cholinergic neurons (28), or decreases receptors of cholinergic system in AD. These may aggravate the reduction of cholinergic neurons already present in AD patients. However, the improvement of cognitive function in alcoholics after abstinence from alcohol suggests that cognitive deficits may reflect neurochemical alterations rather than neuronal loss (24,28). Therefore, without appearing as an accelerator of AD process, alcohol may induce its effects on the cholinergic system, independently from the cholinergic deficits caused by AD (29,30). However, alcohol-related brain damage appears to differ in young and older alcohol consumers (24), although data suggest that alcohol abuse may accelerate aging-related changes in the brain at any age and that older adults may be more vulnerable to the effects of alcohol (8). Apart from cholinergic deficit, another negative effect of heavy alcohol consumption on cognitive function may be attributed to nutritional deficiency or vascular change (31,32), which consist of damage that may be relatively irreversible even after abstinence from heavy alcohol consumption (11).

Therefore, despite the above evidence showing cognitive detrimental effects of heavy alcohol consumption, to the best of our knowledge, no study has established a clear association between alcohol consumption and AD (33). More consistently, recent genetical study on a Japanese population provided evidence that the mitochondrial aldehyde dehydrogenase 2 (ALDH2*2, metabolizes acetaldehyde into acetate, protecting against oxidative stress and playing an important role in the development of AD), and two functional single-nucleotide polymorphisms (SNPs) of the dopamine- β -hydroxylase (*DBH*) gene, involved in the pathophysiology of alcoholism and whose activity is reduced in the neocortex of AD, did not modify the risk for developing AD, suggesting that the polymorphism of the *ALDH2* and *DBH* genes were not associated with AD (34). Nevertheless, future studies must be undertaken on other populations worldwide.

3. Potential benefits of alcohol on AD

Low-to-moderate alcohol intake is considered to protect against neurodegeneration pathology (13,15), dementia (35-39)

and cognitive deterioration (40-43). Of the biologic mechanisms suggested to explain such potential beneficial effects on the brain, there are mainly the antioxidant properties of wine flavonoids (44), the effects against A β (45) and the prevention of ischemia or stroke by alcohol (46). Specifically, polyphenols, members of a large family of plant-derived compounds, are molecules containing one or more phenolic group. There are thousands of polyphenols that have been identified thus far including, bioflavonoids (anthocyanins, flavanols, flavanols, favones, flavanones, isoflavones and proanthocyanins), coumestans, ligans and stilbenoids (47). Polyphenolics are in general antioxidant molecules that reduce the *in vitro* process aggregation of A β , reducing the neuronal death of cortical neurons preventing neurodegeneration (48). The morin for example, a specific flavonoid described in red wine exhibited significant effects in preventing aggregation of A β (49). In agreement with these data, red wine Cabernet Sauvignon significantly reduced the number of A β plaque-induced neuropathology and attenuated a spatial memory decrease in an adult Tg2576 mouse model of AD (50). Another flavonoid that plays an important role in red wine is resveratrol. Resveratrol is known to protect against cardiovascular diseases (which are risk factors for developing AD) and various types of cancer, together with the promotion of the antiaging effect, the modulation of pathomechanisms of debilitating neurological disorder such as strokes, ischemia and Huntington's disease, as well as protection against neuronal degeneration (14,51). Other natural molecules including fulvic acid, altered the aggregation mechanism of τ proteins, a critical protein involved in the stabilization of microtubule and axonal transport, found to be involved in AD pathogenesis (52). However, whatever the facts around the benefits that may follow the potential benefits of low-to-moderate consumption of alcohol, the importance of drinking patterns and specific beverages consumed remain elusive. The operational definition of low or moderate drinking which may vary greatly across studies (53) and the concept of a moderate drinker, which may also be imprecise, comprising a wide range measure that may include those consuming less than one drink a day (54). Red wine consumption appears to promote far more protective effects than the consumption of other ethanol containing beverages (55). This is in keeping with variations that may be introduced by consumers who sometimes associate the consumption of tobacco, or the possible toxicity effects of chronic exposure of the liver to alcohol (56), which can also contribute to brain alteration in regions involved in memory (57).

Taken together, the data provide insufficient evidence to suggest abstainers should initiate alcohol consumption to protect against dementia or AD.

4. Conclusion

Based on the abovementioned research data, no relationship between alcohol consumption and AD exists. In addition, although low-to-moderate consumption of alcohol may protect against AD, leading to benefits on neurodegeneration, A β , oxidative stress and τ neurofibrillary tangle formation suggest bias regarding the definition of low-to-moderate consumption as well as the variability of beverages containing alcohol. In addition, the absence of studies on the possible side effects of

chronic exposure to alcohol on peripheral organs such as liver, and kidney lead to the necessity to delineate global and standard protocols for advanced studies. These studies render the benefits associated with low-to-moderate alcohol consumption against AD. However, the results should be considered as controversial and insufficient to suggest abstainers that initiate alcohol consumption in a preventive manner against AD.

References

- Kumar A, Singh A and Ekavali E: A review on Alzheimer's disease pathophysiology and its management: an update. *Pharmacol Rep* 67: 195-203, 2015.
- Querfurth HW and LaFerla FM: Alzheimer's disease. *N Engl J Med* 362: 329-344, 2010.
- Medeiros R and LaFerla FM: Astrocytes: conductors of the Alzheimer disease neuroinflammatory symphony. *Exp Neurol* 239: 133-138, 2013.
- Mucke L and Selkoe DJ: Neurotoxicity of amyloid β -protein: synaptic and network dysfunction. *Cold Spring Harb Perspect Med* 2: a006338, 2012.
- Doody RS, Thomas RG, Farlow M, Iwatsubo T, Vellas B, Joffe S, Kieburtz K, Raman R, Sun X, Aisen PS, *et al*: Alzheimer's Disease Cooperative Study Steering Committee; Solanezumab Study Group: Phase 3 trials of solanezumab for mild-to-moderate Alzheimer's disease. *N Engl J Med* 370: 311-321, 2014.
- Krstic D and Knuesel I: Deciphering the mechanism underlying late-onset Alzheimer disease. *Nat Rev Neurol* 9: 25-34, 2013.
- Reitz C, Conrad C, Roszkowski K, Rogers RS and Mayeux R: Effect of genetic variation in LRRTM3 on risk of Alzheimer disease. *Arch Neurol* 69: 894-900, 2012.
- Evert DL and Oscar-Berman M: Alcohol-related cognitive impairments. *Alcohol Health Res World* 19: 89, 1995.
- Smith DM and Atkinson RM: Mood disorders secondary to drugs and pharmacologic agents. *Semin Clin Neuropsychiatry* 2: 285-295, 1997.
- Tyas SL: Are tobacco and alcohol use related to Alzheimer's disease? A critical assessment of the evidence and its implications. *Addict Biol* 1: 237-254, 1996.
- Toda A, Tagata Y, Nakada T, Komatsu M, Shibata N and Arai H: Changes in Mini-Mental State Examination score in Alzheimer's disease patients after stopping habitual drinking. *Psychogeriatrics* 13: 94-98, 2013.
- Neafsey EJ and Collins MA: Moderate alcohol consumption and cognitive risk. *Neuropsychiatr Dis Treat* 7: 465-484, 2011.
- Peters R, Peters J, Warner J, Beckett N and Bulpitt C: Alcohol, dementia and cognitive decline in the elderly: a systematic review. *Age Ageing* 37: 505-512, 2008.
- Granzotto A and Zatta P: Resveratrol and Alzheimer's disease: Message in a bottle on red wine and cognition. *Front Aging Neurosci* 6: 95, 2014.
- Anstey KJ, Mack HA and Cherbuin N: Alcohol consumption as a risk factor for dementia and cognitive decline: meta-analysis of prospective studies. *Am J Geriatr Psychiatry* 17: 542-555, 2009.
- Panza F, Frisardi V, Seripa D, Logroscino G, Santamato A, Imbimbo BP, Scafato E, Pilotto A and Solfrizzi V: Alcohol consumption in mild cognitive impairment and dementia: harmful or neuroprotective? *Int J Geriatr Psychiatry* 27: 1218-1238, 2012.
- Mudher A and Lovestone S: Alzheimer's disease - do tauists and baptists finally shake hands? *Trends Neurosci* 25: 22-26, 2002.
- Ridderinkhof KR, de Vlugt Y, Bramlage A, Spaan M, Elton M, Snel J and Band GP: Alcohol consumption impairs detection of performance errors in mediofrontal cortex. *Science* 298: 2209-2211, 2002.
- Zhu W, Volkow ND, Ma Y, Fowler JS and Wang G-J: Relationship between ethanol-induced changes in brain regional metabolism and its motor, behavioural and cognitive effects. *Alcohol* 39: 53-58, 2004.
- Oslin DW and Cary MS: Alcohol-related dementia: validation of diagnostic criteria. *Am J Geriatr Psychiatry* 11: 441-447, 2003.
- Weissenborn R and Duka T: Acute alcohol effects on cognitive function in social drinkers: their relationship to drinking habits. *Psychopharmacology (Berl)* 165: 306-312, 2003.
- Meyer JS, Terayama Y, Konno S, Akiyama H, Margishvili GM and Mortel KF: Risk factors for cerebral degenerative changes and dementia. *Eur Neurol* 39 (Suppl 1): 7-16, 1998.
- Wilkinson DA and Poulos CX: The chronic effects of alcohol on memory. A contrast between a unitary and dual system approach. *Recent Dev Alcohol* 5: 5-26, 1987.
- Krill JJ and Halliday GM: Brain shrinkage in alcoholics: a decade on and what have we learned? *Prog Neurobiol* 58: 381-387, 1999.
- Reed LJ, Lasserson D, Marsden P, Stanhope N, Stevens T, Bello F, Kingsley D, Colchester A and Kopelman MD: FDG-PET findings in the Wernicke-Korsakoff syndrome. *Cortex* 39: 1027-1045, 2003.
- Jensen GB and Pakkenberg B: Do alcoholics drink their neurons away? *Lancet* 342: 1201-1204, 1993.
- Tyas SL: Alcohol use and the risk of developing Alzheimer's disease. *Alcohol Res Health* 25: 299-306, 2001.
- Arendt T: Impairment in memory function and neurodegenerative changes in the cholinergic basal forebrain system induced by chronic intake of ethanol. In: *Cell and Animal Models in Aging and Dementia Research*. Hoyer S, Müller D and Plaschke K (eds). Vol 44. Springer, Vienna, pp173-187, 1994. doi: 10.1007/978-3-7091-9350-1_13.
- Freund G and Ballinger WE Jr: Alzheimer's disease and alcoholism: possible interactions. *Alcohol* 9: 233-240, 1992.
- Diamond I and Gordon AS: Cellular and molecular neuroscience of alcoholism. *Physiol Rev* 77: 1-20, 1997.
- Harper C and Matsumoto I: Ethanol and brain damage. *Curr Opin Pharmacol* 5: 73-78, 2005.
- Schmidt KS, Gallo JL, Ferri C, Giovannetti T, Sestito N, Libon DJ and Schmidt PS: The neuropsychological profile of alcohol-related dementia suggests cortical and subcortical pathology. *Dement Geriatr Cogn Disord* 20: 286-291, 2005.
- Piazza-Gardner AK, Gaffud TJ and Barry AE: The impact of alcohol on Alzheimer's disease: a systematic review. *Aging Ment Health* 17: 133-146, 2013.
- Komatsu M, Shibata N, Ohnuma T, Kuerban B, Tomson K, Toda A, Tagata Y, Nakada T, Shimazaki H and Arai H: Polymorphisms in the aldehyde dehydrogenase 2 and dopamine β hydroxylase genes are not associated with Alzheimer's disease. *J Neural Transm Vienna* 121: 427-432, 2014.
- Orgogozo JM, Dartigues JF, Lafont S, Letenneur L, Commenges D, Salamon R, Renaud S and Breteler MB: Wine consumption and dementia in the elderly: a prospective community study in the Bordeaux area. *Rev Neurol (Paris)* 153: 185-192, 1997.
- Truelsen T, Thudium D and Grønbaek M; Copenhagen City Heart Study: Amount and type of alcohol and risk of dementia: The Copenhagen City Heart Study. *Neurology* 59: 1313-1319, 2002.
- Huang W, Qiu C, Winblad B and Fratiglioni L: Alcohol consumption and incidence of dementia in a community sample aged 75 years and older. *J Clin Epidemiol* 55: 959-964, 2002.
- Luchsinger JA, Tang MX, Siddiqui M, Shea S and Mayeux R: Alcohol intake and risk of dementia. *J Am Geriatr Soc* 52: 540-546, 2004.
- Mukamal KJ, Kuller LH, Fitzpatrick AL, Longstreth WT Jr, Mittleman MA and Siscovick DS: Prospective study of alcohol consumption and risk of dementia in older adults. *JAMA* 289: 1405-1413, 2003.
- Cervilla JA, Prince M and Mann A: Smoking, drinking, and incident cognitive impairment: a cohort community based study included in the Gospel Oak project. *J Neurol Neurosurg Psychiatry* 68: 622-626, 2000.
- Christian JC, Reed T, Carmelli D, Page WF, Norton JA Jr and Breitner JC: Self-reported alcohol intake and cognition in aging twins. *J Stud Alcohol* 56: 414-416, 1995.
- Leroi I, Sheppard J-M and Lyketsof CG: Cognitive function after 11.5 years of alcohol use: relation to alcohol use. *Am J Epidemiol* 156: 747-752, 2002.
- Zuccalà G, Onder G, Pedone C, Cesari M, Landi F, Bernabei R and Cocchi A; Gruppo Italiano di Farmacoepidemiologia nell'Anziano Investigators: Dose-related impact of alcohol consumption on cognitive function in advanced age: results of a multicenter survey. *Alcohol Clin Exp Res* 25: 1743-1748, 2001.
- Commenges D, Scotet V, Renaud S, Jacqmin-Gadda H, Barberger-Gateau P and Dartigues J-F: Intake of flavonoids and risk of dementia. *Eur J Epidemiol* 16: 357-363, 2000.
- Ho L, Chen LH, Wang J, Zhao W, Talcott ST, Ono K, Teplow D, Humala N, Cheng A, Percival SS, *et al*: Heterogeneity in red wine polyphenolic contents differentially influences Alzheimer's disease-type neuropathology and cognitive deterioration. *J Alzheimers Dis* 16: 59-72, 2009.

46. Berger K, Ajani UA, Kase CS, Gaziano JM, Buring JE, Glynn RJ and Hennekens CH: Light-to-moderate alcohol consumption and risk of stroke among U.S. male physicians. *N Engl J Med* 341: 1557-1564, 1999.
47. Pasinetti GM: Novel role of red wine-derived polyphenols in the prevention of Alzheimer's disease dementia and brain pathology: experimental approaches and clinical implications. *Planta Med* 78: 1614-1619, 2012.
48. Kang I-J, Jeon YE, Yin XF, Nam J-S, You SG, Hong MS, Jang BG and Kim M-J: Butanol extract of *Ecklonia cava* prevents production and aggregation of beta-amyloid, and reduces beta-amyloid mediated neuronal death. *Food Chem Toxicol* 49: 2252-2259, 2011.
49. Noor H, Cao P and Raleigh DP: Morin hydrate inhibits amyloid formation by islet amyloid polypeptide and disaggregates amyloid fibers. *Protein Sci* 21: 373-382, 2012.
50. Wang J, Ho L, Zhao Z, Seror I, Humala N, Dickstein DL, Thiyagarajan M, Percival SS, Talcott ST and Pasinetti GM: Moderate consumption of Cabernet Sauvignon attenuates Abeta neuropathology in a mouse model of Alzheimer's disease. *FASEB J* 20: 2313-2320, 2006.
51. Anekonda TS: Resveratrol - a boon for treating Alzheimer's disease? *Brain Res Brain Res Rev* 52: 316-326, 2006.
52. Cornejo A, Jiménez JM, Caballero L, Melo F and Maccioni RB: Fulvic acid inhibits aggregation and promotes disassembly of tau fibrils associated with Alzheimer's disease. *J Alzheimers Dis* 27: 143-153, 2011.
53. Ganguli M, Vander Bilt J, Saxton JA, Shen C and Dodge HH: Alcohol consumption and cognitive function in late life: a longitudinal community study. *Neurology* 65: 1210-1217, 2005.
54. Reid MC, Van Ness PH, Hawkins KA, Towle V, Concato J and Guo Z: Light to moderate alcohol consumption is associated with better cognitive function among older male veterans receiving primary care. *J Geriatr Psychiatry Neurol* 19: 98-105, 2006.
55. Baur JA and Sinclair DA: Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 5: 493-506, 2006.
56. Sun Q, Long Z, Wu H, Liu Y, Wang L, Zhang X, Wang X and Hai C: Effect of alcohol on diethylnitrosamine-induced hepatic toxicity: critical role of ROS, lipid accumulation, and mitochondrial dysfunction. *Exp Toxicol Pathol* 67: 491-498, 2015.
57. Matsumoto H and Matsumoto I: Alcoholism: protein expression profiles in a human hippocampal model. *Expert Rev Proteomics* 5: 321-331, 2008.