Outcome of antegrade intramedullary fixation for juvenile fifth metacarpal neck fracture with titanium elastic nail

KAIYING SHEN^{1*}, YUNLAN XU^{1*} , DAN CAO², ZHIGANG WANG¹ and HAIQING CAI¹

¹Department of Pediatric Orthopedics, Shanghai Children's Medical Center, Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai 200127; ²Department of Gynecology and Obstetrics, International Peace Maternity and Child Health Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai 20030, P.R. China

Received February 1, 2016; Accepted February 1, 2017

DOI: 10.3892/etm.2017.4369

Abstract. The purpose of the current study was to assess the outcome of antegrade intramedullary fixation with titanium elastic nail (TEN) in displaced fifth metacarpal neck fractures. The present study included 69 consecutive juvenile patients with displaced fifth metacarpal fractures. The head-shaft angle of the fifth metacarpal and range of motion (ROM) of the metacarpophalangeal (MCP) joint were evaluated. A disabilities of the arm, shoulder and hand (DASH) questionnaire was used to assess upper arm function. The head-shaft angle of the affected side was significantly improved postoperatively (P<0.05). No marked difference was observed between the affected and unaffected side in head-shaft angle and ROM. The average DASH score was 1.7 (range, 0-6.0). All patients obtained anatomical reduction postoperatively and the average healing time was 5.7±1.09 weeks (range, 5-10 weeks) with no non-union cases. Therefore antegrade intramedullary fixation with TEN is recommended as an easy, reliable and minimally invasive surgical technique for treating displaced fifth metacarpal neck fractures.

Introduction

The term 'fifth metacarpal neck fracture' is known as 'the boxer's fracture'. It is the most common fracture of the hand and is responsible for 9.7-50% of all cases (1-3). It is commonly observed in the dominant hand of male patients (4). The

E-mail: haiqingcaihqc@163.com

E-mail: wzgproscmc@163.com

*Contributed equally

majority of these fractures are nondisplaced or minimally displaced without rotational deformity, and may be treated by conservative treatment with a good functional outcome (5,6). However, operative intervention is recommended for severely displaced fracture or volar angulation with severe rotational deformity.

Previously, numerous surgical techniques have been proposed to treat unstable neck fractures of fifth metacarpal fracture to perform closed/open reduction and achieve stability. These techniques include retrograde pinning (RP) by means of intramedullary K-wires, transverse pinning (TP) using K-wires, external fixation and standard or low-profile plates (7-9). However, complications including marked angulation accompanied by potential reduction in the range of motion (ROM) and reduced grip strength (10,11) may occur. Previous studies have demonstrated that antegrade intramedullary nailing (AIMN) is a reliable and minimally invasive method to treat metacarpal fractures (12-14). Although AIMN was reported to be a safe and reliable option to treat 66 children and juveniles with metacarpal fractures (15), more cases and studies are required to provide clinical guidance for treating the disease in juvenile patients.

Titanium elastic nail (TEN) has received increasing attention in previous years and is considered minimally invasive with excellent functional and cosmetic outcomes in treating fractures (15,16). Therefore the present study investigated the efficacy of treating displaced fifth metacarpal neck fractures in 69 juvenile patients using antegrade intramedullary fixation with TEN. The results of the current study may provide useful information for guiding clinical treatment of neck fractures.

Materials and methods

Patients. The present prospective study was approved by the Ethics committee of Shanghai Children's Medical Center affiliated to Shanghai Jiao Tong University School of Medicine (Shanghai, China) and written informed consent was obtained from the parents of participants prior to the study. It included 69 juvenile patients who received surgery intervention due to fifth metacarpal neck fractures in the Department of Pediatric orthopaedics, Shanghai Children's Medical Center affiliated to Medicine School of Shanghai Jiaotong University (Shanghai, China) between February 2012 and

Correspondence to: Dr Haiqing Cai or Zhigang Wang, Department of Pediatric Orthopedics, Shanghai Children's Medical Center, Affiliated to Shanghai Jiao Tong University School of Medicine, 1678 Dongfang Road, Pudong New Area, Shanghai 200127, P.R. China

Key words: antegrade intramedullary fixation, fifth metacarpal neck fracture, titanium elastic nail, head-shaft angle

January 2014. The inclusion criteria were as follows: i) volar angulation of the fifth metacarpal head >30° (8) or ii) rotational deformity (>15°) (13). Participants were excluded if they had segmental comminuted fractures or intraarticular involvement. The mean age of the patients was 14.9 ± 1.5 years (range, 12-17 years). They included 60 males and 9 females, all of who experienced unilateral limb injury. The right hand was involved in 63 cases and the left hand in 6 cases. The head-shaft angle of fifth metacarpal fracture angle was measured in accordance with a previous study by Picture Archiving and Communication Systems (PACS) (17). Its normal value is ~15°, however, in the current study, the mean head-shaft angle of fifth metacarpal fracture was $48.8\pm3.0^\circ$.

Surgical technique. All surgery was performed under general anesthesia. Anesthesia was induced with midazolam (0.1 mg/kg; Fresenius Kabi AG, Ban Homburg, Germany), propofol (3 mg/kg; Fresenius Kabi AG), fentanyl (2 mg/kg; Fresenius Kabi AG) and rocuronium (0.6 mg/kg Esmeron; MSD, Shanghai, China) and maintained with sevoflurane (3%; Baxter, Deerfield, IL, USA). The surgery was performed by experienced surgeons under fluoroscopy. Following routine skin sterilization and draping, a medial incision (0.5-0.8 cm in length) was made over the dorsal ulnar side of the fifth metacarpal bone. The subcutaneous tissue was separated bluntly to expose bone cortex. A preflex TEN (2 mm in diameter, Synthes GMbH, Umkirch, Germany) was selected based on the pre-operative width of the medullary canal. Entry point of the nail was located on the dorsal ulnar side of the fifth metacarpal bone, which was not covered by nerves and tendons, under constant C-arm radiographic guidance (Siemens AG, Munich, Germany). The entry point was then drilled and the TEN was inserted into medullary canal under constant C-arm radiographic guidance (Fig. 1A) and the awl (Synthes GmbH, Umkirch, Germany) was used to perforate the cortex. The tip of TEN was introduced under direct view, went over the fracture line and was fixed on the edge of the metacarpal articular surface (Fig. 1B). The nail was then rotated through 180° to acquire good reduction of the fracture (Fig. 1C). Re-examination was conducted to confirm good fracture reduction and proper position of the TEN using C-arm fluoroscopy, followed by cutting off the TEN tail (Fig. 1D and E). The remaining nail tail was buried intradermally and the incision was closed. Closed reduction was not conducted in the present study.

Postoperative treatment. All patients received routine antibiotics (cefazolin, 0.5 mg/kg) postoperatively. The affected hand and forearm were immobilized in a short arm plaster cast for 2 weeks. X-ray examination of the hand was performed for review at two days and two weeks after surgery. The plaster was then removed, and the patients performed functional exercise of metacarpophalangeal (MCP) joints gradually as previously described (18). As not all hospitals had rehabilitation departments, patients were not specifically required to complete functional exercises in the hospital. A hand X-ray was taken to evaluate fracture healing 5 weeks postoperatively. If firm fracture union was not observed, the X-ray was taken again 8 weeks postoperatively. When firm fracture union was observed radiographically, the TEN at the medial insertion site Table I. General characteristics of the enrolled patients.

Variable	Mean (Range) 14.9±1.5 (12-17)		
Age, year			
Sex, male/female	60/9		
Operating time, min	15.5±6.5 (5-32)		
Number of fluoroscopy	7.8±2.4 (5-13)		
Healing time, weeks	5.7±1.09 (5-10)		
Follow-up time	16.9±6.3 (7-30)		
DASH	1.7 (0-3.0)		

Data are presented as mean \pm standard deviation. DASH, disabilities of the arm, shoulder and hand.

of the fracture was directly removed under general anesthesia in our hospital 3 months postoperatively.

Patients were asked to return for re-evaluation using X-ray examination during follow-up. The patients were clinically followed up from 7 to 37 months following surgery (average 16.9±6.3 months). From the clinical point of view, operating time, time of C-arm usage and healing time were recorded for each patient. Additionally, the head-shaft angle of the fifth metacarpal and the ROM of the MCP joint for the affected and unaffected sides were evaluated. Disabilities of the arm, shoulder and hand (DASH) scale (0-100 points) (19) was performed to assess upper arm function. The DASH questionnaire consists of 30 items and facilitates comprehensive evaluation of upper-extremity functional activity (20). Although it is not recommended for children, it is suitable for these participants in the present study, as they had mean age of 14.9±1.5 years. Each patient finished the questionnaire.

Statistical analysis. Statistical analysis was performed using SPSS 16.0 software (SPSS, Chicago, IL, USA). The measurement data were expressed as mean \pm standard deviation and analyzed using a paired *t* test. P<0.05 was considered to indicate a statistically significant difference.

Results

Clinical outcomes. The general characteristics of the enrolled patients are presented in Table I. The present study included 60 males and 9 females with an age range of 12-17 years (mean, 14.9±1.5 years). All patients experienced unilateral limb injury, including 63 cases with right hand injury and 6 cases with left hand injury. The operating time ranged from 5-32 min (mean time, 15.5±6.5 min). The frequency of the C-arm usage intraoperatively was 7.8±2.4 times per surgery (range, 5-13 times), and the duration time was 0.5 sec per time. Patient healing time ranged from 5-10 weeks (mean time, 5.7±1.09 weeks). The results of head-shaft angle and ROM of the MCP joint are presented in Table II. The mean ROM of the MCP joint of the affected side was 87.9±4.3° postoperatively, and no significant difference was observed compared with that of the unaffected side (89.1±4.6°, P=0.35). The mean DASH score was 1.7 (range, 0 to 6.0).

Side	ROM of MCP joint, °	Head-shaft angle of the fifth metacarpal		
		Pre-op, °	Postop, °	P-value
Affected	87.9±4.3	48.8±3.0	15.8±1.8	<0.001ª
Unaffected	89.1±4.6	-	15.0±1.6	-
P-value		0.35	0.06	

Table II. Head-shaft angle of the fifth metacarpal and the ROM of the MCP joints.

^aP<0.001. Data are presented as mean ± standard deviation. ROM, range of motion; MCP, metacarpophalangeal; pre-op, pre-operatively; postop, postoperatively.



Figure 1. Intra-operative images of one representative patient. (A) Enter point is localized, (B) the tip of TEN is introduced to the medullary canal and (C) the nail is rotated through 180° to acquire good reduction of the fracture. (D and E) The nail tails were removed.

Radiographic outcomes. The head-shaft angle of the affected side was $48.8\pm3.0^{\circ}$ pre-operatively and $15.8\pm1.8^{\circ}$ postoperatively, indicating a significant improvement (P<0.001) following surgery. However, there were no significant differences in the head-shaft angle between the affected and unaffected sides ($15.8\pm1.8^{\circ}$ vs. $15.0\pm1.6^{\circ}$, respectively; P=0.06). All patients obtained anatomical reduction and presented with complete fracture healing.

Subjective outcomes. Antegrade intramedullary fixation with TEN was successful in each patient with displaced fifth metacarpal neck fracture (Fig. 2). There were no cases of complications, including paralysis of dorsal branch of ulnar nerve, injury of dorsal extensor tendon, avascular necrosis of metacarpal head, nonunion, and abnormal healing. One case (16 year old male) experienced delayed union (>8 weeks, Fig. 3) but acquired good healing after 3 months, at which point the nail was removed. The nail was removed within 3 months of surgery in all patients as fracture union was confirmed radiographically. Excellent functional results were presented during the final follow-up at 37 months post-surgery. A total of four cases developed nail tail irritation but it was tolerated and did not require special management. The symptoms disappeared when the internal fixation was removed.

Discussion

In the present study, the clinical outcome of antegrade intramedullary fixation with TEN in displaced fifth metacarpal neck fractures was evaluated. The results determined that antegrade intramedullary fixation with TEN is a simple technique to perform. In addition, it is also a straightforward, reliable, minimally invasive surgical technique, and seems to achieve excellent functional results.

Initial treatment for the fifth metacarpal neck fracture is nonsurgical and involves certain types of immobilization. Different types of studies demonstrate that treatment of this injury consists of a closed reduction of the fracture and immobilization with cast application (21,22). This treatment is suitable for nondisplaced or minimally displaced fracture without rotational deformity (23). However, long-term cast immobilization limits the activities of the patients and may lead to a stiff joint and pressure necrosis of the skin, thus prolonging rehabilitation. In addition, the nondisplaced



Figure 2. Images of one patient with fifth metacarpal neck fracture of right hand. (A) Pre-operation image presenting dorso-ulnar swelling, (B) X-ray image demonstrating the fifth metacarpal neck fracture, (C) X-ray image at 6 weeks post-operation and (D) X-ray image at final follow-up. (E) Cosmetic and functional image. Post-operatively, ulnar scar is visible.



Figure 3. Images of the case experiencing delayed union. X-ray images (A) Pre-operation, (B) 8 weeks post-operation demonstrating a delayed union, (C) at 3 months post-operation and (D) at final follow-up. (E) Cosmetic and functional images.

fracture may shift or cause mal-union due to cast loss. Therefore, surgical treatment with internal fixation is recommended for severely displaced and unstable fractures to avoid impairment of hand function. Surgery is recommended for injuries with a dorsal angulation $\geq 30^{\circ}$, a shortening ≥ 5 mm and rotatory deformity >15° (8). Percutaneous or cross intramedullary Kirschner wire (K-wire) fixation is a common procedure used in open reduction for operative treatment (24,25). However, patients may experience complications including osteomyelitis, tendon rupture, nerve lesion, pin tract infection, pin loosening and migration (26-28), highlighting the limitations of this procedure.

TEN is widely accepted as the most reliable and valid method for treating paediatric long bone fractures (29-33). TEN is able to maintain biomechanical stability owing to the divergent 'C' configuration, which generates six points of fixation and permits the construct to act as an internal splint (34). Furthermore, TEN offers stable and elastic fixation and permits restricted motion at the fracture site, leading to healing by external callus (35). Additionally, it avoids any growth disturbance by conserving the epiphyseal growth plate and reducing bone impairment or weakening due to the construct elasticity, which offers a load sharing and biocompatible internal splint (29). TEN also allows rapid mobilization with few complications and avoids prolonged immobilization (36). Finally, it involves a minimal risk of bone infection. Antegrade intramedullary fixation with TEN was used in the present study to treat displaced fifth metacarpal neck fractures in 69 juvenile patients. This technique achieved good functional results.

In line with aforementioned previous studies (27,37), the operating time in the present study was relatively short. The ROM of the MCP joint recovered well postoperatively and the average DASH score was 1.7, demonstrating that the function of the affected side underwent good recovery (DASH scores are on a scale from 0-100, with higher scores indicating greater disability). A 2 mm preflex TEN was selected as the material of internal fixation to treat the fifth metacarpal neck fractures. It was not big enough to suppress endosteum of the affected side. Furthermore, no cases required open reduction postoperatively, indicating that the inserted TEN did not injure the periosteum. This suggests that TEN insertions do not impede healing of the fifth metacarpal neck fracture.

There were no cases of serious complications reported in the present study, such as paralysis of dorsal branch of ulnar nerve and injury of dorsal extensor tendon. To avoid these complications, it is worthwhile mentioning a number of important points. Firstly, correct selection of the incision site is of paramount importance for operative success; secondly, superficial sensory nerves should be protected when subcutaneous tissue is separated; thirdly, attention should be focused on managing the nail tails to avoid irritation of the surrounding tissue, thus truncation and short length of nails would be good choices. Finally, it should be stressed that antegrade intramedullary fixation with TEN is suitable only for fractures presenting with an intact metacarpal head. If a comminuted fracture of the metacarpal head is involved, intramedullary nail is insufficient to maintain post-reduction stability and may penetrate into the MCP joint, causing internal fixation failure. In the current study, only one case experienced delayed recovery (>8 weeks), which may be due to a variety of factors, such as the severity of injury or the nutritional status of patients.

The present study should be cautiously interpreted due to certain limitations. It included a small sample size and a short-term follow-up, therefore larger-scale studies are required to confirm the analysis results. Long-term follow-up is in process to evaluate the clinical outcome of antegrade intramedullary fixation using TEN.

Antegrade intramedullary fixation with TEN is a straightforward, reliable and minimally invasive surgical technique. The results may provide good clinical learning experiences for treating juveniles with fifth metacarpal neck fracture.

References

- 1. Feehan LM and Sheps SB: Incidence and demographics of hand fractures in British Columbia, Canada: A population-based study. J Hand Surg Am 31: 1068-1074, 2006.
- 2. Gudmundsen TE and Borgen L: Fractures of the fifth metacarpal. Acta Radiol 50: 296-300, 2009.
- 3. Keenan M: Managing boxer's fracture: A literature review. Emerg Nurse 21: 16, 18-24, 2013.
- Arafa M, Haines J, Noble J and Carden D: Immediate mobilization of fractures of the neck of the fifth metacarpal. Injury 17: 277-278, 1986.

- 5. Harris AR, Beckenbaugh RD, Nettrour JF and Rizzo M: Metacarpal neck fractures: Results of treatment with traction reduction and cast immobilization. Hand (N Y) 4: 161-164, 2009.
- Poolman RW, Goslings JC, Lee JB, Statius Muller M, Steller EP and Struijs PA: Conservative treatment for closed fifth (small finger) metacarpal neck fractures. Cochrane Database Syst Rev: CD003210, 2005.
- 7. Potenza V, Caterini R, De Maio F, Bisicchia S and Farsetti P: Fractures of the neck of the fifth metacarpal bone. Medium-term results in 28 cases treated by percutaneous transverse pinning. Injury 43: 242-245, 2012.
- Rhee SH, Lee SK, Lee SL, Kim J, Baek GH and Lee YH: Prospective multicenter trial of modified retrograde percutaneous intramedullary Kirschner wire fixation for displaced metacarpal neck and shaft fractures. Plast Reconstr Surg 129: 694-703, 2012.
- Calder JD, O'Leary S and Evans SC: Antegrade intramedullary fixation of displaced fifth metacarpal fractures. Injury 31: 47-50, 2000.
- Hofmeister EP, Kim J and Shin AY: Comparison of 2 methods of immobilization of fifth metacarpal neck fractures: A prospective randomized study. J Hand Surg Am 33: 1362-1368, 2008.
- 11. Midgley R and Toemen A: Evaluation of an evidence-based patient pathway for non-surgical and surgically managed metacarpal fractures. Hand Therapy 16: 19-25, 2011.
- 12. Corkum JP, Davison PG and Lalonde DH: Systematic review of the best evidence in intramedullary fixation for metacarpal fractures. Hand (N Y) 8: 253-260, 2013.
- 13. Strub B, Schindele S, Sonderegger J, Sproedt J, von Campe A and Gruenert JG: Intramedullary splinting or conservative treatment for displaced fractures of the little finger metacarpal neck? A prospective study. J Hand Surg Eur Vol 35: 725-729, 2010.
- 14. Sletten IN, Nordsletten L, Hjorthaug GA, Hellund JC, Holme I and Kvernmo HD: Assessment of volar angulation and shortening in 5th metacarpal neck fractures: An inter- and intra-observer validity and reliability study. J Hand Surg Eur Vol 38: 658-666, 2013.
- Lieber J, Härter B, Schmid E, Kirschner HJ and Schmittenbecher PP: Elastic stable intramedullary nailing (ESIN) of pediatric metacarpal fractures: Experiences with 66 cases. Eur J Pediatr Surg 22: 305-310, 2012.
- 16. Chen YF, Wei HF, Zhang C, Zeng BF, Zhang CQ, Xue JF, Xie XT and Lu Y: Retrospective comparison of titanium elastic nail (TEN) and reconstruction plate repair of displaced midshaft clavicular fractures. J Shoulder Elbow Surg 21: 495-501, 2012.
- Theeuwen GA, Lemmens JA and Van Niekerk JL: Conservative treatment of boxer's fracture: A retrospective analysis. Injury 22: 394-396, 1991.
- Stamm TA, Machold KP, Smolen JS, Fischer S, Redlich K, Graninger W, Ebner W and Erlacher L: Joint protection and home hand exercises improve hand function in patients with hand osteoarthritis: A randomized controlled trial. Arthritis Rheum 47: 44-49, 2002.
- Gummesson C, Atroshi I and Ekdahl C: The disabilities of the arm, shoulder and hand (DASH) outcome questionnaire: Longitudinal construct validity and measuring self-rated health change after surgery. BMC Musculoskelet Disord 4: 11, 2003.
- 20. Jester A, Harth A, Wind G, Germann G and Sauerbier M: Disabilities of the arm, shoulder and hand (DASH) questionnaire: Determining functional activity profiles in patients with upper extremity disorders. J Hand Surg Br 30: 23-28, 2005.
- 21. Lundeen J and Shin A: Clinical results of intraarticular fractures of the base of the fifth metacarpal treated by closed reduction and cast immobilization. J Hand Surg Br 25: 258-261, 2000.
- 22. Zwitser EW and Breederveld RS: Fractures of the fifth metatarsal; diagnosis and treatment. Injury 41: 555-562, 2010.
- Kollitz KM, Hammert WC, Vedder NB and Huang JI: Metacarpal fractures: Treatment and complications. Hand 9: 16-23, 2014.
- 24. Khan J, Shrestha SK, Pradhan NM, Acharya BK and Khanal P: Percutaneous K wire fixation of fifth metacarpal neck fracture-new and simple technique. J Nepal Health Res Counc 10: 61-65, 2012.
- Kelsch G and Ulrich C: Intramedullary k-wire fixation of metacarpal fractures. Arch Orthop Trauma Surg 124: 523-526, 2004.
- 26. Stahl S and Schwartz O: Complications of K-wire fixation of fractures and dislocations in the hand and wrist. Arch Orthop Trauma Surg 121: 527-530, 2001.

- 27. Lee SK, Kim KJ and Choy WS: Modified retrograde percutaneous intramedullary multiple Kirschner wire fixation for treatment of unstable displaced metacarpal neck and shaft fractures. Eur J Orthop Surg Traumatol 23: 535-543, 2013.
- Hsu LP, Schwartz EG, Kalainov DM, Chen F and Makowiec RL: Complications of K-wire fixation in procedures involving the hand and wrist. J Hand Surg Am 36: 610-616, 2011.
- 29. El-Adl G, Mostafa MF, Khalil MA and Enan A: Titanium elastic nail fixation for paediatric femoral and tibial fractures. Acta Orthop Belg 75: 512-520, 2009.
- Gwyn DT, Olney BW, Dart BR and Czuwala PJ: Rotational control of various pediatric femur fractures stabilized with titanium elastic intramedullary nails. J Pediatr Orthop 24: 172-177, 2004.
- Aksoy MC, Caglar O, Ayvaz M, Yazici M and Alpaslan AM: Treatment of complicated pediatric femoral fractures with titanium elastic nail. J Pediatr Orthop B 17: 7-10, 2008.
- 32. Pulate A, Jadhav A and Shah N: Study of the outcome of titanium elastic nail system in diaphyseal femoral fractures in children 1. J Maharashtra Orthop Assoc 7: 6-8, 2012.

- 33. Borah S, Kumar A and Saurabh D: Titanium elastic nail system (TENS) or external fixation in paediatric femoral shaft fractures. Bone Joint J Orthopaedic Proceedings Suppl 95: 34, 2013.
- Ligier JN, Metaizeau JP, Prévot J and Lascombes P: Elastic stable intramedullary pinning of long bone shaft fractures in children. Z Kinderchir 40: 209-212, 1985.
- 35. Sankar WN, Jones KJ, David Horn B and Wells L: Titanium elastic nails for pediatric tibial shaft fractures. J Child Orthop 1: 281-286, 2007.
- 36. Flynn JM, Hresko T, Reynolds RA, Blasier RD, Davidson R and Kasser J: Titanium elastic nails for pediatric femur fractures: A multicenter study of early results with analysis of complications. J Pediatr Orthop 21: 4-8, 2001.
- 37. Moon SJ, Yang JW, Roh SY, Lee DC and Kim JS: Comparison between intramedullary nailing and percutaneous K-wire fixation for fractures in the distal third of the metacarpal bone. Arch Plast Surg 41: 768-772, 2014.