

Viscum album (L.) in experimental animal tumors: A meta-analysis

LEONI VILLANO BONAMIN¹, ALOISIO CUNHA DE CARVALHO¹ and SILVIA WAISSE²

¹Research Center, University Paulista, São Paulo 04026-002;

²Department of History of Science, Pontifical Catholic University of São Paulo, São Paulo 01303-000, Brazil

Received September 6, 2016; Accepted January 26, 2017

DOI: 10.3892/etm.2017.4372

Abstract. Mistletoe (*Viscum album* L.) has been used as complementary anticancer treatment for ~100 years. Although the clinical efficacy of mistletoe in cancer and associated survival benefits remain contested, several studies point to its effectiveness and others have reported antitumor and immunomodulatory properties. In the present review, a search was conducted for original articles reporting the outcomes of treatments for experimental animal tumors with mistletoe. The inclusion criteria were: Publication in English, from 1996 onwards and in peer-reviewed journals included in the database PubMed. The parameters analyzed were: Provenance and time of publication, rationale, methods (animal species used, mistletoe preparation, treatment protocol, tumor lineage, blinding, randomization, controls and concomitant treatments), outcomes and investigated mechanisms of action. A total of 37 studies met the inclusion criteria. The quality of the studies was adequate in the terms of sample size and use of controls, and the only animal species employed were mice and rats. However, few studies reported having performed random allocation and none reported blinding. There was wide variation in the type and preparation of mistletoe used, route of administration, regimen, tumor type and the mechanism of action assessed. A temporal trend was identified; earlier studies sought to establish the antitumor effect of mistletoe and its possible mechanisms, cytotoxicity and immunomodulation in particular, whereas the later ones tended to focus more on biologically active principles, genomics and oxidative stress. A total of 32/37 studies reported an antitumor effect, 3 of which had mixed results. A total of 2 studies did not detect any antitumor effect and a further 2 found stimulation of tumor growth in the treated groups. One study did not assess antitumor effects, investigating immunomodulation action instead. The quality of the studies was satisfactory and the majority reported positive outcomes. Nevertheless, there is a great deal

of methodological heterogeneity among the studies, which precludes conclusive comparisons. Based on these results, the present authors strongly suggest developing guidelines for reporting *in vivo* mistletoe cancer treatment experiments.

Introduction

Mistletoe (*Viscum album* L.) has been used for the treatment of various illnesses for centuries (1); however, modern use of mistletoe in cancer treatment was introduced in the 1920s by Rudolf Steiner, the founder of anthroposophy (2). As Kröz *et al* (3) reported, in 2017 mistletoe will have been used for the treatment of cancer for almost a century, and the significance of this is corroborated by >1,200 citations in the PubMed database, which correspond to a multitude of studies demonstrating the immunomodulatory, cytotoxic and proapoptotic, anti-angiogenesis and DNA stabilizing properties of mistletoe. There have also been a number of studies and reviews concerning quality of life and patient perspectives (4,5), evidence-based benefits in human patients (6,7), ongoing clinical studies in humans (8,9) and adverse effects (10).

The mechanism of mistletoe's anticancer action has been of continued interest since its cytotoxic and immunomodulatory qualities were documented in 1990 (11). In the following decade, several studies were concerned with identifying the most effective therapeutic dosage, time and regimen of treatment (12-14). Recently, it has been suggested that mistletoe exerts a potent anti-inflammatory effect via selective inhibition of Cox protein expression, which may contribute to the antitumor action (15,16).

Over time, several biologically active components have been identified, most notably viscotoxins and lectins (ML-I, ML-II, ML-III) (17-21); eventually a recombinant lectin (viscumin, rML) was developed and tested (22-24). Research has also been conducted into lipophilic components, namely triterpenes, which also exhibit marked cytotoxic effects (25-27).

Mistletoe is a semi-parasitic plant that is able to grow on several host trees; the types most frequently used for therapeutic purposes are the ones grown on fir (*A-Abies*), apple (*M-Malus*), pine (*P-Pinus*), poplar (*Po-Populus*) and oak (*Qu-Quercus*) trees (2). Asian subspecies of *V. album* (Korean and Chinese mistletoe, *Viscum album* subsp. *coloratum* Kom.) (28) have also been studied and found to have properties similar to European mistletoe. The concentration of active components and biological activity vary as

Correspondence to: Professor Leoni Villano Bonamin, Research Center, University Paulista, 1212 Rua Doutor Bacelar, São Paulo 04026-002, Brazil
E-mail: leonibonamin@unip.br

Key words: *Viscum album*, experimental oncology, viscotoxins, lectins, review

a function of the host tree, harvesting time and extraction procedure (aqueous extraction with/without fermentation, other extraction procedures, pressing) (29), which also remain a notable target of research (30).

The clinical efficacy of and survival benefit associated with the use of mistletoe remains contested (31). As is known, the intrinsic complexity of cancer and the composition of mistletoe, pose a problem for clinical trials (2-4). *In vivo* tumor models may therefore be particularly elucidative, as features of the tumor microenvironment may be observed and compared with the diverse clinical outcomes. The authors of the present study were able to locate a single review on this topic; however, it was limited to breast and gynecological tumors (29). Another source of relevant information is an NIH/National Cancer Institute overview of the use of mistletoe in anticancer treatment, in which the results of studies involving animal models are discussed (32). As there are no recent reviews of mistletoe treatment in animal models, the current study presents a meta-analysis of the peer-reviewed, published literature on the effect of mistletoe in animal cancer models (*in vivo* models).

Materials and methods

Literature search. A literature search was performed to locate original research articles reporting the outcomes of mistletoe treatments for experimental tumors in animals. The search was limited to articles published from 1996 onwards, in English, in peer-reviewed journals included in the database PubMed (ncbi.nlm.nih.gov/pubmed/) and with relevant references cited. The aim of this was to facilitate access to the raw data for the interested readership. These criteria were adopted as a considerable number of studies are published in German and/or as books or meeting proceedings, and the peer-reviewed criterion ensures *a priori* the quality of articles. Several searches were performed using the following search terms: '(Viscum OR mistletoe) AND cancer'; '(Viscum OR mistletoe) AND tumor'; and 'Viscum album AND tumor AND animal'. Only articles reporting *in vivo* animal models were considered for inclusion.

Parameters of analysis. The parameters of analysis were as follows: i) Provenance and time of publication; ii) rationale underlying the study; iii) methods used [studied animal species, number analyzed, type of mistletoe/mistletoe-derived preparations used, treatment protocol, grafted tumor lineage (epithelial, mesenchymal, melanoma), blinding, random group allocation, controls (placebo, reference drug) and other combined therapies] iv) results (remission, improvement, worsening and survival) and v) mechanisms of action investigated (antiproliferative activity, cytotoxicity, immunomodulatory, tumor leukocyte infiltration, expression of specific phenotypic markers, and other tumor microenvironment aspects likely to correlate with the reported outcomes).

Data presentation. The data were entered in an *ad hoc* form and are presented in two tables. Table I lists the articles reporting *in vivo* studies with detailed descriptions of the substance used, dose and regimen, animal model tested, tumor cell lines, concomitant treatments and outcomes. Table II describes methodological aspects used to determine the quality of studies.

Results and discussion

Initial findings. The literature search located 911 articles. Duplicates were excluded and the remaining studies were analyzed based on their titles and abstracts until 37 remained that met the inclusion criteria and were considered for the present analysis. These 37 studies were written in English, published in peer-reviewed journals from 1996 onward and included in PubMed (mean, 1.85 studies/year; range, 0-7). As mentioned above, only 1 similar review was located for comparison to the present study (29), which assessed 34 studies from 1938 to 2008 (mean, 0.48 studies/year) of which 21 had been published in peer-reviewed journals. The apparently favorable trend detected in the present review should be considered cautiously, as Kienle *et al* (29) considered breast and gynecological tumors only. However, according to those same authors ~50 articles/year are published on mistletoe, implying that the proportion of studies using *in vivo* animal models is rather low. Tables I and II describe the results corresponding to the analyzed parameters.

Mistletoe origin. *Viscum album* (L.) is a parasitic plant whose composition may be influenced by the host plant or the harvest season (2). As a result of this, the commercially available preparations are typically made from mixtures of fresh leafy shoots harvested in both summer and winter, and fruits harvested in winter, when the concentration of the main active principles (lectins and viscotoxins) is highest (28). Only two studies included in the present review reported the harvest season, and this was reported as winter in both (27,33). A total of 32 studies used European mistletoe and 5 studies used Korean/Chinese mistletoe. The host species on which mistletoe was grown were reported as follows: 13 studies on apple, 4 on oak, 4 on pine, 4 on fir, 2 on poplar and 1 on plum (*Prunus*) trees. A total of 12 studies did not report the host species.

Distribution per date of publication. A total of 25 studies were published between 1996 and 2005 (mean, 2.5 studies/year) and 12 from 2006 to 2016 (mean, 1.2 studies/year). This is a ~50% decrease in the period 2006-2016 compared with the number of studies published in the previous decade. One hypothesis that could explain such a decrease in the number of studies is the trend towards reducing animal protocols in cancer research (34) due to their poor translational features, but this remains to be confirmed.

Country. The majority of studies were conducted in Germany (20 in their entirety, >37 including multicenter studies), followed by Croatia, India and Korea (4 in each), Austria, Belarus, Brazil, Canada, China, France, Norway, Romania, Serbia and the UK (1 in each). Compared with the review by Kienle *et al* (29), in which studies originated in Germany, Switzerland, Austria, USA, India, Croatia and Serbia only, the distribution reported here represents a favorable trend, as it seems to denote a more global interest in the use of mistletoe.

Animal species. Mice and rats were used in 32 and 5 studies, respectively. This finding is similar to the one reported by Kienle *et al* (mice, 28; rats, 6) (29); however, as shown herein,

Table I. Study design and outcomes (n=37) according to treatment, animal model, tumor cell line and main outcomes.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
Han <i>et al</i> , 2015	Endragit: Enteric-coated lectin from KM grown on poplar. Toxicity test: 500, 1,000, 2,000 mg/kg, single dose, p.o. Antitumor effect: Added to feed at 1, 2 or 4% concentration.	Toxicity test: ICR mice, both sexes; Antitumor activity: Male GDFl mice	Murine B16.F10 melanoma	POSITIVE Significant inhibition of tumor growth at 4% concentration. Increased survival rate (at 2 or 4%). Significant dose- and time-dependent reduction in tumor cell viability. Mechanisms: Cycle-cell deregulation, apoptosis.	(53)
Delebinski <i>et al</i> , 2015	VA-M; TT (OA 4.0 mg/ml + BA 0.38 mg/ml); ML-I (3.7 µg/ml); TT/ML-I (qualitatively positive); 6 groups: (1) dose escalation 20/40/60 mg/kg TT; (2) 0.3/0.4/0.5 µg/kg ML-I; (3) TT/ML-I combination; VAE given twice every 3rd day, i.v., from day 3 after tumor inoculation; (4) ARA-C 100 mg, i.v., every 3rd day, 4 doses; (5) TT/ML-I + ARA-C; and (6) control, received equal amounts of cyclodextrins.	Female NOD/SCID/IL2rg mice in pathogen-free conditions	Human HL-60 cells	POSITIVE TT/ML-I combination significantly reduced tumor weight, comparable to ARA-C. The greatest effect was observed with TT/ML-I + ARA-C combination.	(25)
Facina <i>et al</i> , 2014	VA-Qu AE (Iscaador Qu; 5 mg with 0.375 ng/ml of lectins), 0.1 ml, s.c. peritumoral, daily for 10 days	Female C57BL/6N mice	B16F10 melanoma	MIXED No effect on tumor volume. Upregulation of apoptosis-related gene expression, particularly the extrinsic apoptotic pathway. Immunomodulatory Th-1-like action, with IL recruitment and cell death activation through caspase 7, associated upstream with <i>Card-10</i> and downstream with CAD.	(44)
Stan <i>et al</i> , 2013	VA-M, ethanolic extract, 50 mg/kg, i.p., days 1, 3, 6. DOXO 2.5 mg/kg, i.p., days 1 and 6; 7 groups (VA-M yes/no, DOXO, yes/no, EAC yes/no)	Swiss female mice	EAC	POSITIVE VA-M enhanced the antitumor effect of DOXO reflected in tumor development (body weight, tumor volume and cell concentration) and oxidative stress-related enzymes.	(58)
Strüh <i>et al</i> , 2013	VA-M AE; STT; CD as control (2.2 g/kg). Treatments and controls injected s.c. peritumorally, every 2nd day starting at day 3 after tumor	Male C57BL/6Nrl mice	Murine B6F10 melanoma cells	POSITIVE Both AE and STT significantly slowed down tumor growth and increased median survival. The effect of STT was greater, but highly toxic. Both AE and STT	(26)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
	injection up to 10 cycles. Experiment 1: AE 12 $\mu\text{g/kg}$ of ML-I; STT 12 $\mu\text{g/kg}$ of ML-I and 93 mg/kg OA. Experiment 2: AE 3.5 $\mu\text{g/kg}$ of ML-I; STT 3.5 $\mu\text{g/kg}$ of ML-I and 71 mg/kg OA; TT 71 mg/kg OA.			significantly increased tumor necrosis and reduced tumor angiogenesis. These effects remained with the lower dose, but were reduced; in turn, they were not associated with toxicity. TT alone had no significant effect on tumor growth. STT and TT alone induced greater tumor necrosis compared with AE. Active caspase-3 expression was weak in all the groups. No treatment had any effect on proliferation of viable melanoma cells (Ki-67 positive).	
Podlech <i>et al.</i> , 2012	Iscaador Qu; (1) 100 $\mu\text{g/ML}$, culture medium then 2 μg , intratumor, (2) 1 μg , s.c., periodically (NR), (3) 20 μl (100 $\mu\text{g/ML}$), intratumor, single dose.	Athymic CD1-deficient NMRI nude mice; VMDk mice	LNT-229; SMA560 glioma cells	MIXED Pretreatment reduced tumor growth, however this was not due to altered proliferation or cell death. Systemic treatment was associated with slight, non-significant inhibition of tumor growth, whereas intratumor administration had significant effect.	(52)
Li <i>et al.</i> , 2011	CM-1 of Chinese mistletoe grown on poplar. Treated groups: 10 $\mu\text{g.kg}^{-1}$ CM-1 every 3rd day or 20 $\mu\text{g/kg}^{-1}$ every 10 days, i.v.	Female athymic nude (Balb/c nu nu) mice	CRC CLY and HT-29 cells	POSITIVE CM-1-treated groups exhibited a dose-dependent reduction in tumor growth and high tumor growth inhibition rate. Downregulation of specific miRNA induced by CM-1 was not due to suppression of host gene transcription, but to direct degradation of their precursors. There was direct correlation between miR-135a&b levels and cellular sensitivity to CM-1.	(69)
Thies <i>et al.</i> , 2008	ML-I from leaves of mistletoe grown on poplar (40%), apple (30%) and red oak (30%) trees collected in the winter. ML stock solution with 963 $\mu\text{g/ML}$ ML-I, ~1% ML-II and <0.5% ML-II. Treatment groups: 30, 150 and 500 ng/kg in 200 μl PBS, i.p., 19 days.	Pathogen-free male Balb/c SCID/SCID mice	Human melanoma MV3	MIXED Primary tumor: Significant reduction of tumor growth with low dose, significant increase of tumor weight with intermediate vs. low dose. The apoptosis rate was significantly higher in all 3 treatment groups, significantly higher with low vs. intermediate and high doses. There was an increased number of infiltrating DC in all treatment groups, significant	(33)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
Seifert <i>et al.</i> , 2008	VA-A (Helixor), 160 ng/ml ML-II/III and 5-10% ML-I; VA-P, 725 ng/ml ML-II/III; treated groups: 1, 50, 100 mg/kg, i.p., 4 d/wk., 3 wks., 1 day after tumor inoculation. Controls: CPA 100 mg/kg on day 3.	Male CB-17 SCID/SCID mice; specific-pathogen free	Human ALL NALM-6 cells	increase of DC apoptosis with intermediate and high doses vs. low dose; significant increase in the number of infiltrating DCs with low dose vs. controls. Lung metastases: Significant reduction in number with low dose vs. control. Mean number increased with intermediate and high dose vs. low dose. Vascular count increase with high dose vs. controls. POSITIVE Both VA extracts exhibited concentration-dependent inhibition of cell proliferation, due to apoptosis; even low concentrations of extracts induced dose-dependent induction of early apoptosis markers. All extracts at all doses increased survival; there were 3 long-term survivors, all treated with VA-A (2 from 50 mg and 1 from 100 mg groups). None of these animals exhibited leukemia or lymphoma at necropsy. POSITIVE Volatile extract includes TT. Reduction of tumor cell number and viability; reduction of antioxidative enzyme activity in tumor bearing animals. POSITIVE Both extracts induced a significant reduction in tumor weight; reduction of tumor cell proliferation rate; increased cell necrosis and apoptosis rates.	(59)
Cebovic <i>et al.</i> , 2008	CO ₂ extract of leaves of mistletoe grown on plum collected in winter; 2.0 ml/kg/d, i.p., duration NR, several doses tested but NR	Male and female Hann: NMRI mice	EAC	POSITIVE Volatile extract includes TT. Reduction of tumor cell number and viability; reduction of antioxidative enzyme activity in tumor bearing animals.	(27)
Beuth <i>et al.</i> , 2006	VA-A (Helixor) 3.6 ng/ml ML-I, 354 ng/ml ML-III; VA-M, 230 ng/ml ML-I, 500 ng/ml ML-III. Treated groups: 5 mg/μl/injection, i.p., of VA-A or VA-M on days 5, 6, 7, 12, 13 and 14	Male Balb/c mice	BT474 human ductal breast carcinoma	POSITIVE Both extracts induced a significant reduction in tumor weight; reduction of tumor cell proliferation rate; increased cell necrosis and apoptosis rates.	(68)
Duong Van Huyen <i>et al.</i> , 2006	Iscaador Qu FrF, 20 μg/dose, once/day, i.p., 7 days starting from the day of tumor inoculation	C57BL/6 and B16.129-II2b (IL-12 deficient) mice	B16F1 melanoma	POSITIVE The chosen dose was 20 μg, which was an effective dose that did not promote lethality. Reduced tumor weight and restoration of proliferative capacity of splenocytes in tumor-bearing animals, associated with increased secretion of IL-12.	(67)
Pryme <i>et al.</i> , 2004	Pure ML-I 10 mg/day as dietary supplement in different concentrations	Female NMRI mice	Krebs II NHL cells adapted to grow	POSITIVE Reduction in the number of mitotic figures;	(70)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
	(2.5, 5.0, 10.0 mg ML-I/d)		as intraperitoneal ascites tumor	intra-tumor lymphoid infiltrate doubled, with a corresponding increase in tumor necrosis; accelerated cell turnover inside tumor, and increased number of apoptotic cells. 11% of treated animals no longer had histological evidence of viable tumors on day 11.	
Yoon <i>et al.</i> , 2003	Aqueous extract (KM-110) and lectin (KML-C) of KM. KM-110 100 µg/animal; KML-C 0.05, 0.02, 0.005 µg/animal, i.v.; lung metastasis: 1 and 2 days before tumor inoculation; liver and spleen metastasis 1 day after tumor inoculation	Experimental lung metastasis: Balb/c; C57BL/6 mice Experimental liver and spleen metastasis: f CDF1 mice Spontaneous lung metastasis: C57BL/6 mice	Experimental lung metastasis: colon26-M3.1 carcinoma; B16.BL6 melanoma, 2.5x10 ⁴ Experimental liver and spleen metastasis: L51784-ML25 lymphoma cells, 4x10 ⁴ Spontaneous lung metastasis: B16.BL6 melanoma, 5x10 ⁵	POSITIVE KML-C prophylactically and therapeutically inhibited tumor metastasis of both lung metastatic tumor lines and reduced tumor metastasis of hematogenous tumor cells. KML-C 5.0 ng/animal, 5 days after inoculation, 5 times at 3-day intervals significantly inhibited spontaneous lung metastasis. KML-C 550 ng/mouse increased NK cell activity. KML-c at various doses given 2 days before inoculation increased macrophage-mediated cytotoxicity anti B16.BL6.	(66)
Braun <i>et al.</i> , 2002	sME Iscador Spezial, 2, 20, 100 µg/animal, s.c., 3 times per wk., 24 h after tumor inoculation	Inbred male Balb/c mice	RAW-117 P (liver); L-1 (lung) lymphosarcoma	POSITIVE Dose-dependent reduction in liver and lung tumor colonies and inhibition of local tumor growth (weight and volume) in all treated groups. Thymocytes upregulation; increased number of leukocytes, lymphocytes and monocytes in peripheral blood.	(49)
Elsässer-Beile <i>et al.</i> , 2001	rML 300 or 1,500 ng/ml in solution (EDTA, NaCl, PBS, PVP); 0.1 ml per intravesical catheter. Control group: NMU on wks. 0,2,4,6 Treated groups: 30 or 150 ng on weeks 8 to 13 or 14 to 19	Inbred female Fischer 344 rats	NMU-induced urinary bladder carcinoma	POSITIVE All 4 treated groups exhibited significant reduction of rates of atypical hyperplasia/neoplastic transformation compared to controls. No change in any of the assessed cytokines.	(22)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
Braun <i>et al</i> , 2001	Helixor ME-A 160 ng/ml, Helixor ME-P® 725 ng/ml of ML-II/III (both >95%), 5 and 50 µg/mouse, s.c., i.p., 3 times/wk, 24 h after tumor inoculation	Inbred male Balb/c mice	RAW-117 P (liver); L-1 (lung) lymphosarcoma	POSITIVE Dose-dependent reduction of liver and lung colonies; i.p. significantly greater effect. Dose-dependent increase of leukocyte, lymphocyte and monocyte count.	(48)
Schaffrath <i>et al</i> , 2001	rML, 0.3, 3, 30, 150 ng/kg, s.c., every 2nd day, starting 2 days after tumor inoculation	Inbred male Balb/c mice	RAW-117 P (liver); L-1 (lung) lymphosarcoma	POSITIVE Increased survival rate; reduced number of liver and lung tumor colonies. Increased number of leukocytes in RAW-117 P	(23)
Burger <i>et al</i> , 2001	Lektinol, 405 µg/ml of ML, from poplar, 0.3, 3, 30, 300 ng/kg/d, i.p. or s.c., once per day, 5 days/wk. for up to 4 wk.; positive controls: DOXO, 8 mg/kg, i.v., on days 0 and 14; 5-FU, 100 mg/kg, i.p., on days 0, 7, 14	C57BL/6, Balb/c mice, both genders	B16.F10 melanoma, C8 38 colon carcinoma, F9 testicular carcinoma, Lewis lung carcinoma or Rencal renal cell carcinoma.	MIXED Dose-dependent tumor growth inhibition (Rencal, colon, testicle) borderline effect; greatest inhibitory effect in Rencal model with dose 300 ng, however, the sensitivity of the 3 tumor types to VA was similar. In Rencal model VA as effective but less toxic than DOXO; colon carcinoma responded better to 5-FU (comparable toxicity). Lewis carcinoma and melanoma did not respond to VA.	(45)
Timoshenko <i>et al</i> , 2001	ML-I 1 ng/kg dissolved in PBS and 50 µg/ml mouse albumin to concentration 0.22 ng/ml, 100 µl injection, s.c., days 7, 10, 13, 16, 19 after tumor inoculation; IL-2, i.p., 10 ⁵ Cetus Units/ml, 100 µl injection, every 8 h, 5 days. Experiment 1: 2 rounds with 4-day interval. Experiment 2: 1 round. 5 groups: (1) healthy mice, (2) tumor inoculated (3) IL-2 alone (4) ML-I alone and (5) IL-2 + ML-I	Female C3He/HeJ mice	C3L5 murine mammary adenocarcinoma, 1 x 10 ⁶	NEGATIVE ML-I alone increased primary tumor growth and development and growth of metastases. In combination, ML-I did not modify the tumor growth suppressive action of IL-2. ML-I did not cause capillary leakage neither inhibited the one caused by IL-2 and associated nitric oxide changes.	(47)
Park <i>et al</i> , 2001	KML, 10, 30, 50 ng/animal; (1) Tumor growth: 10-50 ng/mouse, i.p., 3 times per wk. (2) Antimetastatic effect: 10-50 ng/mouse 2 days before and 1 day after tumor inoculation	Female C57BL/6 mice	Melanoma B16.BL6	MIXED No effect on tumor growth; increased survival. Dose-dependent inhibition of tumor cell proliferation via apoptosis, dose-dependent anti-angiogenesis effect. No effect on cell cycle.	(46)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
Zarkovic <i>et al</i> , 2001	Isorel, 50 μ l, every 2nd day, 3 consecutive wks., s.c. or s.c. contralateral to tumor (local vs. systemic)	Female CBA/HZgr mice	Murine mammary carcinoma	POSITIVE Inhibition of tumor growth with local administration; increased apoptosis and necrosis in both treated groups, reduction of mitosis with local treatment only, reduction of tumor invasiveness in both treated groups. Tumor infiltration without mononuclear cells in controls.	(65)
Mengs <i>et al</i> , 2000	AE, 30 or 300 ng/ml ML, i.v., 3 days/wk., 4 consecutive wks., total volume 0.1 ml	Female C57BL/6J mice.	Urinary bladder carcinoma MB49	POSITIVE Increased survival; disappearance of tumor-both effects statistically significant, non-dose-dependent.	(54)
Schumacher <i>et al</i> , 2000	rML, 20, 150, 500 ng/kg, Monday to Friday until day 83 or 120% of initial body weight	SCID/SCID mice	SoTü 3 human ovarian cystoadenoma	POSITIVE Significantly longer survival with highest dose; symptom-free interval increased among animals that developed malignant ascites.	(24)
Kunze <i>et al</i> , 2000	ML-I, 1 ng/kg, twice/week, s.c., 6 or 15 months	Pathogen-free female Wistar rats	BBN-induced bladder carcinoma	NEGATIVE No difference in tumor growth or spectrum of histological types of carcinomas. In the groups treated for 15 months, tumor size was 2x greater. No significant stimulation of any cell population in lamina propria.	(40)
Antony <i>et al</i> , 1999	Iscaador M: (1) For spleen cell activation, 1.66 mg/dose/animal, i.p., 5 days. (2) For treatment: 1.66 mg/dose, i.p., daily, 10 days	C57BL/6 mice	B16.F10 melanoma	POSITIVE Significant inhibition of lung metastasis (<i>in vivo</i> activated spleen cells), increased with concomitant treatment; similar effect in metastasis-bearing mice. <i>In vitro</i> activation of spleen cells. Increased survival of all treated groups (best effect: <i>In vivo</i> activation + treatment). Markers: Reduction of hydroxyproline, sialic acid and GGTP (cell proliferation).	(64)
Lenartz <i>et al</i> , 1998	ML-1. (1) Systemic: 1, 10 ng/kg, s.c., 2 wk. (2) Local: 10, 100 ng, single dose, i.p.	Adult female Fischer 344 rats	F98 glioma	POSITIVE Reduction in tumor volume in all treated groups, only significant with lowest dose, both routes. Dose-dependent increased tumor necrosis with local administration only.	(50)
Yoon <i>et al</i> , 1998	KM AE grown on oak. Survival: 100 μ g single dose i.v. Metastasis inhibition: 100 μ g i.v., 2 and 4 days	Specific-pathogen free female C57BL/6, Balb/c, CDF ₁ mice	B16.BL6 melanoma; 26-M3.1 colon carcinoma; spleen	POSITIVE Administration 2 days before tumor inoculation significantly inhibited melanoma and colon	(51)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
Weber <i>et al</i> , 1998	before tumor inoculation. Dose/route test: 1-100 μ g/dose, i.v., s.c., p.o., intranasal		and liver metastases of L5178Y-ML25 lymphoma	carcinoma metastasis, 4 days before melanoma metastasis only. Similar pattern in lymphoma; systemic routes had dramatic effect even in low dose. Enhanced NK cell activity. Increased survival.	(55)
	Lektinol, 30 ng/ml of bioactive MLs; 4 groups: (1) Vehicle, (2-4) 3, 30, 150 ng/kg ML, respectively, i.v., 1 hour after tumor injection then 4 days/wk., 3 wk.	Male Balb/c mice	B16.F10 melanoma	POSITIVE Dose-dependent inhibition of metastasis; reduction of melanoma cell count in lungs. <i>In vitro</i> : stimulation of macrophages, CD4 ⁺ CD8 lymphocytes; significant enhancement of anti-CD8, no effect on anti-CD4.	
Zarkovic <i>et al</i> , 1998	Isorel M; pretreatment of tumor cells with whole extract, ML-1 and 2 fractions-high and low molecular weight, 0.1 μ g/ml	Male C57BL/6J mice	Melanoma	POSITIVE All tested preparations inhibited metastasis formation.	(21)
Kunze <i>et al</i> , 1998	Lectin, 1 ng/kg, s.c., twice per wk., 15 months	Female Wistar rats; specific pathogen-free	MNU-induced bladder carcinoma	NO EFFECT The incidence of bladder carcinoma did not differ between treated and control groups. There was no difference in the population of inflammatory cells infiltrating the tumor.	(41)
Zarkovic <i>et al</i> , 1997	Isorel M, 100 mg/kg, single dose, i.p.	Male C57BL/6J mice	B16F10 melanoma	POSITIVE Tumor growth reduced in treated animal, with a reduction of tumor viable cells. Tumor necrosis and abundant inflammatory response. <i>In vitro</i> tests: Strong stimulation of lymphocytes, with increased cytotoxic activity; increased sensitivity of tumor cells to cytotoxic activity of lymphocytes; inhibition of growth of tumor cells more than normal cells.	(63)
Jurin <i>et al</i> , 1997	Isorel M; strength 60, 2 protocols: (1) Single dose, 1 day before, simultaneous, 1 day after sheep RBC injection; 140 or 1,400 mg/kg/animal. (2) Repeated doses: a) normal mice and tumor-bearing	Male and female CBA/Hzgr mice	Murine methylcholantrene-induced fibrosarcoma	IMMUNOMODULATING EFFECT 1) Isorel single dose before sheep RBC injection restored the immune response; 140 mg/kg had greatest effect. 2) Dose-dependent immunostimulating effect; prolonged application (5 wk.) had dose-dependent	(62)

Table I. Continued.

Author/year	Treatments	Animal model	Tumor cell lines	Outcomes	(Refs.)
	mice: 14 or 140 mg/kg, daily, 14 consecutive days; b) normal mice: 14, 140 or 140 mg/kg, daily, 5 wk. (25 doses); route NR			immunosuppressing effect on normal mice only.	
Antony <i>et al</i> , 1997	Iscaador M, prophylactic, 5 doses, 1.66 mg/kg/dose, 4 groups: Control, before tumor inoculation, with tumor then 10 consecutive days, 5 days after tumor, 10 consecutive days	C57BL/6 mice	B16.F10 melanoma	POSITIVE Reduction of lung metastasis formation; increased survival rate of metastasis-bearing animals; lifespan increased in all treated groups. Markers: reduction of hydroxyproline (fibrosis), sialic acid (melanoma development).	(61)
Kunze <i>et al</i> , 1997	Purified lectin, 50 µg/ml, 1 ng/kg, s.c., twice per wk., 7 h after tumor induction and then for 6 months	Pathogen-free adult, female Wistar rats	NMU-induced urinary bladder carcinoma	NO EFFECT No significant differences in tumor rates, characteristics or histology.	(42)
Kuttan <i>et al</i> , 1997	5% Iscaador M, 1.66 mg/dose, route NR; 2 experiments: Carcinogenesis, twice/wk., 15 wks. Metastasis inhibition: 5 days before, simultaneous with and after tumor inoculation, 10 days in each group	Female Swiss albino and C57BL/6 mice	Methylcholantrene-induced sarcoma; B16.F10 melanoma for metastasis induction	POSITIVE Reduction of tumor size and development; reduction of metastasis formation; increased survival. Markers: Reduction of hydroxyproline and sialic acid.	(60)
Kuttan <i>et al</i> , 1996	Iscaador; (1) 1 mg/dose, i.p., twice/wk., 15 wks. (2) dilutions 0 to 1:3,000 (1.66 mg ⁻¹ .66 µg), i.p., twice/wk., 15 wks.	Female Swiss albino mice	Methylcholantrene-induced sarcoma	POSITIVE Iscaador fully inhibited sarcoma development; 100% of animals survived; effect was dose-dependent.	(56)

KM, Korean mistletoe; p.o., oral route; VA-M, mistletoe grown on apple tree; TT, triterpenes; OA, oleanolic acid; BA, betulinic acid; ML-I, lectin-I; VAE, *Viscum album* extract; wk(s), week(s); i.v., intravenous; ARA-C, cytarabine; NOD, non-obese diabetic; SCID, severe combined immunodeficiency; VA-Qu, mistletoe grown on oak tree; AE, aqueous extract; s.c., subcutaneous; Th, T-helper; IL, interleukin; CAD, caspase-activated DNase; i.p., intraperitoneal; DOXO, doxorubicin; EAC, Ehrlich ascites carcinoma; STT, soluble TT; CD, cyclodextrins; NR, not-reported; CM-1, Chinese mistletoe lectin >90% identical with European mistletoe ML-I; CRC, colorectal cancer; miR, microRNA; DC, dendritic cells; VA-A, mistletoe grown on fir; VA-P, mistletoe grown on pine; CPA, cyclophosphamide; ALL, acute lymphoblastic leukemia; NHL, non-Hodgkin lymphoma; FrF, extract prepared without fermentation; NK, natural killer cell; rML, recombinant lectin; 5-FU, 5-fluorouracil; BBN, *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine; GGTP, gamma-glutamyl transpeptidase; PVP, polyvinylpyrrolidone; NMU; *N*-methyl-*N*-nitrosourea; RBC; red blood cells.

Table II. Methodological characteristics of studies (n=37), considering animal species, tumor cell origin, quality criteria (blindness, randomization, sample size) and incidence of toxic effects.

Author/year	Animal species	Tumor type	Blinded study	Control	Random allocation	Number of animals/group	Toxicity	(Refs.)
Han <i>et al</i> , 2015	Mice	Melanoma	NR	Negative: 0.85% saline Positive: VAE + PKA	Yes	Toxicity test: 5; Antitumor effect: 9	Toxicity test: No deaths or clinical morbidity, no significant effect on body weight. No significant change in food intake.	(53)
Delebinski <i>et al</i> , 2015	Mice	Mesenchymal (Leukemia)	NR	Negative: CD; Positive: ARA-C	NR	10	Very good tolerability; no evidence of toxicity measured by body weight and autopsy with liver, spleen histology; some spleen atrophy in group ARA-C.	(25)
Facina <i>et al</i> , 2014	Mice	Melanoma	NR	Negative: Distilled water	NR	6	NR	(44)
Stan <i>et al</i> , 2013	Mice	Epithelial	NR	Negative: Vehicle Tumor control (EAC) Positive control: DOXO	NR	8	NR	(58)
Strüh <i>et al</i> , 2013	Mice	Melanoma	NR	Negative: CD vehicle	NR	8-9 (additional 4 animals/group for histology)	Higher dose: STT caused skin necrosis, apathy and weight loss, with 3 dropouts. All treatments caused local skin inflammation and inflammatory peritumoral infiltrates. Lower dose: No toxic effects; reduced inflammatory infiltrates.	(26)
Podlech <i>et al</i> , 2012	Mice	Mesenchymal (Glioma and astrocytoma)	NR	No treatment; vehicle	NR	6	NR	(52)
Li <i>et al</i> , 2011	Mice	Epithelial	NR	Negative control: PBS	NR	10	NR	(69)
Thies <i>et al</i> , 2008	Mice	Melanoma	NR	Negative control: PBS	NR	20	Vitality score, behavior, physiological response and water and food intake assessed: No animals exhibited changes; no treatment-related lesions were found on biopsy.	(33)
Seifert <i>et al</i> , 2008	Mice	Mesenchymal (Leukemia)	NR	Negative control: PBS Positive control: CPA	NR	8 (treated groups) 10 (control groups)	Tolerability was very good; no changes in body weight or assessed hematological parameters. VA-P was more toxic than VA-A <i>in vitro</i> , especially at the higher concentrations (50 mg/kg).	(59)
Cebovic <i>et al</i> , 2008	Mice	Epithelial	NR	Placebo; tumor control	NR	6	NR	(27)

Table II. Continued.

Author/year	Animal species	Tumor type	Blinded study	Control	Random allocation	Number of animals/group	Toxicity	(Refs.)
Beuth <i>et al</i> , 2006	Mice	Epithelial	NR	Negative control: PBS	NR	8	All animals free from side effects (behavior, appearance).	(68)
Duong Van Huyen <i>et al</i> , 2006	Mice	Melanoma	NR	Untreated; vehicle	NR	6	NR	(67)
Pryme <i>et al</i> , 2004	Mice	Mesenchymal (lymphoma)	NR	Standard diet	NR	5	NR	(70)
Yoon <i>et al</i> , 2003	Mice	Epithelial and mesenchymal	NR	Negative control: PBS	NR	5	NR	(66)
Braun <i>et al</i> , 2002	Mice	Mesenchymal (Lymphoma and sarcoma)	NR	PBS	NR	10	NR	(49)
Elsässer-Beile <i>et al</i> , 2001	Rats	Epithelial	NR	Tumor control	NR	Control: 56 Treated: a) 14; b) 23; c) 22; d) 19; + 15 in group b) for cytokine mRNA assessment	NMU-induced carcinoma was accompanied by high mortality associated with the tumor or procedure (15/71 in the control group and 40/123 in the treated groups). All treated animals exhibited weight loss over the 6 weeks of treatment.	(22)
Braun <i>et al</i> , 2001	Mice	Mesenchymal (Lymphoma and sarcoma)	NR	PBS	NR	8	NR	(48)
Schaffrath <i>et al</i> , 2001	Mice	Mesenchymal (Lymphoma and sarcoma)	NR	PBS	NR	10	No toxic effects (agility, food intake, body weight).	(23)
Burger <i>et al</i> , 2001	Mice	Epithelial and melanoma	NR	Negative control: PBS; positive control: DOXO, 5-FU	NR	5-7	VA less toxic than DOXO (body weight; mortality); not different from 5-FU.	(45)
Timoshenko <i>et al</i> , 2001	Mice	Epithelial	NR	Healthy mice; tumor control	NR	10 (2 rounds), 8 (1 round)	Nephrotoxicity.	(47)
Park <i>et al</i> , 2001	Mice	Melanoma	NR	PBS; no treatment	NR	NR	NR	(46)
Zarcovic <i>et al</i> , 2001	Mice	Epithelial	NR	Saline	NR	7	NR	(65)
Mengs <i>et al</i> , 2000	Mice	Epithelial	NR	Cehicle	Yes	13	NR	(54)

Table II. Continued.

Author/year	Animal species	Tumor type	Blinded study	Control	Random allocation	Number of animals/group	Toxicity	(Refs.)
Schumacher <i>et al</i> , 2000	Mice	Epithelial	NR	PBS + Tween-80	NR	20	NR	(24)
Kunze <i>et al</i> , 2000	Rats	Epithelial	NR	No treatment	Yes	60/70 in tumor-bearing groups; 20 in untreated and non-tumor bearing groups	NR	(40)
Antony <i>et al</i> , 1999	Mice	Melanoma	NR	No treatment	NR	14	NR	(64)
Lenartz <i>et al</i> , 1998	Rats	Mesenchymal (glioma)	NR	PBS	NR	5	NR	(50)
Yoon <i>et al</i> , 1998	Mice	Epithelial, melanoma, mesenchymal	NR	PBS	NR	5	NR	(51)
Weber <i>et al</i> , 1998	Mice	Melanoma	NR	PBS + povidone	Yes	5	No toxic effects (mortality; clinical signs; food intake; body weight; no significant findings on necropsy).	(55)
Zarcovic <i>et al</i> , 1998	Mice	Melanoma	NR	Culture medium	NR	5	NR	(21)
Kunze <i>et al</i> , 1998	Rats	Epithelial	NR	No treatment	NR	75	NR	(41)
Zarkovic <i>et al</i> , 1997	Mice	Melanoma	NR	Saline	NR	14 (total number, allocation NR)	NR	(63)
Jurin <i>et al</i> , 1997	Mice	Mesenchymal	NR	Healthy mice; saline	NR	8-12	NR	(62)
Antony <i>et al</i> , 1997	Mice	Melanoma	NR	No treatment	NR	8	NR	(61)
Kunze <i>et al</i> , 1997	Rats	Epithelial	NR	No treatment	NR	Controls: 57; Treated: 61	43/100 controls and 59/120 treated mice died by causes associated with the tumor-induction procedure.	(42)

Table II. Continued.

Author/year	Animal species	Tumor type	Blinded study	Control	Random allocation	Number of animals/group	Toxicity	(Refs.)
Kuttan <i>et al.</i> , 1997	Mice	Mesenchymal, melanoma	NR	No treatment	NR	15 (carcinogenesis) 8-10 metastasis induction	NR	(60)
Kuttan <i>et al.</i> , 1996	Mice	Mesenchymal	NR	Saline	Yes	15	No toxic effects (body weight, food intake); all animals survived and were healthy at the end of the experiment.	(56)

NR, not reported; VAE, *Viscum album* extract; PKA, polysaccharide K; CD, cyclodextrins; ARA-C, cytarabine; EAC, Ehrlich ascites carcinoma; DOXO, doxorubicin; STT, solubilized triterpenes; CPA, cyclophosphamide; VA-P, mistletoe grown on pine; VA-A, mistletoe grown on fir; NMU; *N*-methyl-*N*-nitrosourea 5-FU, 5-fluorouracil.

in recent decades mice have become the model used in almost all studies of antineoplastic drugs. This fact raises critical questions as to the translational aspects of such studies and the real possibility of extrapolating experimental findings to clinical practice with humans and other animal species (34,35). Such issues notwithstanding, veterinary clinical studies have demonstrated that mistletoe extracts elicit effective responses in various other animal species (36-39).

Number of animals per group. The number of animals included in each study ranged from 5 to 20 (median, 8), excluding 4 studies that induced autochthonous urinary bladder carcinoma (22,40-42), which included a large numbers of animals in the control and treated groups (range, 20-100). The possible reason for this may be the notably high procedure-related mortality, as is discussed below. These findings agree with those of Kienle *et al* (n=5 to 20) (29) and can be rated adequate according to the current standards (43); however, it should be noted that none of the studies presented a justification for the sample size used.

Tumor type. There was an even distribution of tumor types in the studies, including the following: 14 in melanoma cell lines, 15 in epithelial cell lines and 13 in mesenchymal cell lines. Only 7 studies reported negative or no effects relative to models of melanoma (44-46) or epithelial tumors (40-42,45,47). The toxic effects reported were rare and unrelated to the type of tumor. One study reported skin inflammation and necrosis, mainly peritumoral (26) in mice bearing melanoma, whereas 2 studies revealed weight loss in rats and nephrotoxicity in mice bearing epithelial tumors (22,47).

Preparations. A total of 23 studies used whole plant extracts, 13 used lectins alone, 4 used recombinant lectin I, 3 used triterpenes and 1 used a high and low molecular fraction (> and <30 kDa; total number is >37 as some studies tested more than one substance either alone or in combination). These findings differ from the ones reported by Kienle *et al* (29): Whole plant extracts=24; lectins=2; and recombinant lectins=2. There is a clear tendency towards increased use of isolated active components, lectins in particular, and the previous interest in other isolated proteins and polysaccharides (4 each) appears to have been replaced by lipophilic components.

Route of administration. A total of 36 studies tested mistletoe administered via various systemic routes (intraperitoneal, oral, intranasal, intravenous, subcutaneous) and 5 at the local tumor site. Three studies pretreated the tumor cells and 4 did not report the route used. These results show a tendency towards systemic administration compared to the results of the study by Kienle *et al* (29), in which the frequency of systemic and local administration was similar (17 vs. 15 studies, respectively).

A total of 3 studies compared the subcutaneous (SC) and intraperitoneal (IP) routes of administration. In the Braun *et al* study (48), a reduction in the number of lung and liver colonies of RAW-117 P (liver) and L-1 (lung) lymphosarcoma was obtained only with IP administration of aqueous extract standardized for ML-II/III; however, in a later study they used the SC route only and obtained less positive results (49). Lenartz *et al* (50) reported a reduction in F98 glioma volume

for both the SC and IP routes, but tumor necrosis was observed in the animals treated with IP administration only. This supports the hypothesis that IP administration leads to better pharmacokinetics of the main active compounds. Nevertheless, Burger *et al* (45) compared the SC and IP routes in F9 testicular carcinoma and did not find a difference in the anti-tumor effect.

Yoon *et al* (51) compared a variety of dosages (1-100 μ g) of aqueous extract of Korean mistletoe via various routes-systemic (intravenous, subcutaneous) and mucosal (oral and intranasal). Systemic administration induced a dramatic inhibition of colon carcinoma even at a very low dose (2 μ g). Furthermore, mucosal routes were associated with metastasis inhibition at a low dose and significantly, no antitumor effect was detected with higher doses via the intranasal route. One study (52) used lectin-rich extract (Iscaador Qu) to investigate the effects of pretreatment on tumor cells, systemic and local treatment. Pretreatment mitigated tumor growth, indicating that the antitumor effects mistletoe extract were transferred *in vivo* to growing tumors; however, this effect was not due to reduced cell proliferation or increased cell death. While intra-tumor treatment significantly reduced tumor growth, systemic administration had no significant effect.

Taken together, the results are highly variable and divergent, and no definite conclusions can be drawn. With that proviso, there do appear to be indications that local administration may be more effective (see below).

Reported random allocation. Only 5 of the 37 studies included reported having randomly allocated the animals to the experimental groups (40,53-56), whereas in the Kienle *et al* (29) review, 6 of 31 studies reported randomization. Randomization is a crucial step in any experimental protocol, particularly in the case of complementary medicine as the risk of false positive or false negative results is very high (57). It may therefore be beneficial if a methodological guideline were formulated, as it has been for homeopathic studies (57). In the present review, as well as in the Kienle *et al* review (29), none of the included studies reported whether blinding was performed or not. It has been suggested that this is not reported as it is a standard procedure (29). However, unlike in clinical studies, the use of blinded protocols in pharmacological basic research is very rare. For example, in the present review, searching the key works 'animal model' and 'blind protocol' in PubMed only identified 94 articles. This is another point to be considered in a putative guideline.

Controls/combination therapy. All 37 studies included control groups, as follows: 26 studies used a placebo-saline/distilled water (5), vehicle (18) or culture medium (1); 12 compared to untreated (10) and/or healthy mice (3). A total of 5 studies compared mistletoe with standard chemotherapy/adjuvant drugs (polysaccharide K, cytarabine, doxorubicin, cyclophosphamide, 5-fluorouracil) alone or in combination (25,45,53,58,59); and 1 study compared with interleukin-2 alone and in combination (47).

Aims of studies. The objectives of the studies exhibit clear temporal trends. Studies conducted between 1996 and 2005 sought to establish the antitumor effect of mistletoe and its

possible mechanisms, cytotoxicity and immunomodulation in particular (21,46,54-56,60-68), eventually with comparison to or in combination with standard chemotherapeutic agents. By the end of the 1990 s, the focus of studies shifted towards the study of lectins, in particular ML-I (21,50), also including Korean mistletoe (46,51,53,66). Between 2000 and 2001, novel studies concerning recombinant lectin were beginning to be published (22-24); however, the specific effects of lectins II and III began to be explored later (48,49). The overall profile of studies performed in the past decade is notably different to previous studies, tending to investigate biologically active principles (33,53,59,69) including lipophilic ones, in particular triterpenes (25-27). A shift can be also noticed in investigations of the putative mechanism of action, with a stronger focus on genomics (44,52,69) and oxidative stress (27,28).

Side-effects/toxicity. A total of 12 of 37 studies (~30%) specifically investigated side effects as variables of interest, including mortality (45,53,55,56), clinical morbidity (26,53,55), body weight (22,23,25,53,55,56,59), food intake (23,33,53,55,56), water intake (33), histology on necropsy (25,26,33,55), behavior (26,33,68), vitality score (33), physiological response (33), hematological parameters (59), overall appearance (68) and agility (23). The majority of studies reported no significant toxic findings, the exception being a study by Strüh *et al* (26) in which the highest tested dose of solubilized triterpenes administered SC peritumorally resulted in skin necrosis and apathy, which led to 3/13-14 dropouts. Furthermore, animals in all the treated groups exhibited skin inflammation and peritumoral inflammatory infiltrating cells. In the Seifert *et al* study (59), mistletoe grown on pine trees proved to be more toxic compared to that grown on fir. In the Burger *et al* study (45) mistletoe was demonstrated to be considerably less toxic than doxorubicin, but showed no differences compared with 5-fluorouracil.

It is worth noting that in the studies using *N*-methyl-*N*-nitroso-urea (NMU)-induced autochthonous bladder urinary carcinoma the tumor-induction procedure and complications caused high mortality, ranging from 20 to 50% (22,41,42). In a later study, Kunze *et al* (40) employed *N*-butyl-*N*-(4-hydroxybutyl)-nitrosamine (BBN) as a tumor inducer; although the procedure-related mortality was significantly lower (0 to 11%), the number of animals that developed tumors was also low, as is discussed below.

Outcomes. A total of 32/37 studies reported some antitumor effect, 3 of which had mixed results. Facina *et al* (44) did not detect any changes in tumor volume due to mistletoe treatment, however they did demonstrate evidence of genetic regulation of apoptosis. Burger *et al* (45) reported a dose-dependent reduction in tumor growth for 3 of the tested cell lines, but no effect on the other 2 lines tested. In Park *et al* study (46), mistletoe was demonstrated to have no effect on tumor growth; however, it did induce dose-dependent inhibition of metastasis and led to a slight increase in survival rates. Two studies reported no antitumor effect (41,42) and 2 reported an overt negative effect in the treated groups (40,47). Finally, 1 study assessed immunomodulatory activity only, with no investigation of antitumor effects (62); the results from this study indicated that treatment with mistletoe restored the immune response,

however prolonged application also had a dose-dependent immunosuppressing effect on normal cells. The relationship between the concentration of active molecules and the balance between cytotoxicity and immunomodulatory effects requires further investigation; the combination of both factors in the same protocol may be a useful tool to treat aggressive tumors with mistletoe components.

A total of 21 studies assessed tumor growth inhibition. Of these, 17 found a reduction in tumor size, 7 of them in a dose-dependent manner and with eventual inversion of the effect at a higher dose (33), and one of these studies (45) indicated a reduction in tumor size in only 3 out of the 5 tested cell lines. In 1 study, remission was reported when mistletoe was administered inside the tumor, whereas there was no such effect when it was administered via the systemic route (52). Two studies reported an unexpected increase in tumor size in the treated groups (40,47). Future studies investigating the underlying molecular mechanisms may explain these paradoxical results.

Survival was analyzed in a total of 12 studies, and was higher in the treated groups in all cases, most of them in all studied doses but dose-dependence was reported in 2 studies (24,56).

Inhibition of primary or metastatic tumor formation was assessed by 1 (56) and 12 studies, respectively. Tumorigenesis was inhibited in all studies with the exception of Timoshenko *et al* (47). Based on these findings, the authors of the present study suggest that the action of mistletoe active principles in the tumor microenvironment should be investigated further, as the onset of the metastasis-formation process depends on it. Additionally, the variability of results among the different studies may also be attributed to the fact that most protocols were based on transplantable tumors, in which tumorigenesis cannot be evaluated, since tumor cells are inoculated directly in the animals, being immunogenic stimuli. The cases that are most illustrative of this are the 4 studies that assessed the effect of mistletoe lectin on autochthonous urinary bladder carcinoma. Kunze *et al* (41,42) and Elsässer-Beile *et al* (22) induced tumors using NMU, and Kunze *et al* (40) induced tumors with BBN. In all 3 studies by Kunze *et al*, a lectin isolated from unspecified mistletoe was used. Of these, the 2 studies that induced tumors with NMU had a high procedure-related mortality, and no significant difference was found between the control and treated groups. In contrast, the study that used BBN had a very low procedure-mortality rate, but also a low number of induced tumors in the control and treated groups ($n=6$ and 4 , 10.2 and 6.7% , respectively, in the 6-month experiment; $n=16$ and 13 , 25.8 and 19.7% , respectively, in the 15-month experiment). In this case, at 15 months, the tumor size was twice larger, on average, in the lectin-treated group. Elsässer-Beile *et al* (22) used recombinant lectin starting 8 or 14 weeks following tumor induction and reported significantly lower rates of atypical hyperplasia and neoplastic transformation in all treated groups compared with the control. The main difference between these two groups of experiments is that Kunze *et al* used the subcutaneous route for long periods of time (6 and 15 months), whereas Elsässer-Beile *et al* (22) applied shorter 6-week treatments via the local route of administration (intravesical instillation). It could therefore be that the divergence in these results is due to the methodological

differences between the studies. Mengs *et al* (54) used the intravesical route to administer a lectin-rich aqueous mistletoe extract to mice implanted with urinary bladder carcinoma MB49 cells in a short 4-week course of treatment; the results of this study indicated a dose-dependent statistically significant tumor growth inhibition in the treated groups.

Several putative mechanisms for the anticancer action of mistletoe were tested, including tumor cell viability [reduced, (27,53,63)] and number [reduced, (37)]; tumor cell turnover [increased, (70)]; antiproliferative effect [4 studies, no effect in 1 (26)]; proapoptotic action (increased in 8 studies); increased tumor necrosis (6 studies); increased macrophage cytotoxicity (66); reduction of mitotic figures in tumor cells (65,70); cycle cell regulation [deregulated in one study (53), and no effect in another (46)]. Furthermore, 1 study reported degradation of specific miRNAs, this being an indirect effect (affection of precursors) (71) and 2 studies reported a reduction in oxidative stress (27,58).

Reduced tumor invasiveness was reported by 1 study (65), whereas in another (70), 11% of the treated animals exhibited no histological evidence of viable tumors on day 11 after the onset of treatment. Reduced angiogenesis was identified in 2 studies (26,46); however, in 1 study, the vascular count increased when high doses were used (33). The possible underlying antineoplastic mechanisms require further investigation using specific molecular methods in animal models, as developing an effective mistletoe anticancer agent may on the addition of different mechanisms, possibly associated with a mixture of active molecules in different concentrations and their interactions with the tumor microenvironment. Few studies have been performed from this perspective; a total of 5 studies analyzed the role of local cell infiltrates in tumor development, with an increase in absolute numbers of infiltrating leukocytes being found in 3 studies (23,63,70) and no significant effect found in 2 studies (40,41).

For the immunomodulatory action of mistletoe, induction of the Th-1 response was assessed and detected in 1 study (44) and cytokines were assessed in 2, of which 1 reported no effect (22) and the other reported upregulation of IL-12 only (67). An increase in natural killer cell activity was detected in 2 studies (51,66). Macrophage stimulation was reported in 1 study (55) and 2 studies found an increase in white cell count in the peripheral blood, but not of granulocytes (48,49). Upregulation of the number of thymocytes (49), activation of spleen cells (64), lymphocyte stimulation ($CD4^+$ and $CD8^+$, but only anti $CD8^+$ activity was enhanced) (55) and finally stimulation of cytotoxic population (63) were also reported.

In conclusion, the majority of studies investigating the effects of mistletoe in animal cancer models analyzed in the present review reported positive outcomes in terms of the inhibition of tumor formation and growth. However, there is considerable methodological heterogeneity among studies, which precludes performing conclusive comparisons and explains the variation in results. A great deal of research is still required with standardization of mistletoe preparation, dose, concentration, regime of administration, length of treatment, targeted cancer type, and other aspects. Similarly, while it is universally assumed that mistletoe has cytotoxic and immunomodulatory effects, there is a wide variation among studies in terms of the mechanisms assessed. The present review found little or no continuity within

and among research groups, with little progress being made in the past 20 years of studies with experimental animal models. Based on these results, the present authors suggest that the community of interested researchers should develop a guideline for reporting *in vivo* mistletoe cancer treatment experiments.

Acknowledgements

The authors of the present study would like to thank the Library of Hiscia Institute, Arlesheim, Switzerland, and Library of the School of Medicine, University of São Paulo, Brazil, for the help in the retrieval of articles. The authors would also like to thank Stephan Baumgartner, Claudia Scherr and Maria Olga Kokornaczyk for their helpful comments on the manuscript.

References

1. Carpernaros Z: The golden bough: The case for mistletoe. *Eur J Herbal Med* 1: 19-24, 1994.
2. Marvibaigi M, Supriyanto E, Amini N, Abdul Majid FA and Jaganathan SK: Preclinical and clinical effects of mistletoe against breast cancer. *Biomed Res Int* 2014: 785479, 2014.
3. Kröz M, Kienle GS, Feder G, Kaveri S and Rosenzweig S: Mistletoe: From basic research to clinical outcomes in cancer and other indications. *Evid Based Complement Alternat Med*: 987527, 2014.
4. Evans M, Bryant S, Huntley AL and Feder G: Cancer patients' experiences of using mistletoe (*Viscum album*): A qualitative systematic review and synthesis. *J Altern Complement Med* 22: 134-144, 2016.
5. Tröger W, Galun D, Reif M, Schumann A, Stanković N and Miličević M: Quality of life of patients with advanced pancreatic cancer during treatment with mistletoe: A randomized controlled trial. *Dtsch Arztebl Int* 111: 493-502, 2014.
6. Sunjic SB, Gasparovic AC, Vukovic T, Weiss T, Weiss ES, Soldo I, Djakovic N, Zarkovic T and Zarkovic N: Adjuvant cancer biotherapy by *Viscum album* extract Isorel: Overview of evidence based medicine findings. *Coll Antropol* 39: 701-708, 2015.
7. von Schoen-Angerer T, Wilkens J, Kienle GS, Kiene H and Vagedes J: High-dose *Viscum album* extract treatment in the prevention of recurrent bladder cancer: A retrospective case series. *Perm J* 19: 76-83, 2015.
8. Rose A, El-Leithy T, vom Dorp F, Zakaria A, Eisenhardt A, Tschirdewahn S and Rübber H: Mistletoe plant extract in patients with nonmuscle invasive bladder cancer: Results of phase Ib/IIa single group dose escalation study. *J Urol* 194: 939-943, 2015.
9. Rexer H: Study for the treatment of nonmuscle invasive bladder cancer: A phase III efficacy trial for intravesical instillation of mistletoe extract in superficial bladder cancer (TIM) - AB 40/11 of the AUO. *Urologe A*: 55: 963-965, 2016 (In German).
10. Steele ML, Axtner J, Happe A, Kröz M, Matthes H and Schad F: Adverse drug reactions and expected effects to therapy with subcutaneous mistletoe extracts (*Viscum album* L.) in cancer patients. *Evid Based Complement Alternat Med* 2014: 724258, 2014.
11. Kissel D, Jurin M and Zarkovic N: Über die zytostatischen und immunologischen Effekte von *Viscum album*. *Erfahrungsheilkunde* 39: 59-66, 1990.
12. Jurin M, Zarković N, Hrzenjak M and Ilić Z: Antitumorous and immunomodulatory effects of the *Viscum album* L. *Oncology* 50: 393-398, 1993.
13. Beuth J, Ko HL, Gabius HJ and Pulverer G: Influence of treatment with the immunomodulatory effective dose of the beta-galactoside-specific lectin from mistletoe on tumor colonization in BALB/c-mice for two experimental model systems. *In Vivo* 5: 29-32, 1991.
14. Kuttan G, Vasudevan DM and Kuttan R: Effect of a preparation from *Viscum album* on tumor development in vitro and in mice. *J Ethnopharmacol* 29: 35-41, 1990.
15. Saha C, Hegde P, Friboulet A, Bayry J and Kaveri SV: *Viscum album*-mediated COX-2 inhibition implicates destabilization of COX-2 mRNA. *PLoS One* 10: e0114965, 2015.
16. Hegde P, Maddur MS, Friboulet A, Bayry J and Kaveri SV: *Viscum album* exerts anti-inflammatory effect by selectively inhibiting cytokine-induced expression of cyclooxygenase-2. *PLoS One* 6: e26312, 2011.
17. Lyu SY and Park WB: Mistletoe lectin transport by M-cells in follicle-associated epithelium (FAE) and IL-12 secretion in dendritic cells situated below FAE in vitro. *Arch Pharm Res* 33: 1433-1441, 2010.
18. Lyu SY and Park WB: Mistletoe lectin modulates intestinal epithelial cell-derived cytokines and B cell IgA secretion. *Arch Pharm Res* 32: 443-451, 2009.
19. Lavastre V, Cavalli H and Girard D: Anti-inflammatory effect of *Viscum album* agglutinin-I (VAA-I): Induction of apoptosis in activated neutrophils and inhibition of lipopolysaccharide-induced neutrophilic inflammation in vivo. *Clin Exp Immunol* 137: 272-278, 2004.
20. Lavastre V, Pelletier M, Saller R, Hostanska K and Girard D: Mechanisms involved in spontaneous and *Viscum album* agglutinin-I-induced human neutrophil apoptosis: *Viscum album* agglutinin I accelerates the loss of antiapoptotic Mcl-1 expression and the degradation of cytoskeletal paxillin and vimentin proteins via caspases. *J Immunol* 168: 1419-1427, 2002.
21. Zarković N, Kalisnik T, Lončarić I, Borović S, Mang S, Kissel D, Konitzer M, Jurin M and Grainza S: Comparison of the effects of *Viscum album* lectin ML-1 and fresh plant extract (Isorel) on the cell growth in vitro and tumorigenicity of melanoma B16F10. *Cancer Biother Radiopharm* 13: 121-131, 1998.
22. Elsässer-Beile U, Ruhnau T, Freudenberg N, Wetterauer U and Mengs U: Antitumoral effect of recombinant mistletoe lectin on chemically induced urinary bladder carcinogenesis in a rat model. *Cancer* 91: 998-1004, 2001.
23. Schaffrath B, Mengs U, Schwartz T, Hilgers RD, Beuth J, Möckel B, Lentzen H and Gerstmayer B: Anticancer activity of rViscumin (recombinant mistletoe lectin) in tumor colonization models with immunocompetent mice. *Anticancer Res* 21: 3981-3987, 2001.
24. Schumacher U, Feldhaus S and Mengs U: Recombinant mistletoe lectin (rML) is successful in treating human ovarian cancer transplanted into severe combined immunodeficient (SCID) mice. *Cancer Lett* 150: 171-175, 2000.
25. Delebinski CI, Twardziok M, Kleinsimon S, Hoff F, Mulsow K, Rolff J, Jäger S, Eggert A and Seifert G: A natural combination extract of *Viscum album* L. containing both triterpene acids and lectins is highly effective against AML in vivo. *PLoS One* 10: e0133892, 2015.
26. Strüh CM, Jäger S, Kersten A, Schempp CM, Scheffler A and Martin SF: Triterpenoids amplify anti-tumoral effects of mistletoe extracts on murine B16.f10 melanoma in vivo. *PLoS One* 8: e62168, 2013.
27. Cebović T, Spasić S and Popović M: Cytotoxic effects of the *Viscum album* L. Extract on Ehrlich tumour cells in vivo. *Phytother Res* 22: 1097-1103, 2008.
28. Büssing A (ed). Mistletoe: The Genus *Viscum*. Harwood Academic Publishing, Amsterdam, 2000.
29. Kienle GS, Glockmann A, Schink M and Kiene H: *Viscum album* L. extracts in breast and gynaecological cancers: A systematic review of clinical and preclinical research. *J Exp Clin Cancer Res* 28: 79, 2009.
30. Baumgartner S, Flückiger H, Kunz M, Scherr C and Urech K: Evaluation of preclinical assays to investigate an anthroposophic pharmaceutical process applied to mistletoe (*Viscum album* L.) extracts. *Evid Based Complement Alternat Med* 2014: 620974, 2014.
31. Horneber MA, Bueschel G, Huber R, Linde K and Rostock M: Mistletoe therapy in oncology. *Cochrane Database Syst Rev* CD003297, 2008.
32. National Cancer Institute: Mistletoe Extracts (PDQ®). <http://www.cancer.gov/about-cancer/treatment/cam/hp/mistletoe-pdq>. Accessed August 1, 2016.
33. Thies A, Dautel P, Meyer A, Pfüller U and Schumacher U: Low-dose mistletoe lectin-I reduces melanoma growth and spread in a scid mouse xenograft model. *Br J Cancer* 98: 106-112, 2008.
34. Begley CG and Ellis LM: Drug development: Raise standards for preclinical cancer research. *Nature* 483: 531-533, 2012.
35. Mansky PJ, Wallerstedt DB, Sannes TS, Stagl J, Johnson LL, Blackburn MR, Grem JL, Swain SM and Monahan BP: NCCAM/NCI phase 1 study of mistletoe extract and gemcitabine in patients with advanced solid tumors. *Evid Based Complement Alternat Med* 2013: 964592, 2013.

36. Carvalho AC, Porto E and Bonamin L: Canine neurofibrosarcoma treatment with *Viscum album* in serial dilutions. *Int J High Dilution Res* 12: 106, 2013.
37. Christen-Clottu O, Klocke P, Burger D, Straub R and Gerber V: Treatment of clinically diagnosed equine sarcoid with a mistletoe extract (*Viscum album austriacus*). *J Vet Intern Med* 24: 1483-1489, 2010.
38. Blostin R and Faivre C: Bénéfices du gui fermenté chez le chat après exérèse de fibrosarcome: Résultats d'une étude préliminaire. *Phytothérapie* 6: 352-358, 2008.
39. Lefebvre GNF, Bonamin LV and Oliveira CM: Treatment of transmissible venereal tumor (TVT) in dogs with *Viscum album* (Mistletoe) associated to chemotherapy. *Clínica Veterinária* 12: 78-86, 2007.
40. Kunze E, Schulz H, Adamek M and Gabius HJ: Long-term administration of galactoside-specific mistletoe lectin in an animal model: No protection against N-butyl-N-(4-hydroxybutyl)-nitrosamine-induced urinary bladder carcinogenesis in rats and no induction of a relevant local cellular immune response. *J Cancer Res Clin Oncol* 126: 125-138, 2000.
41. Kunze E, Schulz H and Gabius HJ: Inability of galactoside-specific mistletoe lectin to inhibit N-methyl-N-nitrosourea-induced tumor development in the urinary bladder of rats and to mediate a local cellular immune response after long-term administration. *J Cancer Res Clin Oncol* 124: 73-87, 1998.
42. Kunze E, Schulz H, Ahrens H and Gabius HJ: Lack of an antitumoral effect of immunomodulatory galactoside-specific mistletoe lection on N-methyl-N-nitrosourea-induced urinary bladder carcinogenesis in rats. *Exp Toxicol pathol* 49: 167-180, 1997.
43. Charan J and Kantharia ND: How to calculate sample size in animal studies? *J Pharmacol Pharmacother* 4: 303-306, 2013.
44. Facina AS, Facina G, Silva IDCG, Gonçalves GA, Almeida FA, Noronha SMR and Nakamura MU: *Viscum album* modulates apoptotic related genes in melanoma tumor of mice. *Am J Mol Biol* 4: 49-58, 2014.
45. Burger AM, Mengs U, Schüler JB and Fiebig HH: Anticancer activity of an aqueous mistletoe extract (AME) in syngeneic murine tumor models. *Anticancer Res* 21: 1965-1968, 2001.
46. Park WB, Lyu SY, Kim JH, Choi SH, Chung HK, Ahn SH, Hong SY, Yoon TJ and Choi MJ: Inhibition of tumor growth and metastasis by Korean mistletoe lectin is associated with apoptosis and antiangiogenesis. *Cancer Biother Radiopharm* 16: 439-447, 2001.
47. Timoshenko AV, Lan Y, Gabius HJ and Lala PK: Immunotherapy of C3H/HeJ mammary adenocarcinoma with interleukin-2, mistletoe lectin or their combination: Effects on tumour growth, capillary leakage and nitric oxide (NO) production. *Eur J Cancer* 37: 1910-1920, 2001.
48. Braun JM, Ko HL, Schierholz JM, Weir D, Blackwell CC and Beuth J: Application of standardized mistletoe extracts augments immune response and down regulates metastatic organ colonization in murine models. *Cancer Lett* 170: 25-31, 2001.
49. Braun JM, Ko HL, Schierholz JM and Beuth J: Standardized mistletoe extract augments immune response and down-regulates local and metastatic tumor growth in murine models. *Anticancer Res* 22: 4187-4190, 2002.
50. Lenartz D, Andermahr J, Plum G, Menzel J and Beuth J: Efficiency of treatment with galactoside-specific lectin from mistletoe against rat glioma. *Anticancer Res* 18: 1011-1014, 1998.
51. Yoon TJ, Yoo YC, Kang TB, Baek YJ, Huh CS, Song SK, Lee KH, Azuma I and Kim JB: Prophylactic effect of Korean mistletoe (*Viscum album coloratum*) extract on tumor metastasis is mediated by enhancement of NK cell activity. *Int J Immunopharmacol* 20: 163-172, 1998.
52. Podlech O, Harter PN, Mittelbronn M, Pöschel S and Naumann U: Fermented mistletoe extract as a multimodal antitumoral agent in gliomas. *Evid Based Complement Alternat Med* 2012: 501796, 2012.
53. Han SY, Hong CE, Kim HG and Lyu SY: Anti-cancer effects of enteric-coated polymers containing mistletoe lectin in murine melanoma cells in vitro and in vivo. *Mol Cell Biochem* 408: 73-87, 2015.
54. Mengs U, Schwarz T, Bulitta M and Weber K: Antitumoral effects of an intravesically applied aqueous mistletoe extract on urinary bladder carcinoma MB49 in mice. *Anticancer Res* 20: 3565-3568, 2000.
55. Weber K, Mengs U, Schwarz T, Hajto T, Hostanska K, Allen TR, Weyhenmeyer R and Lentzen H: Effects of a standardized mistletoe preparation on metastatic B16 melanoma colonization in murine lungs. *Arzneimittelforschung* 48: 497-502, 1998.
56. Kuttan G, Menon LG and Kuttan R: Prevention of 20-methylcholantrene-induced sarcoma by a mistletoe extract Iscador. *Carcinogenesis* 17: 1107-1109, 1996.
57. Stock-Schröer B, Albrecht H, Betti L, Endler PC, Linde K, Lüdtke R, Musial F, van Wijk R, Witt C and Baumgartner S: Reporting experiments in homeopathic basic research (REHBaR)-a detailed guideline for authors. *Homeopathy* 98: 287-298, 2009.
58. Stan RL, Hangan AC, Dican L, Sevastre B, Hanganu D, Catoi C, Sarpataki O and Ionescu CM: Comparative study concerning mistletoe viscotoxins antitumor activity. *Acta Biol Hung* 64: 279-288, 2013.
59. Seifert G, Jesse P, Laengler A, Reindl T, Lüth M, Lobitz S, Henze G, Prokop A and Lode HN: Molecular mechanisms of mistletoe plant extract-induced apoptosis in acute lymphoblastic leukemia in vivo and in vitro. *Cancer Lett* 264: 218-228, 2008.
60. Kuttan G, Menon LG, Antony S and Kuttan R: Anticarcinogenic and antimetastatic activity of Iscador. *Anticancer Drugs* 8 (Suppl 1): S15-S16, 1997.
61. Antony S, Kuttan R and Kuttan G: Effect of *Viscum album* in the inhibition of lung metastasis in mice induced by B16F10 melanoma cells. *J Exp Clin Cancer Res* 16: 159-162, 1997.
62. Jurin M, Zarkovic N, Borovic S and Kissel D: *Viscum album* L. preparation Isorel modifies the immune response in normal and intumour-bearing mice. *Anticancer Drugs* 8 (Suppl 1): S27-S31, 1997.
63. Zarkovic N, Zarkovic K, Grainca S, Kissel D and Jurin D: The *Viscum album* preparation Isorel inhibits the growth of melanoma B16F10 by influencing the tumour-host relationship. *Anticancer Drugs* 8 (Suppl 1): S17-S22, 1997.
64. Antony S, Kuttan R and Kuttan G: Inhibition of lung metastasis by adoptive immunotherapy using Iscador. *Immunol Invest* 28: 1-8, 1999.
65. Zarkovic N, Vukovic T, Loncaric I, Miletic M, Zarkovic K, Borovic S, Cipak A, Sabolovic S, Konitzer M and Mang S: An overview on anticancer activities of the *Viscum album* extract Isorel. *Cancer Biother Radiopharm* 16: 55-62, 2001.
66. Yoon TJ, Yoo YC, Kang TB, Song SK, Lee KB, Her E, Song KS and Kim JB: Antitumor activity of the Korean mistletoe lectin is attributed to activation of macrophages and NK cells. *Arch Pharm Res* 26: 861-867, 2003.
67. Duong Van Huyen JP, Delignat S, Bayry J, Kazatchkine MD, Bruneval P, Nicoletti A and Kaveri SV: Interleukin-12 is associated with the in vivo anti-tumor effect of mistletoe extracts in B16 mice melanoma. *Cancer Lett* 243: 32-37, 2006.
68. Beuth J, Ko HL, Schneider H, Tawadros S, Kasper HU, Zimst H and Schierholtz JM: Intratumoral application of standardized mistletoe extracts down regulates tumor weight via decreased cell proliferation, increased apoptosis and necrosis in a murine model. *Anticancer Res* 26: 4451-4456, 2006.
69. Li LN, Zhang HD, Zhi R and Yuan SJ: Down-regulation of some miRNAs by degrading their precursors contributes to anti-cancer effect of mistletoe lectin-I. *Br J Pharmacol* 162: 349-364, 2011.
70. Pryme IF, Bardocz S, Pusztai A, Ewen SW and Pfüller U: A mistletoe lectin (ML-I)-containing diet reduces the viability of a murine non-Hodgkin lymphoma tumor. *Cancer Detect Prev* 28: 52-56, 2004.
71. Lee CH, Kim JK, Kim HY, Park SM and Lee SM: Immunomodulating effects of Korean mistletoe lectin in vitro and in vivo. *Int Immunopharmacol* 9: 1555-1561, 2009.