

Correlation between unexplained recurrent spontaneous abortion with CD4⁺CD25⁺ regulatory T-cell and killer cell immunoglobulin-like receptor levels

XIAOZHEN QUAN and XUEZHOU YANG

Department of Obstetrics and Gynecology, Xiangyang Central Hospital,
The Affiliated Hospital of Hubei University of Arts and Science, Xiangyang, Hubei 441021, P.R. China

Received May 12, 2017; Accepted June 8, 2017

DOI: 10.3892/etm.2017.4634

Abstract. The present study investigated the correlation between unexplained recurrent spontaneous abortion (URSA) with CD4⁺CD25⁺ regulatory T-cell (Treg) and killer cell immunoglobulin-like receptor (KIR)-2DL1 levels. A total of 76 URSA patients were enrolled (35 without pregnancy, Group A, and 41 with early abortion, Group B). Additionally, 30 patients who received a regular abortion as planned (Group C) and 30 healthy volunteers (Group D) were selected. Peripheral venous blood and fresh decidual tissue samples were obtained from all the patients, and flow cytometry was performed to detect CD4⁺CD25⁺Treg and Foxp3 transcription factor levels. mRNA and protein KIR-2DL1 expression levels were assayed using quantitative PCR and western blot analysis, respectively. No statistically significant differences in peripheral venous blood CD4⁺CD25⁺Treg/CD4⁺ and Foxp3⁺/CD4⁺CD25⁺Treg cell proportions were found among the groups ($P>0.05$). However, the decidual tissues of Group C presented significantly higher levels of both cell types versus other groups ($P<0.05$). No statistically significant differences were found in comparisons among Groups A, B, and D ($P>0.05$). In peripheral venous blood, mRNA and protein KIR-2DL1 expression levels in Group C were significantly higher than those in the other three groups ($P<0.05$), but again, there were no statistically significant differences among Groups A, B, and D ($P>0.05$). In decidual tissues, KIR-2DL1 levels were significantly higher in Group C relative to Groups A, B, and D ($P<0.05$). Decreased CD4⁺CD25⁺Treg counts and KIR-2DL1 expression levels were closely associated

with the onset of URSA. CD4⁺CD25⁺Tregs mainly exert their effects on decidual tissues, while KIR-2DL1 can act on peripheral venous blood and decidual tissues. These may present new targets for early intervention in URSA.

Introduction

In women of reproductive age, the incidence of spontaneous abortion ranges from 5 to 25%. Previous findings have indicated that spontaneous abortion is associated with gene expression, environmental factors, immune tolerance, inflammatory response, microbiological contamination, endocrine disorders, and diseases in reproductive tract diseases (1,2). However, 30-50% of patients suffer from unexplained recurrent spontaneous abortion (URSA), which is generally considered to be caused by a decrease in immune tolerance (3). Maternal circulatory and placental tissue cell abnormalities, as well as abnormal humoral immune function, can affect the regular transfer and development of the embryo, resulting in early abortion. It has been found that regulatory T-cell (Treg) levels, especially those of the CD4⁺CD25⁺ subtypes, are significantly decreased in the peripheral blood and decidual tissues of patients with spontaneous abortion, and this decrease aligns well with the timing of the abortion (4).

Natural killer (NK) cells account for 65-80% of the lymphocytes in uterine decidual tissues, and in contrast to peripheral blood populations, uterine NK cells are mainly of the CD56^{bright}CD16^{dim} subtype (5). NK cell activity is regulated by killer cell immunoglobulin-like receptors (KIR). KIR-2DL1, as an inhibitor, can specifically identify and combine with HLA-Cw molecules spread on the surface of fetal trophoblastic cells to transmit inhibitory signals to maintain maternal-fetal immune tolerance (6).

In this study, we aimed to analyze the correlation between URSA and CD4⁺CD25⁺Treg and KIR-2DL1 expression levels to identify new potential targets for clinical treatment.

Patients and methods

Patient information. We sequentially selected 76 patients who were diagnosed with URSA at Xiangyang Central Hospital between June 2015 and October 2016. Of these, 35 patients were

Correspondence to: Dr Xuezhou Yang, Department of Obstetrics and Gynecology, Xiangyang Central Hospital, The Affiliated Hospital of Hubei University of Arts and Science, 136 Jingzhou Street, Xiangcheng, Xiangyang, Hubei 441021, P.R. China
E-mail: yangxz258@163.com

Key words: CD4⁺CD25⁺ regulatory T cells, killer cell immunoglobulin-like receptor, unexplained recurrent spontaneous abortion

not pregnant (Group A) and 41 patients suffered early abortion (Group B). We additionally selected 30 patients who received a regular abortion as planned (Group C) and 30 healthy volunteers (Group D). Diagnostic criteria for URSA were as follows: i) patients experiencing spontaneous abortion at least twice; ii) patients with no explicit etiological factors (e.g., abnormal chromosomes, reproductive tract diseases, microbiological infections, endocrine disorders, menstrual disorders, positive test results for antinuclear and anticardiolipin antibodies, negative test results for blocking antibodies, medications, trauma and exercise); and iii) patients whose partners had no semen abnormalities. Written informed consent was obtained from all the subjects.

The average age of Group A was 25.6 ± 6.3 years, the average number of abortions was 2.5 ± 0.4 times, and the average gestational age at abortion was 7.3 ± 1.2 weeks. The average age of Group B was 24.3 ± 5.7 years, the average number of abortions was 2.4 ± 0.5 times, and the average gestational age at abortion was 7.6 ± 0.8 weeks. The average age of Group C was 23.9 ± 5.5 years and the average gestational age at abortion was 6.9 ± 1.6 weeks. The average age of Group D was 23.5 ± 5.9 years. No significant differences in these parameters were observed among groups ($P > 0.05$).

Observation indices. Peripheral venous blood and fresh decidual tissues were obtained from patients to detect CD4⁺CD25⁺Treg and Foxp3 transcription factor levels via flow cytometry. KIR-2DL1 mRNA and protein levels were assayed via quantitative PCR and western blot analysis, respectively.

Flow cytometry. Fasting venous blood (5 ml) was collected and subjected to gradient density centrifugation to isolate peripheral blood mononuclear cells. Fresh decidual tissues were also collected and converted into cellular suspensions. Mouse anti-human CD4-FITC (5 μ l) and CD 25-PE (5 μ l) monoclonal antibodies (Beyotime Biotech Co., Ltd., Jiangsu, China) were added, respectively, to two tubes, which were incubated for 20 min at 37°C in the dark. Buffer (0.5 ml) containing a permeabilization reagent was then added and the tubes were incubated for an additional 30 μ l in the dark at 4°C. The samples were then washed using PBS, centrifuged for 5 min at 1,500 \times g, and then pelleted cells were resuspended after discarding of the supernatant. The cells were then rinsed using permeabilization buffer and incubated with 1 μ l normal mouse serum and 50 μ l permeabilization buffer for 15 min in the dark at 4°C. APC mouse IgG2a isotype antibody (10 μ l; Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd., Beijing, China) was added to one tube as a control, while 10 μ l APC-anti-human Foxp3 antibody (Sigma-Aldrich, St. Louis, MO, USA) was added into the other tube. Both tubes were incubated for 30 min at 4°C in the dark, after which the cells in the two tubes were washed using permeabilization buffer. Cells were finally suspended in PBS for 1 h and detected using a FACS Caliber flow cytometer (BD Biosciences, New Jersey, NY, USA).

qPCR. Regular TRIzol reagent was used to extract total cellular RNA; total concentration and purity were determined via ultraviolet spectrometer and cDNA synthesis was performed using a reverse transcription kit. Primers

were synthesized by Sangon Biotech Co., Ltd. (Shanghai, China) in accordance with GeneBank sequences: KIR-2DL1 forward, 5'-GCAGCACCATGTCGCTCT-3' and reverse, 5'-GTCAGTGGGAGCTGACAC-3', 356 bp; GAPDH forward, 5'-CGCGAGAAGATGACCCAGAT-3' and reverse, 5'-GCACTGTGTTGGCGTACAGG-3', 226 bp. SYBR-Green I was used to detect cDNA amplification using a Prism 7500 Sequence Detector Fluorescence Quantitative PCR Amplifier (both from ABI, Vernon, CA, USA). Reaction system components were as follows: cDNA 3.0 μ l + upstream primer 1.0 μ l + downstream primer 1.0 μ l + 2X SYBR-Green I Mix 5.0 μ l + ddH₂O 10.0 μ l. Reaction conditions were as follows: 95°C for 5 min, 30 cycles of 95°C for 30 sec, 58°C for 30 sec, and 72°C for 60 sec, and a final 10 min step at 72°C. PCR products were identified using 2% agarose gel electrophoresis, ultraviolet imaging was performed using a gel imaging analysis system, and gray value analysis was carried out via digital photos. The detection procedures were repeated three times and the results were presented via the $2^{-\Delta\Delta C_q}$ method.

Western blot analysis. Radio immunoprecipitation assay (RIPA kit from R&D Systems, Minneapolis, MN, USA) lysis buffer was added to samples in order to extract total cellular protein, with a preliminary quantitative assay performed via the Coomassie Brilliant Blue method (kit from Bio-Rad Laboratories, Hercules, CA, USA). Total protein (30 μ g) was subjected to 8% SDS-PAGE to separate proteins, which were then electrically transferred onto polyvinylidene fluoride membranes (4°C, 100 V, 50 min). The membranes were blocked using 5% skimmed milk at room temperature for 2 h. Mouse anti-human KIR-2DL1 monoclonal antibody (dilution, 1:500; cat. no. sc-53595; Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) was added onto the membrane, which was incubated with constant agitation at 4°C overnight. The membrane was then washed using Tris-buffered saline and Tween-20 (TBST) and incubated with horseradish peroxidase-labeled rabbit anti-mouse IgG poly-clonal secondary antibody (dilution, 1:2,000; cat. no. A11008; Invitrogen, Carlsbad, CA, USA) for 4 h with constant agitation at 37°C. Following this, the membranes were washed again using TBST and treated with ECL for protein visualization. Results were scanned and preserved, with semi-quantitative analysis performed using Lab Works 4.5 gel imaging software (Media Cybernetics, Rockville, MD, USA). Results were expressed as integral optical density units. Experiments were carried out three times in order to determine the mean.

Statistical analysis. SPSS 20.0 software (IBM SPSS, Armonk, NY, USA) was used for statistical analysis. Measurement data were presented as mean \pm standard deviation. Single-factor analysis of variance was used for intergroup comparisons, the least significant difference-test was used for paired comparisons, and the independent t-test was employed for comparisons between the two groups. $P < 0.05$ was used to indicate statistical significance.

Results

CD4⁺CD25⁺Treg and Foxp3 levels. We found no statistically significant differences between groups in terms

Table I. CD4⁺CD25⁺Treg and Foxp3 levels (%).

Variables	Group A	Group B	Group C	Group D
Peripheral blood				
CD4 ⁺ CD25 ⁺ Treg	4.77±1.22	4.86±1.32	5.62±1.54	5.33±1.46
Foxp3	19.85±8.63	21.52±8.52	25.69±7.95	20.34±7.84
Decidual tissues				
CD4 ⁺ CD25 ⁺ Treg	8.65±2.52	12.67±6.49	24.52±12.32	10.34±5.87
Foxp3	35.62±16.24	52.95±25.85	89.69±35.57	46.32±24.64

Treg, regulatory T cell.

Table II. KIR-2DL1 mRNA and protein expression levels.

Variables	Group A	Group B	Group C	Group D
Peripheral blood				
mRNA	0.0869±0.0124	0.1127±0.0564	0.2451±0.1324	0.0958±0.0232
Protein	0.07±0.02	0.09±0.03	0.22±0.11	0.07±0.02
Decidual tissues				
mRNA	0.1765±0.0856	0.2546±0.1025	0.4659±0.1654	0.1854±0.0965
Protein	0.17±0.08	0.23±0.09	0.44±0.21	0.16±0.05

KIR, killer cell immunoglobulin-like receptor.

of peripheral venous blood CD4⁺CD25⁺Treg/CD4⁺ and Foxp3⁺/CD4⁺CD25⁺Treg percentage ($P>0.05$). Group C decidual tissues showed elevated levels of both indices compared to the other groups ($P<0.05$), but no statistically significant differences were found between Groups A, B, and D ($P>0.05$; Table I).

KIR-2DL1 mRNA and protein expression levels. Peripheral venous blood KIR-2DL1 mRNA and protein expression levels in Group C were significantly higher than in the other three groups ($P<0.05$), but there were no statistically significant differences between Groups A, B, and D ($P>0.05$). In decidual tissues, Group C levels were significantly higher in Groups A, B, and D ($P<0.05$; Table II).

Discussion

Research has suggested that CD4⁺CD25⁺Treg cells play an extremely important role in cellular immune function. During immunological rejection of organ transplantation, there is a decrease in the amount of CD4⁺CD25⁺Treg cells and CD4⁺CD25⁺ T cells are activated to generate cytotoxicity, causing immune injuries and assisting B-cell involvement in the antigen-antibody combination reaction (7). After T-cell receptor stimulation and activation, Tregs can suppress the transcription and expression of the *IL-2* gene in CD4⁺ and CD8⁺ cells in order to inhibit the proliferation and activation of T cells, thus regulating immune tolerance. Foxp3, a characteristic molecule expressed by CD4⁺CD25⁺Tregs, can regulate the development and maintain the function of cells.

Foxp3 is highly expressed in CD4⁺CD25^{high}, lowly expressed in CD4⁺CD25^{low}, and not expressed in CD4⁺CD25⁻ cells (8). A study by Khattri *et al* revealed that the elimination of Foxp3⁺CD4⁺CD25⁺Treg cells in the mouse using a specific antibody can cause severe autoimmune responses (9). Aluvihare *et al* found that Foxp3 expression levels in the uteruses of intrabreeding BALB/c pregnant mice and interbreeding BALB/c and C57BL/6 pregnant mice were 1,000-fold greater than those in non-pregnant mice of the same age (10). However, no statistically significant differences were found in Foxp3 expression level in uteruses between intrabreeding BALB/c pregnant mice and interbreeding BALB/c and C57BL/6 pregnant mice (10). This result suggested that the increase in Foxp3 expression was caused by pregnancy but not correlated with homoantigen stimulation. In the present study, we found no differences in the proportion of peripheral venous blood CD4⁺CD25⁺Treg/CD4⁺ and Foxp3⁺/CD4⁺CD25⁺Treg cells. We also found that Group C decidual tissues showed significantly higher levels of both indices, but no statistically significant differences were found among Groups A, B, and D. This finding suggested that decreased CD4⁺CD25⁺Treg levels and Foxp3 expression were closely correlated with the onset of URSA, and CD4⁺CD25⁺Treg-mediated immune tolerance was mainly generated in local decidual tissues. However, this result did not coincide with results found in a previous study (11). Local immune regulation in decidual tissues lay the foundation for immune identification and responses between the matrix and the embryo. Formation, uterine settlement, nutrition, and embryo development can significantly affect the local decidual tissues and induce increased levels of CD4⁺CD25⁺Treg cells.

These, however, seldomly enter the peripheral circulation, where there are insufficient corresponding antigens or receptors to interact with CD4⁺CD25⁺Treg (12).

KIR-2DL1, a transmembrane glycoprotein, can specifically identify the 77th amino acid, aspartic acid, and 80th amino acid, lysine, in the HLA-Cw molecule. Immunoreceptor tyrosine-based inhibitory motifs inside the phosphorylated structure can transmit negative regulation signals to inhibit the killing activity of NK cells (13). Ntrivalas *et al* (14) and Yamada *et al* (15) have confirmed that in women with recurrent spontaneous abortion, peripheral blood NK expression of KIR-2DL1 was significantly lower than in healthy women. In this study, we found that peripheral venous blood KIR-2DL1 mRNA and protein expression levels in Group C were significantly higher than those in the other three groups, but there were no statistically significant differences among Groups A, B, and D. In decidual tissues, Group C expression levels were significantly higher than those in Groups A, B, and D. This result suggested that a decreased expression of KIR-2DL1 was closely correlated with the onset of URSA, and can occur in both peripheral blood and decidual tissues. NK cells can simultaneously mediate cellular and humoral immunity. In addition to high expression on the surface of fetal trophoblast cells, HLA-Cw is also expressed on the surface of T and B cells (16). In conclusion, CD4⁺CD25⁺Treg and KIR-2DL1 may serve as new targets for the early intervention of URSA.

Acknowledgements

The present study was supported by the Natural Science Foundation of Hubei Province of China (grant no. 2014CFB251).

References

1. Işık G, Demirezen Ş, Dönmez HG and Beksaç MS: Bacterial vaginosis in association with spontaneous abortion and recurrent pregnancy losses. *J Cytol* 33: 135-140, 2016.
2. Wu T, Yin B, Zhu Y, Li G, Ye L, Chen C, Zeng Y and Liang D: Molecular cytogenetic analysis of early spontaneous abortions conceived from varying assisted reproductive technology procedures. *Mol Cytogenet* 9: 79, 2016.
3. Zhu L, Chen H, Liu M, Yuan Y, Wang Z, Chen Y, Wei J, Su F and Zhang J: Treg/Th17 cell imbalance and IL-6 profile in patients with unexplained recurrent spontaneous abortion. *Reprod Sci* 24: 882-890, 2016.
4. Ebina Y, Shimada S, Deguchi M, Maesawa Y, Iijima N and Yamada H: Divergence of helper, cytotoxic, and regulatory T cells in the decidua from miscarriage. *Am J Reprod Immunol* 76: 199-204, 2016.
5. Moffett A, Chazara O, Colucci F and Johnson MH: Variation of maternal KIR and fetal HLA-C genes in reproductive failure: Too early for clinical intervention. *Reprod Biomed Online* 33: 763-769, 2016.
6. Dambaeva SV, Lee DH, Sung N, Chen CY, Bao S, Gilman-Sachs A, Kwak-Kim J and Beaman KD: Recurrent pregnancy loss in women with killer cell immunoglobulin-like receptor KIR2DS1 is associated with an increased HLA-C2 allelic frequency. *Am J Reprod Immunol* 75: 94-103, 2016.
7. Safinia N, Vaikunthanathan T, Fraser H, Thirkell S, Lowe K, Blackmore L, Whitehouse G, Martinez-Llordella M, Jassem W, Sanchez-Fueyo A, *et al*: Successful expansion of functional and stable regulatory T cells for immunotherapy in liver transplantation. *Oncotarget* 7: 7563-7577, 2016.
8. Winger EE and Reed JL: Low circulating CD4(+) CD25(+) Foxp3(+) T regulatory cell levels predict miscarriage risk in newly pregnant women with a history of failure. *Am J Reprod Immunol* 66: 320-328, 2011.
9. Khattri R, Cox T, Yasayko SA and Ramsdell F: Pillars article: An essential role for Scurfin in CD4⁺CD25⁺ T regulatory cells. *Nat Immunol* 4: 337-342, 2003.
10. Aluvihare VR, Kallikourdis M and Betz AG: Tolerance, suppression and the fetal allograft. *J Mol Med (Berl)* 83: 88-96, 2005.
11. Alijotas-Reig J, Melnychuk T and Gris JM: Regulatory T cells, maternal-fetal immune tolerance and recurrent miscarriage: New therapeutic challenging opportunities. *Med Clin (Barc)* 144: 265-268, 2015.
12. Lu QB and Zhu S: Modulation of an aqueous extract of Chinese medicine prescription Anzi Heji on ratio of CD4(+)CD25(+) FOXP3(+) regulatory T cells in anticardiolipin antibody-positive patients with threatened abortion. *Chin J Integr Med* 26: 654-655, 2016.
13. Ozturk OG, Sahin G, Karacor ED and Kucukgoz U: Evaluation of KIR genes in recurrent miscarriage. *J Assist Reprod Genet* 29: 933-938, 2012.
14. Ntrivalas EI, Bowser CR, Kwak-Kim J, Beaman KD and Gilman-Sachs A: Expression of killer immunoglobulin-like receptors on peripheral blood NK cell subsets of women with recurrent spontaneous abortions or implantation failures. *Am J Reprod Immunol* 53: 215-221, 2005.
15. Yamada H, Shimada S, Kato EH, Morikawa M, Iwabuchi K, Kishi R, Onoé K and Minakami H: Decrease in a specific killer cell immunoglobulin-like receptor on peripheral natural killer cells in women with recurrent spontaneous abortion of unexplained etiology. *Am J Reprod Immunol* 51: 241-247, 2004.
16. Dakhly DM, Bayoumi YA, Sharkawy M, Gad Allah SH, Hassan MA, Gouda HM, Hashem AT, Hatem DL, Ahmed MF and El-Khayat W: Intralipid supplementation in women with recurrent spontaneous abortion and elevated levels of natural killer cells. *Int J Gynaecol Obstet* 135: 324-327, 2016.