

IMB-6G induces endoplasmic reticulum stress-mediated apoptosis in human nasopharyngeal carcinoma cells

YETING PAN, YANNI ZHANG, LIANG GONG, JIANDING ZOU, BOXIA HU and SICONG ZHANG

Department of Otorhinolaryngology, Affiliated Cixi Hospital of Wenzhou Medical College, Cixi, Zhejiang 315300, P.R. China

Received March 22, 2018; Accepted June 29, 2018

DOI: 10.3892/etm.2018.6724

Abstract. IMB-6G is a novel N-substituted sophoridine acid that has been reported to have anticancer effects. The purpose of the present study was to investigate the effect and underlying mechanism of IMB-6G on human nasopharyngeal carcinoma (NPC) cells. The NPC cell line C666-1 was used in the present study and treated with different concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h. Subsequently, cell viability was determined using the Cell Counting kit-8 assay and cell apoptosis was analyzed by performing flow cytometry. The expression levels of genes and proteins in the current study were determined using reverse transcription-quantitative polymerase chain reaction and western blot analysis, respectively. Results indicated that IMB-6G dose-dependently inhibited C666-1 cell viability and induced apoptosis. It was also revealed that IMB-6G induced apoptosis via inducing endoplasmic reticulum (ER) stress activation. Notably, IMB-6G administration enhanced the expression levels of Binding immunoglobulin protein and CCAAT-enhancer-binding protein homologous protein in C666-1 cells. Further analysis suggested that IMB-6G treatment activated inositol-requiring enzyme 1 α (IRE1 α) and PKR-like ER kinase (PERK) signaling pathways in C666-1 cells. In addition, silencing of IRE1 α and PERK significantly reversed IMB-6G-induced cell growth inhibition and apoptosis. In conclusion, the present findings indicated that IMB-6G induced ER stress-mediated apoptosis through activating IRE1 α and PERK signaling pathways. The present study suggests that IMB-6G may be a promising agent for NPC treatment.

Introduction

As the most common malignant epithelial tumor of the head and neck in Southeast Asia and Southern China, nasopharyngeal carcinoma (NPC) seriously affects the quality of human life (1,2). Various factors, including genetic, environmental and viral factors are involved in the development of NPC (3-5). Notably, early detection of NPC remains difficult in the clinic because of a lack of NPC-identifiable symptoms and high metastatic potential. Approximately 75% of the patients were reported with the late stage NPC and local node metastases and/or distant metastasis at first diagnosis (6). Once recurrence or metastasis occurs following treatment, the prognosis of NPC is poor, which makes it the leading cause of fatality (7). Because of the frequent metastasis and poor prognosis, NPC has been identified as a highly malignant tumor (8). The standard approach for the treatment of NPC is chemoradiotherapy (9). Currently, although great progress has been made in the development of therapeutic strategies for NPC, the therapeutic efficacy remains unsatisfactory. Therefore, studies for novel treatment therapies for NPC are urgently required.

Sophoridine is a major bioactive component that is isolated from the traditional medicinal herb *Sophora alopecuroides* L (10). Studies have reported that sophoridine provides an anti-tumor effect against malignant trophoblastic tumors (11,12). As a novel N-substituted sophoridine acid derivative, IMB-6G has been indicated to have an anti-proliferation effect on several human tumor cells through inducing apoptosis and G₀/G₁ cell cycle arrest (13). Zhang *et al* (14) reported that IMB-6G induces endoplasmic reticulum (ER) stress-mediated apoptosis through the activation of inositol-requiring enzyme 1 α (IRE1 α) and PKR-like ER kinase (PERK) signaling pathways. Furthermore, IMB-6G also exhibits reasonable bio-availability and has been demonstrated to possess positive pharmacokinetic properties and exhibits good safety profile *in vivo* (13,15). However, to the best of our knowledge, no study has investigated the influence of IMB-6G in NPC cells.

The purpose of the present study was to investigate the anticancer activity and the underlying molecular mechanisms of IMB-6G against human NPC cells. The present findings investigated whether IMB-6G induces human NPC cell apoptosis by activating ER stress. Additionally, the study explored whether IRE1 α - and PERK-mediated ER stress may be involved in IMB-6G-induced apoptosis.

Correspondence to: Dr Jianding Zou, Department of Otorhinolaryngology, Affiliated Cixi Hospital of Wenzhou Medical College, 999 South Second Ring Road, Cixi, Zhejiang 315300, P.R. China
E-mail: zoujiand1803@163.com

Key words: IMB-6G, nasopharyngeal carcinoma, apoptosis, endoplasmic reticulum stress

Materials and methods

Cell culture and treatment. The NPC cell line C666-1 was obtained from the Chinese Academy of Sciences Cell Bank (Shanghai, China) and cultured in RPMI-1640 medium supplemented with 10% fetal bovine serum (HyClone, Logan, UT, USA), 1% penicillin/streptomycin solution at 37°C in an atmosphere containing 5% CO₂. Cells were treated with various concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h at 37°C as described previously (14) prior to experiments described below.

Cell transfection. The C666-1 cells were transfected with 50 nM IRE1 α -small interfering (si)RNA (5'-GCGUCUUU ACUACGUAUUCU-3'; Shanghai GenePharma Co., Ltd., Shanghai, China), 50 nM PERK-siRNA (cat no. sc-36213) or 50 nM control-siRNA (cat no. sc-37007; both Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) using Lipofectamine 2000 (Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA) according to the manufacturer's protocol. Following 24 h of transfection, the cells were treated with or without 5 μ M IMB-6G for 24 h prior to the subsequent experiments.

Cell viability detection. For cell viability detection, the Cell Counting kit-8 (CCK-8) assay was used. In brief, C666-1 cells were plated into a 96-well plate (5x10³ cells/well) and incubated for 24 h at 37°C. Subsequently, the cells were treated with various concentrations (0.000, 0.625, 1.250, 2.500, 5.000, 10.000 and 15.000 μ M) of IMB-6G for 24 h. Each well was treated with 10 μ l CCK-8 assay solution (Dojindo Molecular Technologies, Inc., Kumamoto, Japan). Cells were subsequently incubated for a further 1 h at 37°C. To assess the cell viability, the optical density at 570 nm was measured using a microplate reader.

Apoptosis analysis. To detect C666-1 cell apoptosis, an Annexin V-fluorescein isothiocyanate (FITC) apoptosis detection kit (cat no. 45115; Cell Signaling Technology, Inc., Danvers, MA, USA) was utilized. C666-1 cells were treated with various concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h. C666-1 cells (5x10⁵ cells per well) were subsequently labeled with Annexin V-FITC and propidium iodide per as the manufacturer's protocol. Flow cytometry (BD Biosciences, Franklin Lakes, NJ, USA) was performed to analyze cell apoptosis, and data were analyzed using WinMDI (version 2.5; Purdue University Cytometry Laboratories; www.cyto.purdue.edu/flowcyt/software/Catalog.htm).

Western blot analysis. Total cellular protein from C666-1 cells was extracted using radioimmunoprecipitation assay buffer (cat. no. P0013B; Beyotime Institute of Biotechnology, Nanjing, China). A bicinchoninic protein assay kit was used to detect protein concentration. Protein samples (25 μ g per lane) were separated using 12% SDS-PAGE and then transferred to a polyvinylidene membrane. Following this, the membranes were blocked with 5% non-fat milk for 1 h at room temperature and blotted overnight at 4°C with the following primary antibodies: Cleaved caspase-9 (cat. no. 20750; 1:1,000 dilution), cleaved caspase-3 (cat. no. 9579; 1:1,000 dilution),

binding immunoglobulin protein (Bip; cat. no. 3177; 1:1,000 dilution), CCAAT-enhancer-binding protein homologous protein (CHOP; cat. no. 5554; 1:1,000 dilution), IRE1 α (cat. no. 3294; 1:1,000 dilution; all Cell Signaling Technology, Inc.), p-IRE1 α (ab48187; 1:1,000 dilution; Abcam, Cambridge, MA, USA), PERK (cat. no. 5683; 1:1,000 dilution), p-PERK (cat. no. 3179; 1:1,000 dilution), p-eukaryotic initiation factor 2 α (eIF2 α ; cat. no. 3597; 1:1,000 dilution), eIF2 α (cat. no. 5324; 1:1,000 dilution) and β -actin (cat. no. 4970; 1:1,000 dilution; all Cell Signaling Technology, Inc.). The membranes were subsequently incubated with the anti-rabbit IgG horseradish peroxidase-linked antibody (cat. no. 7074; 1:5,000 dilution; Cell Signaling Technology, Inc.) at room temperature for 2.5 h. Protein bands were visualized using enhanced chemiluminescence reagent (EMD Millipore, Billerica, MA, USA).

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Total RNA was isolated from C666-1 cells using TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol. Total RNA was reverse transcribed into cDNA using PrimeScript RT Reagent kit (Takara Bio Inc., Otsu, Japan) according to the manufacturer's protocol. SYBR Premix Ex Taq II (Takara Bio Inc.) was used for qPCR analysis. Amplification conditions were as follows: 95°C for 10 min, followed by 40 cycles at 95°C for 15 sec, 72°C for 30 sec and 78°C for 1.5 min. Samples were stored at 4°C. GAPDH was used as an internal control. All primer sequences for PCR were listed in Table I. Relative gene expression was analyzed using the 2^{- $\Delta\Delta$ C_q} method (16).

Statistical analysis. Experiments were performed in triplicate. Data were presented as the mean \pm standard deviation. SPSS 16.0 statistical software (SPSS, Inc., Chicago, IL, USA) was applied for all statistical analyses. One-way analysis of variance followed by a post hoc Tukey's test or a Student's t-test was used to analyze the differences between groups. P<0.05 was considered to indicate a statistically significant difference.

Results

IMB-6G inhibits C666-1 cell viability and induces cell apoptosis. To investigate the anticancer effect of IMB-6G on NPC, human NPC cells (C666-1) were treated with a variety of concentrations of IMB-6G (0.000, 0.625, 1.250, 2.500, 5.000, 10.000 and 15.000 μ M) for 24 h, and assessed via CCK-8 assay analysis to measure C666-1 cell viability. Fig. 1A revealed the molecular structure of IMB-6G. As indicated in Fig. 1B, IMB-6G inhibited the cell viability of C666-1 cells in a dose-dependent manner. To determine whether cell apoptosis participates in IMB-6G-induced NPC cell death, C666-1 cell apoptosis was analyzed by flow cytometry following treatment with different concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h (Fig. 1C and D). Furthermore, western blot analysis indicated that IMB-6G promoted the expression levels of cleaved caspase-9 and cleaved caspase-3 (Fig. 1E). The findings suggested that IMB-6G dose-dependently induced C666-1 cell apoptosis and inhibited cell growth.

IMB-6G activates ER stress in C666-1 cells. ER stress has been identified to serve an important role in regulating

Table I. Primer sequence for polymerase chain reaction.

Gene	Direction	Sequence (5'-3')
Bip	Forward	5'-TAGCGTATGGTGCTGCTGTC-3'
	Reverse	5'-TTTGTCAAGGGTCTTTCACC-3'
CHOP	Forward	5'-GAGGAGAGAGTGTTCAGAAGG-3'
	Reverse	5'-TCTGGGAGGTGCTTGTGAC-3'
IRE1 α	Forward	5'-TAGTCAGTTCTGCGTCCGCT-3'
	Reverse	5'-TTCCAAAAATCCCAGGGCCG-3'
PERK	Forward	5'-AGGACAGAGGGGACAGAGTTG-3'
	Reverse	5'-TAATGACCTTTTCTTCCCTGCTCC-3'
GAPDH	Forward	5'-CTTTGGTATCGTGAAGGACTC-3'
	Reverse	5'-GTAGAGGCAGGGATGATGTTCT-3'

Bip, binding immunoglobulin protein; CHOP, CCAAT-enhancer-binding protein homologous protein; IRE1 α , inositol-requiring enzyme 1 α ; PERK, PKR-like ER kinase.




Figure 1. IMB-6G inhibits cell viability and induces apoptosis in C666-1 cells. (A) Structure of the IMB-6G molecule. (B) C666-1 cells were treated with various concentrations of IMB-6G (0.000, 0.625, 1.250, 2.500, 5.000, 10.000 and 15.000 μ M) for 24 h and the cell viability was assessed using the Cell Counting kit-8 assay. (C) C666-1 cells were treated with various concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h and cell apoptosis was determined by flow cytometry. (D) Statistical analysis results of flow cytometric analysis of apoptosis were indicated. (E) Protein expression levels of cleaved caspase-9 and cleaved caspase-3 were measured using western blot analysis. Data were presented as the mean \pm standard deviation. * $P < 0.05$, ** $P < 0.01$ vs. 0 μ M IMB-6G. PI, propidium iodide; FITC, fluorescein isothiocyanate.

apoptosis (17,18). Thus, it was determined whether ER stress was involved in IMB-6G-mediated apoptosis in the present study. Following treatment with different concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h, the protein and mRNA expression levels of Bip and CHOP in C666-1 cells were detected by western blot analysis and RT-qPCR, respectively. Results indicated that IMB-6G dose-dependently

increased the protein and mRNA expression levels of Bip and CHOP (Fig. 2).

IRE1 α and PERK are involved in the IMB-6G-activated ER stress signaling pathway. To identify whether the ER stress signaling pathway is activated by IMB-6G, the phosphorylation of PERK, IRE1 α and eIF2 α was measured in C666-1 cells




Figure 2. IMB-6G triggers ER stress in C666-1 cells. C666-1 cells were treated with various concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h and the expression levels of Bip and CHOP were measured by (A) western blot analysis and (B and C) reverse transcription-quantitative polymerase chain reaction, respectively. Data were presented as the mean \pm standard deviation. * P <0.05, ** P <0.01 vs. 0 μ M IMB-6G. Bip, binding immunoglobulin protein; CHOP, CCAAT-enhancer-binding protein homologous protein.




Figure 3. IMB-6G activates the ER stress signaling pathway via IRE1 α and PERK. (A) C666-1 cells were treated with various concentrations of IMB-6G (0, 1, 2 and 5 μ M) for 24 h and the phosphorylation of PERK (Thr980), IRE1 α (Ser724) and eIF2 α (Ser51) was measured by western blot analysis. (B-D) The protein expression levels of p-PERK (Thr980), p-IRE1 α (Ser724) and p-eIF2 α (Ser51) were analyzed. (E) The protein levels of total PERK, total IRE1 α and total eIF2 α in C666-1 cells were measured by western blotting. Data were presented as the mean \pm standard deviation. * P <0.05, ** P <0.01 vs. 0 μ M IMB-6G. ER, endoplasmic reticulum; IRE1 α , inositol-requiring enzyme 1 α ; PERK, PKR-like ER kinase; eIF2 α , eukaryotic initiation factor 2 α .

following treatment with IMB-6G at various concentrations of IMB-6G (0, 1, 2, 5 μ M) for 24 h. Western blot analysis demonstrated that IMB-6G administration dose-dependently enhanced p-PERK, p-IRE1 α and p-eIF2 α expression levels (Fig. 3). Furthermore, the current study determined that IMB-6G administration exhibited no effect on the expression of PERK, IRE1 α and eIF2 α total protein in C666-1 cells.

IMB-6G induces C666-1 cell apoptosis via IRE1 α and PERK pathways. To confirm whether the IRE1 α and PERK signaling pathways are involved in IMB-6G-induced apoptosis of NPC cells, the effect of IRE1 α or PERK inhibition on IMB-6G-induced C666-1 cell apoptosis was investigated. Following transfection with control-siRNA, PERK-siRNA or IRE1 α -siRNA for 24 h, C666-1 cells were treated with

0 or 5 μ M IMB-6G for another 24 h. The results demonstrated that IRE1 α or PERK knockdown significantly reversed the IMB-6G-induced cell growth inhibition and apoptosis when compared with the control group (Fig. 4).

Discussion

NPC is considered to be relatively sensitive to chemotherapy due to the fact that >90% of NPCs are of the undifferentiated variety (19). Although some of the known therapies for NPC have demonstrated high efficacy, the survival of patients with advanced NPC has not been significantly improved (20). Therefore, there is an urgency to identify novel and efficient agents for the treatment of patients with advanced NPC. IMB-6G, a novel N-substituted sophoridinic acid derivative, has




Figure 4. IRE1 α and PERK signaling pathways are critical for IMB-6G-induced NPC cell viability inhibition and apoptosis. C666-1 cells were transfected with IRE1 α -siRNA, PERK-siRNA or control-siRNA for 24 h and then treated with or without 5 μ M IMB-6G for 24 h. (A) Protein expression levels of IRE1 α and PERK were measured by western blot analysis. (B and C) mRNA expression levels of IRE1 α and PERK were measured by reverse transcription-quantitative polymerase chain reaction. (D) Cell viability was assayed using a Cell Counting kit-8 assay. (E) Cell apoptosis was detected by flow cytometry. Data were presented as the mean \pm standard deviation. **P<0.01 vs. si-control group; #P<0.05 and ##P<0.01 vs. IMB-6G+si-control group. siRNA, small interfering RNA; IRE1 α , inositol-requiring enzyme 1 α ; PERK, PKR-like ER kinase.

been demonstrated to have anti-tumor effect (13,14); however, little is known about its effect on NPC. In the present study, the effect of IMB-6G and any associated possible signaling pathways was investigated on human NPC C666-1 cells *in vitro*.

Firstly, the impact of IMB-6G on C666-1 cell viability was investigated. The results suggested that IMB-6G dose-dependently inhibited C666-1 cell viability and induced cell apoptosis, indicating IMB-6G induced cytotoxicity and apoptosis in human NPC cells. Mounting evidence has revealed that ER stress is critical in the regulation of apoptosis (17,18). ER serves an important role in regulating protein folding and trafficking, calcium homeostasis and lipid synthesis (21). Notably, a variety of factors can induce ER stress and trigger an adaptive response called the unfolded protein response (22). To investigate whether ER stress was involved in IMB-6G-mediated apoptosis, the ER stress responses in IMB-6G-treated NPC C666-1 cells were analyzed in the present study. The findings indicated that ER stress activation may be involved in the induced cytotoxicity of IMB-6G, as suggested by the increased expression levels of Bip and CHOP in IMB-6G-treated C666-1 cells. Subsequently, the signaling pathways that participated in the effect of IMB-6G on human NPC cells were explored. B-cell lymphoma 2 (Bcl-2)/Bcl-2-associated X protein-regulated Ca²⁺ release from the ER, IRE1 α -mediated tumor necrosis factor receptor-associated factor 2 activation and PERK/eIF2 α -dependent induction of the proapoptotic transcriptional factor CHOP (C/EBP homologous protein) are all associated with ER stress-mediated apoptosis (23-25). A previous study reported that IMB-6G induces ER stress-mediated apoptosis in human hepatocellular carcinoma cells by activating

IRE1 α and PERK signaling pathways (14). Therefore, it was investigated whether IRE1 α and PERK signaling pathways were involved in ER stress-mediated apoptosis in human NPC C666-1 cells in the present study. The findings indicated that IMB-6G dose-dependently enhanced the expression levels of p-PERK and p-IRE1 α , indicating the involvement of IRE1 α and PERK signaling pathways in ER stress-mediated apoptosis in human NPC cells. In addition, it was revealed that IRE1 α and PERK knockdown significantly reversed IMB-6G-induced cell viability inhibition and apoptosis, which indicated the activation of IRE1 α and PERK signaling pathways served an important role in human NPC cell apoptosis induced by IMB-6G.

Taken together, the present study suggested that IMB-6G induced cytotoxicity and apoptosis in human NPC C666-1 cells through activating ER stress via the activation of IRE1 α and PERK signaling pathways. The data indicated that IMB-6G is a potential anticancer agent that acts via inducing ER stress associated apoptosis in human NPC cells. Therefore, targeting ER stress by IMB-6G may be a promising treatment strategy for the treatment of NPC.

Acknowledgements

Not applicable.

Funding

The present study was supported by the Project of Zhejiang traditional Chinese Medicine Research (grant no. 2013ZB115) and the Scientific Project of CiXi (grant no. CN2013013).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YP designed the current study. YP, YZ, and LG accessed and analyzed the data. JZ, BH and SZ interpreted results. All authors collaborated to write the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Chang ET and Adami HO: The enigmatic epidemiology of nasopharyngeal carcinoma. *Cancer Epidemiol Biomark Prev* 15: 1765-1777, 2006.
- Cao SM, Simons MJ and Qian CN: The prevalence and prevention of nasopharyngeal carcinoma in China. *Chin J Cancer* 30: 114-119, 2011.
- Lo KW and Huang DP: Genetic and epigenetic changes in nasopharyngeal carcinoma. *Semin Cancer Biol* 12: 451-462, 2002.
- Tao Q and Chan AT: Nasopharyngeal carcinoma: Molecular pathogenesis and therapeutic developments. *Expert Rev Mol Med* 9: 1-24, 2007.
- Young LS and Murray PG: Epstein-Barr virus and oncogenesis: From latent genes to tumours. *Oncogene* 22: 5108-5121, 2003.
- Katano A, Takahashi W, Yamashita H, Yamamoto K, Ando M, Yoshida M, Saito Y, Abe O and Nakagawa K: Radiotherapy alone and with concurrent chemotherapy for nasopharyngeal carcinoma: A retrospective study. *Medicine (Baltimore)* 97: e0502, 2018.
- Chiang AK, Mak NK and Ng WT: Translational research in nasopharyngeal carcinoma. *Oral Oncol* 50: 345-352, 2014.
- Wang J, Guo LP, Chen LZ, Zeng YX and Lu SH: Identification of cancer stem cell-like side population cells in human nasopharyngeal carcinoma cell line. *Cancer Res* 67: 3716-3724, 2007.
- Sze H, Blanchard P, Ng WT, Pignon JP and Lee AW: Chemotherapy for nasopharyngeal carcinoma-current recommendation and controversies. *Hematol Oncol Clin North Am* 29: 1107-1122, 2015.
- Wang WX, Sun ZH, Chen HM, Xu BN and Wang FY: Role and mechanism of Sophoridine on proliferation inhibition in human glioma U87MG cell line. *Int J Clin Exp Med* 8: 464-471, 2015.
- Liang L, Wang XY, Zhang XH, Ji B, Yan HC, Deng HZ and Wu XR: Sophoridine exerts an anti-colorectal carcinoma effect through apoptosis induction in vitro and in vivo. *Life Sci* 91: 1295-1303, 2012.
- Zhao WC, Song LJ and Deng HZ: Effect of sophoridine on dextran sulfate sodium-induced colitis in C57BL/6 mice. *J Asian Nat Prod Res* 12: 925-933, 2010.
- Bi CW, Zhang CX, Li YH, Tang S, Deng HB, Zhao WL, Wang Z, Shao RG and Song DQ: Novel N-substituted sophoridinol derivatives as anticancer agents. *Eur J Med Chem* 81: 95-105, 2014.
- Zhang N, Bi C, Liu L, Dou Y, Tang S, Pang W, Deng H and Song D: IMB-6G, a novel N-substituted sophoridinic acid derivative, induces endoplasmic reticulum stress-mediated apoptosis via activation of IRE1 α and PERK signaling. *Oncotarget* 7: 23860-23873, 2016.
- Bi C, Zhang C, Li Y, Tang S, Wang S, Shao R, Fu H, Su F and Song D: Synthesis and biological evaluation of sophoridinol derivatives as a novel family of potential anticancer agents. *ACS Med Chem Lett* 5: 1225-1229, 2014.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2011.
- Herold MJ, O'Reilly LA, Lin A, Srivastava R, Doerflinger M, Bouillet P, Strasser A and Puthalakath H: Evidence against upstream regulation of the unfolded protein response (UPR) by pro-apoptotic BIM and PUMA. *Cell Death Dis* 5: e1354, 2014.
- Verfaillie T, Garg AD and Agostinis P: Targeting ER stress induced apoptosis and inflammation in cancer. *Cancer Lett* 332: 249-264, 2013.
- Gao W, Chan JY and Wong TS: Curcumin exerts inhibitory effects on undifferentiated nasopharyngeal carcinoma by inhibiting the expression of miR-125a-5p. *Clin Sci (Lond)* 127: 571-579, 2014.
- Peng H, Chen L, Chen YP, Li WF, Tang LL, Lin AH, Sun Y and Ma J: The current status of clinical trials focusing on nasopharyngeal carcinoma: A comprehensive analysis of clinical trials. *gov database. PLoS One* 13: e0196730, 2018.
- Wang M and Kaufman RJ: The impact of the endoplasmic reticulum protein-folding environment on cancer development. *Nat Rev Cancer* 14: 581-597, 2014.
- Walter P and Ron D: The unfolded protein response: From stress pathway to homeostatic regulation. *Science* 334: 1081-1086, 2011.
- Ron D and Hubbard SR: How IRE1 reacts to ER stress. *Cell* 132: 24-26, 2008.
- Ghosh AP, Klocke BJ, Ballester ME and Roth KA: CHOP potentially co-operates with FOXO3a in neuronal cells to regulate PUMA and BIM expression in response to ER stress. *PLoS One* 7: e39586, 2012.
- Han J, Back SH, Hur J, Lin YH, Gildersleeve R, Shan J, Yuan CL, Krokowski D, Wang S, Hatzoglou M, *et al*: ER-stress-induced transcriptional regulation increases protein synthesis leading to cell death. *Nat Cell Biol* 15: 481-490, 2013.