

Alteration of vaginal microbiota in patients with unexplained recurrent miscarriage

FENG ZHANG¹, TAO ZHANG², YINGYING MA³, ZHANGQIAN HUANG¹,
YAO HE¹, HAITAO PAN², MIN FANG² and HAIGANG DING¹

¹Department of Gynecology; ²Genetic Laboratory; ³Reproductive Center,
Shaoxing Women and Children's Hospital, Shaoxing, Zhejiang 312000, P.R. China

Received February 26, 2018; Accepted October 12, 2018

DOI: 10.3892/etm.2019.7337

Abstract. The way in which a balanced vaginal microbiome helps prevent gynecological diseases in women and maintain health remains to be fully elucidated. In the present study, the potential effect of aberrations in the vaginal flora on unexplained recurrent miscarriage (RM) was investigated. The vaginal bacterial communities of 10 patients with unexplained RM and 10 healthy volunteers were sampled and subjected to sequencing analysis of the V3-V4 regions of the bacterial 16S ribosomal RNA gene using the Illumina MiSeq platform. Beta diversity analysis/principal component analysis indicated that bacterial community structures were different between the RM and control groups. A lower microbiota diversity in samples from RM patients was revealed by alpha diversity estimation. Taxonomic analysis demonstrated that abundance of three types of phyla (*Firmicutes*, *Actinobacteria* and *Bacteroidetes*) was significantly different between the RM and the normal control group. Furthermore, at the genus level, *Lactobacillus* was the most dominant genus in the two groups. Statistically significant differences were observed in 5 genera between the two groups. In the RM group, 3 bacterial taxa (*Atopobium*, *Prevotella* and *Streptococcus*) were significantly more abundant, while only 2 taxa were overrepresented in the control group (*Lactobacillus* and *Gardnerella*). In conclusion, the present results provide experimental evidence supporting dysbiosis of the vaginal flora in women with RM.

Introduction

Unexplained recurrent miscarriage (RM) is defined as ≥ 3 consecutive idiopathic miscarriages prior to the gestational age of 12 weeks of pregnancy. RM affects 1-2% of females

attempting to conceive and represents a considerable challenge for physicians (1,2). RM may be explained by genetic abnormalities, maternal immunological causes and endocrine dysfunction only in a minority of cases, while the majority of cases are diagnosed as unexplained RM (3). The vagina harbors different species of bacteria in variable quantities and relative proportions that have important effects on the health of women (4). A healthy vaginal flora, dominated by *Lactobacillus* species, has an important role in the protection against genital infections, which are considered as a common cause of miscarriage (5). A change in the bacterial flora of the vagina is represented by a decrease of *Lactobacillus* and overgrowth of several anaerobic or facultative bacteria, including *Mobiluncus* species, *Prevotella* species, *Gardnerella vaginalis* and genital *Mycoplasma*: *Mycoplasma hominis* and *Ureaplasma urealyticum* (6). At present, bacterial vaginosis is thought to be mainly caused by vaginal dysbacteriosis, which may lead to premature birth, premature rupture of membranes, low birth weight at premature birth, RM, chorioamnionitis and a series of diseases (7).

Advances in sequencing technologies and analytical methods have enabled the exploration of the human microbiota (8). At the end of 2007, the US National Institutes of Health invested hundreds of millions of dollars to start a five-year human microbial genome project, mainly using high-throughput sequencing technology analysis of different parts of the body's bacterial community structure and the female vaginal micro-ecological environment is an important part of this program (9,10). Theoretically, 16S ribosomal (r) RNA gene sequencing is designed to detect most pathogenic bacteria. Via this method, bacteria that are not detectable by cultivation and empirical antibiotic treatments are identifiable. In addition, this method may be utilized to identify novel pathogens that have not been previously recognized as etiologic agents (11).

The diversity of the microflora in the female reproductive tract is affected by various drugs, as well as changes in the environment, and the host's hormone levels and immune system (12). Numerous studies have demonstrated that the microbial balance in the reproductive tract is closely associated with a poor outcome of pregnancy (13-16). However, due to the complex etiology and pathogenesis of RM, the association of aberrations in the vaginal flora and RM has remained to

Correspondence to: Mr. Haigang Ding, Department of Gynecology, Shaoxing Women and Children's Hospital, 305 East Street, Yuecheng, Shaoxing, Zhejiang 312000, P.R. China
E-mail: haigangding1234@sina.com

Key words: unexplained recurrent miscarriage, vaginal microbiome, bacterial diversity, 16S ribosomal RNA

be sufficiently elucidated. In the present study, the diversity of the vaginal microbial community in patients with unexplained RM was analyzed in order to determine the involvement of the vaginal flora in the mechanisms of RM.

Materials and methods

Sample collection. Vaginal secretions of 10 women with RM (case group) and 10 healthy volunteers (control group) were collected at Shaoxing Women and Children's Hospital (Shaoxing, China) between September 2016 and March 2017. The samples were collected in accordance with the relevant guidelines and all subjects provided written informed consent for use of their samples in the present study. The present study was approved by the Ethics Committee of Shaoxing Women and Children's Hospital (Shaoxing, China). None of the women were pregnant, while they were of reproductive age (case group, 20–35 years, median age, 28 years; control group, 26–34 years; median age, 30 years; $P < 0.05$), had a regular menstrual cycle and a history of sexual activity, and had not taken any antibiotics or antimycotic drugs in the past 30 days. Women were asked to refrain from sexual activity for 3 days prior to sampling. Women were excluded from the study if they had used douches, vaginal medications or suppositories, feminine sprays, genital wipes or contraceptive spermicides, or had reported vaginal discharge in the past 48 h (12). The secretions were scraped from the vaginal walls of the RM patients and healthy controls using aseptic swabs. From each subject, two vaginal swabs were collected. The sample centrifuge tubes were immediately placed in a prepared ice box or in a foam box filled with ice packs to maintain a low temperature. The samples were then transferred to the laboratory and stored at -80°C .

DNA extraction and polymerase chain reaction (PCR) amplification. Microbial DNA was extracted from vaginal discharge using the QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) according to manufacturer's protocols. The V3–V4 region of the bacterial 16S rRNA gene was amplified by PCR (ABI GeneAmp[®] 9700). The thermocycling protocol was as follows: 95°C for 2 min, followed by 25 cycles at 95°C for 30 sec, 55°C for 30 sec and 72°C for 30 sec and a final extension at 72°C for 5 min. The primers used were 341F 5'-barcode-CCTACG GGGGCGWGCAG-3' and 805R 5'-barcode-GACTACHVGG GTATCTAATCC-3', where the barcode is an eight-base sequence unique to each sample. PCR was performed in triplicate in 20 μl mixtures containing 4 μl 5X Fast Pfu Buffer, 2 μl 2.5 mM deoxynucleoside triphosphates, 0.8 μl of each primer (5 μM), 0.4 μl FastPfu Polymerase and 10 ng template DNA (17). All samples were subjected to the same conditions and each sample was assessed in duplicate. The PCR products of the same sample were mixed and examined by 2% agarose gel electrophoresis. The PCR products were recovered using AxyPrepDNA Gel Recovery Kit (Axygen; Corning, Inc., Corning, NY, USA) Tris-HCl elution (18).

Illumina MiSeq sequencing. The PCR products were quantified using the QuantiFluor[™]-ST Blue Fluorescence Quantitative System (Promega Corp., Madison, WI, USA) after obtaining the initial quantitative results of the electrophoresis, followed

by mixing the corresponding proportions according to the sequencing requirements of each sample. Amplicons were extracted from 2% agarose gels and purified using the AxyPrep DNA Gel Extraction kit (Axygen; Corning, Inc.) according to the manufacturer's instructions and quantified using QuantiFluor[™]-ST (Promega Corp.). Purified amplicons were pooled in equimolar amounts and paired-end sequenced (2x250) on an Illumina MiSeq platform according to the standard protocols.

Processing of sequencing data. Sequences produced by Miseq sequencing contain barcode sequences, and the primers and linker sequences are added at the time of sequencing. First, the primer sequences were removed using Cutadapt and the pairs were merged into a sequence according to the overlap between paired-end (PE) reads by using The PE reAd merger (19). Next the sample data were identified and distinguished according to the barcode tag sequences. Finally, quality control and filtration of each sample was performed using Prinseq (20) to obtain data.

Raw fastq files were demultiplexed and quality-filtered using FLASH (21) with the following criteria: i) The 3' end of the sequencing primer was removed and the Read1 3' end sequencing link was TGGAATTCTCGGGTGCCAAGGAAC TC; ii) according to the overlap between PE reads, pairs of reads were merged into a sequence, and the allowed maximum mismatch ratio of the splicing sequence overlap area was 0.1; iii) data for each sample were divided from the merged data according to each sample's barcode sequence; iv) the reads were truncated at any site receiving an average quality score of <20 over a 10-bp sliding window; if the average quality value of the window was <20 , the back-end base from the window was cut. v) Reads containing N sequences were removed along with the short sequence in the data, with a length threshold of 200 bp. vi) The low-complexity sequences were filtered.

Bacterial community characterization. According to the barcodes, the high-throughput pyrosequencing reads from 20 vaginal secretion samples were reassigned to samples. Operational taxonomic units (OTUs) were clustered with a 97% similarity cutoff through Usearch (version 7.0; <http://drive5.com/uparse/>) (22), and chimeric sequences were identified and removed using UCHIME (23). Alpha diversity analysis was performed using the OTUs that reached a similarity level of 97%, and the species diversity in a single sample was analyzed by evaluation of Chao, abundance-based coverage estimators (ACE) and Simpon parameters using Mothur (24). A Venn diagram was generated to illustrate the shared and unique OTUs among the groups, based on the occurrence of OTUs in a group of samples regardless of their relative abundance, and this was analyzed using the Venn Diagram package of R language tools (25).

Beta diversity was analyzed to investigate the similarity of the bacterial community structure among groups using Weighted Unifrac distances and visualized analysis via principal component analysis (PCoA). Circos, a sample and species association map, is a visual circle diagram that describes the correspondence between sample and species. The functional profiles of microbial communities were predicted by using PICRUSt (26). The 10,506 OTUs were normalized to the 16S




Figure 1. Alpha diversity of the bacterial flora in the two group. (A) ACE diversity, (B) Chao diversity, (C) Richness diversity and (D) Simpon diversity. Box edges represent the quartiles, horizontal lines represent the medians, bars indicate edge values and circles indicate outliers. ACE, abundance-based coverage estimators.

rRNA copy number and their metagenomic contributions were predicted according to the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways. Based on the PICRUSt functional secondary classification results, the differences in abundance between samples were compared.

Statistical analysis. All statistical analyses were performed using R packages (V.3.2) (25). The screening criterion for analyzing differences in bacterial abundance between samples was $P \leq 0.05$. Fisher's exact test was used for comparison between samples and Welch's t-test was used for comparison between groups (27). Finally, the P-value was verified by using the false discovery rate to obtain the q-value.

Results

Alpha diversity analysis between the case and control groups. To characterize the vaginal microbiota and potential variations associated with RM, the vaginal secretions from 10 cases of RM were collected, and those from 10 healthy women were used as a control group. In total, 745,971 usable sequences from 20 samples were obtained. A total of 10,506 OTUs were delineated at a 97% similarity level. On average, >90% of reads in a sample were retained. At a 97% sequence similarity level, Good's coverage values for all sampled bacterial communities ranged from 0.989 to 0.995 when estimated from all reads (Table I).

Alpha diversity was applied to analyze the complexity of the bacterial diversity of the samples. In Fig. 1, a boxplot

of richness estimators was generated using the QIIME toolkit, providing a clear visualization of the diversities in the different sample groups. No significant difference in Simpson, ACE and Chao diversity was identified between the control and case groups in the box plots ($P > 0.05$; Fig. 1). Community richness comparison indicated that the case group had a significantly increased number of observed and estimated OTUs compared with those in the control group ($P = 0.037$). As displayed in the Venn diagram, the groups had 1,924 OTUs in common, and the proportion of unique OTUs among the total OTUs was 43.8 and 41.9% in the case and control group, respectively. However, the case group had 63 more OTUs than the control group (Fig. 2). This result demonstrated a significantly higher diversity in the case group compared with that in the control group. As presented in Fig. 3, the Simpson rarefaction curves for all samples were saturated, which indicated that the 16S rRNA gene sequence was highly abundant in the database, and that the present analysis had an adequate depth to retrieve most of the information on microbial diversity.

Characteristics of beta diversity analysis between the case and control groups. In the PCoA plot (Fig. 4), each dot represents one sample. PCoA based on 10,506 OTUs (grouped at a sequence identity of 97%) revealed a slight separation of the two different groups on the first two PCo scores, PCo1 and PCo2, which accounted for 36.8 and 29.9% of the total variations, respectively. This indicated that the development of RM may contribute to microbiota imbalance. Furthermore,

Table I. Summary of pyrosequencing data.

Sample_ID	Sequence_no.	OTUs	Filtered OTUs	Shannon_index	ACE_index	Chao1_index	Coverage	Simpon
RM_control1	34948	447	37322	1.2204	641.11437	580.913043	0.994964	0.56009
RM_control2	39002	463	41410	0.421023	1672.661269	971.683544	0.992718	0.905217
RM_control3	36163	551	38664	0.883533	885.603703	787.014925	0.993032	0.717632
RM_control4	38540	336	39876	0.384966	448.529403	407.90625	0.996938	0.919171
RM_control5	33366	637	35693	1.586938	874.401789	812.506329	0.992927	0.478691
RM_control6	38787	377	39998	1.193842	691.127976	618.325	0.994921	0.460157
RM_control7	33174	421	34615	1.90784	639.519629	559.125	0.994876	0.282237
RM_control8	38476	604	40815	0.605582	1702.873689	1098.161017	0.991111	0.872647
RM_control9	38193	418	40387	0.349364	1476.004523	827.296296	0.993245	0.927695
RM_control10	39055	556	41136	0.430429	1525.391993	1033.097087	0.99196	0.913746
RM_case1	38861	530	40899	0.47373	1411.786593	889.486486	0.992718	0.901358
RM_case2	37973	495	39464	0.969683	688.77094	613.106383	0.995181	0.737364
RM_case3	32766	659	34398	2.121407	1435.948065	1057.649635	0.989898	0.263954
RM_case4	38107	633	40429	0.559294	1226.25393	1100.12782	0.990737	0.882253
RM_case5	36180	553	37780	0.964259	1496.239405	983	0.99168	0.62994
RM_case6	39523	692	42178	0.492288	1817.054442	1243.338129	0.990082	0.904096
RM_case7	34778	478	36739	1.20537	766.093643	675.280992	0.993703	0.446528
RM_case8	39885	510	41821	0.325184	1492.01625	904.345794	0.992704	0.94037
RM_case9	39854	537	41768	0.576914	1407.226829	936.794393	0.992648	0.866492
RM_case10	38340	609	41200	0.486695	1252.493401	1152.128205	0.990689	0.900224

The OTU, Chao I, Shannon and Simpon data were calculated at a 3% distance. RM_case, disease group; RM_control, control group; OTUs, operational taxonomic units; ACE, abundance-based coverage estimators.




Figure 2. Venn diagram of OTUs for different group samples. The yellow area was used to represent the case group and light blue area was used to represent the control group, while dark blue was used to represent the common OTUs of these two different groups. OTUs, operational taxonomic units.




Figure 3. Simpon rarefaction of all samples. The horizontal axis was used to represent the number of sequences, the vertical axis was used to represent the Simpon index and the curves of different colors were used to represent different samples.

microbiota disturbances may further promote the development of RM (Fig. 4).

Comparison of bacterial communities in the case and control groups at the phylum level. When comparing the alpha diversity between the groups, no significant differences in diversity were observed except for a higher richness index in the case group compared with that in the control group. However, the PCoA plot based on Weighted Unifrac data displayed distinct bacterial community structures in the case and the control group. To investigate the specific changes of the microbiota in the samples from the case group, the relative abundance of taxa in the different groups was assessed. Fig. 5 presents taxonomic distributions of the predominant bacteria (relative abundance, >1% of the total sequences) at the phylum level. The bacterial flora analysis indicated that *Firmicutes* was the most predominant phylum, accounting for 79.57 and 73.12% of the microbiota in case and control group, respectively. The second most dominant phylum was *Actinobacteria* (18.86% in the case group and 22.95% in the




Figure 4. PCoA plots based on Bray-Curtis metrics for the control and case groups. The X-axis represents the PCo1 (accounting for 36.8% of the total variations) and the Y-axis represents the PCo2 (accounting for 29.9% of the total variations). Each data-point represents one sample. PCoA, principal component analysis.

control group). Changes at the phylum level were mainly in the three most common types of bacteria (*Firmicutes*, *Bacteroidetes* and *Actinobacteria*). Compared with that in the control group, the relative abundance of *Firmicutes* in the case group was increased by 5.72%, while the *Actinobacteria* and the *Bacteroidetes* were reduced by 3.08 and 3.07%, respectively.

Comparison of bacterial communities between case and control groups at the genus level. The bacterial diversity and relative abundance in all samples at the genus level are presented in Fig. 6. Fig. 6A displays the composition of the vaginal flora of the patients with RM and healthy women. In the stacked bar chart, each bar represents the average relative abundance of each bacterial taxon. The top 50 taxa with high relative abundance are illustrated. All core bacteria in the vagina, including *Lactobacillus*, *Gardnerella*, *Atopobium* and *Prevotella*, were identified in the present study. Of all the detected genera, *Lactobacillus* of the phylum *Firmicutes* were the most abundant ones. Circos was then used to identify bacterial taxa that were significantly different between the groups. A Circos presentation of bacterial taxa that are differentially represented between the different groups is provided in Fig. 6B. In the case group, 3 bacterial taxa were significantly more abundant (*Atopobium*, *Prevotella* and *Streptococcus*), while only 2 taxa were overrepresented in the control group (*Lactobacillus* and *Gardnerella*).

Functional annotation. As displayed in Fig. 7, a total of 14 KEGG pathways (e.g. 'Signal Transduction' and 'Cell Motility') were more abundant in the case group and 27 pathways (e.g. 'Metabolism of Cofactors and Vitamins' and 'Carbohydrate Metabolism') were more abundant in the control group. However, the differences in the mean proportion of the abundance of the functional pathways between the groups were not significant ($P > 0.05$). In the future, differences in the functional pathways specific for the vaginal microbial flora of the two groups will be further experimentally verified.




Figure 5. Relative abundance of the predominant bacteria at the phylum level. The predominant taxa (>1% relative abundance) is presented. (A) Distribution of community components at the phylum level in each individual sample. (B) Group merge of community composition at the phylum level in all samples. The X-axis represents the different samples and the Y-axis represents the relative abundance ratio. Different colors represent the different phyla, and the different color block heights indicate the relative abundance ratio of different species.

Discussion

Previous studies on vaginal bacterial communities suggested that the human vaginal microbiota has a key role in preventing a number of urogenital diseases, including bacterial vaginosis, yeast infections, sexually transmitted infections and urinary tract infections (12,28). However, studies indicating a direct association between RM and vaginal microbiota are currently lacking. Compared with the intestinal flora, the vaginal flora is less diverse, and women of childbearing age may have ~40 species of bacteria and facultative anaerobic bacteria, including *Lactobacillus*, *Bacteroides*, *Coccidioides*, *Corynebacterium*,

Escherichia coli, *Velveti* and *Gardneria* (29). A previous study sampled and analyzed 396 women from four ethnic groups and identified that the common constituents of the vaginal flora may be roughly divided into five categories, of which four are able to produce a large quantity of lactic acid to generate the acidic environment in the vagina (30). Bacterial vaginosis is characterized by a complete loss of lactobacilli and a concomitant increase in Gram-variable and Gram-negative rods, with *Gardnerella vaginalis*, as well as *Bacteroidetes*, *Prevotella* and *Mobiluncus* species being primary among them. The presence of an abnormal vaginal microbiota in early pregnancy is a recognized risk factor for preterm delivery and low birth weight (31).




Figure 6. Relative abundance of the predominant bacteria at the genus level. (A) Distribution of microbial community components in each sample at the genus level. The height of the bars for each genus indicates its relative abundance in the sample. (B) The data were visualized in a Circos presentation.

In the present study, the vaginal microbiome of the RM group exhibited an increased richness in species as well as a significant shift in the overall microbial diversity. A statistically significant increase in richness was observed in the RM group ($P=0.037$). These results suggested the presence of disorders in the profile of the vaginal microbiome in patients with RM and miscarriage may be either a cause or an effect of the altered composition of the vaginal microbiome. The PCo coefficients obtained in the PCoA analysis of the present study indicated that RM may be the most important factor contributing to changes in the vaginal bacterial composition. It was suggested that RM may be associated with imbalances in the microbiota. Local microbiota disturbances may further promote the development of RM (32).

When comparing the alpha diversity between samples, significant differences in diversity were observed in richness indices. Furthermore, PCoA indicated distinct bacterial community structures between the two groups. To further clarify the specific differences in the intestinal microflora between the case and control groups, taxonomic analysis at

the phylum and genus level, as well as beta diversity analysis were performed. At the phylum level, *Firmicutes* was the most predominant phylum, followed by *Actinobacteria* and then *Bacteroidetes*. Furthermore, at the genus level, *Lactobacillus* was the most dominant genus. Statistically significant differences in five genera were observed. In the case group, three bacterial taxa were significantly more abundant (*Atopobium*, *Prevotella* and *Streptococcus*), while only two taxa were overrepresented in the control group (*Lactobacillus* and *Gardnerella*). Kuon *et al* (33) reported that RM patients with elevated peripheral and uterine natural killer cells suffer more frequently from colonization by *Gardnerella vaginalis* and gram-negative anaerobes. McDonald *et al* (34) demonstrated that group B streptococcus is a key pathogen in intrauterine infection and a frequent cause of spontaneous midgestation abortions. *Lactobacillus* is one of the diverse and phylogenetically heterogeneous orders of lactic acid-producing bacteria that includes the genus *Lactobacillus*, as well as the genera *Facklamia*, *Granulicatella*, *Leuconostoc*, *Pediococcus* and *Streptococcus* (35). The healthy human vagina is dominated




Figure 7. Enriched Kyoto Encyclopedia of Genes and Genomes pathways of bacteria in the two groups. The left panel indicates the abundance ratio of the different functional pathways in the two groups, and the right panel indicates the difference in the mean proportion of the abundance of the functional pathways and the 95% confidence interval, as well as the P-values.

by a variety of *Lactobacillus* species, which have an essential role in protecting women from genital infection. A deficiency in *Lactobacilli* may disturb the microbial balance in the vagina, frequently resulting in the syndrome of bacterial vaginosis, which is associated with a quantitative and qualitative shift from normally occurring *Lactobacilli* to a mixed microflora dominated by anaerobic bacteria (31). In the present study, the three anaerobic bacteria of *Atopobium*, *Prevotella* and *Streptococcus* were more abundant in the case group compared with those in the control group, which was consistent with the results of previous studies.

In conclusion, the present study defined a structural imbalance in the vaginal microbiota of patients with RM, represented by an increased incidence of *Atopobium*, *Prevotella* and *Streptococcus* and reduction of *Lactobacillus*, which was identified by comparing the vaginal microbiota composition of RM patients with that of healthy individuals. PCoA analysis suggested that changes in vaginal flora may be the cause of/associated with RM. The present results point towards a novel strategy aimed at preventing the development of RM through restoring the homeostasis of the vaginal microbiome, by improvement in lifestyle or early intervention with drugs or probiotics.

Acknowledgements

Not applicable.

Funding

The present study was supported by the National Natural Science Foundation of China (grant no. 81701522); the National Natural Science Foundation of Zhejiang Province, China (grant nos. LY15H040001 and LY16H040008); the Science Technology Department of Zhejiang Province, China (grant no. 2015C33280, 2016C33222, 2016C33223 and 2018C37102); the Medical and Health Project of Zhejiang Province, China (grant nos. 2014KYA275, 2014KYA276, 2017KY669, 2017KY670, 2017KY672, 2018KY844, 2018KY845, 2018KY846 and 2018KY848); the Health and Family Planning Commission of Shaoxing, China (grant nos. 2017QN008 and 2017CX011); and the Science Technology Department of Shaoxing, China (grant no. 2017B70004).

Availability of data and material

The datasets used and/or analyzed during the present study are available from the corresponding author on reasonable request.

Authors' contributions

FZ, TZ and HD conceived and designed the current study. YM and ZH obtained samples; YH performed the experiments; HP collected the data; MF and HD analysed and interpreted the data. All authors approved the final version of the manuscript for publication.

Ethical approval and consent to participate

All subjects provided written informed consent for use of their samples in the present study. The present study was approved by the Ethics Committee of Shaoxing Women and Children's Hospital (Shaoxing, China) and was performed in compliance with the Declaration of Helsinki.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Sugiura-Ogasawara M, Ozaki Y and Suzumori N: Management of recurrent miscarriage. *J Obstet Gynaecol Res* 40: 1174-1179, 2014.
2. Dempsey MA, Flood K, Burke N, Murray A, Cotter B, Mullers S, Dicker P, Fletcher P, Geary M, Kenny D and Malone FD: Platelet function in patients with a history of unexplained recurrent miscarriage who subsequently miscarry again. *Eur J Obstet Gynecol Reprod Biol* 188: 61-65, 2015.
3. Christiansen OB, Larsen EC, Egerup P, Lunoe L, Egestad L and Nielsen HS: Intravenous immunoglobulin treatment for secondary recurrent miscarriage: A randomised, double-blind, placebo-controlled trial. *BJOG* 122: 500-508, 2015.
4. Obiero JA, Waititu KK, Mulei I, Omar FI, Jaoko W and Mwethera PG: Baboon vaginal microbial flora. *J Med Primatol* 45: 147-155, 2016.
5. Farr A, Kiss H, Hagmann M, Machal S, Holzer I, Kueronya V, Husslein PW and Petricevic L: Role of lactobacillus species in the intermediate vaginal flora in early pregnancy: A retrospective cohort study. *PLoS One* 10: e0144181, 2015.
6. Romero R, Chaiworapongsa T, Kuivaniemi H and Tromp G: Bacterial vaginosis, the inflammatory response and the risk of preterm birth: A role for genetic epidemiology in the prevention of preterm birth. *Am J Obstet Gynecol* 190: 1509-1519, 2004.
7. Romero HD and Andreu DA: Bacterial vaginosis. *Enfermedades Infecciosas Y Microbiol Clin* 34 (Suppl 3): 14-18, 2016.
8. Tseng CH, Lin JT, Ho HJ, Lai ZL, Wang CB, Tang SL and Wu CY: Gastric microbiota and predicted gene functions are altered after subtotal gastrectomy in patients with gastric cancer. *Sci Rep* 6: 20701, 2016.
9. NIH HMP Working Group, Peterson J, Garges S, Giovanni M, McInnes P, Wang L, Schloss JA, Bonazzi V, McEwen JE, Wetterstrand KA, *et al*: The NIH human microbiome project. *Genome Res* 19: 2317-2323, 2009.
10. Human Microbiome Jumpstart Reference Strains Consortium, Nelson KE, Weinstock GM, Highlander SK, Worley KC, Creasy HH, Wortman JR, Rusch DB, Mitreva M, Sodergren E, *et al*: A catalog of reference genomes from the human microbiome. *Science* 328: 994-999, 2010.
11. Ling Z, Liu X, Chen X, Zhu H, Nelson KE, Xia Y, Li L and Xiang C: Diversity of cervicovaginal microbiota associated with female lower genital tract infections. *Microb Ecol* 61: 704-714, 2011.
12. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket CO, *et al*: Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci USA* 108 (Suppl 1): 4680-4687, 2011.
13. Leitch H and Kiss H: Asymptomatic bacterial vaginosis and intermediate flora as risk factors for adverse pregnancy outcome. *Best Pract Res Clin Obstet Gynaecol* 21: 375-390, 2007.
14. Allsworth JE and Peipert JF: Prevalence of bacterial vaginosis: 2001-2004 National Health and Nutrition Examination Survey data. *Obstet Gynecol* 109: 114-120, 2007.
15. Trabert B and Misra DP: Risk factors for bacterial vaginosis during pregnancy among African American women. *Am J Obstet Gynecol* 197: 477.e1-e8, 2007.
16. Donders GG, Veerecken A, Bosmans E, Dekersmaecker A, Salembier G and Spitz B: Definition of a type of abnormal vaginal flora that is distinct from bacterial vaginosis: Aerobic vaginitis. *BJOG* 109: 34-43, 2002.
17. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N and Knight R: Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci USA* 108 (Suppl 1): 4516-4522, 2011.
18. Aguirre E, Galiana A, Mira A, Guardiola R, Sánchez-Guillén L, García-Pachón E, Santibañez M, Royo G and Rodríguez JC: Analysis of microbiota in stable patients with chronic obstructive pulmonary disease. *APMIS* 123: 427-432, 2015.
19. Zhang J, Kobert K, Flouri T and Stamatakis A: PEAR: A fast and accurate Illumina Paired-End reAd mergeR. *Bioinformatics* 30: 614-620, 2014.
20. Schmieder R and Edwards R: Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27: 863-864, 2011.
21. Magoč T and Salzberg SL: FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27: 2957-2963, 2011.
22. Edgar RC: Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460-2461, 2010.
23. Edgar RC, Haas BJ, Clemente JC, Quince C and Knight R: UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27: 2194-2200, 2011.
24. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, *et al*: Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75: 7537-7541, 2009.
25. Snipen L and Liland KH: micropan: An R-package for microbial pan-genomics. *BMC Bioinformatics* 16: 79, 2015.
26. Langille MG, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkepille DE, Vega Thurber RL, Knight R, *et al*: Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nat Biotechnol* 31: 814-821, 2013.

27. Parks DH, Tyson GW, Hugenholtz P and Beiko RG: STAMP: Statistical analysis of taxonomic and functional profiles. *Bioinformatics* 30: 3123-3124, 2014.
28. Hernándezrodríguez C, Romerogonzález R, Albanicampanario M, Figueroa-Damián R and Meraz-Cruz N and Hernández-Guerrero C: Vaginal microbiota of healthy pregnant mexican women is constituted by four lactobacillus species and, several vaginosis-associated bacteria. *Infect Dis Obstet Gynecol* 2011: 851485, 2011.
29. Wells CL, Jechorek RP and Maddaus MA: The translocation of intestinal facultative and anaerobic bacteria in defined flora mice. *Microb Eco Health Dis* 1: 227-235, 1988.
30. Mendes-Soares H, Suzuki H, Hickey RJ and Forney LJ: Comparative functional genomics of *Lactobacillus* spp. reveals possible mechanisms for specialization of vaginal lactobacilli to their environment. *J Bacteriol* 196: 1458-1470, 2014.
31. Petricevic L, Domig KJ, Nierscher FJ, Sandhofer MJ, Fidesser M, Krondorfer I, Husslein P, Kneifel W and Kiss H: Characterisation of the vaginal *Lactobacillus* microbiota associated with preterm delivery. *Sci Rep* 4: 5136, 2014.
32. Lu Y, Chen J, Zheng J, Hu G, Wang J, Huang C, Lou L, Wang X and Zeng Y: Mucosal adherent bacterial dysbiosis in patients with colorectal adenomas. *Sci Rep* 6: 26337, 2016.
33. Kuon RJ, Togawa R, Vomstein K, Weber M, Goeggel T, Strowitzki T, Markert UR, Zimmermann S, Daniel V, Dalpke AH and Toth B: Higher prevalence of colonization with *gardnerella vaginalis* and gram-negative anaerobes in patients with recurrent miscarriage and elevated peripheral natural killer cells. *J Reprod Immunol* 120, 15-19, 2017.
34. McDonald HM and Chambers HM: Intrauterine infection and spontaneous midgestation abortion: Is the spectrum of microorganisms similar to that in preterm labor? *Infect Dis Obstet Gynecol* 8: 220-227, 2000.
35. Goldstein EJ, Tyrrell KL and Citron DM: *Lactobacillus* species: Taxonomic complexity and controversial susceptibilities. *Clin Infect Dis* 60 (Suppl 2): S98-S107, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.