

MicroRNA-138 attenuates myocardial ischemia reperfusion injury through inhibiting mitochondria-mediated apoptosis by targeting HIF1- α

YAN LIU¹, JIANFENG ZOU², XIAOYAN LIU¹ and QUAN ZHANG³

¹The First Ward; ²The Third Ward, Department of Cardiology, Rizhao People's Hospital, Rizhao, Shandong 276800;

³Department of Cardiology, Affiliated Hospital of Weifang Medical University, Weifang, Shandong 260141, P.R. China

Received July 24, 2018; Accepted March 26, 2019

DOI: 10.3892/etm.2019.7976

Abstract. Myocardial ischemia-reperfusion (I/R) injury is considered to have a detrimental role in coronary heart disease, which is considered to be the leading cause of death worldwide. However, the molecular mechanism involved in the progression of myocardial I/R injury is still unclear. The current study aimed to investigate the expression and function of microRNA (miR)-138 in the process of myocardial I/R injury. First, miR-138 expression levels were analyzed both in myocardium with I/R injury and control myocardium using reverse transcription-quantitative polymerase chain reaction analysis. Then, the relationship between the levels of miR-138 and hypoxia-inducible factor (HIF)1- α was also investigated using a luciferase reporter assay. Assessment of myocardial infarct size, measurements of serum myocardial enzymes and electron microscopy analysis were all utilized to analyse the effect of miR-138 on myocardial I/R injury. The authors of current study also used western blotting to examine the expression levels of the mitochondrial fission-related proteins dynamin-1-like protein and mitochondrial fission 1 protein. It was found that miR-138 is downregulated and HIF1- α is upregulated after myocardial ischemia reperfusion injury. Overexpression of miR-138 reduced myocardial I/R injury-induced infarct sizes and myocardial enzyme levels, and it also inhibited the expression of proteins related to mitochondrial morphology and myocardial I/R-induced mitochondrial apoptosis by targeting HIF1- α . Taken together, these findings provide a novel insight into the molecular mechanism of miR-138 and HIF1- α in the progression of myocardial

I/R injury. miR-138 has the potential to become a promising therapeutic target for treating myocardial I/R injury.

Introduction

Myocardial ischemia-reperfusion (I/R) injury is considered to have a detrimental role in coronary heart disease (CHD), which is considered to be the leading cause of death worldwide (1-3). Myocardial infarction is caused by coronary occlusion and the subsequent insufficient supply of blood to the myocardium, which may then cause irreversible necrosis to occur (4,5). Restoring blood flow to the ischemic myocardium is the most effective treatment to rescue ischemic myocardium cells and to save the life of the patient (6). However, this treatment can cause an abrupt restoration of the oxygen supply and the reperfusion of myocardium may aggravate myocardial injury, leading to a reperfusion injury (7,8). Therefore, it is necessary to explore and to better understand the potential molecular mechanism of myocardial I/R injury.

MicroRNAs (miRNAs or miRs) are small, single-stranded, non-coding RNAs (20-22 nucleotides) that are involved in numerous biological processes (9). miRNAs participate in many biological processes, such as cell apoptosis, cell differentiation and cell development (10). More and more evidence has indicated that several miRNAs are expressed abnormally during myocardial I/R injury, suggesting the involvement of miRNAs in myocardial I/R injury development (11-13). Furthermore, the inhibition of mitochondria-mediated apoptosis to reduce cardiomyocyte apoptosis is considered to be an important mechanism (14,15). However, little evidence is available regarding the role of miRNAs and mitochondria-mediated apoptosis in myocardial I/R injury.

Hypoxia-inducible factor (HIF) 1 is known as a heterodimeric transcription factor composed of an oxygen-labile α subunit (HIF1- α) and a constitutive β subunit (HIF1- β), and HIF1 can bind to the hypoxia response element to regulate gene expression (16,17). Furthermore, HIF1- α is the regulatory subunit that senses tissue oxygen level, responds to various types of cellular stimulation and exerts a vast array of physiological functions, enabling cells to adapt to temporary hypoxia (18). Some studies found that HIF1- α activity showed some effects in preventing diabetic cardiomyopathy

Correspondence to: Professor Quan Zhang, Department of Cardiology, Affiliated Hospital of Weifang Medical University, 2428 Yuhe Road, Weifang, Shandong 260141, P.R. China
E-mail: quan00208dia@126.com

Key words: myocardial ischemia-reperfusion injury, coronary heart disease, microRNA-138, mitochondria-mediated apoptosis, hypoxia-inducible factor 1- α

and cardiac remodeling (19,20). However, the role of HIF1- α in myocardial I/R injury is incompletely understood.

The aim of the current study was to investigate the effect of miRNAs and mitochondria-mediated apoptosis in a myocardial I/R injury model. In the current study, the authors uncovered the pivotal role of miR-138 in the myocardial I/R injury model, which may be connected with the inhibition of myocardial I/R-induced mitochondrial apoptosis.

Materials and methods

Ethical statement. The experiments involving animals conform to local and national Guide for the Care and Use of Laboratory Animal guidelines. Furthermore, these experimental methods are also approved by the Clinical Ethics Committee of Affiliated Hospital of Weifang Medical University (Shandong, China).

Mouse model of myocardial I/R Injury. A total of 60 mice (30 males and 30 females; age, 8-10 weeks; weight, 18-25 g) were obtained from the Animal Center of Weifang Medical University (Weifang, China). Animals were housed at a temperature of 23-25°C and a relative humidity of 40-60%, under a 14/10 h light/dark cycle and free access to water and food. Housing conditions also included a light intensity of ~40 lux at the position of the animal in cage. Mice were divided into five groups (each, n=12): A sham group (mice without injury), a control group (I/R injury with miR-138 negative control injection), an I/R injury group, a miR-138 mimic group (I/R injury with miR-138 mimic injection) and a miR-138 inhibitor group (I/R injury with miR-138 inhibitor injection). The mouse model of myocardial I/R injury was established as previously described (21). In short, pentobarbital (50 mg/kg) was used to anesthetize the mice. Then, a left horizontal incision was made at the third intercostal space. Subsequently, a silk suture we used to tie around the left anterior descending artery and a silicon tube (1 mm outside diameter). In the sham group, the mouse only received the identical surgical procedure without ligation. The silicon tube was then removed to achieve reperfusion for 6 h after 30 min of ischemia. After reperfusion, the heart samples were collected as quickly as possible. For some experiments, miR-138 mimic (20 nmol/l), miR-138 inhibitor (20 nmol/l), HIF1- α siRNA (100 nmol/l) or miR-138 negative control (20 nmol/l) were administered intraperitoneally for seven days before modeling. miR-138 mimic, miR-138 inhibitor, HIF1- α siRNA were all purchased from Genscript (Piscataway, NJ, USA). The miR-138 mimic sequence was 5'-AGCUGGUGUUGUGAAUCAGGCCG-3', the miR-138 inhibitor sequence was 5'-CGGCCUGAUUCACAACACCAGCU-3', the HIF1- α siRNA sequence was 5'-AUCCAGAGU CACUGGAACU-3' and the NC sequence was 5'-UGAAUC CUUGAAUAGGUGUGUU-3'. Carbon dioxide was used to euthanize the mice with a fill rate of 10-30% of the volume of the cage per minute. The authors of the current study followed the American Veterinary Medical Association guidelines' recommendations to confirm the death of the mice, including heartbeat, breathing, corneal reflex and responses to firm toe pinch, graying of the mucous membranes and rigor mortis.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) analysis. RNA from heart samples from each

group was extracted using the TRIzol® reagent (Invitrogen). Subsequently, the RNA was reverse transcribed into cDNA using the cDNA Reverse Transcription kit (Applied Biosystems; Thermo Fisher Scientific, Inc., Waltham, MA, USA), and we perform RT-qPCR with SYBR Premix Ex Taq (Takara Biotechnology Co., Ltd.). GAPDH was used as the internal normalized reference. The relative levels of miR-138 and HIF1- α were calculated using the $2^{-\Delta\Delta C_q}$ method (22). The thermocycling conditions were as follows: 45°C for 10 min and one cycle of 95°C for 10 min, followed by 40 cycles at 95°C for 15 sec and 60°C for 45 sec. The primers sequences were as follows: miR-138: Forward, 5'-TCCGAGCCTGACTAAGTG TTGTGGTCTGA-3' and reverse, 5'-GTGCAGGGTCCGAGG T-3'; GAPDH: Forward, 5'-TGGTATCGTGGAAGGACTC-3' and reverse 5'-AGTAGAGGCAGGGATGATG-3'; HIF1- α : Forward, 5'-CTCAGCCCCAGTGCATTGTA-3' and reverse 5'-GAACCTCCTATAGCCACCGC-3'.

Western blot analysis. Myocardial tissues from each group were lysed using radioimmunoprecipitation assay buffer. Subsequently, a bicinchoninic acid protein assay to detect the concentration of protein. Protein lysates (20 μ l) were then separated using 10% SDS-PAGE and transferred to polyvinylidene fluoride membranes (EMD Millipore, Billerica, MA, USA). Membranes were then blocked with 5% skim milk in TBST for 1 h at room temperature. The membranes were incubated with the following antibodies obtained from Abcam (Cambridge, UK): HIF1- α (1:2,000; cat. no. ab51608), cleaved caspase-9 (1:2,000; cat. no. ab2324), cleaved caspase-3 (1:2,000; cat. no. ab2302), Drp1 (1:2,000; cat. no. ab184247), Fis1 (1:2,000; cat. no. ab71498) and GAPDH (1:5,000; Santa Cruz Biotechnology, Inc.) at 4°C overnight. Then, they were washed using TBST and incubated with secondary antibodies (horseradish peroxidase conjugated Goat Anti-Rabbit IgG; 1:5,000; cat. no. ab205718, Abcam) at 4°C for 1 h. ECL Western blotting substrate (Pierce; Thermo Fisher Scientific, Inc.) was used to visualize and detect the results. ImageJ software 1.43v (National Institutes of Health, Bethesda, MD, USA) was used to quantify protein levels.

Assessment of myocardial infarct size. After sacrificing the mice in each group, Evans blue dye was used to demarcate the ischemic area-at-risk. Subsequently, heart tissues were excised and sliced (50 μ m). These samples were stained with 1% triphenyltetrazolium chloride at 37°C for 20 min and then fixed using 4% paraformaldehyde for 8 h at 4°C. Infarcted myocardium was separated from the non-infarcted myocardium and weighed carefully. Heart tissue from each group were dissected and weighed. The infarct size was presented as a percentage of the total ischemic area.

Measurement of serum myocardial enzymes. After 6 h of reperfusion, blood samples from each group were collected, centrifuged at 2,000 \times g for 30 min at 4°C and transferred to Eppendorf tubes. Subsequently, ELISA was performed to measure the serum levels of troponin I (cat. no. BEK-2212-1P/2P; Biosensis), cardiac muscle (cTn I; cat. no. EKC40439, R&D Systems Inc.) and creatine kinase M-type/B-type (CK-MB; cat. no. ABIN415661; R&D Systems Inc.) according to the manufacturer's protocol.




Figure 1. Expression of miR-138 and HIF1- α in the myocardium with I/R injury. (A) miR-138 mRNA expression measured by RT-qPCR. (B) HIF1- α mRNA expression measured by RT-qPCR. (C) Western blotting and (D) densitometric quantification of proteins. (***) $P < 0.001$. HIF1- α , hypoxia-inducible factor 1- α ; miR, microRNA; I/R, ischemia-reperfusion; RT-qPCR, reverse transcription-quantitative polymerase chain reaction.

Electron microscopy analysis. The method of electron microscopy analysis was described as previously (23). The electron microscopy images of mitochondria were analyzed using PhotoshopCS5.0 software (Adobe, Inc.). For the analysis, the number of the myocardial mitochondria in each group were measured.

Luciferase reporter assay. TargetScan (www.targetscan.org/) was used to predict the target gene of miR-138. Following, a wild-type (WT) 3'-untranslated region (UTR) fragment of HIF1- α containing the putative miR-138 binding sequence was inserted into a pmirGLO Dual-luciferase miRNA Target Expression Vector (Promega Corporation, Madison, WI, USA), while mutant (MUT) 3'-UTR was also cloned into the vector to generate a mutated binding site. Subsequently, the cells obtained from the American Type Culture Collection were co-transfected with HIF1- α -WT or HIF1- α -MUT and miR-138 mimics (20 nmol/l) using Lipofectamine™ 2000 (Thermo Fisher Scientific, Inc.). Dual Luciferase reporter assay system (DLR® Assay, Promega Corporation) was used to evaluate the luciferase activity after 48 h. Renilla Luciferase was used as a normalizing transfection control.

Statistical analysis. GraphPad Prism 4 software was used to analyze the experimental data. The data were presented as mean \pm standard deviation. Statistical analyses were performed using Student's t-test and one-way analysis of variance, followed by a Tukey's post-hoc test. $P < 0.05$ indicated that the difference between groups was statistically significant.

Results

miR-138 is downregulated and HIF1- α is upregulated after myocardial I/R injury. To identify whether miR-138 performed important functions in myocardial I/R injury, its expression level in adult mouse myocardium with and without I/R injury was examined. The results indicated that the miR-138 expression level was significantly downregulated in the myocardium of the myocardial I/R injury model compared with the control myocardium (Fig. 1A). Moreover, the expression level of HIF1- α was significantly upregulated in the myocardium of the myocardial I/R injury model compared with the control myocardium (Fig. 1B). The western blotting results also confirmed the alteration of HIF1- α expression level as HIF1- α protein expression significantly upregulated in the myocardium of the I/R injury group compared with the control group (Fig. 1C and D).

Overexpression of miR-138 reduces the myocardial I/R-induced increase in infarct size and myocardial enzymes. As mentioned previously, compared with the control group, the miR-138 expression level was significantly decreased in the myocardial I/R injury group. Subsequently, to further explore the role of miR-138 in the myocardial I/R injury model, miR-138 mimic, miR-138 inhibitor or a negative control were administered intraperitoneally before modeling. Compared with the I/R injury group, the infarct size of, and serum CTn I and CK-MB levels in the myocardium with I/R injury were significantly decreased in the miR-138 mimic group and increased in the miR-138 inhibitor group (Fig. 2).




Figure 2. miR-138 influences the infarct size and myocardial enzymes in the myocardium with I/R injury. Effect of miR-138 on (A) infarct size, (B) cTn I concentration and (C) CK-MB activity. (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). miR, microRNA; I/R, ischemia-reperfusion; cTn I, Troponin I, cardiac muscle; CK-MB, creatine kinase M-type/B-type.




Figure 3. miR-138 inhibited the expression of cleaved caspase-9 and -3 in myocardial I/R injury. Effect of miR-138 on (A) the number of mitochondria and (B) the expression of cleaved caspase-3 and caspase-9 in the myocardium. Densitometric quantification of (C) caspase-3 and (D) caspase-9 in the myocardium. (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). miR, microRNA; I/R, ischemia-reperfusion.

miR-138 inhibits the expression of cleaved caspase-9 and -3 in myocardial I/R injury. The number of the mitochondria in the myocardial I/R injury group per area was significantly decreased compared with the sham group. However, in the miR-138 mimic group, the number of mitochondria per area were increased significantly, while in the miR-138 inhibitor group, the number of mitochondria was significantly reduced

compared with the I/R injury group (Fig. 3A). Notably, the results of the western blot analysis indicated that the expression levels of cleaved caspase-9 and -3 were significantly increased in the myocardial I/R injury group compared with sham mice, and after the injection of the miR-138 mimic, expression levels were decreased compared with the I/R injury group (Fig. 3B-D). In the miR-138 inhibitor group, cleaved caspase-9




Figure 4. miR-138 inhibited the expression of proteins related to mitochondrial morphology (A) Effect of miR-138 on the expression of Drp1 and Fis1 in the myocardium. Densitometric quantification of (B) Drp1 and (C) Fis1 in the myocardium. (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). miR, microRNA; Drp1, dynamin-1-like protein; Fis1, mitochondrial fission 1 protein.




Figure 5. miR-138 directly targeted the 3'-UTR of HIF1- α mRNA. (A) Schematic representation of the mature miR-138 sequence, putative miR-138 target site in the 3'-UTR of HIF1- α mRNA. (B) The relative luciferase activity in the WT 3'-UTR of HIF1- α mRNA and MUT 3'-UTR of HIF1- α after miR-138 overexpression. (C) HIF1- α protein expression after miR-138 mimic transfection. (** $P < 0.01$, *** $P < 0.001$). miR, microRNA; UTR, untranslated region; HIF1- α , hypoxia-inducible factor 1- α ; WT, wild-type; MUT, mutated.

and -3 expression levels were significantly increased reduced compared with the I/R injury group.

miR-138 inhibits the expression of proteins related to mitochondrial morphology. To explore the mechanism of miR-138 in relation to morphological alterations of mitochondria in myocardial I/R injury, the expression levels of the mitochondrial fission-related proteins, Drp1 and Fis1 (24,25), were examined using western blot analysis. The results indicated that the expression levels of Drp1 and Fis1 in the myocardial I/R injury group were both increased compared with the sham group (Fig. 4). When treated with miR-138 mimic, however, Drp1 and Fis1 expression levels were significantly decreased, while their expression levels were both increased in the miR-138 inhibitor group compared with the I/R injury group.

HIF1- α is the target of miR-138. The bioinformatic results were confirmed by a luciferase reporter assay. Fig. 5A shows the predicted miR-138 binding sequence in HIF1- α . The luciferase activity of the construct with the WT 3'-UTR was significantly inhibited after transfection with the miR-138 mimic (Fig. 5B). Western blotting results indicated that miR-138 mimic transfection markedly decreased the expression of HIF1- α compared with NC mimic transfection (Fig. 5C).

Downregulation of HIF1- α inhibits the expression of cleaved caspase-9 and caspase-3 in myocardial I/R injury. To explore the role of HIF1- α in the model of myocardial I/R injury, siRNA of HIF1- α were administered intraperitoneally before modeling. HIF1- α siRNA was shown to significantly decrease the expression of HIF1- α in the myocardium compared with the control group (Fig. 6A). The number of mitochondria per




Figure 6. Downregulation of HIF1- α inhibited myocardial I/R-induced mitochondrial apoptosis. (A) HIF1- α mRNA expression measured by RT-qPCR. (B) Effect of HIF1- α siRNA on the number of mitochondria and (C) the expression of cleaved caspase-3 and caspase-9 in the myocardium. Densitometric quantification of (D) caspase-3 and (E) caspase-9 in the myocardium. (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). miR, microRNA; siRNA, small interfering RNA; I/R, ischemia-reperfusion; HIF1- α , hypoxia-inducible factor 1- α ; RT-qPCR, reverse transcription-quantitative polymerase chain reaction.

area was increased significantly in the mouse myocardium of the HIF1- α siRNA group compared with the I/R injury group (Fig. 6B). Notably, western blotting results indicated that the protein expression levels of cleaved caspase-9 and -3 were decreased significantly in the myocardium after HIF1- α siRNA administration compared with the I/R injury group (Fig. 6C-E).

Discussion

CHD is considered to be the leading cause of death worldwide, with approximately 17.5 million people dying because of cardiovascular disease according to the estimates from the World Health Organization (26-28). Acute myocardial I/R usually causes the detrimental effects of CHD (29). Therefore, to explore the molecular mechanism involved in myocardial I/R injury, the authors of the current study conducted functional experiments and identified the critical role of miR-138 and mitochondria-mediated apoptosis in the myocardial I/R injury model. The results indicated that low expression levels of miR-138 were found in the myocardium with I/R injury compared with that of control myocardium. Furthermore, the level of HIF1- α was significantly upregulated in the myocardium of the myocardial I/R injury model compared with control myocardium. Overexpression of miR-138 reduced the myocardial I/R-induced increase in infarct size and myocardial enzymes by targeting HIF1- α to inhibit myocardial I/R-induced mitochondrial apoptosis. These findings suggest that miR-138 may prevent damage of the myocardium after I/R injury.

miRNAs are a type of small non-coding RNAs that bind to their target mRNAs specifically, and subsequently cause the downregulation of the target gene by repressing degradation

or translation (30). In recent years, miRNAs have been implicated in many processes related to the heart, such as cardiac hypertrophy, cardiac development and heart failure (31). miRNAs function as endogenous intracellular regulators of mRNA translation, but the significance of miR-138 in the process of myocardial I/R injury has not been reported previously, particularly its role in myocardial I/R-induced mitochondrial apoptosis. In the current study, the results indicated that miR-138 was downregulated in myocardium with I/R injury compared with control myocardium. The results also showed that the infarct size and serum CTn I and CK-MB levels of the myocardium with I/R injury were significantly decreased in the miR-138 mimic group, but were significantly increased in the miR-138 inhibitor group compared with the control group.

Apoptosis is known as a critical pathological process in the course of myocardial I/R injury and the amount of apoptosis determines the severity of myocardial I/R injury (32,33). Therefore, understanding the mechanism involved in cardiomyocyte apoptosis in myocardial I/R injury is critically important in the development of effective treatment methods. Prior research has shown that myocardial I/R-induced mitochondrial dysfunction and apoptosis are responsible for the exacerbation of cardiac ischemic injury in diabetic patients (15). In the present study, the authors found that in the miR-138 mimic group, the mitochondria per area were significantly increased, while the number of mitochondria was significantly decreased in the miR-138 inhibitor group. Furthermore, the expression levels of cleaved caspase-9 and -3 were both significantly increased in the myocardial I/R injury model compared with control mice. Additionally, after injection of the miR-138 mimic, their expression levels were

decreased. After treatment with the miR-138 mimic, protein expression levels of Drp1 and Fis1, proteins related to mitochondrial morphology, were both significantly decreased, while their expression levels were increased in the miR-138 inhibitor group. Moreover, HIF1- α was confirmed as the target of miR-138. Therefore, the aforementioned results showed that miR-138 might lessen myocardial ischemia reperfusion injury by inhibiting mitochondria-mediated apoptosis, due to its associated targeting of HIF1- α .

There are some limitations in the current study. The mitochondrial membrane potential generated by proton pumps is an essential component in the process of energy storage during oxidative phosphorylation, and it is associated with cells' capacity to generate ATP by oxidative phosphorylation (34,35). Several fluorescent lipophilic cationic dyes (including, tetramethylrhodamine methyl ester and tetramethylrhodamine ethyl ester, Rhodamine 123, 3,3'-dihexyloxacarbocyanine iodide and 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide) have become important tools for directly measuring the mitochondrial membrane potential (36). Furthermore, apoptosis inducing factor (AIF), such as Cyt-C, Smac and Apaf-1, are proteins that trigger chromatin condensation and DNA fragmentation in a cell in order to induce programmed cell death (37,38). The mitochondrial AIF protein was found to be a caspase-independent death effector that can allow independent nuclei to undergo apoptotic changes (39). Therefore, future studies should detect the change of mitochondrial membrane potential and apoptosis factors to detect mitochondrial apoptosis.

This study indicated that overexpression of miR-138 reduces the myocardial I/R-induced increase in infarct size and myocardial enzymes by targeting HIF1- α to inhibit myocardial I/R-induced mitochondrial apoptosis. These results demonstrated a cardioprotective effect of miR-138 and suggested the potential to become a promising target to alleviate myocardial I/R injury.

Acknowledgements

Not applicable.

Funding

The present study was supported by Shandong Province Natural Science Foundation (grant no. ZR2015HL011) and Shandong Medical and Health Technology development Project (grant no. 2016WS0687).

Availability of data and materials

The datasets used and/or analyzed during the present study are available from the corresponding author on reasonable request.

Authors' contributions

QZ conceptualized the idea; YL and JFZ performed the experiments; YL and XYL searched the literature; YL, XYL and JFZ analyzed the data; YL and JFZ designed and made the figures; YL created the tables; YL and QZ wrote the manuscript. QZ reviewed the paper.

Ethics approval and consent to participate

This current study was approved by the Clinical Ethics Committee of Affiliated Hospital of Weifang Medical University (Shandong, China).

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Nichols M, Townsend N, Scarborough P and Rayner M: Cardiovascular disease in europe 2014: Epidemiological update. *Eur Heart J* 35: 2950-2959, 2014.
- Cuevas P, Carceller F and Gimenez-Gallego G: Fibroblast growth factors in myocardial ischemia/reperfusion injury and ischemic preconditioning. *J Cell Mol Med* 5: 132-142, 2001.
- Song CL, Liu B, Wang JP, Zhang BL, Zhang JC, Zhao LY, Shi YF, Li YX, Wang G, Diao HY, *et al*: Anti-apoptotic effect of microRNA-30b in early phase of rat myocardial ischemia-reperfusion injury model. *J Cell Biochem* 116: 2610-2619, 2015.
- Jennings RB and Reimer KA: The cell biology of acute myocardial ischemia. *Annu Rev Med* 42: 225-246, 1991.
- Mu F, Duan J, Bian H, Yin Y, Zhu Y, Wei G, Guan Y, Wang Y, Guo C, Wen A, *et al*: Cardioprotective effects and mechanism of radix salviae miltiorrhizae and lignum dalbergiae odoriferae on rat myocardial ischemia/reperfusion injury. *Mol Med Rep* 16: 1759-1770, 2017.
- Yellon DM and Hausenloy DJ: Myocardial reperfusion injury. *New Engl J Med* 357: 1121-1135, 2007.
- Prompt E, Sanit J, Barrere-Lemaire S, Nargeot J, Noordali H, Madhani M and Kumphune S: The cardioprotective effects of secretory leukocyte protease inhibitor against myocardial ischemia/reperfusion injury. *Exp Ther Med* 15: 5231-5242, 2018.
- Li X, Liu M, Sun R, Zeng Y, Chen S and Zhang P: Protective approaches against myocardial ischemia reperfusion injury. *Experimental and therapeutic medicine* 12: 3823-3829, 2016.
- Shi Z, Zhou H, Lu L, Li X, Fu Z, Liu J, Kang Y, Wei Z, Pan B, Liu L, *et al*: The roles of microRNAs in spinal cord injury. *Int J Neurosci* 127: 1104-1115, 2017.
- Chen C, Zhou Y, Wang J, Yan Y, Peng L and Qiu W: Dysregulated microRNA involvement in multiple sclerosis by induction of T helper 17 cell differentiation. *Front Immunol* 9: 1256, 2018.
- Yang J, Chen L, Yang J, Ding J, Li S, Wu H, Zhang J, Fan Z, Dong W and Li X: MicroRNA-22 targeting CBP protects against myocardial ischemia-reperfusion injury through anti-apoptosis in rats. *Mol Biol Rep* 41: 555-561, 2014.
- Zhang SB, Liu TJ, Pu GH, Li BY, Gao XZ and Han XL: MicroRNA-374 exerts protective effects by inhibiting SP1 through activating the PI3K/Akt pathway in rat models of myocardial ischemia-reperfusion after sevoflurane preconditioning. *Cell Physiol Biochem* 46: 1455-1470, 2018.
- Zhu ZD, Ye JY, Niu H, Ma YM, Fu XM, Xia ZH and Zhang X: Effects of microRNA-292-5p on myocardial ischemia-reperfusion injury through the peroxisome proliferator-activated receptor- α /gamma signaling pathway. *Gene Ther* 25: 234-248, 2018.
- Huang LH, Li J, Gu JP, Qu MX, Yu J and Wang ZY: Butorphanol attenuates myocardial ischemia reperfusion injury through inhibiting mitochondria-mediated apoptosis in mice. *Eur Rev Med Pharmacol Sci* 22: 1819-1824, 2018.
- Wu Y, Leng Y, Meng Q, Xue R, Zhao B, Zhan L and Xia Z: Suppression of excessive histone deacetylases activity in diabetic hearts attenuates myocardial ischemia/reperfusion injury via mitochondria apoptosis pathway. *J Diabetes Res* 2017: 8208065, 2017.
- Liu J, Zhang C, Zhao Y, Yue X, Wu H, Huang S, Chen J, Tomskey K, Xie H, Khella CA, *et al*: Parkin targets HIF-1 α for ubiquitination and degradation to inhibit breast tumor progression. *Nat Commun* 8: 1823, 2017.

17. Semenza GL, Shimoda LA and Prabhakar NR: Regulation of gene expression by HIF-1. *Novartis Found Symp* 272: 2-8, 2006.
18. Semenza GL, Jiang BH, Leung SW, Passantino R, Concordet JP, Maire P and Giallongo A: Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1. *J Biol Chem* 271: 32529-32537, 1996.
19. Chen JX and Stinnett A: Ang-1 gene therapy inhibits hypoxia-inducible factor-1 α (HIF-1 α)-prolyl-4-hydroxylase-2, stabilizes HIF-1 α expression, and normalizes immature vasculature in db/db mice. *Diabetes* 57: 3335-3343, 2008.
20. Xue W, Cai L, Tan Y, Thistlethwaite P, Kang YJ, Li X and Feng W: Cardiac-specific overexpression of HIF-1 α prevents deterioration of glycolytic pathway and cardiac remodeling in streptozotocin-induced diabetic mice. *Am J Pathol* 177: 97-105, 2010.
21. Elrod JW, Calvert JW, Morrison J, Doeller JE, Kraus DW, Tao L, Jiao X, Scalia R, Kiss L, Szabo C, *et al*: Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function. *Proc Natl Acad Sci USA* 104: 15560-15565, 2007.
22. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
23. Gupta S and Knowlton AA: HSP60, Bax, apoptosis and the heart. *J Cell Mol Med* 9: 51-58, 2005.
24. Frank S, Gaume B, Bergmann-Leitner ES, Leitner WW, Robert EG, Catez F, Smith CL and Youle RJ: The role of dynamin-related protein 1, a mediator of mitochondrial fission, in apoptosis. *Dev Cell* 1: 515-525, 2001.
25. Perrelli MG, Pagliaro P and Penna C: Ischemia/reperfusion injury and cardioprotective mechanisms: Role of mitochondria and reactive oxygen species. *World J Cardiol* 3: 186-200, 2011.
26. Zhang W, Li Y and Wang P: Long non-coding RNA-ROR aggravates myocardial ischemia/reperfusion injury. *Braz J Med Biol Res* 51: e6555, 2018.
27. Arora M, Kaul D and Sharma YP: Human coronary heart disease: Importance of blood cellular miR-2909 RNomics. *Mol Cell Biochem* 392: 49-63, 2014.
28. Venardos KM, Zatta AJ, Marshall T, Ritchie R and Kaye DM: Reduced L-arginine transport contributes to the pathogenesis of myocardial ischemia-reperfusion injury. *J Cell Biochem* 108: 156-168, 2009.
29. Hausenloy DJ and Yellon DM: Myocardial ischemia-reperfusion injury: A neglected therapeutic target. *J Clin Invest* 123: 92-100, 2013.
30. Esteller M: Non-coding RNAs in human disease. *Nat Rev Genet* 12: 861-874, 2011.
31. Cheng J, Wu Q, Lv R, Huang L, Xu B, Wang X, Chen A and He F: MicroRNA-449a inhibition protects H9C2 cells against hypoxia/reoxygenation-induced injury by targeting the notch-1 signaling pathway. *Cell Physiol Biochem* 46: 2587-2600, 2018.
32. Konstantinidis K, Whelan RS and Kitsis RN: Mechanisms of cell death in heart disease. *Arterioscler Thromb Vasc Biol* 32: 1552-1562, 2012.
33. Geng YJ: Molecular mechanisms for cardiovascular stem cell apoptosis and growth in the hearts with atherosclerotic coronary disease and ischemic heart failure. *Ann N Y Acad Sci* 1010: 687-697, 2003.
34. Basheer WA, Fu Y, Shimura D, Xiao S, Agvastian S, Hernandez DM, Hitzeman TC, Hong T and Shaw RM: Stress response protein GJA1-20k promotes mitochondrial biogenesis, metabolic quiescence, and cardioprotection against ischemia/reperfusion injury. *JCI Insight* 3, 2018.
35. Kadenbach B, Ramzan R, Moosdorf R and Vogt S: The role of mitochondrial membrane potential in ischemic heart failure. *Mitochondrion* 11: 700-706, 2011.
36. Solaini G, Sgarbi G, Lenaz G and Baracca A: Evaluating mitochondrial membrane potential in cells. *Biosci Rep* 27: 11-21, 2007.
37. Muzaffar S and Chattoo BB: Apoptosis-inducing factor (Aif1) mediates anacardic acid-induced apoptosis in *Saccharomyces cerevisiae*. *Apoptosis* 22: 463-474, 2017.
38. Zhang X, Chen J, Graham SH, Du L, Kochanek PM, Draviam R, Guo F, Nathaniel PD, Szabó C, Watkins SC and Clark RS: Intracellular localization of apoptosis-inducing factor (AIF) and large scale DNA fragmentation after traumatic brain injury in rats and in neuronal cultures exposed to peroxynitrite. *J Neurochem* 82: 181-191, 2002.
39. Joza N, Susin SA, Daugas E, Stanford WL, Cho SK, Li CY, Sasaki T, Elia AJ, Cheng HY, Ravagnan L, *et al*: Essential role of the mitochondrial apoptosis-inducing factor in programmed cell death. *Nature* 410: 549-554, 2001.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.