Hydrogen sulfide modulates high glucose-induced NLRP3 inflammasome activation in 3T3-L1 adipocytes

TIAN-XIAO HU^{1*}, NING-NING ZHANG^{2*}, YUN RUAN¹, QING-YING TAN¹ and JING WANG¹

¹Department of Endocrinology, Chinese People's Liberation Army 903 Hospital, Hangzhou, Zhejiang 310013;

²Department of Physiology, Naval Medical University, Shanghai 200433, P.R. China

Received March 5, 2019; Accepted August 23, 2019

DOI: 10.3892/etm.2019.8242

Abstract. Activation of the NACHT leucine rich repeat and pyd domains-containing 3 (NLRP3) inflammasome plays an important role in the initiation of inflammation in adipose tissue in diabetic patients. However, the mechanisms underlying this are not fully understood. Hydrogen sulfide (H₂S) has been shown to have anti-inflammatory properties in various cell types. The present study aimed to investigate the effect of H₂S on high glucose (HG)-induced NLRP3 inflammasome activation in adipocytes. Adipocytes were differentiated from 3T3-L1 cells and treated with low glucose (LG), HG, H₂S donor sodium hydrosulfide (NaHS) or N-acetyl-tyrosyl-valyl-alanyl-aspartyl chloromethyl ketone, an inhibitor of the cysteine protease caspase-1. The expression levels of NLRP3, apoptosis-associated speck-like protein containing A CARD (ASC) and caspase-1, and the release of interleukin (IL)-1 β and IL-18 were measured. The results of the present study indicated that HG increased the expression levels of NLRP3, ASC and cleaved caspase-1, and the release of IL-1 β and IL-18 in adipocytes. Caspase-1 inhibition abolished HG-induced production of IL-1 β and IL-18 in adipocytes. Furthermore, NaHS inhibited the expression of NLRP3, ASC and cleaved caspase-1, and the production of IL-1 β and IL-18 in adipocytes treated with HG. In conclusion, HG may increase and exogenous H₂S may inhibit HG-induced NLRP3 inflammasome activation in adipocytes.

Introduction

Type 2 diabetes mellitus (T2DM) presents a major and growing health problem throughout the world. Microvascular and macrovascular complications often occur in T2DM, and chronic low-grade inflammation plays a central role in this process (1-3). The inflammatory response is associated with both insulin resistance and the development of vascular complications in T2DM patients (4,5). T2DM patients with higher body mass index (BMI) and a large amount of adipose tissue are considered to be at greater risk of developing more severe insulin resistance (6,7) and cardiovascular disease (8,9).

Adipose tissue is regarded as an important endocrine organ and is known to be involved in regulating inflammation (10-12). Studies have shown that the NACHT leucine rich repeat and pyd domains-containing 3 (NLRP3) inflammasome plays a role in the initiation of inflammation in adipose tissue (3,13). A wide range of pathogens and cellular damage can activate the NLRP3 inflammasome, and its activation leads to caspase-1 activation and the release of the inflammatory cytokines interleukin (IL)-1 and IL-18 (14,15). It has been shown that the expression of NLRP3 is significantly elevated in adipose tissues of patients with obesity, dyslipidemia and diabetes, and is positively correlated with the severity of atherosclerosis (16). It is hypothesized that a high glucose (HG) environment may induce aberrant expression of the NLRP3 inflammasome in adipose tissue of T2DM patients, and that this may contribute to the development of atherosclerosis. However, the effects of HG on the expression of NLRP3 in adipocytes and adipose tissues remain to be elucidated.

Hydrogen sulfide (H₂S), a gaseous signaling transmitter, can be produced by a wide spectrum of mammalian tissues (17). Previous studies have indicated that H₂S plays numerous regulatory effects in the cardiovascular, gastrointestinal and neurological systems (18-21). The anti-inflammatory properties of H₂S have been identified in a range of cell types (22-24). In the cardiovascular system, H₂S is suggested to inhibit the development of atherosclerosis through suppression of inflammation (25,26). Cystathionine- γ -lyase (CSE) and cystathionine- β -synthase (CBS) are enzymes responsible for the synthesis of H₂S. It has been reported that both CSE and CBS are expressed in adipose tissues and adipocytes (27,28). Pan *et al* (29) demonstrated that HG inhibits expression of

Correspondence to: Dr Jing Wang or Dr Qing-Ying Tan, Department of Endocrinology, Chinese People's Liberation Army 903 Hospital, 14 Lingyin Road, Hangzhou, Zhejiang 310013, P.R. China

E-mail: wangjing200710@126.com

E-mail: tanqy1973@163.com

*Contributed equally

Key words: diabetes, NACHT leucine rich repeat and pyd domains-containing 3 inflammasome, adipocyte, hydrogen sulfide

CSE, and thus the production of H₂S, in adipocytes. This was confirmed by our previous study, which additionally revealed that HG did not affect the expression level of CBS (30). Several studies have also suggested that HG significantly increases the secretion of cytokines, including tumor necrosis factor- α (TNF- α), IL-6, monocyte chemoattractant protein-1 and adiponectin (31,32). However, the effect that this HG-induced reduction in CSE expression and H₂S production has on the expression of the NLRP3 inflammasome remains unknown. The hypothesis of the present study was that H₂S may play a role in HG regulation of NLRP3 expression in adipocytes. To test this hypothesis the expression of the NLRP3 inflammasome in adipocytes exposed to HG was investigated and an H₂S donor was used to try to reverse the HG effects.

Materials and methods

Cell culture and treatment. Adipocytes were cultured at 37°C with 5% CO₂ and differentiated from 3T3-L1 cells (American Type Culture Collection) as previously described (30). After disassociation using 0.125% trypsin, 1x10⁶/ml adipocytes were seeded and grouped for treatment. To determine the effect of HG, cells were treated with either low glucose (LG) DMEM (5.5 mmol/l glucose; HyClone; GE Healthcare Life Sciences; cat. no. SH30021.01) or HG DMEM (25.0 mmol/l glucose; HyClone; GE Healthcare Life Sciences; cat. no. SH30022.01) for 24 h. To determine the effect of H₂S, cells were treated with HG DMEM containing increasing concentrations of sodium hydrosulfide (NaHS; Sigma-Aldrich; Merck KGaA; cat. no. 161527) or HG DMEM without NaHS for 24 h. Our previous study revealed that 10, 25 and 50 nmol/l are effective concentrations of NaHS for treatment of adipocytes, so those concentrations were used in the present study (30). To inhibit the activity of the NLRP3 inflammasome, cells were treated with HG DMEM containing 10 μ g/ml N-acetyl-tyrosyl-valyl-alanyl-aspartyl chloromethyl ketone (Ac-YVAD-CMK; Sigma-Aldrich; Merck KGaA; cat. no. SML0429) or DMSO for vehicle for 24 h.

Western blot analysis. Proteins were extracted from adipocytes using radioimmunoprecipitation assay lysis buffer (Beyotime Institute of Biotechnology). The protein concentration was assayed using a BCA Protein Assay kit (Beyotime Institute of Biotechnology). Protein samples (50 μ g) were used for SDS-PAGE (4 and 10% gel) and subsequently transferred to nitrocellulose membranes. After blocking with 5% skim milk for 2 h at 37°C, membranes were incubated with primary antibodies against NLRP3 (Abcam; cat. no. ab10931; 1:1,000), apoptosis-associated speck-like protein containing A CARD (CST Biological Reagents Co., Ltd.; cat. no. 4628; 1:1,000), caspase-1 (Santa Cruz Biotechnology, Inc.; cat. no. SC-514; 1:1,000) and β -actin (Sigma-Aldrich; Merck KGaA; cat. no. A5441; 1:8,000) for 12 h at 4°C. The nitrocellulose membranes were then incubated with a secondary HRP-conjugated antibody (1:2,000). Immunoreactive proteins were then visualized using Immobilon Western Chemiluminescent HRP Substrate (Merck KGaA) and a Tanon 5200 Multi scanner (Tanon Science and Technology Co., Ltd.). The band intensities were calculated by ImageJ (version 1.51b; National Institutes of Health). Then the ratio of

band intensities to β -actin was obtained to quantify the relative protein expression levels.

ELISA analysis of IL-1 β and IL-18 release. After treatment, culture media were collected and the concentrations of IL-1 β and IL-18 released into culture media were determined using IL-1 β (cat. no. F10770) and IL-18 (cat. no. F10920) ELISA kits (Shanghai Westang Biotech Co., Ltd.) according to the manufacturer's instructions. All assays were performed in duplicate.

Statistical analysis. The data are presented as the mean $\pm$ SEM and were analyzed by one-way ANOVA followed by LSD-t test using SPSS (version 20; IBM Corp.). P<0.05 was considered to indicate a statistically significant difference.

Results

HG significantly upregulates the expression of NLRP3 inflammasome in adipocytes. Activation of the NLRP3 inflammasome initiates inflammation in adipose tissues and adipocytes. To investigate whether HG is associated with activation of the NLRP3 inflammasome, its expression in adipocytes was observed. The expression levels of NLRP3 inflammasome components NLRP3, ASC and caspase-1 were determined by western blot analysis. Compared with the LG group, HG significantly increased the level of NLRP3 and ASC, and the ratio of cleaved caspase-1 to pro-caspase-1 (Fig. 1A-D).

HG stimulates the release of IL-1 β and IL-18 by adipocytes via activation of the NLRP3 inflammasome. In the present study, mature IL-1 β and IL-18 levels in culture media were determined by ELISA. The data revealed that IL-1 β and IL-18 concentrations were significantly higher in the culture media of adipocytes treated with HG compared with those treated with LG. In order to confirm the role of NLRP3 inflammasome activation in this HG-induced IL-1 β and IL-18 release, adipocytes were treated with Ac-YVAD-CMK. As shown in Fig. 2, the IL-1 β and IL-18 concentrations in the culture media of adipocytes treated with HG + Ac-YVAD-CMK were significantly lower compared with HG DMEM containing vehicle.

Role of H₂S in the regulation of NLRP3 inflammasome activation in adipocytes. To confirm the role of H₂S in regulation of NLRP3 inflammasome activation, adipocytes were treated with HG DMEM containing increasing doses of the H₂S donor NaHS. The treatment of adipocytes with NaHS resulted in decreased HG-induced expression levels of NLRP3 and ASC, a reduced cleaved caspase-1 to pro caspae-1 ratio (Fig. 3), and a reduced HG-induced release of IL-1 β and IL-18 (Fig. 4).

Discussion

The present study demonstrated that HG increased the activity of the NLRP3 inflammasome in adipocytes. Additionally, NaHS, an H₂S donor, inhibited HG-induced expression of the NLRP3 inflammasome and the release of IL-1 and IL-18.

Figure 1. Effect of HG on the expression levels of NLRP3, ASC and caspase-1 in adipocytes. (A) Western blotting was performed to determine the expression levels of (B) NLRP3, (C) ASC and (D) the ratio of caspase-1 to pro caspase-1 in adipocytes treated with LG and HG. Data are presented as the mean $\pm$ SEM of four replicates (n=4 cultures). **P<0.01 as indicated. ASC, apoptosis-associated speck-like protein containing A CARD; HG, high glucose; LG, low glucose; NLRP3, NACHT leucine rich repeat and pyd domains-containing 3.

Figure 2. Effect of caspase-1 inhibitor Ac-YVAD-CMK on HG-induced IL-1 β and IL-18 release in adipocytes. ELISA was performed to determine the concentrations of (A) IL-1 β and (B) IL-18 in culture media from adipocytes treated with LG and HG. Data are presented as the mean $\pm$ SEM of four replicates. *P<0.05 and **P<0.01 as indicated. HG, high glucose; IL, interleukin; LG, low glucose; Ac-YVAD-CMK, N-acetyl-tyrosyl-valyl-alanyl-aspartyl chloromethyl ketone.

The link between T2DM and inflammation is well established. T2DM is considered to be, in part, a consequence of subclinical chronic low-grade inflammation (1-3). Several studies have reported that circulating inflammatory cytokines, such as c-reactive protein, TNF- α , IL-1 β , IL-6 and IL-18, are significantly elevated in T2DM patients (33-38). Elevated levels of these inflammatory cytokines directly induce insulin resistance and impair

glucose homeostasis (39-41). Cytokines of the IL-1 family are critical regulators of inflammation and control numerous inflammatory processes. Both IL-1 β and IL-18, which are classic pro-inflammatory cytokines of the IL-1 family, contribute to insulin resistance and islet β -cell damage in T2DM (39,42,43). Additionally, chronic inflammation is also a major feature of atherosclerosis. In the progression of T2DM, IL-1 β and IL-18 increase the risk of microvascular

Figure 3. Effect of H₂S donor NaHS on HG-induced NLRP3 inflammasome expression in adipocytes. (A) Western blot analysis was performed to determine the expression levels of (B) NLRP3 and (C) ASC and (D) the ratio of caspase-1 to pro caspase-1 in adipocytes treated with HG DMEM containing increasing concentrations of NaHS (0, 10, 25, 50 nM). Data are presented as the mean $\pm$ SEM of four replicates. * P <0.05 and ** P <0.01 as indicated. ASC, apoptosis-associated speck-like protein containing A CARD; H₂S, hydrogen sulfide; HG, high glucose; LG, low glucose; NaHS, sodium hydrosulfide; NLRP3, NACHT leucine rich repeat and pyd domains-containing 3.

Figure 4. Effect of H₂S donor NaHS on HG-induced IL-1 β and IL-18 release in adipocytes. ELISA was performed to determine the concentrations of (A) IL-1 β and (B) IL-18 in culture media from adipocytes treated with HG DMEM containing increasing concentrations of NaHS (0, 10, 25, 50 nM). Data are presented as the mean $\pm$ SEM of four replicates. * P <0.05 and ** P <0.01 as indicated. HG, high glucose; IL, interleukin; LG, low glucose; NaHS, sodium hydrosulfide.

and macrovascular complications by accelerating atherosclerosis (44). However, the underlying molecular mechanism behind the elevation of IL-1 β and IL-18 levels in T2DM patients has not been fully elucidated.

The majority of cytokines in the IL-1 family have been linked to obesity (14,15). Accumulating evidence has indicated that the NLRP3 inflammasome plays a critical role in regulating IL-1 β and IL-18 production in adipose tissues (3,13).

Once NLRP3 is activated, the inflammasome recruits pro-caspase-1. The clustering of pro-caspase-1 subunits at the inflammasome complex results in auto-cleavage and formation of active caspase-1. Active caspase-1 converts pro-IL-1 β and pro-IL-18 into their mature forms, IL-1 β and IL-18. In the adipose tissues of obese individuals, compared with lean individuals, activity of the NLRP3 inflammasome and expression levels of IL-1 β and IL-18 are significantly elevated (16,45,46).

T2DM patients with higher BMI and a greater amount of adipose tissue are found to have higher serum IL-1 β and IL-18 levels, which results in more severe insulin resistance and increased risk of cardiovascular disease (35,47). Despite this, whether an HG environment effects NLRP3 inflammasome expression and IL-1 β and IL-18 release in the adipose tissue of T2DM patients is unknown and merits further investigation. Results of the present study indicated that HG significantly increased the expression levels of NLRP3 and ASC, and caspase-1/pro caspase-1 ratio in adipocytes. The results also suggested that HG significantly increased IL-1 β and IL-18 release in the media of cultured adipocytes. In order to identify whether NLRP3 inflammasome activation was involved in HG-induced IL-1 β and IL-18 release, cells were treated with HG DMEM containing an NLRP3 inflammasome inhibitor. Results indicated that inhibition of the NLRP3 inflammasome abolished HG-induced IL-1 β and IL-18 release. These data suggest that HG increased the production of IL-1 β and IL-18 in adipocytes via activation of the NLRP3 inflammasome.

H₂S, a gaseous signaling transmitter, is reportedly involved in inflammation in various tissues (17). Adipocytes have been shown to express both CBS and CSE, and the expression of CSE and the generation of H₂S have been shown to be suppressed by HG (29,48). As previously discussed, activation of the NLRP3 inflammasome and release of IL-1 β and IL-18 increased in adipocytes exposed to HG. Therefore, it was hypothesized that a reduction in H₂S in HG DMEM may increase NLRP3 inflammasome expression and IL-1 β and IL-18 release. In order to verify this hypothesis, adipocytes were treated with HG DMEM in the presence of increasing concentrations of H₂S donor NaHS. The findings indicated that NaHS significantly suppressed NLRP3 inflammasome expression and IL-1 β and IL-18 release in adipocytes. These data suggest that exogenous H₂S can inhibit HG-induced NLRP3 inflammasome activation in adipocytes.

In summary, the results of the present study suggest that HG increased activation of the NLRP3 inflammasome in adipocytes. Exogenous H₂S donor NaHS significantly inhibited NLRP3 inflammasome expression, and IL-1 β and IL-18 production in adipocytes.

Acknowledgements

The authors thank Dr Xin Ni from the Department of Physiology, Second Military Medical University (Shanghai, China) for valuable comments on the manuscript.

Funding

This work was supported by the Natural Science Foundation of China (grant no. 81701481) and the Health Science and Technology Plan (General) Project of Hangzhou (grant no. 2016B56).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

TXH and NNZ were involved in drafting the manuscript. TXH, NNZ and YR collected and analyzed the data. QYT and JW interpreted the data, and all authors gave final approval of the version to be published. All authors reviewed the initial manuscript and revised it critically for important intellectual content.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Calle MC and Fernandez ML: Inflammation and type 2 diabetes. *Diabetes Metab* 38: 183-191, 2012.
2. Prattichizzo F, De Nigris V, La Sala L, Procopio AD, Olivieri F and Ceriello A: 'Inflammaging' as a druggable target: A senescence-associated secretory phenotype-centered view of type 2 diabetes. *Oxid Med Cell Longev* 2016: 1810327, 2016.
3. Esser N, Legrand-Poels S, Piette J, Scheen AJ and Paquot N: Inflammation as a link between obesity, metabolic syndrome and type 2 diabetes. *Diabetes Res Clin Pract* 105: 141-150, 2014.
4. Assar ME, Angulo J and Rodriguez-Manas L: Diabetes and ageing-induced vascular inflammation. *J Physiol* 594: 2125-2146, 2016.
5. Bessueille L and Magne D: Inflammation: A culprit for vascular calcification in atherosclerosis and diabetes. *Cell Mol Life Sci* 72: 2475-2489, 2015.
6. Kelsey MM, Forster JE, Van Pelt RE, Reusch JE and Nadeau KJ: Adipose tissue insulin resistance in adolescents with and without type 2 diabetes. *Pediatr Obes* 9: 373-380, 2014.
7. Lee J: Adipose tissue macrophages in the development of obesity-induced inflammation, insulin resistance and type 2 diabetes. *Arch Pharm Res* 36: 208-222, 2013.
8. Silaghi CA, Silaghi H, Crăciun AE, Fărcaș A, Colosi HA, Cosma DT, Pais R, Hâncu N and Georgescu CE: Age, abdominal obesity, and glycated hemoglobin are associated with carotid atherosclerosis in type 2 diabetes patients with nonalcoholic fatty liver disease. *Med Ultrason* 17: 300-307, 2015.
9. Jung CH, Kim BY, Kim KJ, Jung SH, Kim CH, Kang SK and Mok JO: Contribution of subcutaneous abdominal fat on ultrasonography to carotid atherosclerosis in patients with type 2 diabetes mellitus. *Cardiovasc Diabetol* 13: 67, 2014.
10. McGown C, Biredine A and Younossi ZM: Adipose tissue as an endocrine organ. *Clinics Liver Dis* 18: 41-58, 2014.
11. Coelho M, Oliveira T and Fernandes R: Biochemistry of adipose tissue: An endocrine organ. *Arch Med Sci* 9: 191-200, 2013.
12. Adamczak M and Wiecek A: The adipose tissue as an endocrine organ. *Semin Nephrol* 33: 2-13, 2013.
13. Yin Z, Deng T, Peterson LE, Yu R, Lin J, Hamilton DJ, Reardon PR, Sherman V, Winnier GE, Zhan M, *et al*: Transcriptome analysis of human adipocytes implicates the NOD-like receptor pathway in obesity-induced adipose inflammation. *Mol Cell Endocrinol* 394: 80-87, 2014.
14. Kursawe R, Dixit VD, Scherer PE, Santoro N, Narayan D, Gordillo R, Giannini C, Lopez X, Pierpont B, Nouws J, *et al*: A role of the inflammasome in the low storage capacity of the abdominal subcutaneous adipose tissue in obese adolescents. *Diabetes* 65: 610-618, 2016.

15. Murphy AM, Lyons CL, Finucane OM and Roche HM: Interactions between differential fatty acids and inflammatory stressors-impact on metabolic health. *Prostaglandins Leukot Essent Fatty Acids* 92: 49-55, 2015.
16. Bando S, Fukuda D, Soeki T, Nishimoto S, Uematsu E, Matsuura T, Ise T, Tobiume T, Yamaguchi K, Yagi S, *et al*: Expression of NLRP3 in subcutaneous adipose tissue is associated with coronary atherosclerosis. *Atherosclerosis* 242: 407-414, 2015.
17. Renga B: Hydrogen sulfide generation in mammals: The molecular biology of cystathionine- β -synthase (CBS) and cystathionine-gamma-lyase (CSE). *Inflamm Allergy Drug Targets* 10: 85-91, 2011.
18. Tripatara P, Patel NS, Collino M, Gallicchio M, Kieswich J, Castiglia S, Benetti E, Stewart KN, Brown PA, Yaqoob MM, *et al*: Generation of endogenous hydrogen sulfide by cystathionine gamma-lyase limits renal ischemia/reperfusion injury and dysfunction. *Lab Invest* 88: 1038-1048, 2008.
19. Panthi S, Chung HJ, Jung J and Jeong NY: Physiological importance of hydrogen sulfide: Emerging potent neuroprotector and neuromodulator. *Oxid Med Cell Longev* 2016: 9049782, 2016.
20. Meng XM, Huang X, Zhang CM, Liu DH, Lu HL, Kim YC and Xu WX: Hydrogen sulfide-induced enhancement of gastric fundus smooth muscle tone is mediated by voltage-dependent potassium and calcium channels in mice. *World J Gastroenterol* 21: 4840-4851, 2015.
21. Das A, Samidurai A, Hoke NN, Kukreja RC and Salloum FN: Hydrogen sulfide mediates the cardioprotective effects of gene therapy with PKG-I α . *Basic Res Cardiol* 110: 42, 2015.
22. Vandiver M and Snyder SH: Hydrogen sulfide: A gasotransmitter of clinical relevance. *J Mol Med (Berl)* 90: 255-263, 2012.
23. Wang XH, Wang F, You SJ, Cao YJ, Cao LD, Han Q, Liu CF and Hu LF: Dysregulation of cystathionine γ -lyase (CSE)/hydrogen sulfide pathway contributes to ox-LDL-induced inflammation in macrophage. *Cell Signal* 25: 2255-2262, 2013.
24. Hirata I, Naito Y, Takagi T, Mizushima K, Suzuki T, Omatsu T, Handa O, Ichikawa H, Ueda H and Yoshikawa T: Endogenous hydrogen sulfide is an anti-inflammatory molecule in dextran sodium sulfate-induced colitis in mice. *Dig Dis Sci* 56: 1379-1386, 2011.
25. Yu XH, Cui LB, Wu K, Zheng XL, Cayabyab FS, Chen ZW and Tang CK: Hydrogen sulfide as a potent cardiovascular protective agent. *Clin Chim Acta* 437: 78-87, 2014.
26. Xu S, Liu Z and Liu P: Targeting hydrogen sulfide as a promising therapeutic strategy for atherosclerosis. *Int J Cardiol* 172: 313-317, 2014.
27. Tsai CY, Peh MT, Feng W, Dymock BW and Moore PK: Hydrogen sulfide promotes adipogenesis in 3T3L1 cells. *PKLoS One* 10: e0119511, 2015.
28. Beltowski J: Endogenous hydrogen sulfide in perivascular adipose tissue: Role in the regulation of vascular tone in physiology and pathology. *Can J Physiol Pharmacol* 91: 889-898, 2013.
29. Pan Z, Wang H, Liu Y, Yu C, Zhang Y, Chen J, Wang X and Guan Q: Involvement of CSE/H₂S in high glucose induced aberrant secretion of adipokines in 3T3-L1 adipocytes. *Lipids Health Dis* 13: 155, 2014.
30. Hu TX, Wang G, Wu W, Gao L, Tan QY and Wang J: Hydrogen sulfide inhibits high glucose-induced sFlt-1 production via decreasing ADAM17 expression in 3T3-L1 adipocytes. *Int J Endocrinol* 2017: 9501792, 2017.
31. Grosick R, Alvarado-Vazquez PA, Messersmith AR and Romero-Sandoval EA: High glucose induces a priming effect in macrophages and exacerbates the production of pro-inflammatory cytokines after a challenge. *J Pain Res* 11: 1769-1778, 2018.
32. Briones L, Andrews M, Pizarro F and Arredondo-Olguin M: Expression of genes associated with inflammation and iron metabolism in 3T3-L1 cells induced with macrophages-conditioned medium, glucose and iron. *Biomaterials* 31: 595-604, 2018.
33. Akbarzadeh M, Eftekhari MH, Dabbaghmanesh MH, Hasanazadeh J and Bakhshayeshkaram M: Serum IL-18 and hsCRP correlate with insulin resistance without effect of calcitriol treatment on type 2 diabetes. *Iran J Immunol* 10: 167-176, 2013.
34. Mir M, Rostami A and Hormozi M: Comparison of serum levels of IL-18 in peripheral blood of patients with type II diabetes with nephropathy clinical protests and patients with type II diabetes without nephropathy clinical protests. *Diabetes Metab Syndr* 11: 245-250, 2017.
35. Herder C, Dalmás E, Boni-Schnetzler M and Donath MY: The IL-1 pathway in type 2 diabetes and cardiovascular complications. *Trends Endocrinol Metab* 26: 551-563, 2015.
36. Banerjee M and Saxena M: Interleukin-1 (IL-1) family of cytokines: Role in type 2 diabetes. *Clin Chim Acta* 413: 1163-1170, 2012.
37. Daniele G, Guardado Mendoza R, Winnier D, Fiorentino TV, Pengou Z, Cornell J, Andreozzi F, Jenkinson C, Cersosimo E, Federici M, *et al*: The inflammatory status score including IL-6, TNF- α , osteopontin, fractalkine, MCP-1 and adiponectin underlies whole-body insulin resistance and hyperglycemia in type 2 diabetes mellitus. *Acta Diabetol* 51: 123-131, 2014.
38. Hussain G, Rizvi SA, Singhal S, Zubair M and Ahmad J: Serum levels of TNF- α in peripheral neuropathy patients and its correlation with nerve conduction velocity in type 2 diabetes mellitus. *Diabetes Metab Syndr* 7: 238-242, 2013.
39. Hardaway AL and Podgorski I: IL-1 β , RAGE and FABP4: Targeting the dynamic trio in metabolic inflammation and related pathologies. *Future Med Chem* 5: 1089-1108, 2013.
40. Dinarello CA: Interleukin-18 and the pathogenesis of inflammatory diseases. *Semin Nephrol* 27: 98-114, 2007.
41. Lee CC, Lorenzo C, Haffner SM, Wagenknecht LE, Festa A, Goodarzi MO, Stefanovski D, Olson NC, Norris JM, Rewers MJ and Hanley AJ: The association of inflammatory and fibrinolytic proteins with 5 year change in insulin clearance: The Insulin resistance atherosclerosis study (IRAS). *Diabetologia* 56: 112-120, 2013.
42. Maedler K, Dharmadhikari G, Schumann DM and Storling J: Interleukin-1 beta targeted therapy for type 2 diabetes. *Expert Opin Biol Ther* 9: 1177-1188, 2009.
43. Bosch M, Lopez-Bermejo A, Vendrell J, Musri M, Ricart W and Fernandez-Real JM: Circulating IL-18 concentration is associated with insulin sensitivity and glucose tolerance through increased fat-free mass. *Diabetologia* 48: 1841-1843, 2005.
44. Frostegard J: Immune mechanisms in atherosclerosis, especially in diabetes type 2. *Front Endocrinol (Lausanne)* 4: 162, 2013.
45. Vandanmagsar B, Youm YH, Ravussin A, Galgani JE, Stadler K, Mynatt RL, Ravussin E, Stephens JM and Dixit VD: The NLRP3 inflammasome instigates obesity-induced inflammation and insulin resistance. *Nat Med* 17: 179-188, 2011.
46. Esser N, L'Homme L, De Roover A, Kohnen L, Scheen AJ, Moutschen M, Piette J, Legrand-Poels S and Paquot N: Obesity phenotype is related to NLRP3 inflammasome activity and immunological profile of visceral adipose tissue. *Diabetologia* 56: 2487-2497, 2013.
47. Zilverschoon GR, Tack CJ, Joosten LA, Kullberg BJ, van der Meer JW and Netea MG: Interleukin-18 resistance in patients with obesity and type 2 diabetes mellitus. *Int J Obes (Lond)* 32: 1407-1414, 2008.
48. Feng X, Chen Y, Zhao J, Tang C, Jiang Z and Geng B: Hydrogen sulfide from adipose tissue is a novel insulin resistance regulator. *Biochem Biophys Res Commun* 380: 153-159, 2009.