

Effects of various catheter fix sites on catheter-associated lower urinary tract symptoms

LIKUN ZHU^{1,2}, RUI JIANG^{1,2}, XIANGJUN KONG^{1,2}, XINWEI WANG^{1,2},
LIJUN PEI^{1,2}, QINGFU DENG^{1,2} and XU LI^{1,2}

¹Department of Urology Surgery, The Affiliated Hospital of Southwest Medical University,

²Sichuan Clinical Medical Research Center for Nephropathy, Luzhou, Sichuan 646000, P.R. China

Received March 5, 2020; Accepted July 17, 2020

DOI: 10.3892/etm.2020.9478

Abstract. The present study aimed to compare the effects of various catheter fix sites on catheter-associated lower urinary tract symptoms (CALUTS) in 450 patients who underwent surgical removal of upper urinary calculi 24 h earlier. All patients had 16 French Foley catheters inserted and the balloons were filled. In group A, the catheters were fixed on the top one-third of the thigh. In group B, the catheters were fixed on the abdominal wall. Patients in whom the catheters were neither fixed on the thigh nor abdominal wall were designated as controls. There were 150 patients in each group. CALUTS, such as frequency, urgency, burning during micturition, odynuria, bladder pain and other symptoms, including urethral discharge, a red and swollen external urethral orifice, catheter traction or blockage and catheter-associated discomfort were recorded. Patients in group A compared with the control group had a significantly lower incidence of frequency, urgency, odynuria, urethral discharge, catheter traction and catheter-associated discomfort ($P<0.05$). Patients in group B were observed to have a significantly lower incidence of urgency, urethral discharge, catheter traction and catheter-associated discomfort compared with the control group ($P<0.05$), but a higher incidence of odynuria, urethral pain, urethral discharge and a red and swollen external urethral orifice compared with group A ($P<0.05$). An additional catheter fixation site for bedridden patients was necessary and an additional fix site on the thigh was preferred to the abdominal wall, which may further reduce catheter-associated lower urinary tract symptoms.

Introduction

Indwelling urethral catheters are a mainstay in the care of hospitalized patients. An estimated 30 million indwelling

urethral (Foley) catheters are used annually in the United States (1) and ~20% of hospitalized patients have a urethral catheter at any given time (2,3). Catheter-associated lower urinary tract symptoms (CALUTS) refer to a range of symptoms, including frequency, urgency, burning during micturition, odynuria and suprapubic pain (3). These symptoms may be caused by involuntary detrusor contractions induced by activation of muscarinic type III receptors, also known as M_3mAChR , and the increased release of acetylcholine, which are caused by reactions of the mucosa in the trigone and urethra after insertion of a catheter (1-8). An indwelling catheter enables the continuous drainage of urine, which maintains an empty bladder, enhances the stimulatory effects of the catheter on the bladder walls and causes an increased incidence of CALUTS (6). In addition, severe CALUTS may also be associated with higher the volume of the balloon and the weight of the catheter and/or urine bag (9). Previous studies have reported that CALUTS may aggravate post-operative pain, increase the occurrence of post-operative complications, including burning during micturition, odynuria and suprapubic pain or even prolong the length of hospital stay (7,10,11). Previous studies in CALUTS prevention have focused on the use of post-operative medications, including anticholinergic drugs, a receptor blockers (2-4,12-14). However, the side effects, such as nausea, vomiting, dizziness and conscious state changes may have adverse effects on patient recovery (11).

The present study aimed to compare the effects of various fixed catheter sites on preventing CALUTS in patients who underwent surgical removal of upper urinary calculi 24 h after surgery.

Materials and methods

Study design and subjects. A total of 450 patients who underwent surgical removal of upper urinary calculi at The Affiliated Hospital of Southwest Medical University (Luzhou, China) between January 2018 and September 2018 were included in this study. A total of 450 patients were selected in the present study, however, later 4 patients were excluded due to breakage of their catheter balloons. A final of 446 patients were included in the present study, which were control group ($n=148$) and groups A and B ($n=149$ /group). A 16 French double-cava Foley urethral catheter (Bardex) was inserted

Correspondence to: Dr Likun Zhu, Department of Urology Surgery, The Affiliated Hospital of Southwest Medical University, 25 Taiping Road, Luzhou, Sichuan 646000, P.R. China
E-mail: zhulikun2019@163.com

Key words: catheter-associated lower urinary tract symptoms, additional fixation, abdominal wall, thigh

into the bladders of each patient following surgery (Fig. 1). Patients >18 years of age with normal cognitive ability and renal function (serum creatinine $\leq 44.00 \mu\text{mol/l}$, urea nitrogen $\leq 1.70 \text{ mmol/l}$) met the inclusion criteria for participation in the present study. Patients who were diagnosed with benign prostate hyperplasia or urethrostenosis, had a history of bladder dysfunction (overactive bladder, nocturia >3, or micturition >8 for 24 h before the surgery (7) neurogenic bladder, bladder inflammation, incontinence, lower urinary tract infection, urinary tuberculosis or cognitive disorders were excluded. One hundred and fifty patients were included in each group. The present study was approved by the Ethics Committee of the Affiliated Hospital of Southwest University (Luzhou, China; approval no. K2019002-R). Informed written consent was obtained from all the patients.

All patients were randomly and evenly divided into 3 groups (control, group A and group B), 150 patients in each group. The present study was not double blinded as the fixed site for the catheter on patients was visible. Patients with renal calculi who underwent percutaneous nephrolithotomy were treated with 100 ml of 0.9% normal saline (NS) + 50 mg of flurbiprofen axetil (Beijing Forte Meditec Co., Ltd.) in an intravenous drip twice on the day of surgery and the first day after surgery, while patients with ureteral calculi were not treated with post-operative analgesics. Ureteral stents (double J tube; C.R. Bard, Inc.) were retained for one month in all patients post-operatively.

A highly trained nurse was responsible for catheter fixation and all patients were followed-up for 24 h after catheter insertion. Catheters were inserted into the bladders of all patients and the balloons were filled. The catheter was pulled back after filling the balloon and stopped when the balloon was in contact with the bladder neck. In group A, the catheters were fixed on the top one-third of the thigh at the same level as the external urethral orifice. In doing so the extra-urethral part of the catheter was U-shaped (Fig. 2A). In group B, the catheters were fixed to the lower abdominal wall adjacent to the groin (Fig. 2B). The patients in whom the catheters were neither fixed on the thigh nor the abdominal wall were designated as controls (Fig. 2C).

Observation indices were recorded 24 h following catheter insertion, including the incidence of various CALUTS, such as frequency, urgency, odynuria, pain in the urethra or bladder, an increase in urethral discharge, a red and swollen external urethral orifice and other discomfort associated with the indwelling catheter or catheter obstruction.

Pain in the urethra and bladder were evaluated according to a previous study as follows: 0, no pain; 1-3, mild pain; 4-6, moderate pain; and 7-10, severe pain (15). The catheter-associated level of discomfort was quantified as follows: 0, no discomfort; 1, mild discomfort; 2, moderate discomfort, but able to tolerate the indwelling catheter; and 3, severe discomfort requiring removal of the catheter (6).

Statistical analysis. The data were analyzed using the SPSS 18.0 (SPSS, Inc.). The normality of data distribution of the continuous variables was assessed using the Shapiro-Wilk test. Continuous variables with normal distribution are presented as the mean \pm standard deviation (SD). The means of normally distributed continuous variables were compared




Figure 1. 3M tape clipping. Foley urethral catheter was inserted into the bladders of each patient following surgery and fixed using 3M tape clipping cut into this shape.

using Tukey's test and one-way ANOVA (Respectively, if the results of ANOVA was significant, Tukey's test was used to compare the differences between the two means.). Tukey's test was used to compare between groups. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Basic characteristics. No significant differences were observed in indices, such as age, sex, education level, types of surgery, including percutaneous nephroscopy (PCNL) and ureteroscopy (URL) and the length of the operation between the control group and groups A and B ($P > 0.05$; Table I).

Comparison of urinary catheter related LUTS. Patients in group A had significantly lower incidences of frequency, urgency and odynuria compared with patients in the control group ($P < 0.05$). The proportion of patients who complained of bladder pain in groups A or B was significantly lower compared with patients in the control group ($P < 0.05$) (Table II), while the number of patients with mild bladder pain in group A was significantly higher compared with group B ($P < 0.05$) (Table II). The incidence of dysuria and total urethral pain scores of group A was significantly lower than group B ($P < 0.05$). In addition, patients in group B had a significantly lower incidence of urgency when compared with the control group ($P < 0.05$) (Table II). Significantly lower incidences of urethral discharge and catheter traction were observed in patients in group A compared with patients in the group B ($P < 0.05$; Table III). In group B, a significantly higher incidence of urethral discharge was observed compared with group A and the control group, and a significantly higher incidence of red and swollen external urethral orifices in group B compared with group A was observed ($P < 0.05$; Table III). The incidence of urethrorrhea in the control group was significantly higher than group A and B (10 vs. 1 and 4, respectively), while there were only two cases with catheter blockage in group A.

Comparison of catheter associated discomfort. There were 51 cases with mild of catheter-associated discomfort in control group, while it was 40 and 41 cases in group A and group B,




Figure 2. Catheter fixture. (A) Catheters were fixed on the abdominal wall or thigh. (B) Catheters were fixed on the abdominal wall. (C) Catheters were fixed on neither the thigh nor the abdominal wall.

respectively. The total cases with catheter associated discomfort was 63 in control group, while it was 42 and 46 in group A and group B, respectively. Patients in groups A and B had a lower incidence of mild discomfort when compared with the control group (Table IV).

Discussion

In the present study the effects of various fixed sites of catheters on preventing CALUTS in patients who underwent surgical removal of upper urinary calculi were assessed, and it was demonstrated that catheters fixed by filling the balloon then fixing the catheter to the top one-third of the thigh were most beneficial for patients.

In the present study, patients in group A complained of significantly less frequency and urgency, odynuria and bladder pain compared with the control group, which may be due to the additional fixed site of the catheter. The additional fixed site on the thigh changed the pressure of the catheter and urine bag weight on the additional fixed position. Hence, this fixation site could decrease the pull force of the catheter and urine bag on the neck of the bladder and urethra. In the control group, the pull force of the catheter and urine bag may directly press on the neck of the bladder and urethra, especially when the patients were in bed. The aforementioned reason may explain the complaints of majority of the patients in the control group who reported more serious CALUTS when in bed, and may also be responsible for the significantly higher incidence of frequency, urgency and odynuria in the control group compared with group A and B.

In the present study, patients in group B also had an additional fixed site of the catheter on the abdominal wall; however, they had a significantly lower incidence of frequency and a significantly decreased composition ratio of bladder pain when compared with the control group ($P < 0.05$). Patients in group B had a significantly increased incidence of odynuria, urethral pain and proportion of severe bladder pain compared with group A ($P < 0.05$). The findings of the present study were consistent with those of Zhang and Zhang (16) who reported that the additional fixation of the catheter on the inside of the thigh may lower the incidence of external urethral orifice pain, ulcers and necrosis in patients with neurosurgical crises. The present study hypothesized that the difference of CALUTS between groups A and B in the present study may be due to the different fixed position of urethral catheter resulting in different changes in normal physiological structure of the urethra. Although, in the present study the additional fixation of the catheter on the abdominal wall decreased the pull forces from the catheter and urine bag on the urethra, the additional fixation site was still at a higher level compared with the external urethral orifice, which caused a reverse angle to be formed between the extra-urethral part of the catheter and the urethra (the size of the angle depends on the fat thickness beneath the abdominal wall). This reverse angle causes the catheter to pull and press on the urethra and therefore increases the incidence rate of odynuria. In addition, the additional fixation on the abdominal wall affects the physio anatomy of the urethra and causes the disappearance of the urethral ante curvature (6,17). The urethral ante curvature was therefore passive straightened in the present study which increased the urethral pain. In

Table I. Parameters of patients (n=446) with different catheter fixation sites who underwent surgical removal of upper urinary calculi.

Patients	Age, years $\bar{X} \pm S$	Length of the operation, min $\bar{X} \pm S$	Sex, n (%)		Education level, n (%)				Type of surgery, n (%)		
			Male	Female	Illiterate	Primary	Junior	High	University	PCNL	URL
Control (n=148)	54.14 \pm 18.97	45.53 \pm 17.89	91 (61.49)	57 (38.51)	20 (13.51)	53 (35.81)	45 (30.41)	20 (13.51)	10 (6.76)	64 (43.24)	84 (56.76)
Group A (n=149)	54.27 \pm 18.54	45.79 \pm 17.58	95 (63.76)	54 (36.24)	21 (14.09)	50 (33.56)	47 (31.54)	22 (14.77)	9 (6.04)	67 (44.97)	82 (55.03)
Group B (n=149)	53.99 \pm 19.12	46.01 \pm 17.64	93 (62.42)	56 (37.58)	20 (13.42)	51 (34.23)	48 (32.21)	23 (15.44)	7 (4.70)	65 (43.62)	84 (56.38)
χ^2 value ^a	0.06	0.13	0.164		0.3					0.085	
P-value ^a	>0.05	>0.05	0.686		0.99 ^d					0.765 ^e	
χ^2 value ^b	t=0.07	0.23	0.027		0.871					0.004	
P-value ^b	>0.05	>0.05	0.869		0.929 ^d					0.947 ^e	
χ^2 value ^c	t=0.13	0.11	0.058		0.317					0.054	
P-value ^c	>0.05	>0.05	0.810		0.989 ^d					0.816 ^e	

^aRepresents the comparison between group A and control group; ^brepresents comparison between group B and control group; and ^crepresents comparison between group A and group B. PCNL, percutaneous nephrolithotripsy; URL, transurethral ureteroscopic lithotripsy; ^dcomparison of constituent ratio of different education levels; ^ecomparison of different surgical types.

Table II. Comparison of catheter-associated lower urinary tract symptoms between the 3 groups of patients.

Patients	Frequency n (%)	Urgency	Odynuria	Urethral pain scores (15) n (%)					Bladder pain scores n (%)				
				1-3	4-6	7-10	Total		1-3	4-6	7-10	Total	
Control (n=148)	86 (58.11)	36 (24.32)	18 (12.16)	54 (36.49)	3 (2.03)	2 (1.35)	59 (39.87)		23 (15.54)	7 (4.73)	7 (4.73)	37 (25.00)	
Group A (n=149)	68 (45.64)	16 (10.74)	13 (8.72)	33 (22.15)	0	2 (1.34)	35 (23.49)		23 (15.44)	1 (0.67)	0 (0.00)	24 (16.11)	
Group B (n=149)	73 (48.99)	18 (12.08)	26 (17.45)	55 (36.91)	2 (1.34)	2 (1.34)	59 (39.59)		20 (13.42)	0 (0.00)	5 (3.36)	5 (3.36)	
χ^2 value ^a	4.635	9.489	0.938	3.082			9.203		11.974			3.598	
P-value ^a	0.032	0.002	0.333	0.214 ^e			0.002		0.003 ^d			0.058	
χ^2 value ^b	2.480	7.482	1.645	0.211			0.002		7.911			3.038	
P-value ^b	0.115	0.006	0.200	0.900 ^e			0.962		0.019 ^d			0.081	
χ^2 value ^c	0.337	0.133	4.986	2.135			8.951		4.073			0.024	
P-value ^c	0.562	0.716	0.026	0.344 ^e			0.003		0.044 ^d			0.876	

^aRepresents the comparison between group A and control group; ^brepresents comparison between group B and control group; ^crepresents comparison between group A and group B; ^drepresents the comparison constituent ratio of urethra pain; and ^erepresents the comparison of bladder pain with different severity.

Table III. Comparison of other catheter-associated lower urinary tract symptoms between the 3 groups of patients.

Patients	Urethral discharge n (%)	Urethrorrhea n (%)	Catheter traction, n (%)	Red and swollen urethra, n (%)	Catheter blockage n (%)
Control (n=148)	77 (52.03)	10 (6.76)	71 (47.97)	12 (8.11)	0 (0.00)
Group A (n=149)	94 (63.09)	1 (0.67)	14 (9.40)	7 (4.70)	2 (1.34)
Group B (n=149)	111 (74.50)	4 (2.68)	13 (8.72)	20 (13.42)	0 (0.00)
χ^2 value ^a	3.718	7.710	54.089	1.442	0.497
P-value ^a	0.054	0.005	0.000	0.230	0.481
χ^2 value ^b	16.137	2.741	56.388	2.182	-
P-value ^b	0.000	0.098	0.000	0.140	-
χ^2 value ^c	4.517	0.814	0.041	6.883	0.503
P-value ^c	0.034	0.367	0.840	0.009	0.478

^arepresents the comparison between group A and control group; ^brepresents comparison between group B and control group; and ^crepresents comparison between group A and group B.

Table IV. Comparison of catheter associated discomfort between the 3 groups of patients (6).

Patients	Level 1 (mild) n (%)	Level 2 (moderate) n (%)	Level 3 (severe) n (%)	Total n (%)
Control (n=148)	51 (34.46)	8 (5.41)	4 (2.70)	63 (42.57)
Group A (n=149)	40 (26.85)	0 (0.00)	2 (1.34)	42 (28.19)
Group B (n=149)	41 (27.52)	0 (0.00)	5 (3.36)	46 (30.88)

group A, the additional fixation of the catheter on the thigh enabled the catheter and external urethral orifice to be at the same level which thereby maintaining the physiological curve of the urethra, decreased the press force of the catheter on the urethra and decreased the incidence of odynuria.

In the present study, patients in groups A and B were observed to have a significantly lower incidence of catheter traction compared with the control group ($P<0.05$); however, no significant difference in the incidence of catheter traction or blockage was observed between groups A and B ($P>0.05$). The findings of the present study were not consistent with those of Zheng and Li (18), which considered that catheter traction was more likely to occur in the thigh fixation than in the abdominal wall fixation. The present study hypothesized that the different additional fixation sites and methodology utilized may be responsible for this inconsistency. In the present study, in patients from group A the pull force of the catheter and urine bag on the urethra and bladder neck could be released when the patients abducted their thigh, bend their knees and rolled over in bed. Patients in groups A or B demonstrated no significant difference in incidence of a red and swollen external urethral orifice compared with the control group ($P>0.05$), while a significantly lower incidence was found in group A when compared with group B ($P<0.05$). The aforementioned findings were consistent with the results obtained by Zhang and Zhang (16). The present study hypothesized that the additional fixation on the abdominal wall may increase the pull force of the catheter on the urethra, form the reverse angle and cause the passive straight urethral antecurvature. In addition, the significant increase in urethral discharge in

group B compared with group A and the control group may also be responsible for the difference observed between the groups ($P<0.05$).

In the present study, the incidence of catheter-associated discomfort and the proportion of severe catheter-associated discomfort in groups A and B was significantly lower compared with the control group. In groups A and B, the majority of patients complained of mild catheter-associated discomfort, while in the control group, the majority of patients complained of moderate and severe discomfort. The different incidence rates of pull force by the catheter may be responsible for this finding. The pull force from the catheter may have a crucial effect on CALUTS; however, in the present study the incidence of odynuria, urethral and bladder pain in group A was significantly lower compared with group B, while no significant difference was observed in the incidence of catheter-associated discomfort and the proportion of severe catheter-associated discomfort between groups A and B. The present study hypothesized that catheter-associated discomfort may involve the incidence of CALUTS and the pull force of the catheter, and might also depend on the tolerance of the patient. This may explain inconsistent results between the present study and those of Sun and Wang (19) who compared the different comfort levels between various catheter fixation sites and concluded that there was no significant difference of the comfort of female patients between the abdominal wall fixation and thigh fixation.

The present study had some limitations. First, whether fixation of the catheter could be applied to urological surgery of the lower urinary tract was not evaluated. Further studies with clinical efficacy were needed. Secondly, the observation

time was only 24 h. With the extension of catheter indwelling time, whether different fixation methods have different effects on the occurrence of CALUTS and other accompanying symptoms needs further observation and confirmation.

The present study demonstrated that additional fixation of the catheter for bedridden patients was necessary and decreased the incidence of CALUTS and other concomitant symptoms, such as catheter traction and urethral discharge. An additional fixation site on the thigh is recommended based on the results of the present study.

Acknowledgements

Not applicable.

Funding

The present study was funded by a grant from the Health Department of Sichuan Province for the Research of Continuing Nursing Health Promoting Education Model for Patients with Urinary Stones (grant no. 16158).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

The authors declare that all the listed authors have participated actively in the study and all met the requirements of the authorship. LZ and RJ conducted study design and experimental supervision. LZ wrote the first draft of the manuscript, RJ edited the manuscript and reviewed it, XW conducted secondary fixation and data collection of urinary catheters in all patients, LP and QD were responsible for intraoperative catheter indwelling and airbag fixation, XK and XL were responsible for statistical analysis of data. All authors read and approved the publication of the manuscript.

Ethics approval and consent to participate

This study was approved by the Ethics Committee of The Affiliated Hospital of Southwest University (Luzhou, China; approval no. K2019002-R). Informed written consent was obtained from each patient.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Binhas M, Motamed C, Hawajri N, Yiou R and Marty J: Predictors of catheter-related bladder discomfort in the post-anaesthesia care unit. *Ann Fr Anesth Reanim* 30: 122-125, 2011.
2. Agarwal A, Dhiraaj S, Singhal V, Kapoor R and Tandon M: Comparison of efficacy of oxybutynin and tolterodine for prevention of catheter related bladder discomfort: A prospective, randomized, placebo-controlled, double-blind study. *Brit J Anaesth* 96: 377-380, 2006.
3. Ryu JH, Hwang JW, Lee JW, Seo JH, Park HP, Oh AY, Jeon YT and Do SH: Efficacy of butylscopolamine for the treatment of catheter-related bladder discomfort: A prospective, randomized, placebo-controlled, double-blind study. *Brit J Anaesth* 111: 932-937, 2013.
4. Shi H, Zhang H, Pan W, Lu Y, Zhang Y, Chen L, Wei C and Cui Y: Pooled analysis of the efficacy and safety of intraoperative dexmedetomidine on postoperative catheter-related bladder discomfort. *Low Urin Tract Symptoms*: Jun 4, 2020 (Epub ahead of print). doi: 10.1111/luts.12322.
5. Kim HC, Lee YH, Jeon YT, Hwang JW, Lim YJ, Park JE and Park HP: The effect of intraoperative dexmedetomidine on postoperative catheter-related bladder discomfort in patients undergoing transurethral bladder tumor resection: A double-blind randomised study. *Eur J Anaesthesiol* 32: 596-601, 2015.
6. Wilson M: Causes and management of dwelling urinary catheter-related pain. *Br J Nurs* 17: 232-239, 2008.
7. Anderson KE and Wein AJ: Pharmacology of the lower urinary tract: basis for current and future treatments of urinary incontinence. *Pharmacol Rev* 56: 581-631, 2004.
8. Li JY and Ren Liao R: Prevention of catheter-related bladder discomfort-pudendal nerve block with ropivacaine versus intravenous tramadol: Study protocol for a randomized controlled trial. *Trials* 17: 448, 2016.
9. Hanchett M: Techniques for stabilizing urinary catheters. *Am J Nurs* 102: 44-48, 2002.
10. Bai Y, Wang X, Li X, Pu C, Yuan H, Tang Y, Li J, Wei Q and Han P: Management of catheter-related bladder discomfort in patients who underwent elective surgery. *J Endourol* 29: 640-649, 2015.
11. Lepousé C, Lautner CA, Liu L, Gomis P and Leon A: Emergence delirium in adults in the post-anaesthesia care unit. *Br J Anaesth* 96: 747-753, 2006.
12. Srivastava VK, Agrawal S, Kadiyala VN, Ahmed M, Sharma S and Kumar R: The efficacy of pregabalin for prevention of catheter-related bladder discomfort: A prospective, randomized, placebo-controlled double-blind study. *J Anesth* 29: 212-216, 2015.
13. Maro S, Zarattin D, Baron T, Bourez S, de la Taille A and Salomon L: Catheter-related bladder discomfort after urological surgery: Importance of the type of surgery and efficiency of treatment by clonazepam. *Prog Urol* 24: 628-633, 2014 (In French).
14. Shariat Moharari R, Lajevardi M, Khajavi M, Najafi A, Shariat Moharari G and Etezadi F: Effects of intra-operative ketamine administration on postoperative catheter-related bladder discomfort: A double-blind clinical trial. *Pain Pract* 14: 146-150, 2014.
15. Breivik H, Borchgrevink PC, Allen SM, Rosseland LA, Romundstad L, Hals EK, Kvarstein G and Stubhaug A: Assessment of pain. *Br J Anaesth* 101: 17-24, 2008.
16. Zhang J and Zhang X: Effect of external urinary catheterization on indwelling catheterization in critically ill patients with neurosurgery. *Nurs Res* 19: 1577-1578, 2005.
17. Na YQ, Guo ZH: Practical urology. First Edition. Beijing: People's Medical Publishing House, 2009.
18. Zheng LI and Li D: Comparison of the effects of different methods of fixation of indwelling catheters in elderly patients. *Gen Nurs* 10: 1570-1571, 2012.
19. Sun Y and Wang X: Improvement and effect of female indwelling catheter fixation method. *Nurs Res* 22: 624, 2008.