

Endogenous antisense *walR* RNA modulates biofilm organization and pathogenicity of *Enterococcus faecalis*

SHIZHOU WU¹⁻³, YUNJIE LIU⁴, LEI LEI² and HUI ZHANG¹

¹Department of Orthopedics, West China Hospital; ²State Key Laboratory of Oral Diseases, Department of Preventive Dentistry, West China Hospital of Stomatology, Sichuan University, Chengdu, Sichuan 610041, P.R. China; ³Department of Advanced Oral Sciences and Therapeutics, University of Maryland School of Dentistry, Baltimore, MD 21201, USA; ⁴West China School of Public Health, Sichuan University, Chengdu, Sichuan 610041, P.R. China

Received November 7, 2019; Accepted June 10, 2020

DOI: 10.3892/etm.2020.9501

Abstract. *Enterococcus faecalis* (*E. faecalis*) is regarded as the major pathogen for persistent periapical periodontitis. The aim of the present study was to investigate the role of antisense *walR* RNA in the regulation of adjacent downstream genes. Reverse transcription-PCR assays were performed to validate *walR*. Adjacent downstream genes *walK*, EF1195, EF1196, and EF1197 were co-transcribed and detect antisense *walR* RNA. Northern blotting and 5'-rapid amplification of cDNA ends (5'-RACE) assays were conducted to detect and confirm a novel *walR* antisense (AS*walR*) RNA. AS*walR* overexpression mutants were constructed, and the biofilm biomass was determined using a crystal violet microtiter assay. The present study detected and confirmed a 550-bp noncoding antisense RNA with the potential to attenuate the activities of the essential response regulator *WalR*. The levels of antisense *walR* RNA transcripts were inversely associated with the production of *WalR* protein. It was showed that overexpression of AS*walR* leads to reduced biofilm formation and exopolysaccharide synthesis. Furthermore, the pathogenicity of *E. faecalis* was markedly decreased by AS*walR* overexpression in an *in vivo* periapical periodontitis model. In summary, the present study detected a novel antisense *walR* RNA that leads to a reduction in biofilm formation and the pathogenicity of *E. faecalis*. Collectively, the data suggest a role for AS*walR* as a post-transcriptional modulator of the *WalR* regulator in *E. faecalis*.

Introduction

Enterococcus faecalis (*E. faecalis*), a generally commensal organism, has emerged as the major pathogen for persistent periapical periodontitis (1). Nonpathogenic (or commensal) bacterial pathogens can switch from the commensal stage to a pathogenic state (2). The virulence of *E. faecalis* is derived from its ability to sense and adapt to varying environmental stresses during colonization and in the host (3). During long-term interaction with the host, the two-component signal transduction system (TCS) of *E. faecalis* regulates the expression of virulence genes in response to microenvironmental conditions, including the host environment, and binds to the regulatory regions of target genes accordingly (4). These two-component signal transduction systems are involved in various cellular processes, including cell viability, virulence, biofilm formation, quorum sensing and antibiotic resistance (5). A typical TCS consists of a histidine sensory kinase membrane receptor and its cognate response regulator (6). The histidine kinase is activated by recognition of environmental stimuli, including pH, oxidative stress, antibiotic pressure and nutrient starvation, and relays the activated phosphoryl group to the response regulator for gene expression modulation by binding at the regulatory regions of target genes (4).

The WalRK (also known as VicRK or YycFG) TCS originally from *Bacillus subtilis* is highly conserved in Gram-positive bacteria and is annotated as VicRK or YycFG in some genera, such as *Staphylococcus aureus*, *Enterococcus faecalis*, and *Streptococcus mutans* (*S. mutans*) (6). The system WalRK has been proposed because of its major roles in the regulation of genes associated with cell wall synthesis and biofilm formation (7). The essential gene *walR*, is closely associated with bacterial growth, virulence and biofilm extracellular polysaccharide synthesis and aggregation (1). Bacterial growth and biofilm aggregation are strongly associated with the process of human infections (4). Consistently, the ability of *E. faecalis* to form the three-dimensional biofilm scaffold contributes to the failure of persistent infected root canal treatments (8). In *E. faecalis*, it was reported that the inability to construct *walR* deletion mutants indicated that this regulatory gene was essential for bacterial viability (4).

Correspondence to: Dr Lei Lei, State Key Laboratory of Oral Diseases, Department of Preventive Dentistry, West China Hospital of Stomatology, Sichuan University, 14 Renmin South Road, Chengdu, Sichuan 610041, P.R. China
E-mail: leilei0504@126.com

Dr Hui Zhang, Department of Orthopedics, West China Hospital, Sichuan University, 37 Guoxue Alley, Chengdu, Sichuan 610041, P.R. China
E-mail: caesarzh@163.com

Key words: *WalR*, antisense RNA, *Enterococcus faecalis*, pathogenicity, post-transcriptional control

A noncoding antisense RNA (AS RNA) can bind to the target messenger RNA (mRNA) by base-pairing, and their interaction forms an RNA duplex structure (9). This AS RNA-induced duplex complex generally inhibits mRNA transcription or translation and performs regulatory functions (10). Using high-throughput transcriptomics analyses, AS RNA regulators can be identified in bacteria (11). An endogenous *vicR* antisense RNA transcript has been identified in *S. mutans* (12). In addition, it was found that the production of VicR protein associated inversely with different levels of *vicR* antisense RNA and that the biofilm biomass decreased in the *vicR* antisense overexpression (12). Considering the close proximity of *E. faecalis* and *S. mutans* in related Gram-positive cocci, it was suggested that there were similar structural and functional relationships between *vicR* in *S. mutans* and *walR* in *E. faecalis*. Additionally, by mapping the *E. faecalis* *walR* gene with BLAST searches, a high similarity (the DNA sequences are 70% identical) with the homologous *vicR* gene in *S. mutans* was demonstrated. In the present study, a potential AS*walR* was hypothesized, and whether the potential AS*walR* was specifically associated with regulation of WalR function was investigated in *E. faecalis*.

Materials and methods

Bacterial strains and growth conditions. The bacterial strains and plasmids used in the present study are listed in Table SI. *E. faecalis* strains purchased from Guangdong Huankai Microbial Science & Technology Co., Ltd. were grown in brain heart infusion (BHI) broth (BD Biosciences) at 37°C in a 5% CO₂ atmosphere with 500 µg/ml spectinomycin when AS*walR* overexpression strains were incubated at 37°C overnight. *E. faecalis* strains were cultured to mid-exponential phase (optical density at a wavelength of 600 nm of OD value=0.5). For antisense overexpression strain construction, the shuttle vector pDL278 (Novagen) was used to overexpress AS*walR* or *walR* sequence under the control of the *walR* gene promoter region. First, the antisense *walR* or *walR* sequences were obtained by oligonucleotide synthesis (Sangon Biotech Co., Ltd.). It was reported that *E. faecalis* V583 carrying the empty vector pDL278 has no effects of the vector itself on the phenotypes examined (13). Next, antisense AS*walR* or *walR* sequences were cloned into the pDL278 plasmid at *Bam*HI and *Eco*RI restriction sites. All recombinant AS*walR* RNA or *walR* gene overexpression plasmids were verified by PCR and DNA sequencing analyses. The recombinant plasmid pDL278AS-*walR* or pDL278*walR* was transformed into *E. faecalis* V583 strains, as previously described (1,12).

Reverse transcription-quantitative PCR (RT-qPCR). Total RNAs were purified from *E. faecalis* strains using a MasterPure RNA purification kit (Epicentre; Illumina Inc.). Residual genomic DNA was removed using Turbo RNase-free DNase I (Ambion; Thermo Fisher Scientific, Inc.). The quality and integrity of RNAs were assessed by 1% agarose gel electrophoresis. The concentration and purity of RNAs were determined using a NanoDrop 8000 spectrophotometer (Thermo Fisher Scientific, Inc.). Total RNAs were extracted from *E. faecalis* V583, AS*walR* and *walR* overexpression strains as aforementioned. Subsequently, purified RNA was

reverse transcribed to cDNA using a PrimeScript RT Reagent kit (Takara Bio, Inc). RT-PCR was conducted for identification and detection of antisense RNA. For AS*walR* detection, a gene-specific primer (PCR1) was used for the first strand cDNA synthesis, and another gene-specific primer (AS2) was used for RT-qPCR analysis (Supplementary Table II). The expression of *walR* and *walK* and virulent factor genes including *ace* (adhesin of collagen), *esp* (protein surface), *epal*, *epaA* (enterococcal polysaccharide antigen), and *gel* (gelatinase) of all *E. faecalis* strains were also examined. RT-qPCR was performed using primers listed in Supplementary Table II on a LightCycler 480 (Roche Diagnostics), following the instructions of the PrimeScript RT Reagent kit (Takara Bio, Inc.). RT-qPCR conditions were: 95°C for 30s, followed by 40 cycles at 95°C for 5s; 60°C for 30s, and then dissociation. The threshold cycle values (CT) were quantified and transcription levels of each gene were compared to the expression of the 16sR gene, which was used as a reference gene (1).

Reverse transcription-PCR (RT-PCR) assays for *walR* co-transcribed operator and antisense RNA detection. Conditions for PCR reactions were: 94°C, 3 min (initial denaturation); 32 cycles at 94°C, 30 sec (denaturation); 55°C, 40 sec (primer annealing); and 72°C, 1 min (primer extension). For potential antisense *walR* RNA detection, total RNAs were purified from *E. faecalis* strains after planktonic growth using a MasterPure RNA purification kit (Epicentre; Illumina Inc.). Residual genomic DNA was removed using Turbo RNase-free DNase I (Ambion; Thermo Fisher Scientific, Inc.). The quality and integrity of RNAs were assessed by 1% agarose gel electrophoresis. The concentration and purity of RNAs were determined using a NanoDrop 8000 spectrophotometer (Thermo Fisher Scientific, Inc.).

Total RNA was prepared from strain *E. faecalis* V583 grown as planktonic culture in BHI and used as templates to validate *walR* and adjacent downstream genes, *walK*, EF1195, EF1196, and EF1197 were co-transcribed (12). The primers for PCR amplification were presented in Table SIII. For first-strand DNA synthesis, *walR* antisense-(PCREf) and sense-specific primers (ASEf) were used. The AS*walR* RT-PCR product was purified with QIAquick PCR Purification Kit according to the manufacturer's instructions and subsequently sequenced. Detection of antisense RNA was performed by 1% agarose gel electrophoresis and RT-PCR assays were performed using a first strand-specific primer for synthesis of cDNA transcripts using the First Strand cDNA synthesis kit (Thermo Fisher Scientific, Inc.) (14). The primers (PCREf) used for first strand synthesis PCR are listed in Table SII. Following the reverse transcription reaction, the first-stand cDNA synthesis products were used as templates. The sense strand primer (ASEf) for PCR amplification is shown in Table SIV.

Northern blotting. Northern blotting assays were performed as previously described with minor modifications (12,15). In brief, 10 µg of total RNA was purified from *E. faecalis* planktonic cultures. The RNA samples were detected by electrophoresis in 1% formaldehyde-agarose gels and transferred to nylon membranes (Amersham; Cytiva). The probe was made specifically (listed in Table SIV) and labeled using the DIG High Prime DNA Labeling and Detection Starter Kit II

(Roche Diagnostics). Blots were incubated 16 h with 20 ng/ml probes at 50°C, and signals were detected using the CSPD Star substrate (Roche Diagnostics).

5'-rapid amplification of cDNA ends (5'-RACE) assay. To find the transcription initiation site and probable termination site of *ASwalR*, total RNA (20 µg) from *E. faecalis* strains was ligated to the 5'-RACE outer adapter from the FirstChoice RLM-RACE Kit (Ambion; Thermo Fisher Scientific, Inc.) according to the manufacturer's instructions (12). The primers used for PCR assays for 5'-RLM-RACE included 5'-RACE gene-specific outer primer and 5'-RACE gene-specific inner primer as listed in Table SIV. The nested PCR thermocycling conditions were as follows: Initial activation of 95°C for 3 min; followed by 35 cycles of denaturation at 94°C for 30 sec, annealing at 56°C for 30 sec and extension at 72°C for 1 min; and a final extension at 72°C for 7 min was performed thereafter. An aliquot (6 µl) of the PCR mixture was assessed by 1% agarose gel electrophoresis using reactants without a cDNA template as the control and sequencing by Sangon Biotech Co., Ltd..

Western blotting. Protein extraction and western blotting were proceeded as previously described (12,13). *E. faecalis* cells grown as planktonic cultures of mid-exponential phase were washed and resuspended in 10 mM Tris-HCl buffer (pH 8.0). Cells were mechanically disrupted by ultrasonication with glass beads (diameter, 0.1 mm) for three cycles of 15 sec with 1 min rest on ice. Clear supernatants were collected by centrifugation (12,000 x g, 2 min, 4°C) and protein concentrations were determined using a Bradford assay (BioRad Laboratories, Inc.). For western blotting, equal amounts of protein (30 µg) were mixed with Laemmli sample buffer (BioRad Laboratories, Inc.) in boiling water for 10 min. The protein samples were loaded on precast 4-20%, gradient gels (Beijing Solarbio Science & Technology Co., Ltd.) and electrotransferred to PVDF membranes (Thermo Fisher Scientific, Inc.). Polyclonal antibodies against r-WalR were produced using the standard 70 days Rabbit Protocol (AbMax Biotechnology Co., Ltd.). The membranes were blocked, then probed with purified WalR-specific rabbit antibody (AbMax Biotechnology Co., Ltd.; <http://www.antibodychina.com/>, cat. no. scu001; 1:1,000) for 2 h at room temperature and incubated with horseradish peroxidase-conjugated goat anti-rabbit secondary antibody (HRP conjugated, cat. no. SSA004; 1:10,000) for 2 h at room temperature. A BioRad GS-700 Imaging Densitometer (BioRad Laboratories, Inc.) was used to determine the signal density of protein signals.

Biofilm assessment. *E. faecalis* V583, *ASwalR* and *walR+* strains were cultured in BHI for 24 h at 37°C as previously described (1,12,13). Crystal violet assay was performed to measure the biomass of each *E. faecalis* biofilm. *E. faecalis* V583 biofilms were set as controls. Briefly, the biofilms were washed three times with PBS. Subsequently, the biofilms were stained with 0.1% (w/v) crystal violet for 15 min at 37°C and destained with 1 ml ethanol/acetone (at a ratio of 8:2) solution. The collected solution was removed to a new plate and measured with a microplate reader (ELX800; Gene Company Ltd.) at a wavelength of 600 nm. For confocal laser scanning

microscopy (CLSM), the *E. faecalis* stains in biofilm were labeled with SYTO9 (Invitrogen; Thermo Fisher Scientific, Inc.) and the extracellular polymeric substance (EPS) matrix was stained with an Alexa Fluor 647-labeled dextran conjugate (Invitrogen; Thermo Fisher Scientific, Inc.) (1). Three-dimensional reconstruction of the biofilms and the EPS/bacterial ratio were analyzed using Imaris 7.0 software (Bitplane; Oxford Instruments).

Periapical periodontitis lesions in animal experiments. Animal experiments were approved by the Biomedical Research Ethics Committee of West China Hospital (approval no. 2019128A) and conducted according to the guidelines of Institutional Animal Care and Use Committee (IACUC) for animal care and use of laboratory animals (16). A total of 20 6-week-old female Sprague-Dawley rats (260-280 g) were included in the study and anesthetized with ketamine/xylazine (90 and 10 mg/kg, respectively) by intraperitoneal injection. On the occlusal surface of the left mandible first molars, an access opening was made using a high-speed round bur (Aseptico Inc.). The right first mandible molar was used as a blank control. Next, the root canal preparation was performed with sterile K-files (0.02 taper, sizes #15-#35). Subsequently, 50 µl log-phased *E. faecalis* V583 bacterial suspensions were inoculated into the pulp chamber and sealed with light-cured flowable resin (3M Filtek™ Z350XT; 3M) in the V583 group (n=10). For the *ASwalR* group (n=10), 50 µl log-phased *ASwalR E. faecalis* bacterial suspensions were applied for establishment of an infection model. After 4 weeks of operation, the rats were scarified by euthanasia under deep anesthesia using ketamine/xylazine by cervical dislocation, and the rats were imaged using the Quantum GX Micro-CT System (PerkinElmer, Inc.). The scanning conditions were as follows: kV=90; CT µA=72; 360° scan time=8 sec. The reconstructed images were analyzed with Analyze 12.0 (PerkinElmer, Inc.).

HE and Gram staining and PNA-FISH. For histological evaluation and peptide nucleic acid fluorescence *in situ* hybridization (PNA-FISH), the mandibular bone was longitudinally split into two parts. The specimens were prepared for histological evaluation. The mandibular bone was fixed in 10% neutral buffered formalin for 72 h at room temperature, decalcified in 10% EDTA and embedded in paraffin. The 5-µm slices were prepared for hematoxylin and eosin (HE) and Gram staining for tissue assessment with a light microscope at x40 magnification. For PNA-FISH, bone specimens were also fixed for 72 h in 10% neutral buffered formalin and decalcified in 10% EDTA decalcified for 3 weeks before being embedded in paraffin blocks and sectioned to 3 µm thickness. Smears were deparaffinized in xylene (2x5 mins) and rehydrated by a graded ethanol series (100, 95, 80, 70 and 50%) for 5 min each time and permeated with proteinase K (Wuhan ServiceBio Technology Co., Ltd.) at 37°C for 25 min. Finally, all smears were washed up with distilled water for 10 min and allowed to air dry.

The above slides were fixed in 100% methanol for 10 min at 37°C, followed by immersion in 80% ethanol for 10 min. For pre-hybridization, one drop of hybridization solution without probe containing 10% (w/v) dextran sulfate (Thermo Fisher Scientific, Inc.), 10 mM NaCl (Panreac), 0.2% (w/v)

polyvinylpyrrolidone (Sigma-Adrich; Merck KGaA), 0.2% (w/v) Ficoll (Thermo Fisher Scientific, Inc), 5 mM disodium EDTA (PanReac AppliChem), 0.1% (v/v) Triton X-100 (PanReac AppliChem), 50 mM Tris-HCl (pH 7.5; Thermo Fisher Scientific, Inc.) was applied for 1 h at 37°C. From the template of genomic DNA a fluorescein amidite-labeled PNA probe (5'-GGTGTGTTAGCATTTTCG-3') targeting *E. faecalis* 16S rRNA (Wuhan Servicebio Technology Co., Ltd.) was applied. The slides were then incubated with hybridization solution containing 200 nM PNA probe overnight at 37°C. Subsequently, the slides were co-cultured in saline sodium citrate washing buffer (Wuhan ServiceBio Technology Co., Ltd.) at 37°C for 30 min and dried in air. Bone smears were then stained with 100 μ l DAPI (Thermo Fisher Scientific, Inc.) in the dark for 30 min at room temperature. The samples were washed with PBS buffer twice and dried in air. The slides were mounted with Gold Antifade reagent (Thermo Fisher Scientific, Inc.) and stored at 4°C for further investigation with fluorescent microscope at 40 magnification.

Statistical analysis. Bartlett's test was conducted to analyze the homogeneity of data variances and Shapiro-Wilk test was applied to determine the normal distribution of data. For parametric testing, one-way ANOVA was performed followed by pairwise multiple comparisons of Tukey's test using SPSS software 18.0 (SPSS, Inc.). Data are presented as the mean \pm SD. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

WalR gene operon contains an antisense walR RNA. Downstream of the *walR* (EF1193) in *E. faecalis* V583, a cluster of four genes (EF1194 to EF1197) is co-transcribed (Fig. 1A). Using a series of overlapping primers, RT-PCR indicated co-transcription comprising *walR* and the four downstream genes (Fig. 1B), indicating that *walR*, *walK*, EF1195, EF1196, and EF1197 are part of the WalR regulon. RNA agarose gel electrophoresis for northern blotting was included in Fig. S1.

To investigate whether a potential antisense RNA was specifically associated with *walR*, first strand cDNA synthesis was performed to detect the transcript in *E. faecalis* V583. The sense and antisense sequences of the *walR* gene (EF1193) are shown in Fig. 2A. To determine the 5'-terminus of AS*walR* RNA, 5'-RACE assays of AS*walR* were conducted. The primers used and the design and position of primers for 5'-RACE is indicated in Fig. 2A. The sequencing analyses indicated that the potential AS*walR* contained 550 bp of antisense sequence in the *E. faecalis* V583 genome by RT-PCR assays (Fig. 2B). Western blotting showed that the expression of WalR protein was inversely associated with different levels of AS*walR* transcripts (Fig. 2C). The size of the 5'-RACE PCR product was ~120 bp (Fig. 3A). Additionally, the differences in AS*walR* abundance in *E. faecalis* planktonic growth were confirmed by northern blotting (Fig. 3B). The sequence predicted that the 5'-terminus of AS*walR* begins at and covers most of the *walR* coding sequence (Fig. 3C).

Antisense walR RNA negatively affects the production of WalR and suppresses biofilm formation. Using CLSM visualization

of double-stained biofilms, it was found that overproducing AS*walR* markedly decreased EPS production compared with *E. faecalis* V583 strain, while biofilm cells were packed within the enriched EPS matrix, which was stained red in the *walR*+ strains, similar to the results obtained with the V583 parent strain (Fig. 4A). These findings were further confirmed by quantitation of data revealing that AS*walR* exhibited the lowest EPS/bacterial biomass volume ratio, indicating a role for the *walR* gene in EPS architecture development (Fig. 4B). Next, the effects of AS*walR* overexpression on the ability to form biofilms were evaluated. *E. faecalis* V583, AS*walR* and *walR*+ strains were allowed to form biofilms in BHI for 24 h. The biomass was quantified by a crystal violet microtiter assay, and the results indicated that overexpression of AS*walR* resulted in a 45% decrease in biofilm growth compared with the V583 parent strain (Fig. 4C). However, the biomass of the *walR*+ strain was slightly but non-significantly increased compared with the *E. faecalis* V583 biofilm (Fig. 4C). The expression levels of AS*walR* and *walR* were quantified, which indicated that the transcripts of AS*walR* in the AS*walR* strain were 2.7-fold higher compared with V583 parent cells, while the transformation of a pDL278 empty vector did not affect the levels of *walR* mRNA (1). The results showed that the levels of transcripts of two glycosyltransferase genes, *epal* and *epaA*, were significantly decreased in the AS*walR* strain when compared with *E. faecalis* V583 strain (Fig. 4D).

WalR antisense inhibited the pathogenicity and reduced Periapical Lesion Size. The micro-CT results showed that the severity of bone destruction was markedly higher in the *E. faecalis* V583-infected group compared with the AS*walR*-treated group (Fig. 5A). Quantitatively, the average relative ratio of the periapical cavity was ~2.5% in the AS*walR*-treated group compared with the right first mandible molar as a blank control, which was significantly lower than that of *E. faecalis* V583-treated group (Fig. 5B). This trend indicated that AS*walR*-transformed strains presented a limited capability to infarct infected bone tissues. Histological assessments in HE-stained samples showed bone tissue absorption and inflammatory infiltration in the *E. faecalis* V583-colonized groups, which were more severe compared with control and AS*walR* groups (Fig. 5C; left column). Gram staining results showed that more Gram-positive cells were identified within the bone of the *E. faecalis* V583 group compared with the AS*walR* group (Fig. 5C; middle column). After labeling with PNA fluorescent probes, bacterial cells were specifically detected by green fluorescence. The control group indicated no Gram-positive cells were inoculated (Fig. 5C; right column). The fluorescence intensity of the *E. faecalis* group was higher compared with AS*walR* strain-infected samples. The control group indicated no fluorescence intensity of the *E. faecalis* were observed.

Discussion

The WalRK (also known as VicRK or YycFG) two-component signal transduction system is highly conserved in Gram-positive bacteria, which originated from *Bacillus subtilis* (6). In *E. faecalis*, the downregulation of *walR* led to a reduction in the dextran-dependent aggregation in the biofilm and




Figure 1. WalR regulon and downstream genes in *Enterococcus faecalis* V583. (A) Schematic of the gene locus and RT-PCR primer sets. Downstream of *walR*, a cluster of five genes (EF1194 to EF1197) is similarly transcribed in the same orientation. (B) RT-PCR with lanes numbered 1-4 according to primer sets used. Total RNA was used as the negative control for all the PCR reactions, indicating RT was not contaminated with genomic DNA. RT, reverse transcription; F, forward; R, reverse. RNA agarose gel electrophoresis for Northern blotting was included in the Supplementary Fig. 1.

A *Enterococcus faecalis* *walR* [1159770 – 1160474 bp] *walR* sense/antisense strands

5' GTGAAGAAAA TTTTAGTAGT TGATGACGAG AAGCCAATTT CAGAGATCGT TAAATATAAT TTGGTTAAAG AAGGATATGA
3' CACTTCTTTT AAAATCATCA ACTACTGCTC TTCGGTTAAA GTCTCTAGCA ATTTATATTA AACCAATTC TTCCTATACT
→ PCREf

AGTATTTACT GCTTATGATG GAGAAGAAGC ACTTGAAAAA GTGAAGAAG TGAACACAGA CTTAATTATT TTAGACTTAA
TCATAAATGA CGAATACTAC CTCTTCTCG TGAACTTTTT CACCTTCTTC ACCTTGGTCT GAATTAATAA AATCTGAATT

TGCTCCCTAA AATGGATGGC TTAGAAGTCG CGCGAGAAGT GCGCAAAACA CATGATATGC CAATCATTAT GGTGACTGCC
ACGAGGGATT TTACCTACCG AATCTTCAGC GCGCTCTTCA CGCGTTTTGT GTACTATACG GTTAGTAATA CCACTGACGG

AAAGATTCTG AAATTGATAA GGTTTTAGGA TTGAATTAG GAGCCGATGA CTATGTAACG AAACCATTTT CAAATCGTGA
TTTCTAAGAC TTTAACTATT CCAAAATCCT AACCTTAATC CTCGGCTACT GATACATTGC TTTGGTAAAA GTTTAGCACT

ATTAGTTGCT CGTGTAAGA CCAATTTACG GCGAGGTGCA ACCAATGCGA AAGAAGCCGA GGTGACAACA CAATCTGAAT
TAATCAACGA GCACATTTTC GGTAAATGC CGCTCCAGT TGGTTACGCT TTCTTCGGCT CCACTGTTGT GTTAGACTTA

TAACGATTGG TGATTAAAC ATTATCCTG ATGCATACAT GGTCTCAAAA CGGGGTGAAA AAATTGAATT AACCCACCGT
ATTGCTAACC ACTAAATTGG TAAGTAGGAC TACGTATGTA CCAGAGTTTT GCCCCTTT TTTAACTTAA TTGGGTGGCA

GAATTGAGT TACTTTATTA CTTAGCAAAA CATATCGGAC AAGTGATGAC TCGTGAACAT TTATTACAAA CCGTTTGGGG
CTTAAATCA ATGAATAAT GAATCGTTTT GTATAGCCTG TTCACTACTG AGCACTTGTA AATAATGTTT GGCAAAACCC

TTATGATTAT TTTGGGGATG TCGGACAGT GGACGTAACC GTACGTCGTT TAAGAGAAAA AATTGAAGAT AGTCCAAGTC
AATACTAATA AAACCCCTAC AGCCTGTCA CCTGCATTGG CATGCAGCAA ATTCTCTTTT TTAACCTCTA TCAGGTTTCAG

ATCCAACGTA TTTAGTTACT CGTCGTGGG TTGGTTATTA TCTAAGAAAT CCTGAACAGG AGTAA
TAGGTTGCAT AAATCAATGA GCAGCACCCC AACCAATAAT AGATTCTTTA GGACTTGTC TCATT


→ 5'-RACE gene-specific outer primer

→ 5'-RACE gene-specific inner primer

ASEf

Initial site for AS RNA by 5'RACE

B RT-PCR for AS*walR*



C




Figure 2. Detection of *walR* antisense RNA. (A) Primer design for detection of antisense *walR* RNA. (B) Differences in AS*walR* RNA abundance in *Enterococcus faecalis* cultures by first cDNA strand synthesis and RT-PCR using *walR* gene-specific anti-sense and sense strand primers. The 16S gene was used as an internal standard. (C) WalR production blot was quantified in the cells grown as aforementioned from western blots probed with anti-WalR antibody. PCREf, *walR* antisense-specific primer; ASEf, *walR* sense-specific primer; 5'-RACE, 5'-rapid amplification of cDNA ends; AS, antisense; RT-PCR, reverse transcription-PCR; AS*walR*, *walR* antisense RNA.




Figure 3. Structure of the ASwalR transcript and association with *walR* mRNA. (A) Detection of the 5'-terminus of the ASwalR transcript by 5'-RACE. The PCR product size was ~120 bp. (B) Transcript amounts of ASwalR in *Enterococcus faecalis* cultures were observed in a representative northern blot. The loading sample without RNA was used as the blank control. (C) Schematic of the ASwalR 5'-terminus showing that transcription starts within the *walR* ORF and is complementary to most of the *walR* ORF. The sequence of the first strand synthesis ASwalR RT-PCR amplicon was from 1159871-1160384 bp. The overlapping sequence from first strand-specific ASwalR RT-PCR and 5'-RACE product (89 bp) was from 1159871-1159960 bp in the genome. The full length for ASwalR RNA is 550 bp (1159834-1160384 bp). ASwalR, *walR* antisense RNA; ORF, open reading frame; RT-PCR, reverse transcription-PCR; 5'-RACE, 5'-rapid amplification of cDNA ends; AS, antisense.

inhibited the transcription of virulence genes (1). Several genes in the region of the *walR* locus were assigned to the WalR operon as RT-PCR results indicated co-transcription comprising *walR* and the four downstream genes, including *walK*, EF1195, EF1196, and EF1197. The proximity of *walK* to *walR* suggested a close interaction and role in the regulation of WalRK TCS (17). The function annotated for the EF1195 gene, a hypothetical protein belonging to the conserved protein domain family YchH, is associated with the regulatory protein YchH in the two-component signal transduction system YycFG (18). The most distal genes, EF1196 and EF1197, in the extended WalR regulon encoded a putative Ycl protein and a metallo-beta-lactamase, YycJ, which probably plays a role in regulating protein function by lactamization (19). Although the information of these genes regulated by WalR remains limited, the close proximity of the other genes in the extended WalR regulon suggested a functional relationship, including cell wall biogenesis and biofilm formation, which also needs further investigation.

The evidence has revealed that substantial antisense transcripts, including intergenic and intragenic transcription, exist in the bacteria (11). A cis antisense noncoding RNA to the *walR* gene (ASwalR) was identified and confirmed by northern blotting and 5'-RACE. The effect of ASwalR on decreasing and inhibiting *walR* gene expression and associated production of WalR was further detected using following RT-PCR and western blotting. Similarly, it was shown that antisense RNA could combine with its reverse complementary sequences and restricted the expression of targeted mRNA (12). Based on this, the present study hypothesized an inverse association between ASwalR levels and *walR* transcripts and WalR production. The inverse association observed is consistent with the ASwalR-*walR* mRNA complex blocking its transcription and translation (20).

To determine whether ASwalR RNA activity played a role in biofilm formation, the present study constructed ASwalR and *walR* overexpression mutant strains to inhibit ASwalR




Figure 4. ASwalR overexpression suppresses *E. faecalis* biofilm aggregation and EPS production. (A) Double labeling of the biofilms in the *E. faecalis* V583, ASwalR and walR+ strains. Green, bacterial cells (SYTO9); red, EPS matrix (propidium iodide, PI). Scale bar, 100 μ m. (B) Percentage of EPS matrix/bacteria ratio in biofilm growth. n=10. *P<0.05. (C) Biomass was quantified by crystal violet staining. n=10. *P<0.05. (D) Reverse transcription-PCR analysis showed the gene transcripts in *E. faecalis* V583, ASwalR, and walR+ strains. *E. faecalis* gene expression was quantified using 16sR as an internal control and calculated based on the *E. faecalis* V583 expression, which was set as 1.0. n=10. *P<0.05 when compared to the *E. faecalis* V583. ASwalR, walR antisense RNA; EPS, exopolysaccharide; *E. faecalis*, *Enterococcus faecalis*; OD, optical density; Ace, adhesin of collagen; esp, protein surface; epal, epaA, enterococcal polysaccharide antigen; gel, gelatinase.

levels. Our previous results suggested that the interference of the walR gene suppressed the expression of virulent genes associated with cellular adhesion and biofilm growth (1). In *E. faecalis*, glycosyltransferase is a type of enzyme involved in the synthesis of polysaccharides by the transfer of a sugar precursor onto sugar residues (21). Biofilm is one of major factors for infection development and processes (1,13). By overexpression of ASwalR, the levels of two glycosyltransferase

genes, epal and epaA were significantly reduced, which contributed to impaired EPS production in the ASwalR biofilm and decreased biomass of ASwalR overproducing strains. In addition, the transcripts of virulence genes were lower in all ASwalR-treated groups. Therefore, the ability of ASwalR strains to induce periapical infections was reduced compared to the parent V583 strains, demonstrated by histological evidence from HE and Gram staining (1). The effects of walR




Figure 5. *ASwalR* overexpression reduces the periapical lesion size and inhibits bacterial aggregation. (A) Reconstructed images for micro-CT scanning of rat periapical lesions. (B) Relative periapical cavity levels (%) were calculated $n=10$. * $P<0.05$. (C) Samples were obtained from infected periapical lesions at 4 weeks. HE-stained histological slices (left lane; scale bars, 100 μm) and Gram-stained samples (middle lane; scale bars, 100 μm). The red boxes indicated inflammatory cells in the HE-staining or bacteria in the Gram-staining. The presence of fluorescent *Enterococcus faecalis* was identified with a PNA-FISH probe for bacterial 16S rRNA (right lane; scale bars, 100 μm). *ASwalR*, *walR* antisense RNA; HE, hematoxylin-eosin; PNA-FISH, peptide nucleic acid-fluorescence *in situ* hybridization.

overexpression on both bacterial growth and EPS production were similar to the V583 parental strain levels, justifying additional studies.

These analyses of biofilm formation again confirm that the *ASwalR* RNA functions in biofilm formation by *E. faecalis*. Next, the role of *ASwalR* RNA in the pathogenicity of

E. faecalis was validated using a Sprague-Dawley rat model. The results indicated that AS*walR* overexpression markedly reduced the periapical lesion size at 4 weeks compared to the *E. faecalis* V583 group, which presented large periapical lesions. In the present study, restricting the periapical lesions in AS*walR*-treated group was considered an important step for stimulating the periapical repair process (22). Additionally, histological evidence from HE and Gram staining demonstrated that the ability of AS*walR* strains to cause periapical infections was reduced compared with the parent V583 strains. Furthermore, the presence of *E. faecalis* bacterial cells involved in periapical tissues was identified by PNA-FISH, which showed higher fluorescence intensity in the parent V583 strains. It was hypothesized that the pathogenicity of *E. faecalis* was markedly decreased by AS*walR* overexpression in periapical periodontitis. However, one limitation of the present study is the lack of direct evidence showing whether AS*walR*-mRNA duplexes can be disrupted by additional RNase activities. Bioinformatic studies for the AS*walR* transcript started in the opposite direction to *walR*, which was reverse complementary to the *walR* transcript (BPRM at <http://Linux1.softberry.com/berry.phtml>). A putative-35 box, TTCACA, and a-10 box, TTCTAATCT. Future research is needed to construct a vector which contains the entire AS*walR* gene and the promoter of the AS*walR* to identify the transcriptional initial site of AS*walR*.

In summary, the downstream genes *walK*, EF1195, EF1196, and EF1197 together with the *walR* gene are co-transcribed in *E. faecalis* V583. The present study detected and confirmed a 550-bp noncoding antisense RNA with the potential to attenuate the activities of the essential response regulator WalR. The levels of antisense *walR* RNA transcripts were inversely associated with the production of WalR protein. In addition, the present study showed that overexpression of AS*walR* led to a reduction in biofilm formation and EPS synthesis. Furthermore, the results of animal experiments revealed that the pathogenicity of *E. faecalis* was markedly decreased by AS*walR* overexpression in periapical periodontitis. Collectively, the data suggested a role for AS*walR* as a post-transcriptional modulator of the WalR regulator in *E. faecalis* and revealed that preserving the novel antisense *walR* RNA will be a potential substitutive therapy for *E. faecalis* infections, especially in the treatment of periapical periodontitis.

Acknowledgements

Not applicable.

Funding

This study was supported by the Natural Science Foundation of China (grant no. 81800964), the Sichuan Provincial Natural Science Foundation of China (grant nos. 2018SZ0125 and 2019YFS0270).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

SW contributed to conception, design and interpretation of data, drafted and critically revised the manuscript. YL contributed to design, and data interpretation, drafting manuscript and critical revision the manuscript. LL contributed to conception, design, and interpretation of data, drafted and critically revised the manuscript. HZ contributed to conception, design, and interpretation of data, drafted and critically revised the manuscript. All authors gave their final approval and agree to be accountable for all aspects of the work. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The West China Hospital of Sichuan University Biomedical Research Ethics Committee approved the animal experiments for this investigation (approval no. 2019128A).

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Wu S, Liu Y, Zhang H and Lei L: The susceptibility to calcium hydroxide modulated by the essential *walR* gene reveals the role for *Enterococcus faecalis* biofilm aggregation. *J Endod* 45: 295-301.e2, 2019.
2. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, Lakshmanan A and Wade WG: The human oral microbiome. *J Bacteriol* 192: 5002-5017, 2010.
3. Colomer-Winter C, Flores-Mireles AL, Baker SP, Frank KL, Lynch AJL, Hultgren SJ, Kitten T and Lemos JA: Manganese acquisition is essential for virulence of *Enterococcus faecalis*. *PLoS Pathog* 14: e1007102, 2018.
4. Hancock LE and Perego M: Systematic inactivation and phenotypic characterization of two-component signal transduction systems of *Enterococcus faecalis* V583. *J Bacteriol* 186: 7951-7958, 2004.
5. Gotoh Y, Eguchi Y, Watanabe T, Okamoto S, Doi A and Utsumi R: Two-component signal transduction as potential drug targets in pathogenic bacteria. *Curr Opin Microbiol* 13: 232-239, 2010.
6. Fukuchi K, Kasahara Y, Asai K, Kobayashi K, Moriya S and Ogasawara N: The essential two-component regulatory system encoded by *ycfF* and *ycfG* modulates expression of the *ftsAZ* operon in *Bacillus subtilis*. *Microbiology* 146: 1573-1583, 2000.
7. Dubrac S, Bisicchia P, Devine KM and Msadek T: A matter of life and death: Cell wall homeostasis and the *WalKR* (YycGF) essential signal transduction pathway. *Mol Microbiol* 70: 1307-1322, 2008.
8. Fisher K and Phillips C: The ecology, epidemiology and virulence of *Enterococcus*. *Microbiology* 155: 1749-1757, 2009.
9. Krinke L and Wulff DL: RNase III-dependent hydrolysis of lambda cII-O gene mRNA mediated by lambda OOP antisense RNA. *Genes Dev* 4: 2223-2233, 1990.
10. Saberi F, Kamali M, Najafi A, Yazdanparast A and Moghaddam MM: Natural antisense RNAs as mRNA regulatory elements in bacteria: A review on function and applications. *Cell Mol Biol Lett* 21: 6, 2016.
11. Rasmussen S, Nielsen HB and Jarmer H: The transcriptionally active regions in the genome of *Bacillus subtilis*. *Mol Microbiol* 73: 1043-1057, 2009.
12. Lei L, Stipp RN, Chen T, Wu SZ, Hu T and Duncan MJ: Activity of *Streptococcus mutans* VicR is modulated by antisense RNA. *J Dent Res* 97: 1477-1484, 2018.

13. Ho EC, Donaldson ME and Saville BJ: Detection of antisense RNA transcripts by strand-specific RT-PCR. *Methods Mol Biol* 630: 125-138, 2010.
14. Stipp RN, Boisvert H, Smith DJ, Höfling JF, Duncan MJ and Mattos-Graner RO: CovR and VicRK regulate cell surface biogenesis genes required for biofilm formation in *Streptococcus mutans*. *PLoS One* 8: e58271, 2013.
15. Wu S, Huang F, Zhang H and Lei L: *Staphylococcus aureus* biofilm organization modulated by YycFG two-component regulatory pathway. *J Orthop Surg Res* 14: 10, 2019.
16. Couto M and Cates C: Laboratory guidelines for animal care. *Methods Mol Biol* 1920: 407-430, 2019.
17. Aakra A, Vebø H, Snipen L, Hirt H, Aastveit A, Kapur V, Dunne G, Murray BE and Nes IF: Transcriptional response of *Enterococcus faecalis* V583 to erythromycin. *Antimicrob Agents Chemother* 49: 2246-2259, 2005.
18. Paulsen IT, Banerjee L, Myers GS, Nelson KE, Seshadri R, Read TD, Fouts DE, Eisen JA, Gill SR, Heidelberg JF, *et al*: Role of mobile DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. *Science* 299: 2071-2074, 2003.
19. Wecker P, Klockow C, Ellrott A, Quast C, Langhammer P, Harder J and Glöckner FO: Transcriptional response of the model planctomycete *Rhodopirellula baltica* SH1(T) to changing environmental conditions. *BMC Genomics* 10: 410, 2009.
20. Krinke L and Wulff DL: OOP RNA, produced from multicopy plasmids, inhibits lambda cII gene expression through an RNase III-dependent mechanism. *Genes Dev* 1: 1005-1013, 1987.
21. Dale JL, Cagnazzo J, Phan CQ, Barnes AM and Dunne GM: Multiple roles for *Enterococcus faecalis* glycosyltransferases in biofilm-associated antibiotic resistance, cell envelope integrity, and conjugative transfer. *Antimicrob Agents Chemother* 59: 4094-4105, 2015.
22. Siddiqui YD, Omori K, Ito T, Yamashiro K, Nakamura S, Okamoto K, Ono M, Yamamoto T, Van Dyke TE and Takashiba S: Resolvin D2 induces resolution of periapical inflammation and promotes healing of periapical lesions in rat periapical periodontitis. *Front Immunol* 10: 307, 2019.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.