

Association of vitamin D receptor gene TaqI polymorphism with Alzheimer's disease in a Southeastern European Caucasian population

EFTHIMIOS DIMITRAKIS¹, MARTHA-SPYRIDOULA KATSAROU¹, MARIA LAGIOU¹,
VASILIKI PAPASTEFANOPOULOU², EVANGELIA STANITSA³, DEMETRIOS A. SPANDIDOS⁴,
ARISTIDIS TSATSAKIS⁵, SOCRATIS PAPAGEORGIOU³, PARASKEVI MOUTSATSOU²,
KATERINA ANTONIOU⁶, CHRISTOS KROUPIS² and NIKOLAOS DRAKOULIS¹

¹Research Group of Clinical Pharmacology and Pharmacogenomics, Faculty of Pharmacy, School of Health Sciences, National and Kapodistrian University of Athens, 15784 Athens; ²Department of Clinical Biochemistry and
³Second Department of Neurology, University General Hospital 'ATTIKON', 12462 Athens;
⁴Laboratory of Clinical Virology and ⁵Department of Forensic Sciences and Toxicology, Medical School, University of Crete, 71003 Heraklion; ⁶Department of Pharmacology, Faculty of Medicine, School of Health Sciences, University of Ioannina, 45110 Ioannina, Greece

Received January 29, 2022; Accepted March 2, 2022

DOI: 10.3892/etm.2022.11271

Abstract. The role of vitamin D in Alzheimer's Disease (AD) has been studied over the past years. The results from numerous studies have indicated that the molecular pathways involved in the development of AD are closely related to the molecular pathways of the mechanisms of action of vitamin D. However, only a limited number of studies have described the key role of vitamin D receptor (VDR) in the regulation of the functions of vitamin D and the potential effect of single nucleotide polymorphisms (SNPs) of the *VDR* gene. Thus, the aim of the present study was to investigate the *VDR* TaqI polymorphism in relation to AD in a Southeastern European Caucasian (SEC) cohort. Further, the present study aimed to compare the results obtained with those of other AD populations. For this purpose, blood samples from 90 confirmed patients with AD [median age, 74 years; median mini-mental state examination (MMSE) score of 21; median frontal assessment battery (FAB) score of 10] and 103 healthy controls (median age, 57 years) were

analyzed to determine the genotypes of TaqI (rs731236) using quantitative PCR. The frequencies (%) of the TaqI TT, TC and CC genotypes in the controls/patients were 34/48.9, 47.6/41.1 and 18.4/10.0, respectively. Statistically significant differences were observed for the TaqI C allele [odds ratio (OR), 0.54; 95% confidence interval (CI), 0.30-0.96; $P=0.035$], the TaqI TT genotype (OR, 1.86; 95% CI, 1.04-3.32; $P=0.035$) and the TaqI CC genotype (OR, 0.119; 95% CI, 0.014-0.995; $P=0.032$), in relation to the MMSE score <21 in the patient's group. The TaqI TT allele was found to increase the risk of developing AD by 1.86-fold in the SEC population, while the TaqI C allele may act protectively, with a 46% lower risk of developing the disease. Patients with the TaqI CC genotype were found to have an 88% less likelihood of developing severe cognitive impairment based on the MMSE score. On the whole, the present study did not confirm the results of previous studies on the *VDR* TaqI C allele in patients with AD.

Introduction

Alzheimer's disease (AD) is a neurodegenerative disorder contributing to 60-70% of cases of dementia globally, based on the estimations of the World Health Organization (WHO) (1). Vitamin D is an established molecule which is crucial for the development of neural cells and the nervous system (2,3). A number of studies have reported that vitamin D deficiency is associated with a higher risk of developing dementia and neurodegenerative disorders, such as AD (4-8). In parallel, the beneficial effects of vitamin D in neural cells have been reported in a number of studies that have investigated its functions in relation to nerve growth factor, inducible nitric oxide synthase, amyloid β -peptide clearance and calcium homeostasis (9-14). Vitamin D receptor (VDR) is the critical protein involved in the mediation of the functions of vitamin D (15).

Correspondence to: Professor Nikolaos Drakoulis, Research Group of Clinical Pharmacology and Pharmacogenomics, Faculty of Pharmacy, School of Health Sciences, National and Kapodistrian University of Athens, Panepistimioupolis, 15784 Athens, Greece
E-mail: drakoulis@pharm.uoa.gr

Abbreviations: VDR, vitamin D receptor; AD, Alzheimer's disease; MMSE score, mini-mental state examination score; LOAD, late-onset Alzheimer's disease

Key words: Alzheimer's disease, vitamin D receptor, polymorphism, MMSE, TaqI

Over the past few years, a number of studies have investigated the potential association of vitamin D with the development of late-onset AD (LOAD) in various populations, focusing on the *VDR* gene single nucleotide polymorphisms (SNPs) (16-22). One reason for which the *VDR* gene has received increasing attention is that it is included in the risk locus in chromosome 12, which was previously reported by genetic studies to be linked to LOAD (23,24). Vitamin D functions as a transcription factor after binding to VDR and the large number of transcription binding sites, that have been estimated up to 10,000 per cell, are another good basis for researching the potential effects of *VDR* variants in association with the functions of vitamin D in the nervous system and in the development of LOAD (6,25). However, the initial indication for a potential association of the disease with *VDR* was described in an older study, which reported lower *VDR* mRNA levels in hippocampal and pyramidal cells of patients with AD (26).

In addition, to date, only a limited number of studies have been able to provide knowledge regarding the molecular mechanisms or pathways through which vitamin D and VDR affect the development of AD. A previous study concluded that vitamin D and VDR regulated the production of amyloid β , and therefore the development of the disease through the expression of the proteins involved in secretases (25). Information regarding the potential pathways and genes affected by vitamin D and VDR was also provided through an *in vivo* animal study, which concluded that there was a close connection between vitamin D molecular pathways and Alzheimer's molecular pathways through certain groups of genes affecting amyloid β production, neuroinflammation or other AD-related molecules (27,28). The molecular pathways involved in the mechanisms of action of vitamin D affect proteins that play a key role in the development of a diverse group of diseases, including diabetes, autoimmune, cardiovascular and malignant diseases, Parkinson's disease and AD. One indicative example can be described for the protein kinase B (Akt) pathway. Akt phosphorylation is decreased by PTEN in AD, leading to the progression of the disease (29). In parallel, it has been reported that vitamin D through VDR, stimulates the Akt signaling pathway, leading to a potential attenuation of the progression of the disease (29,30). Overall, the molecular pathways of vitamin D and VDR associated with the development of LOAD have not yet been fully investigated. Moreover, the potential effects of the *VDR* gene SNPs on these pathways have not yet been fully described. *VDR* polymorphisms have been also investigated in specific neurological diseases and conditions, such as multiple sclerosis, Parkinson's disease and cluster headaches, with the results indicating possible associations (31-33).

Thus far, the results from studies investigating *VDR* SNPs and LOAD in various populations have been contradictory. One of the most extensively studied *VDR* SNP is TaqI (rs731236, c.1056T>C, ATT>ATC, p.Ile, NM_000376.3) which is located in exon 9 and is included in the ligand binding site of the gene. The TaqI polymorphism does not result in an amino acid change of the receptor protein, and has been proposed as a genetic biomarker for AD; however, the results differ according to the ethnic population reported in each study (16,34).

The aim of the present study was to investigate potential associations of the single nucleotide variation TaqI of the

VDR gene with the development of LOAD in a Southeastern European Caucasian (SEC) cohort. The present study also wished to compare the observed results with the data from other populations and to evaluate the potential association of this variation with neuropsychology mini-mental state examination (MMSE) and frontal assessment battery (FAB) assessments.

Subjects and methods

Study subjects. The study sample included 90 patients with well-ascertained LOAD (median age, 74 years; range, 51-92 years; males, 48.9%; females, 51.1%; median MMSE score of 21; median FAB score of 10) and 103 healthy controls (median age, 57 years; range, 51-90 years; males, 49.5%; females, 50.5%). LOAD diagnosis was based on current diagnostic criteria for AD, including a physical examination, MMSE/FAB score, a brain CT/MRI scan and amyloid β , tau and p-tau levels in the cerebrospinal fluid. Patients or healthy control subjects not belonging to the SEC population were not included in the study. Patients were recruited from the Outpatient Clinic of the Cognitive Disorder-Dementia Unit of the Second Department of Neurology at the University General Hospital 'ATTIKON' (Athens, Greece). Sample collection took place from January, 2018 to February, 2019. The demographic data of the patient and control groups are presented in Table I. The present study has been approved by the Scientific Council and Bioethics Committee of the University General Hospital 'ATTIKON' (Reg. no. 312; December 21, 2017). Written informed consent for participation in the study and the use of their genetic data was obtained from all participants.

DNA isolation and quantitative PCR (qPCR). Blood samples from the patients and controls were analyzed to determine the genotypes of the SNP TaqI (rs731236) of the *VDR* gene. DNA extraction was performed from 200 μ l whole blood samples using the NucleoSpin® Genomic DNA from Tissue kit (Macherey-Nagel GmbH & Co. KG). Genotypes of the TaqI (rs731236) polymorphism were determined using qPCR (using the LightCycler® 480 system; Roche Diagnostics) with the LightSNiP (SimpleProbe®) assay (TIB Molbiol). For qPCR, an initial polymerase activation and denaturation step at 95°C for 10 min was followed by 45 amplification cycles for each sample in the LightCycler instrument. Cycles included denaturation (95°C for 10 sec), annealing (60°C for 10 sec) and extension (72°C for 15 sec). Fluorescence was measured at the end of the annealing period of each cycle to monitor the progress of amplification. Following amplification, melting curves were created by cooling/holding temperature at 40°C for 30 sec and gradually increasing the temperature up to 95°C. Melting peaks were produced accordingly based on the fluorescence signal using LightCycler® 480 software, version 1.5. Differences in the hybridization between the SimpleProbe® oligonucleotide and the DNA sequence (due to the presence of the polymorphism) result in different melting temperatures, allowing the detection of the alleles. In rs731236, the C allele with three hydrogen bonds is detected at higher temperatures than the T allele with two hydrogen bonds. Homozygous samples (TT or CC) were detected, providing a single melting peak, while heterozygous samples (TC) provided two melting peaks each one corresponding to each allele.

Table I. Demographic data of the study groups.

Parameter	Patients	Controls
Number (n)	90	103
Age, median (years)	74	57
Age, mean (years)	73	60
Max value (years)	92	90
Min value (years)	51	51
Range	41	39
Males, n (%)	44 (48.9)	51 (49.5)
Females, n (%)	46 (51.1)	52 (50.5)
MMSE score, median	21	-
FAB score, mean	10	-

MMSE, mini-mental state examination; FAB, frontal assessment battery.

Statistical analysis. Data from the genotyping results were analyzed using SNPstats web-based software, developed by the Catalan Institute of Oncology, 2006 and SPSS® 17.0.0 software (SPSS, Inc.). Logistic regression analysis was applied to analyze the association of the genotypes in each heredity model with the disease and odds ratios (ORs) with 95% confidence intervals (CIs) were calculated. Fisher's exact test was performed to compare the genotyping results with the MMSE and FAB score within the group of patients. A value of $P < 0.05$ was considered to indicate a statistically significant difference. The results were consistent with the Hardy-Weinberg equilibrium.

Results

The patient group had a median MMSE score of 21 (Fig. 1) and a FAB score of 10. The frequencies (%) of the TaqI TT, TC and CC genotypes in the controls/patients were 34.0/48.9, 47.6/41.1 and 18.4/10.0, respectively (Fig. 2). A statistically significant difference was observed for the TaqI C allele in the dominant model of inheritance TT vs. CT + CC (OR, 0.54; 95% CI, 0.30-0.96; $P = 0.035$) (Table II). On the other hand, a statistically significant difference was observed for the TT genotype in the recessive model of inheritance CC + TC vs. TT (OR, 1.86; 95% CI, 1.04-3.32; $P = 0.035$) (Table II). In the patient group, when analyzing TaqI genotypes in relation to the median value of the MMSE score (21), a statistically significant difference was observed for the TaqI CC genotype and MMSE score < 21 , CC vs. TC + TT (OR, 0.119; 95% CI, 0.014-0.995; $P = 0.032$) (Table III). No statistically significant difference was observed when analyzing TaqI genotype frequencies and FAB score in the patient group (data not shown).

Discussion

In the SEC cohort examined in the present study, the TaqI TT genotype was associated with a higher risk of developing AD by 1.8-fold. In addition, the TaqI C allele was associated with a potential protective effect against the disease, since it was calculated that TaqI C carriers had a 46% less likelihood of developing AD (Table II). Moreover, in the patient group, the




Figure 1. MMSE scores in the patient cohort presented in box and whisker plot. The median MMSE value was 21. MMSE, mini-mental state examination.




Figure 2. Bar chart of frequencies of rs731236 genotypes in the SEC population. SEC, Southeastern European Caucasian.

TaqI CC genotype was associated with a 88% less likelihood of developing severe cognitive impairment measured using the MMSE score (Table III). One of the assuming effects of the VDR polymorphism on the molecular level is the affinity of the ligand site of the receptor to vitamin D. It has been previously reported that VDR polymorphisms may decrease the affinity of VDR to vitamin D and may therefore affect the beneficial effects of vitamin D in neural cells as regards calcium homeostasis, neurotrophin levels and inflammation (16,17). Another potential effect of the TaqI polymorphism is on the stability of VDR mRNA expression (17). It can be assumed that the insufficient effects of vitamin D, associated with the TaqI TT genotype, either by decreased affinity to the VDR or reduced mRNA levels in the nervous system may lead to neurodegeneration and cognitive impairment in SECs. On the contrary, the TaqI C allele appears to be associated with enhanced effects of vitamin D in the nervous system in the specific population studied.

The results of the present study in the SEC cohort are not in agreement with those produced by a previous study in the Turkish population, in which the TaqI TT genotype was more frequent in the control group with a statistically significant

Table II. Frequencies of TaqI genotypes in the different inheritance models.

Model	Genotype	TaqI rs731236 association with AD (n=193)			P-value
		Controls, n (%)	AD, n (%)	OR (95% CI)	
Codominant	TT	35 (34%)	44 (48.9)	1.00	0.064
	TC	49 (47.6)	37 (41.1)	0.60 (0.32-1.11)	
	CC	19 (18.4)	9 (10)	0.38 (0.15-0.94)	
Dominant	TT	35 (34)	44 (48.9)	1.00	0.035
	TC/CC	68 (66)	46 (51.1)	0.54 (0.30-0.96) ^a	
Recessive	TT/TC	84 (81.5)	81 (90)	1.00	0.092
	CC	19 (18.4)	9 (10)	0.49 (0.21-1.15)	
Recessive (for T allele)	CC/TC	68 (66%)	46 (51.1%)	1.00	0.035
	TT	35 (34%)	44 (48.9%)	1.86 (1.04-3.32) ^a	


^aP<0.05. AD, Alzheimer's disease; OR, odds ratio; CI, confidence interval.

Figure 3. Bar chart of frequencies of rs731236 genotypes from the available published studies (16-18,20-22,35).

difference (16). However, another study in the Asian population reported an association of the TaqI TC genotype with AD with an increased risk of 2.8-fold for the heterozygous carriers (35). By contrast, the TaqI TC frequency in the present study was higher in the control group and no statistically significant association with the disease was observed (Table II). In another study on Northwestern European Caucasians, it was reported that the presence of TaqI C allele in adults <75 years of age resulted in a 3-fold higher risk of developing AD in comparison to the control group with a statistically significant difference (17). On the contrary, the TaqI C allele in the SEC cohort of the present study was associated with a potential protective effect against the disease and a lower likelihood of developing AD (Table II). The VDR TaqI polymorphism has also been investigated in the Iranian population; however, the results of that

study did not reveal any statistically significant difference for TaqI genotypes between the patient and control group, and the polymorphism was not associated with the disease (18,20). The results of the present study differ from those reported for the Iranian population in which no risk or protective TaqI allele for AD was detected. The TaqI polymorphism was also previously investigated in a Northeastern European Caucasian cohort and no association with AD was reported (21). Another study on the Spanish population investigated the TaqI polymorphism in relation to serum vitamin D levels (22). However, it was reported that TaqI did not result in any statistically significant difference in the serum vitamin D levels nor was it associated with a higher risk of developing AD (22). These data provided from different studies are presented in the graph depicted in Fig. 3 (16-18,20-22,35).

Table III. TaqI CC genotype vs. MMSE score <21 crosstabulation in the group of patients.

MMSE score	TaqI		OR	95% CI	P-value
	CT + TT (%)	CC (%)			
>21	38 (48.7)	8 (88.9)	0.119	0.014-0.995	0.032
<21	40 (51.3)	1 (11.1)			

MMSE, mini-mental state examination; OR, odds ratio; CI, confidence interval.

Table IV. TaqI genotype frequencies of published studies.

Country	Population	Author	n _{AD} /n _{CO}	Mean age (years) ±SD AD/control group	TaqI TT %		TaqI TC %		TaqI CC %		(Refs.)
					AD	Control group	AD	Control group	AD	Control group	
Turkey	Turkish	Gezen-Ak <i>et al</i>	104/109	75.1±5.7/73.6±7.3	36.5	48.6	48.1	35.8	15.4	15.6	(16)
Iran	Iranian	Esfehiani <i>et al</i>	101/109	>65/>65	9.9	11.9	45.5	42.2	44.6	45.9	(18)
Iran	Iranian	Khorshid <i>et al</i>	145/162	78±8/77±7	11.8	13	44.1	40.1	44.1	46.9	(20)
Poland	Northwestern European Caucasian	Łaczmański <i>et al</i>	108/77	73.3±8.6/64.5±7.8	38.9	40.2	50.9	49.4	10.2	10.4	(21)
UK	Northwestern European Caucasian	Lehmann <i>et al</i>	255/260	78.8±8.5/78.1±8.8	26.7	38.8	53.3	45	20	16.2	(17)
Brazil	Spanish	Oliveira <i>et al</i>	32/24	69.8±9/74±7.2	31.2	54.2	34.4	25	34.4	20.8	(22)
Korea	Asian	Mun <i>et al</i>	144/335	79.82±7.02/68.94±6.10	86.8	90.0	13.2	9.7	0	0.3	(35)

AD, Alzheimer's disease; SD, standard deviation; n_{AD}, number of AD patients; n_{CO}, number of individuals in the control group.

Overall, the results from published studies for each population are contradictory and do not allow for a definite conclusion regarding the association of TaqI with AD (Table IV). The TaqI C allele that appears to increase the risk of developing AD in Northwestern European Caucasians (17) has been shown to exert a protective effect in SECs from the present study. In another study on Northwestern European Caucasians, the TaqI C allele was reported to be associated with the decline in the scoring of tests measuring cognitive performance (36). However, that study did not include patients with AD and therefore a direct comparison to the studies with patients with AD is not applicable. However, the indication of the TaqI C allele that increases the risk of developing cognitive impairment is not confirmed by the results of the present study, in which the TaqI C allele appeared to decrease the risk of developing AD (Table II); homozygous TaqI CC patients also had higher MMSE scores (Table III). The interpretation of the results from different populations should take into account a number of factors, two of which are sun exposure and vitamin D intake. A previous study on Northwestern European Caucasians reported that the activity of VDR may be related to the level of sun exposure of each population (21). That study concluded that in subjects with a lower sun exposure or lower vitamin D intake, the amount of VDR produced was the crucial factor associated with the development of AD (21). In addition, the same study

described that VDR polymorphisms (TaqI and ApaI) were not associated with AD in populations with a high sun exposure due to the increased endogenous vitamin D production, which renders the activity of VDR less dependent to its amount (21). The results of the present study on the SEC population cannot confirm the aforementioned assumption, since sun exposure can be considered high in the geographical area of Greece and the TaqI TT genotype was associated with AD.

The potentially protective effects of the TaqI C allele or the higher MMSE scores of the TaqI genotype CC in patients with AD that have been described in the present study in the SEC population is an interesting finding that creates further questions to be answered regarding the molecular pathways affected by this polymorphism. It appears that the TaqI C allele is associated with the more efficient utilization of vitamin D and therefore, in a more potent effect as regards neurotrophic factor production or amyloid β clearance (6).

The present study concluded that the TaqI polymorphism is associated with AD in the SEC population, both as a risk and protective factor, and may thus be considered a potential biomarker. However, further studies on larger SEC sample sizes are required to confirm these results. Of note, the present study has provided an initial indication as regards the association of the VDR TaqI polymorphism with AD in the SEC population.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

ND conceived study and provided the control samples and data. MSK and ML performed the sample analysis. ED obtained the ethics committee study approval, performed the literature review and the statistical and data analyses, and was responsible for the manuscript composition under the supervision of ND, CK and KA. DAS and AT contributed to the editing of the final manuscript. KA and CK reviewed and edited the statistical analysis. DAS, AT, SP, VP and ES contributed to the collection of the clinical data and patient scores. SP, PM and CK also provided the patient samples. All authors discussed the results and agreed on the conclusions of the study and all authors have read and approved the final manuscript. All authors confirm the authenticity of the raw data.

Ethics approval and consent to participate

The present study was approved by the Scientific Council and Bioethics Committee of the University General Hospital 'ATTIKON' (Reg. No 312; December 21, 2017). Written informed consent for participation in the study and use of their genetic data was obtained from all participants.

Patient consent for publication

Not applicable.

Competing interests

DAS is the Editor-in-Chief for the journal, but had no personal involvement in the reviewing process, or any influence in terms of adjudicating on the final decision, for this article. All the other authors declare that they have no competing interests.

References

- Global Action Plan on the Public Health Response to Dementia 2017-2025. World Health Organization, Geneva, 2017.
- Orme RP, Middleditch C, Waite L and Fricker RA: The Role of Vitamin D₃ in the development and neuroprotection of midbrain dopamine neurons. *Vitam Horm* 100: 273-297, 2016.
- Wrzosek M, Łukaszkiwicz J, Wrzosek M, Jakubczyk A, Matsumoto H, Piątkiewicz P, Radziwoń-Zaleska M, Wojnar M and Nowicka G: Vitamin D and the central nervous system. *Pharmacol Rep* 65: 271-278, 2013.
- Littlejohns TJ, Henley WE, Lang IA, Annweiler C, Beauchet O, Chaves PH, Fried L, Kestenbaum BR, Kuller LH, Langa KM, *et al*: Vitamin D and the risk of dementia and Alzheimer disease. *Neurology* 83: 920-928, 2014.
- Afzal S, Bojesen SE and Nordestgaard BG: Reduced 25-hydroxyvitamin D and risk of Alzheimer's disease and vascular dementia. *Alzheimers Dement* 10: 296-302, 2014.
- Gezen-Ak D, Yilmazer S and Dursun E: Why Vitamin D in Alzheimer's Disease? The Hypothesis. *J Alzheimers Dis* 40: 257-269, 2014.
- Landel V, Annweiler C, Millet P, Morello M and Féron F: Vitamin D, Cognition and Alzheimer's Disease: The Therapeutic benefit is in the D-Tails. *J Alzheimers Dis* 53: 419-444, 2016.
- Mavraki E, Ioannidis P, Tripsianis G, Gioka T, Kolousi M and Vadikolias K: Vitamin D in mild cognitive impairment and Alzheimer's disease. A study in older Greek adults. *Hippokratia* 24: 120-126, 2020.
- Gezen-Ak D, Dursun E and Yilmazer S: The effect of Vitamin D treatment On Nerve Growth Factor (NGF) Release from Hippocampal Neurons. *Noro Psikiyatr Ars* 51: 157-162, 2014.
- Dursun E, Gezen-Ak D and Yilmazer S: The Influence of Vitamin D treatment on the Inducible Nitric Oxide Synthase (iNOS) Expression in Primary Hippocampal Neurons. *Noro Psikiyatr Ars* 51: 163-168, 2014.
- Grădinaru D, Borsă C, Ionescu C, Margina D, Prada GI and Jansen E: Vitamin D status and oxidative stress markers in the elderly with impaired fasting glucose and type 2 diabetes mellitus. *Aging Clin Exp Res* 24: 595-602, 2012.
- Mizwicki MT, Menegaz D, Zhang J, Barrientos-Durán A, Tse S, Cashman JR, Griffin PR and Fiala M: Genomic and nongenomic signaling induced by 1 α ,25(OH)₂-vitamin D₃ Promotes the Recovery of Amyloid- β Phagocytosis by Alzheimer's Disease Macrophages. *J Alzheimers Dis* 29: 51-62, 2012.
- Mizwicki MT, Liu G, Fiala M, Magpantay L, Sayre J, Siani A, Mahanian M, Weitzman R, Hayden EY, Rosenthal MJ, *et al*: 1 α ,25-dihydroxyvitamin D₃ and resolvin D1 retune the balance between amyloid- β phagocytosis and inflammation in Alzheimer's disease patients. *J Alzheimers Dis* 34: 155-170, 2013.
- Gezen-Ak D, Dursun E and Yilmazer S: The effects of vitamin D receptor silencing on the expression of LVSCC-A1C and LVSCC-A1D and the release of NGF in cortical neurons. *PLoS One* 6: e17553, 2011.
- Cui X, Gooch H, Petty A, McGrath JJ and Eyles D: Vitamin D and the brain: Genomic and non-genomic actions. *Mol Cell Endocrinol* 453: 131-143, 2017.
- Gezen-Ak D, Dursun E, Ertan T, Hanagasi H, Gurvit H, Emre M, Eker E, Ozturk M, Engin F and Yilmazer S: Association between Vitamin D receptor gene polymorphism and Alzheimer's disease. *Tohoku J Exp Med* 212: 275-282, 2007.
- Lehmann DJ, Refsum H, Warden DR, Medway C, Wilcock GK and Smith AD: The vitamin D receptor gene is associated with Alzheimer's disease. *Neurosci Lett* 504: 79-82, 2011.
- Esfehani NT, Rahgozar M and Biglarian A: Identification of genetic polymorphism interactions in sporadic Alzheimer's disease using logic regression. *Iran Rehabil J* 9: 45-50, 2011.
- Gezen-Ak D, Dursun E, Bilgic B, Hanagasi H, Ertan T, Gurvit H, Emre M, Eker E, Ulutin T, Uysal O and Yilmazer S: Vitamin D receptor gene haplotype is associated with late onset Alzheimer's disease. *Tohoku J Exp Med* 228: 189-196, 2012.
- Khorram Khorshid HR, Gozalpour E, Saliminejad K, Karimloo M, Ohadi M and Kamali K: Vitamin D receptor (VDR) polymorphisms and late-onset Alzheimer's disease: An association study. *Iran J Publ Health* 42: 1253-1258, 2013.
- Łączmański L, Jakubik M, Bednarek-Tupikowska G, Rymaszewska J, Słoka N and Lwow F: Vitamin D receptor gene polymorphisms in Alzheimer's disease patients. *Exp Gerontol* 69: 142-147, 2015.
- Oliveira ACR, Magalhães CA, Loures CMG, Fraga VG, de Souza LC, Guimarães HC, Cintra MTG, Bicalho MA, Sousa MCR, Silveira JN, *et al*: BsmI polymorphism in the vitamin D receptor gene is associated with 25-hydroxy vitamin D levels in individuals with cognitive decline. *Arq Neuropsiquiatr* 76: 760-766, 2018.
- Poduslo SE and Yin X: Chromosome 12 and late-onset Alzheimer's disease. *Neurosci Lett* 310: 188-190, 2001.
- Gezen-Ak D and Dursun E: Molecular basis of vitamin D action in neurodegeneration: The story of a team perspective. *Hormones (Athens)* 18: 17-21, 2019.
- Gezen-Ak D, Atasoy IL, Candaş E, Alaylioglu M, Yilmazer S and Dursun E: Vitamin D receptor regulates amyloid beta 1-42 production with protein disulfide isomerase A3. *ACS Chem Neurosci* 8: 2335-2346, 2017.
- Sutherland MK, Somerville MJ, Yoong LK, Bergeron C, Haussler MR and McLachlan DR: Reduction of vitamin D hormone receptor mRNA levels in Alzheimer as compared to huntington hippocampus: Correlation with calbindin-28k mRNA levels. *Brain Res Mol Brain Res* 13: 239-250, 1992.

27. Landel V, Millet P, Baranger K, Lloriod B and Féron F: Vitamin D interacts with Esr1 and Igf1 to regulate molecular pathways relevant to Alzheimer's disease. *Mol Neurodegener* 11: 22, 2016.
28. Gubandru M, Margina D, Tsitsimpikou C, Goutzourelas N, Tsarouhas K, Ilie M, Tsatsakis AM and Kouretas D: Alzheimer's disease treated patients showed different patterns for oxidative stress and inflammation markers. *Food Chem Toxicol* 61: 209-214, 2013.
29. Nitulescu GM, Van De Venter M, Nitulescu G, Ungurianu A, Juzenas P, Peng Q, Olaru OT, Gradinaru D, Tsatsakis A, Tsoukalas D, *et al*: The Akt pathway in oncology therapy and beyond (Review). *Int J Oncol* 53: 2319-2331, 2018.
30. Zaulkffali AS, Md Razip NN, Syed Alwi SS, Abd Jalil A, Abd Mutalib MS, Gopalsamy B, Chang SK, Zainal Z, Ibrahim NN, Zakaria ZA and Khaza'ai H: Vitamins D and E stimulate the PI3K-AKT signalling pathway in insulin-resistant SK-N-SH neuronal cells. *Nutrients* 11: 2525, 2019.
31. Zhang D, Wang L, Zhang R and Li S: Association of vitamin D receptor gene polymorphisms and the risk of multiple sclerosis: A meta-analysis. *Arch Med Res* 50: 350-361, 2019.
32. Niu MY, Wang L and Xie AM: ApaI, BsmI, FokI, and TaqI polymorphisms in the vitamin D receptor gene and Parkinson's disease. *Chin Med J (Engl)* 128: 1809-1814, 2015.
33. Papasavva M, Vikelis M, Siokas V, Katsarou MS, Dermizakis E, Raptis A, Dardiotis E and Drakoulis N: VDR gene polymorphisms and cluster headache susceptibility: Case-control study in a Southeastern European caucasian population. *J Mol Neurosci* 72: 382-392, 2022.
34. Geng J, Zhang J, Yao F, Liu X, Liu J and Huang Y: A systematic review and meta-analysis of the associations of vitamin D receptor genetic variants with two types of most common neurodegenerative disorders. *Aging Clin Exp Res* 32: 21-27, 2020.
35. Mun MJ, Kim MS, Kim JH and Jang WC: A TaqI polymorphism of vitamin D receptor is associated with Alzheimer's disease in Korean population: A case-control study. *Int J Clin Exp Med* 9: 19268-19279, 2016.
36. Kuningas M, Mooijaart SP, Jolles J, Slagboom PE, Westendorp RG and Van Heemst D: VDR gene variants associate with cognitive function and depressive symptoms in old age. *Neurobiol Aging* 30: 466-473, 2009.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.