

Paeoniflorin-6'O-benzene sulfonate suppresses fibroblast-like synoviocytes proliferation and migration in rheumatoid arthritis through regulating GRK2-G $\beta\gamma$ interaction

YUWEN ZHANG^{1,2}, XUEZHI YANG¹, CHENCHEN HAN¹, DANDAN WANG¹, YANG MA¹ and WEI WEI¹

¹Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine, Ministry of Education, Collaborative Innovation Center of Anti-inflammatory and Immune Medicine; ²Department of Anesthesiology, The First Affiliated Hospital of Anhui Medical University, Hefei, Anhui 230032, P.R. China

Received March 19, 2022; Accepted May 19, 2022

DOI: 10.3892/etm.2022.11450

Abstract. Rheumatoid arthritis (RA) is a chronic autoimmune disease. Enhanced G protein coupled receptor kinase 2 (GRK2) translocation and prostaglandin E4 receptor (EP4) desensitization play a critical role in fibroblast-like synoviocytes (FLS) dysfunction. Paeoniflorin-6'O-benzene sulfonate (CP-25) exerts a protective effect in arthritis in the RA animal models. To demonstrate the role of G $\beta\gamma$ in EP4 desensitization and the mechanisms of CP-25 that protects FLS in RA, RA-FLS and adjuvant-induced arthritis (AA-FLS) were isolated from synovium of RA patients and AA rats. RA-FLS, AA-FLS and MH7A were treated with CP-25, G $\beta\gamma$ agonist and antagonist. The cell membrane expression of EP4, GRK2, and G $\beta\gamma$ were detected using western blot analysis. Co-immunoprecipitation (Co-IP) and immunofluorescence were adopted to detect the interactions of GRK2-G $\beta\gamma$, GRK2-EP4, and EP4-G $\beta\gamma$. Cell Counting Kit-8 and Transwell assay were used to analyze the proliferation and migration of the FLS. An increased membrane expression of GRK2 and G $\beta\gamma$, enhanced GRK2-G $\beta\gamma$ interaction and decreased EP4 membrane expression in the RA synovial tissue were identified. *In vitro*, prostaglandin E2 (PGE₂) enhanced the proliferation and migration of FLS. CP-25 exhibited an inhibition effect similar to G $\beta\gamma$ inhibitor, which downregulated GRK2-EP4 interaction, blocked the translocation of GRK2, and reversed EP4 desensitization, leading to the suppression of the proliferation and migration induced by PGE₂. These results elucidated that an enhanced

GRK2-G $\beta\gamma$ interaction was involved in the EP4 desensitization and dysfunction. CP-25 regulated EP4-GRK2-G $\beta\gamma$ signaling and re-sensitized EP4 by inhibiting GRK2-G $\beta\gamma$ interaction. The regulation of EP4-G $\beta\gamma$ -GRK2 signaling may be a novel potential therapeutic target in RA.

Introduction

Rheumatoid arthritis (RA) is a chronic, autoimmune, inflammatory disease. The pathological hallmarks of synovium in RA are synovial hyperplasia, pannus formation, cartilage damage and bone erosion (1). Abnormal expansion of fibroblast-like synoviocytes (FLS) leads to synovial hyperplasia, which is the leading pathophysiology of RA. Making up two thirds of the hyperplastic synovial membrane, FLS secrete several adhesion molecules and inflammatory cytokines, and this also leads to bone erosion and cartilage damage. Migration of FLS to unaffected cartilage drives the progression of polyarthritis in RA (2). It has been reported that there is a large amount of prostaglandin E2 (PGE₂) in the synovial fluid of RA patients that triggers anomalous proliferation and migration of FLS (3).

Guanine nucleotide binding protein, also known as the G protein, is a ubiquitous signal molecule that plays an important role in transmembrane signal transduction. G protein couples with the corresponding G protein-coupled receptor (GPCR). The binding of ligand to GPCR induces the dissociation of G protein into G α and G $\beta\gamma$ dimers. G $\beta\gamma$ subunit plays multifunctional roles, such as anchoring the G protein to the cell membrane and conducting signal transduction. GPCRs and their effector molecules bind G protein subunits before targeting to the cell surface. Interacting with various effectors, including G protein-coupled receptor kinase 2 (GRK2), adenylyl cyclase II, and PI-3 kinase (4), G $\beta\gamma$ is important in recruiting GRKs to activated receptors and regulating cellular functions.

GRK2 is associated with GPCR desensitization. PGE₂ binds to prostaglandin receptors (EPs). In the physiologic state, EP4 primarily couples to the G α s subunit to mediate increases in the cAMP concentrations. In a pathological environment, EP4 perceives continuous stimulation from abnormal levels of inflammatory cytokines such as PGE₂, causing an overactive

Correspondence to: Professor Wei Wei, Institute of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine, Ministry of Education, Collaborative Innovation Center of Anti-inflammatory and Immune Medicine, 81 Meishan Road, Hefei, Anhui 230032, P.R. China
E-mail: wwei@ahmu.edu.cn

Key words: CP-25, G $\beta\gamma$, G protein coupled receptor kinase 2, rheumatoid arthritis, fibroblast-like synoviocytes, prostaglandin E4 receptor

inflammatory response. Previous studies have shown that in RA animal models, the PGE₂-EP4-GRK2 signaling pathway plays an important role in the occurrence and development of arthritis. PGE₂ binds to the EP4 receptor, inducing excessive GRK2 translocation to the cell membrane and EP4 desensitization, thereby promoting downregulation of cAMP, FLS dysfunction and synovium hyperplasia with the mechanism of decreased protective Gas signaling (5,6).

Paeoniflorin-6'-O-benzene sulfonate (CP-25; patent number in China: ZL201210030616.4) is a structurally modified compound from paeoniflorin with anti-inflammatory and immunomodulatory effects in RA animal models and other immune diseases (7-10). It was found that CP-25 ameliorated FLS abnormal proliferation and migration by inhibiting GRK2 translocation (11) and re-sensitizing EP4 (6), which was an important mechanism of CP-25 in improving FLS dysfunction.

The GPCR-G $\beta\gamma$ -GRK complex is crucial for the regulation of receptor function and GRK catalysis, and this has been studied in adrenergic and opioid receptors, but has not been reported in EP4 receptors. GRK2 binds both delta opioid receptor and G $\beta\gamma$ to form a receptor-GRK2-G $\beta\gamma$ complex, which is required for GRK-mediated receptor phosphorylation and desensitization (12). In heart failure animal model, GRK2 translocates to β -adrenergic receptor after binding with G $\beta\gamma$ to phosphorylate the agonist-occupied receptor, thus cardiac contractile function collapses as β -adrenergic receptor signaling is attenuated (13). It remains unclear whether G $\beta\gamma$ is involved in PGE₂-EP4-GRK2 signaling. Hence, in the present study, the role of G $\beta\gamma$ and the GRK2-G $\beta\gamma$ interaction in regulating FLS proliferation and migration was mainly explored. In addition, the specific mechanism of CP-25 in restoring dysfunction of EP4 was investigated.

Materials and methods

Sample collection. RA synovial tissue samples were obtained from six patients (4 females, 2 males; age range, 52-73 years) with RA undergoing surgical synovectomy and knee replacement. Trauma synovium samples were obtained from six patients (3 females, 3 males; age range, 28-50 years) with traumatic synovitis undergoing arthroscopic synovectomy. The sample collection period was from December 2017 to December 2018. Patients had been diagnosed with RA at least three years prior to surgery and were treated using the standard of care (non-steroidal anti-inflammatory drugs and disease-modifying anti-rheumatic drugs DMARDs). Informed consent was obtained from all patients. The study protocol was approved (approval no. 20131321) by the Biomedical Ethics Committee of Anhui Medical University (Hefei, China).

Animals. A total of 12 male Sprague Dawley rats (body weight, 180-200 g) were obtained from SPF Animal Laboratory of Anhui Medical University (Hefei, China). All rats were housed in pathogen-free animal houses at 20-26°C under a 12-h light/dark cycle and had free access to sterile water and food. All experiments were conducted in accordance with the principles of the laboratory animal care guidelines and were approved (approval no. 20131321) by Biomedical Ethics Committee of Anhui Medical University (Hefei, China).

Drugs and reagents. PGE₂ was purchased from Cayman Chemical Company. Gallein was purchased from Santa Cruz Biotechnology, Inc. mSIRK was purchased from Sigma-Aldrich; Merck KGaA. GRK2i was purchased from Tocris Bioscience. The Cell Counting Kit-8 (CCK-8) kit was purchased from Beyotime Institute of Biotechnology. The Transwell inserts (8- μ m) were purchased from Corning, Inc. CP-25 (purity >98%) was provided by the Institute of Clinical Pharmacology, Anhui Medical University (Hefei, China). G β 1 siRNA(h) and the control siRNA(h) were purchased from Santa Cruz Biotechnology, Inc. Lipofectamine® 2000 and fetal bovine serum (FBS) were purchased from Invitrogen; Thermo Fisher Scientific, Inc. Dulbecco's Modified Eagle's Medium (DMEM) was purchased from Hyclone; Cytiva.

Induction of adjuvant-induced arthritis (AA). There were six rats in model group, and the other six rats were in control group. The method of AA rat preparation was consistent with that described in a previous study (5). A total of 0.1 ml of complete Freund's adjuvant (CFA) emulsion was injected into the right hind metatarsal footpad. The rats were assessed every three days for the signs of arthritis by two independent observers. If tissue necrosis or ulcerated masses were present (no signs of healing within 3 days) and signs of deterioration such as inability to reach food or water for ~12 h, weight loss >20% of starting weight or any sign of suffering not relieved by pharmacologic analgesia, the rat would be euthanized. The rats in model group were sacrificed for the AA-FLS at day 30 after immunization under anesthesia using isoflurane (induction 3-4%; maintenance 1-2%). Subsequently, all the rats were euthanized via the inhalation of carbon dioxide. The displacement of the euthanasia chamber was between 30-70% of the chamber volume per min. Death would be confirmed by an appropriate method, such as ascertaining cardiac and respiratory arrest or noting fixed and dilated pupils of an animal.

Cell culture and identification. The RA-FLS and AA-FLS were obtained from the knee synovium of the RA patients and AA rats. Human rheumatoid FLS (MH7A) were purchased from Jennio Biotech Co., Ltd. All cells were cultured in DMEM containing 20% FBS, 1% penicillin and 1% streptomycin at 37°C and 5%-CO₂ containing incubator. The RA-FLS and AA-FLS were cultured using Anti-Vimentin (cat. no. sc-373717; Santa Cruz Biotechnology, Inc.) and Anti-CD55 (cat. no. 26580-1-AP; Proteintech) at a dilution ratio of 1:1,000 for 30 min and then detected using immunofluorescence. Mycoplasma testing was carried out for the cell lines used.

Histological analysis. Synovial tissue was fixed with 4% paraformaldehyde solution at room temperature for 24 h and embedded in paraffin. The paraffin sections (4 μ m) were dewaxed in xylene, stained with hematoxylin and eosin (H&E) at room temperature for 3 min and 30 sec, respectively, and were examined using standard light microscopy.

Immunofluorescence assay. The synovial tissue samples were snap-frozen in OCT medium. Cryostat sections (5 μ m) were cut and mounted on adhesive glass slides. Slides of the tissue

samples and cells were fixed with 4% paraformaldehyde at 4°C for 10 min, permeabilized with 0.5% Triton X-100 for 7 min at room temperature and blocked with 5% bovine serum albumin (MilliporeSigma) for 1 h at 37°C. After incubating with primary antibodies (1:100 dilution) at 4°C overnight, the slides were incubated with anti-mouse-Alexa Fluor 594 (1:200 dilution; cat. no. ab150116; Abcam) and anti-rabbit-Alexa Fluor 488 (1:200 dilution; cat. no. ab150073; Abcam) for 1 h at 37°C. DAPI (20 μ l) was used to stain the nuclei for 5 min at room temperature. The slides were then observed under a fluorescence microscope (BX53; Olympus Corporation). The primary antibodies were as follows: G β (cat. no. sc-166123), EP4 (1:1,000 dilution; cat. no. sc-55596; both from Santa Cruz Biotechnology, Inc.) and GRK2 (cat. no. ab228705; Abcam).

Isolation of membranes from synovial tissue. The synovial tissue was washed using PBS, cut into pieces and added with a cold homogenization buffer (0.32 M sucrose, 5 mM Tris-HCl, pH 7.5, 120 mM KCl, 1 mM EDTA and 0.2 mM PMSF). Samples were sonicated and centrifuged at 100,000 x g for 1 h at 4°C. The supernatants were pooled and centrifuged at 100,000 x g for 1 h. The pellets were resuspended in a cold extraction buffer (20 mM HEPES, pH 7.5, 10% glycerol, 2% Triton X-100, 1 mM EDTA, 1 mM EGTA and 0.2 mM PMSF) and incubated at 4°C overnight. The supernatants containing the cell membrane were collected after centrifuging at 7,800 x g for 30 min at 4°C.

Isolation of proteins from the cell cytoplasm and the membrane. RA-FLS, AA-FLS and MH7A cells were lysed with lysis buffer (cat. no. P0013; Beyotime Institute of Biotechnology) and centrifuged at 12,000 x g for 10 min at 4°C. The supernatants were centrifuged at 100,000 x g for 1 h at 4°C. The precipitate was kept as the membrane fraction, while the supernatant was collected as the cytoplasm fraction.

Cell proliferation assay. Cell proliferation was examined using a CCK-8 assay. RA-FLS, AA-FLA and MH7A cells were counted and seeded on to 96-well plates at a concentration of 5×10^6 /ml and incubated in DMEM at 37°C for 48 h. The cells were then treated with PGE₂ (2 μ m) with or without different agonists or antagonists and CP-25 (10^{-7} , 10^{-6} and 10^{-5} mol/l) for 24 h. After treatment, the cells were added to a 10- μ l CCK-8 solution for each well. The plate was incubated for 1 h at 37°C. The optical density values of the plate were measured at 490 nm on a Tecan Infinite M200 Microplate Reader (Tecan Group, Ltd).

Cell migration assay. Cells were seeded into the upper wells of the Transwell chambers at a concentration of 1×10^5 /ml and maintained in a medium containing 5% FBS. The lower chamber was filled with DMEM containing 20% FBS. After 24 h, the upper chamber was washed with PBS and fixed with 4% paraformaldehyde for 20 min at room temperature and stained with 0.1% crystal violet for 15 min at room temperature. All images were captured using a light microscope. Cells were counted from at least four random microscopic fields for each sample in three independent experiments.

Small interfering RNA (siRNA). The G β 1 siRNA (cat. no. sc-41762) and negative control siRNA (cat. no. sc-37007)

were used to knockdown G β 1 in MH7A cells. The MH7A were transfected with 50 nM siRNA at 70% confluency in 24-well plates using Lipofectamine 2000 at room temperature according to the manufacturer's protocol. After 48 h, silencing of G β 1 was confirmed using western blot analysis.

Western blot analysis. Protein samples were collected and boiled in a loading buffer for 10 min at 100°C. The protein concentration was detected using a BCA protein assay kit (Thermo Fisher Scientific, Inc.). Equal amounts of protein (20 μ g) were electrophoretically separated using 10% SDS-PAGE and transferred to a PVDF membrane (MilliporeSigma). The PVDF membrane was then incubated in blocking solution (cat. no. P0205; Beyotime Institute of Biotechnology) at room temperature for 1 h. Subsequently, the membrane was incubated with primary antibodies at 4°C overnight, followed by the horseradish peroxidase-conjugated goat anti-mouse antibody (cat. no. STAR207P; or goat anti-rabbit antibody (cat. no. STAR208P; both 1:10,000 dilution; Bio-Rad Laboratories, Inc.) at room temperature for 2 h. The bands were visualized using chemiluminescence detection reagents (MilliporeSigma) according to the manufacturer's protocol. The following primary antibodies were used: G β (1:1,000; cat. no. sc-166123), G β 1 (1:1,000; cat. no. sc-515764; both from Santa Cruz Biotechnology, Inc.), ATP1A1 (1:3,000; cat. no. 14418-1-AP; ProteinTech Group, Inc.), GRK2 (1:1,000; cat. no. sc-13143; Santa Cruz Biotechnology, Inc.), GRK2 (1:1,000; cat. no. ab228705; Abcam), EP4 (1:1,000; cat. no. sc-55596), β -arrestin2 (1:1,000; cat. no. sc-13140; both from Santa Cruz Biotechnology, Inc.), β -actin (1:5,000; cat. no. 60008-1-Ig) and GAPDH (1:5,000; cat. no. 60004-1-Ig; both from ProteinTech Group, Inc.). The protein bands were visualized using an enhanced chemiluminescence detection system and quantified using ImageJ software (version 1.46; National Institutes of Health).

Coimmunoprecipitation (Co-IP) assay. To examine the association between GRK2 and G β in the synovial tissue or in FLS, the tissue or cell lysate was immunoprecipitated using an anti-GRK2, anti-EP4 or anti-G β antibody and analyzed using western blot analysis. Briefly, the sample was precleared for 2 h at 4°C with 20 μ l of protein A + G agarose beads and subsequently incubated with 4 μ l primary antibody for 2 h at 4°C. Then the precleared lysates were associated with 20 μ l of protein A + G agarose beads and the primary antibody and incubated overnight at 4°C on the rotation shaker. Tubes were then centrifuged at 14,000 x g for 5 sec at 4°C. The pellets were washed with lysis buffer for 10 min. The resin-bound immune complexes were boiled and the following steps were according to the western blot protocol.

Reverse transcription-quantitative PCR (RT-qPCR). Total RNA was extracted from synovial tissue using TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.), which was then used to synthesize cDNA by RT using the PrimeScript™ RT Master Mix Kit (TaKaRa Bio, Inc.) in a reaction volume of 20 μ l according to the manufacturer's protocol. Real-time qPCR assays were performed with an Applied Biosystems 7500 fast real-time PCR system with TB Green Premix Ex Taq II (TaKaRa Bio, Inc.). The conditions of amplification were as

follows: Initial denaturation at 95°C for 30 sec, followed by 40 cycles of annealing and extension at 95°C for 3 sec and 60°C for 30 sec. The oligonucleotide primer sequences were as follows: GNB1 forward, 5'-TCGTCTCTGGTGCTTGTGATGC-3' and reverse, 5'-GTCGTCTGAGCCAGTGGCAAA T-3'; and GNB2 forward, 5'-TGTTGCCGCTTCCTGGATGACA-3' and GNB2 reverse, 5'-CACTGTGTCCAGCAA AACCAC-3'; GAPDH forward, 5'-GGAGCGAGATCCCTC CAAAAT-3' and reverse, 5'-GGCTGTTGTCATACTTCT CATGG-3'. GAPDH was used as an internal reference and the 2^{- $\Delta\Delta C_q$} method was utilized to calculate the relative expression levels (14).

Statistical analysis. Data were analyzed using the GraphPad Prism v7 statistical software (GraphPad Software, Inc.). Values are expressed as the mean \pm standard deviation. An Unpaired Student's t-test was used to analyze the data between two independent groups. A one-way analysis of variance followed by Bonferroni's post hoc test was used to analyze the data from multiple groups. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

The GRK2-G β γ interaction is overactivated in the synovium tissue of patients with RA. It was previously revealed that PGE₂ levels were increased in the RA synovium and AA rats. It was also found that after PGE₂ stimulation, the membrane expression of GRK2 was increased in AA-FLS, which was related to FLS dysfunction (5). In an attempt to identify whether G β γ was involved in the GRK2 translocation to the membrane, the expression of G β and the interaction between GRK2 and G β in the synovial tissues were evaluated using western blot analysis, Co-IP and double-stained immunofluorescence. The pathological feature of the synovial tissue in patients with RA was first assessed using H&E staining and expression of relevant proteins was detected. Compared with the trauma patients, the RA patients had synovial hyperplasia and inflammatory cell infiltration (Fig. 1A). The RA group showed a significantly increased expression of GRK2 (Fig. S1). In the RA group, membrane expression of GRK2 and G β was increased, whereas EP4 was decreased (Fig. 1B). The aforementioned results indicated GRK2 and G β translocation to the membrane and EP4 desensitization in the RA synovium. The results of the Co-IP and immunofluorescence demonstrated that GRK2-G β , EP4-G β and GRK2-EP4 interactions were enhanced in the RA group compared with those in the trauma group (Fig. 1C-F). These results suggested that G β γ was involved in GRK2 translocation and EP4 desensitization in RA.

CP-25 inhibits proliferation and migration of RA-FLS and AA-FLS by downregulating the GRK2-G β γ interaction. Since FLS is the major component of hyperplastic pannus in the RA synovium, the primary synoviocytes from RA patients and AA rats were isolated and identified as cells positive for labeling with Vimentin and CD55 by immunofluorescence (Fig. S2). Before the experimental endpoint, none of the rats succumbed. After the stimulation of PGE₂ (2 μ M) for 24 h, the proliferation and migration in both RA-FLS and

AA-FLS was enhanced. CP-25 (10⁻⁶ and 10⁻⁵ mol/l) significantly inhibited proliferation and migration induced by PGE₂ in both primary cells. Equivalent effects were observed in the G β γ inhibitor and the gallein (10 μ M) group (Fig. 2A and B), which indicated that inhibition of both G β γ and CP-25 may decrease the abnormal FLS proliferation migration in the RA condition. To identify whether G β γ subunits were involved in the protection of CP-25, the membrane expression of G β and its interaction with GRK2 were examined. It was revealed that membrane expression of GRK2 and G β was increased, whereas EP4 was decreased in the PGE₂ group. CP-25 (10⁻⁶ mol/l) restored membrane expression of GRK2, G β and EP4 (Fig. 2C). The Co-IP assay showed enhanced EP4-G β , GRK2-G β , and GRK2-EP4 interactions in the PGE₂ group. CP-25 inhibited translocation of GRK2 and G β , and reduced the binding of GRK2 and G β (Fig. 2D). These results suggested that CP-25 inhibited proliferation and migration of RA-FLS and AA-FLS by downregulating GRK2-G β γ interaction.

CP-25 inhibits EP4-GRK2-G β γ signaling and promotes G β γ -dependent EP4 re-sensitization in MH7A cells. In the next few experiments, drugs that interfered with G β γ in the immortal human synovial fibroblast cell line MH7A were used to investigate whether G β γ controlled the interaction between GRK2 and EP4 and the role of CP-25.

First, excessive proliferation and migration induced by PGE₂ was observed in MH7A cells (Fig. 3A and B). GRK2i, gallein and CP-25 reversed the abnormalities. The results were similar to that in the RA-FLS and the AA-FLS. The proliferation and migration rate were further enhanced in the mSIRK (G β γ agonist) group compared with the PGE₂ group. PGE₂ upregulated the membrane expression of GRK2 and downregulated the membrane expression of EP4. β -arrestin2, related to GPCR desensitization, increased in the PGE₂ group. G β γ inhibitor, GRK2i, and gallein decreased the membrane expression of GRK2 and β -arrestin2 and restored the membrane expression of EP4. Membrane GRK2 was significantly increased in the mSIRK group, whereas β -arrestin2 and EP4 remained unchanged compared with the PGE₂ group (Fig. 3C). Subsequently, it was identified that the GRK2-EP4, EP4-G β , GRK2-G β and EP4- β -arrestin2 interactions were enhanced in the PGE₂ group, while they were inhibited in the gallein group (Fig. 3E and G). Compared with the PGE₂ group, GRK2i decreased the combination of GRK2 and EP4, while mSIRK enhanced the GRK2-EP4 interaction (Fig. 3D and F). CP-25 suppressed the GRK2-G β interaction (Fig. 3E and G), decreased membrane expression of GRK2 and β -arrestin2 (Fig. 3C) and blocked EP4 binding with GRK2 or β -arrestin2 (Fig. 3E), and thus restored membrane EP4 expression (Fig. 3C). These results demonstrated that CP-25 inhibited PGE₂-induced EP4-GRK2-G β γ signaling activation in MH7A. Exogenous stimulation of PGE₂ overactivated EP4-GRK2-G β γ signaling and promoted EP4 desensitization and dysfunction. CP-25 abrogated the binding of GRK2 with G β γ and caused G β γ -dependent re-sensitization of EP4, thus inhibiting abnormal proliferation and migration in the RA condition. It is worthy to note that the total expression of G β nearly remained unchanged between the control, PGE₂, CP-25 and gallein groups (Fig. S3).




Figure 1. Enhanced GRK2-G $\beta\gamma$ interaction in the synovial tissue of patients with RA. (A) H&E-stained histological sections. Scale bars=100 and 20 μ m. (B) Membrane protein expression of GRK2, EP4 and G β detected using western blotting. (C and D) Interactions of (C) GRK2-G β , EP4-G β and (D) GRK2-EP4 by Co-immunoprecipitation. (E and F) The co-localization of GRK2-EP4 and GRK2-G β in frozen section of synovial tissue of patients with RA was detected using immunofluorescence. Scale bars, 20 μ m. * P <0.05, ** P <0.01 and *** P <0.001 vs. trauma group (n=6). GRK2, G protein coupled receptor kinase 2; RA, rheumatoid arthritis; EP4, prostaglandin E4 receptor.

G $\beta\gamma$ knockdown inhibits the PGE₂-induced proliferation and migration by possibly interfering with the GRK2-EP4 interaction. The mRNA expression of G β 1 and G β 2 in synovial tissue of patients with trauma and RA was compared. It was observed that the expression of G β 1 mRNA was higher in the synovial tissue of patients with RA (Fig. S4). To further confirm the functions of G $\beta\gamma$ in MH7A cells, the G β 1 siRNA was transfected to knock down the endogenous G β 1. It was found that the knockdown of G β 1 significantly decreased PGE₂-induced proliferation (Fig. 4A) and migration (Fig. 4B). In untransfected cells, there was increased total expression of GRK2 (Fig. 4C), decreased membrane expression of EP4 (Fig. 4D) and enhanced GRK2-EP4 interaction after the PGE₂ induction (Fig. 4E). These results suggested that G $\beta\gamma$

knockdown inhibits PGE₂-induced proliferation and migration of MH7A cells, possibly by interfering with the GRK2-EP4 interaction. Collectively, G $\beta\gamma$ was involved in the anomalous proliferation and migration by hindering the interaction between GRK2 and EP4.

Discussion

FLS is a prominent component of the hyperplastic synovium. Inflammatory environments induce FLS to migrate, and this results in bone damage and cartilage erosion. Compared with normal synovial tissue, there are high expression levels of GRK2 in patients with RA and in RA animal models (6,7). Accordingly, GRK2 upregulation and translocation to the cell




Figure 2. CP-25 inhibits proliferation and migration of RA-FLS and AA-FLS by downregulating GRK2-G β γ interaction. (A and B) Inhibition of abnormal (A) proliferation and (B) migration of RA-FLS and AA-FLS by CP-25 and gallein was detected using Cell Counting Kit-8 and Transwell assays. (C) Membrane expression of GRK2, EP4 and G β in RA-FLS and AA-FLS was evaluated by western blotting. (D) Combination of EP4-G β , GRK2-G β , and GRK2-EP4 in RA-FLS and AA-FLS using Co-immunoprecipitation. Scale bars, 100 μ m. ** P <0.01 and *** P <0.001 vs. control group; * P <0.05, ** P <0.01 and *** P <0.001 vs. PGE₂ group (n=6). RA, rheumatoid arthritis; AA, adjuvant-induced arthritis; FLS, fibroblast-like synoviocytes; GRK2, G protein coupled receptor kinase 2; EP4, prostaglandin E4 receptor; PGE₂, prostaglandin E2.




Figure 3. Gβγ on proliferation and migration of MH7A cells and the involvement of EP4-GRK2-Gβγ signaling. MH7A cells were treated with Gβγ antagonist (5 μM GRK2i and 10 μM gallein), Gβγ agonist (1 μM mSIRK), CP-25 (10⁻⁶ mol/l) and stimulated by PGE₂ (2 μM). (A) Cell proliferation was detected using Cell Counting Kit-8. (B) Cell migration was detected using Transwell assay. Scale bars, 100 μm. (C) Membrane expression of GRK2, EP4 and Gβ in PGE₂-stimulated MH7A cells was detected using western blotting. (D) Co-expression of GRK2 and EP4 with GRK2i and mSIRK treatment detected by Co-IP. (E) Co-expression of EP4-β-arrestin2, EP4-GRK2 and EP4-Gβ with CP-25 and gallein treatment was detected by Co-IP. (F) Co-localization of GRK2 and EP4 with GRK2i and mSIRK treatment was evaluated by immunofluorescence. Scale bars, 50 μm. (G) Co-localization of GRK2 and Gβ, EP4 and GRK2 in PGE₂-stimulated MH7A cells was evaluated by immunofluorescence. Scale bars=50 μm. *P<0.05, **P<0.01 and ***P<0.001 vs. control group; #P<0.05, ##P<0.01 and ###P<0.001 vs. PGE₂ group (n=5). EP4, prostaglandin E4 receptor; GRK2, G protein coupled receptor kinase 2; PGE₂, prostaglandin E2; Co-IP, Co-immunoprecipitation.

membrane play a role in the mediation of the pathological progression of RA. In the present study, the protein expression of EP4-GRK2-Gβγ signaling was detected using western blot analysis and protein-protein interaction was evaluated using Co-IP and immunofluorescence. Increased GRK2 membrane

expression and enhanced interactions of GRK2-Gβ, EP4-Gβ, and GRK2-EP4 were revealed in the collected synovial tissue of patients with RA.

To mimic the chronic inflammatory environment of RA, RA-FLS, AA-FLS and MH7A cells were stimulated with




Figure 4. G $\beta\gamma$ knockdown inhibits the PGE₂-induced proliferation and migration of MH7A cells. (A and B) PGE₂-induced (A) proliferation and (B) migration of MH7A cells after silencing G $\beta 1$ was detected by Cell Counting Kit-8 and Transwell assays. Scale bars=100 μ m. (C) Total expression of GRK2, G $\beta 1$ and (D) membrane expression of EP4 stimulated by PGE₂ in G $\beta 1$ siRNA-transfected MH7A cells was determined by western blotting. (E) Co-localization of GRK2 and EP4 in G $\beta 1$ siRNA-transfected MH7A cells. Scale bars, 100 μ m. **P<0.01 and ***P<0.001 vs. control group (n=3). PGE₂, prostaglandin E₂; GRK2, G protein coupled receptor kinase 2; siRNA, small interfering RNA.

PGE₂ for 24 h. Similar results as in the synovial tissue of patients with RA were observed, suggesting that G $\beta\gamma$ was involved in synovial hyperplasia. The enhanced GRK2-G $\beta\gamma$ and the downregulated expression of EP4 in the FLS were significantly associated with synovium hypertrophy. The aforementioned mechanisms promoted the occurrence and development of RA. To further identify the role of G $\beta\gamma$ in RA, G $\beta\gamma$ interference with pharmacological and genetic methods was conducted. G $\beta\gamma$ agonist, mSIRK, promoted abnormalities by PGE₂. However, the membrane expression of EP4 and β -arrestin2 were unchanged compared with the PGE₂ alone

group. It was hypothesized that the desensitized EP4 reached a saturated state. There may be other mechanisms independent of EP4 signaling that account for the extra elevated proliferation and migration after the mSIRK treatment. For example, the PI3K/AKT pathway was activated by G $\beta\gamma$ (15). Intervention abrogating GRK2-G $\beta\gamma$ or a G $\beta\gamma$ blocker and G $\beta\gamma$ knockdown mitigated the GRK2 translocation and resulted in a decreased membrane expression of EP4. This finally alleviated the abnormal proliferation and migration of FLS in the RA model. As revealed by the present data, the activation of G $\beta\gamma$ was involved in FLS proliferation and migration.

In the present study, FLS were treated with CP-25 at a concentration of 10^{-7} , 10^{-6} and 10^{-5} mol/l. The results showed that the 10^{-6} and 10^{-5} mol/l CP-25 significantly inhibited the proliferation and migration after the stimulation of PGE_2 , suggesting that CP-25 may provide a potential novel strategy for RA treatment in terms of synovial hyperplasia. In consistency with previous studies, CP-25 suppressed proliferation (5) and migration (15) in the AA-FLS and MH7A cells. In addition, CP-25 inhibits the progress of AA rats by reducing inflammation, immunity and joint injury (16). All of the previous studies by our group proved that CP-25 may be an effective drug for RA (5-11,15,16). In the present study, the effect of CP-25 on the interaction between GRK2 and $\text{G}\beta\gamma$ was evaluated. A decreased GRK2- $\text{G}\beta\gamma$ interaction and less EP4 desensitization was observed under the CP-25 treatment. The results supported the hypotheses that the interaction of $\text{G}\beta\gamma$ and GRK2 contributes to the effect of CP-25 on relieving FLS dysfunction.

GRK2 is a family of protein kinases that regulates the activity of GPCR by phosphorylating the intracellular domain. The pleckstrin homology (PH) domain in the C-terminal of GRK2 is considered to interact with free $\text{G}\beta\gamma$ subunits (17). Phosphorylation at Ser⁶⁸⁵ of GRK2 by PKA facilitates the GRK2- $\text{G}\beta\gamma$ interaction and enhances the kinase activity of GRK2 (18). It was recently found that CP-25 downregulates phosphorylation at Ser⁶⁸⁵ of GRK2 and directly binds to the kinase domain of GRK2 *in vitro* and inhibits GRK2 activity by controlling the key amino acid residue of Ala³²¹ of GRK2 (7,19). Other studies have revealed that when Ser⁶⁷⁰ in the C-terminal domain of GRK2 was phosphorylated by MAPK, the binding of $\text{G}\beta\gamma$ with GRK2 was impaired. Thus, GRK2 translocation to the membrane and subsequent GPCR regulation are inhibited (20,21). Manipulation of GRK2 represents a promising molecular basis for treating RA. Previous studies by our group emphasized the importance of inhibiting GRK2 in maintaining receptor function (6,7). Approaches such as the genetic deletion of *Grk2* and systemic inhibition of GRK2 showed some drawbacks or potential side effects differentiated from a pharmacological approach such as a selective GRK2 inhibitor. For instance, *Grk2*-knockout mice succumbed on day 15 of embryonic development from cardiac hypoplasia and cardiac dysfunction (21). GRK2 knockdown also caused glomerular injury and renal damage when treating heart disease (22). It is worth exploring whether there exists a safer and efficacious influence on the balanced receptor function and kinase activity with minimal side effects, but does not just completely abolish the expression of the receptor or kinase.

At present, small molecule targeting GPCR- $\text{G}\beta\gamma$ -GRK2 signaling is an area of increased research attention in pathologic conditions such as heart failure and kidney dysfunction (23,24). In this pathway, one approach is the pharmacological inhibition of GRK2, such as paroxetine. The other strategy is to interfere with GRK2- $\text{G}\beta\gamma$ or inhibit $\text{G}\beta\gamma$. Preclinical evaluations have revealed a cardioprotective effect for GRK2 genetic deletion or pharmacological inhibition in cardiovascular diseases by binding to and stabilizing GRK2, thus inhibiting downstream β -adrenergic receptor (β -AR) signaling (25,26). Furthermore, M119 and gallein hindered $\text{G}\beta\gamma$ binding to GRK2, leading to decreased translocation to

the membrane of GRK2 and thus preserving the normal function of the β -adrenaline receptor so as to mitigate heart failure progression and cardiac hypertrophy (27). In the present study, gallein and CP-25 suppressed the abnormal proliferation and migration in PGE_2 -stimulated FLS by interfering with the binding of $\text{G}\beta\gamma$ with GRK2. The present results are in a manner analogous to those observed of the β -AR.

There are five different $\text{G}\beta$ and 12 $\text{G}\gamma$ in humans. An investigation of the binding preferences revealed that GRK2 binds preferentially to $\text{G}\beta 1$ and $\text{G}\beta 2$ (28). It was revealed that the expression of $\text{G}\beta 1$ mRNA was higher than $\text{G}\beta 2$ in the synovial tissue of patients with RA; hence, $\text{G}\beta 1$ was silenced in MH7A cells. It has become clear that $\text{G}\beta\gamma$ is a multifunctional protein complex that interacts not only with the $\text{G}\alpha$ subunit and GPCRs but also with intracellular proteins such as PI3K, AC and GIRK (29). Targeting $\text{G}\beta\gamma$ signaling appears to be a growing potential pharmacological intervention for treating cancer cell migration (30). It has been reported that the EP4 antagonist suppresses cancer cell invasion and migration in prostate cancer (31). However, a balance should be maintained in suppressing cancer cell growth and motility without destroying the normal functions of cells. FLS displays cancerous properties of proliferation and migration in RA synovial tissue that resemble cancer-associated fibroblasts in tumors (32). FLS has its own physiological function to maintain the homeostasis of joints (33). Excessive inhibition of the normal function of FLS in rheumatoid or non-rheumatoid joints would cause detrimental outcomes. This notion may be related to the soft regulation of the inflammatory immune response (SRIIR) proposed by our group. This emphasized that the normal function as well as gene and protein expression or activity of the cell should not be completely disabled by drugs. The most appropriate drug is characterized by regulating and restoring abnormal activity to achieve physiological levels with minimal adverse reactions. In the present study, CP-25 likely is a SRIIR drug due to its ability to re-sensitize EP4 and reduce the excessive proliferation and migration in PGE_2 -stimulated FLS.

The long-term activation of GPCRs after ligand exposure has been revealed to desensitize receptors and downregulate downstream signals. GPCR desensitization is generally explained by paradigms proposed in studies on the $\beta 2$ -AR (34,35). Once the activated receptor is phosphorylated by GRK2/3, its affinity for β -arrestin is enhanced. β -arrestin2 binds GPCRs that are targeted for endocytosis and receptor internalization (35). Agonist-induced GRK2- $\text{G}\beta\gamma$ interaction is a prerequisite for GRK2-mediated receptor desensitization and downregulation. The PH domain of GRK2 competitively binds to $\text{G}\beta\gamma$ and sequesters it from the $\text{G}\alpha$ subunit. Evidence has demonstrated that the $\text{G}\beta\gamma$ -mediated recruitment of PI3K in complex with GRK2 is directly implicated in β -AR desensitization (36). In the present study, in response to PGE_2 , GRK2 was passively pulled to the cell membrane by the binding of $\text{G}\beta\gamma$ subunits, and then both simultaneously translocated to the cell membrane. GRK2 was phosphorylated on the membrane. β -arrestin2 was recruited to the membrane simultaneously, inducing excessive desensitization and internalization of EP4. Blocking $\text{G}\beta\gamma$ signaling, either with the small molecule $\text{G}\beta\gamma$ selective inhibitor, gallein and GRK2i, or with $\text{G}\beta 1$ siRNA or CP-25, led to the decreased translocation of GRK2 and EP4

re-sensitization. In this regard, EP4 desensitization via the GRK2-G $\beta\gamma$ interaction could be a novel therapeutic target to treat RA. Although the present study was focused primarily on the interaction between GRK2 and G $\beta\gamma$, it was found that GRK2, G $\beta\gamma$ and EP4 interacted with each other. It is a possibility that these three proteins bind directly to form a ternary complex. Further research is required to demonstrate the complex formed by EP4, GRK2 and G $\beta\gamma$ in a sequential order, as well as the association and disassociation of it. Several issues, including the details of G $\beta\gamma$ releases from G α -GDP, the binding site of G $\beta\gamma$ with GRK2, and the G $\beta\gamma$ downstream signal pathways, remain to be further confirmed.

The limitation of the present study is that G β antibody was selected instead of G $\beta\gamma$. Although distinct G β and G γ antibodies may reveal in an improved way the activity of G $\beta\gamma$, in the present study, main focus was addressed on the interaction between G $\beta\gamma$ and GRK2; it is considered that one kind of G β antibodies may not be optimal, but should be sufficient to draw the conclusion of the present study. It is considered that future work is needed to explore changes of different subunits of G β and G γ .

Collectively, it was revealed that GRK2-G $\beta\gamma$ interaction was enhanced in RA synovial tissue and FLS. PGE₂-induced abnormal proliferation and migration in FLS were associated with strengthened GRK2-G $\beta\gamma$ interaction and EP4 desensitization. Regulating EP4-GRK2-G $\beta\gamma$ signaling may be one of the mechanisms that underlie the salutary effect of CP-25. Additional *in vivo* studies are required to confirm the therapeutic potential of the interfering GRK2-G $\beta\gamma$ interaction in RA.

Acknowledgements

Not applicable.

Funding

The present study was supported by the National Natural Science Foundation of China (grant no. 81330081).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YZ participated in the design of the study, performed most of the experiments and wrote the manuscript. XY, CH and DW performed experiments. YM and WW conceived the study and revised the manuscript. YZ and DW confirm the authenticity of all the raw data. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The study protocol was approved (approval no. 20131321) by the Biomedical Ethics Committee of Anhui Medical University (Hefei, China). Informed consent was obtained from all patients for the use of their samples in scientific research.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Benson RA, McInnes IB, Brewer JM and Garside P: Cellular imaging in rheumatic disease. *Nat Rev Rheumatol* 11: 357-367, 2015.
- Guo Q, Wang Y, Xu D, Nossent J, Pavlos NJ and Xu J: Rheumatoid arthritis: Pathological mechanisms and modern pharmacologic therapies. *Bone Res* 6: 15, 2018.
- Ma Y, Hong FF and Yang SL: Role of prostaglandins in rheumatoid arthritis. *Clin Exp Rheumatol* 39: 162-172, 2021.
- Dupré DJ, Robitaille M, Rebois RV and Hébert TE: The role of Gbetagamma subunits in the organization assembly and function of GPCR signaling complexes. *Annu Rev Pharmacol Toxicol* 49: 31-56, 2009.
- Jia X, Chang Y, Wei F, Dai X, Wu YJ, Sun XJ, Xu S, Wu HX, Wang C, Yang XZ and Wei W: CP-25 reverses prostaglandin E4 receptor desensitization-induced fibroblast-like synoviocyte dysfunction via the G protein-coupled receptor kinase 2 in autoimmune arthritis. *Acta Pharmacol Sin* 40: 1029-1039, 2019.
- Yang X, Zhao Y, Jia X, Wang C, Wu Y, Zhang L, Chang Y and Wei W: CP-25 combined with MTX/ LEF ameliorates the progression of adjuvant-induced arthritis by the inhibition on GRK2 translocation. *Biomed Pharmacother* 110: 834-843, 2019.
- Han C, Li Y, Zhang Y, Wang Y, Cui D, Luo T, Zhang Y, Liu Q, Li H, Wang C, *et al*: Targeted inhibition of GRK2 kinase domain by CP-25 to reverse fibroblast-like synoviocytes dysfunction and improve collagen-induced arthritis in rats. *Acta Pharm Sin B* 11: 1835-1852, 2021.
- Chen J, Wang Y, Wu H, Yan S, Chang Y and Wei W: A modified compound from paeoniflorin, CP-25, suppressed immune responses and synovium inflammation in collagen-induced arthritis mice. *Front Pharmacol* 9: 563, 2018.
- Tu J, Guo Y, Hong W, Fang Y, Han D, Zhang P, Wang X, Körner H and Wei W: The regulatory effects of paeoniflorin and its derivative paeoniflorin-6'-O-benzene sulfonate CP-25 on inflammation and immune diseases. *Front Pharmacol* 10: 57, 2019.
- Shu J, Zhang XZ, Han L, Zhang F, Wu Y, Tang XY, Wang C, Tai Y, Wang QT, Chen JY, *et al*: Paeoniflorin-6'-O-benzene sulfonate alleviates collagen-induced arthritis in mice by down-regulating BAFF-TRAF2-NF- κ B signaling: Comparison with biological agents. *Acta Pharmacol Sin* 40: 801-813, 2019.
- Zhang BJ, Wang YY, Jia CY, Li SS, Wang XW, Xu Y, Chen AY, Xu HP, Wang C, Yang ZY, *et al*: Paeoniflorin-6'-o-benzene sulfonate ameliorates the progression of adjuvant-induced arthritis by inhibiting the interaction between Ahr and GRK2 of fibroblast-like synoviocytes. *Int Immunopharmacol* 108: 108678, 2022.
- Li J, Xiang B, Su W, Zhang X, Huang Y and Ma L: Agonist-induced formation of opioid receptor-G protein-coupled receptor kinase (GRK)-G beta gamma complex on membrane is required for GRK2 function in vivo. *J Biol Chem* 278: 30219-30226, 2003.
- Rudomanova V and Blaxall BC: Targeting GPCR-G $\beta\gamma$ -GRK2 signaling as a novel strategy for treating cardiorenal pathologies. *Biochim Biophys Acta Mol Basis Dis* 1863: 1883-1892, 2017.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Wang DD, Jiang MY, Wang W, Zhou WJ, Zhang YW, Yang M, Chen JY and Wei W: Paeoniflorin-6'-O-benzene sulfonate down-regulates CXCR4-G $\beta\gamma$ -PI3K/AKT mediated migration in fibroblast-like synoviocytes of rheumatoid arthritis by inhibiting GRK2 translocation. *Biochem Biophys Res Commun* 526: 805-812, 2020.
- Chang Y, Jia X, Wei F, Wang C, Sun X, Xu S, Yang X, Zhao Y, Chen J, Wu H, *et al*: CP-25, a novel compound, protects against autoimmune arthritis by modulating immune mediators of inflammation and bone damage. *Sci Rep* 6: 26239, 2016.

17. Homan KT and Tesmer JG: Molecular basis for small molecule inhibition of G protein-coupled receptor kinases. *ACS Chem Biol* 10: 246-256, 2015.
18. Murthy KS, Mahavadi S, Huang J, Zhou H and Sriwai W: Phosphorylation of GRK2 by PKA augments GRK2-mediated phosphorylation, internalization, and desensitization of VPAC2 receptors in smooth muscle. *Am J Physiol Cell Physiol* 294: C477-C487, 2008.
19. Han CC, Liu Q, Zhang Y, Li YF, Cui DQ, Luo TT, Zhang YW, Wang XM, Wang C, Ma Y and Wei W: CP-25 inhibits PGE₂-induced angiogenesis by down-regulating EP4/AC/cAMP/PKA-mediated GRK2 translocation. *Clin Sci (Lond)* 134: 331-347, 2020.
20. Penela P, Ribas C and Mayor F Jr: Mechanisms of regulation of the expression and function of G protein-coupled receptor kinases. *Cell Signal* 15: 973-981, 2003.
21. Jaber M, Koch WJ, Rockman H, Smith B, Bond RA, Sulik KK, Ross J Jr, Lefkowitz RJ, Caron MG and Giros B: Essential role of beta-adrenergic receptor kinase 1 in cardiac development and function. *Proc Natl Acad Sci USA* 93: 12974-12979, 1996.
22. Tutunea-Fatan E, Abd-Elrahman KS, Thibodeau JF, Holterman CE, Holleran BJ, Leduc R, Kennedy CRJ, Gros R and Ferguson SSG: GRK2 knockdown in mice exacerbates kidney injury and alters renal mechanisms of blood pressure regulation. *Sci Rep* 8: 11415, 2018.
23. Hullmann J, Traynham CJ, Coleman RC and Koch WJ: The expanding GRK interactome: Implications in cardiovascular disease and potential for therapeutic development. *Pharmacol Res* 110: 52-64, 2016.
24. Kamal FA, Travers JG, Schafer AE, Ma Q, Devarajan P and Blaxall BC: G protein-coupled receptor-G-protein $\beta\gamma$ -subunit signaling mediates renal dysfunction and fibrosis in heart failure. *J Am Soc Nephrol* 28: 197-208, 2017.
25. Schumacher SM, Gao E, Zhu W, Chen X, Chuprun JK, Feldman AM, Tesmer JJ and Koch WJ: Paroxetine-mediated GRK2 inhibition reverses cardiac dysfunction and remodeling after myocardial infarction. *Sci Transl Med* 7: 277ra31, 2015.
26. Murga C, Arcones AC, Cruces-Sande M, Briones AM, Salaices M and Mayor Jr F: G protein-coupled receptor kinase 2 (GRK2) as a potential therapeutic target in cardiovascular and metabolic diseases. *Front Pharmacol* 10: 112, 2019.
27. Casey LM, Pistner AR, Belmonte SL, Migdalovich D, Stolpnik O, Nwakanma FE, Vorobiof G, Dunaevsky O, Matavel A, Lopes CMB, *et al*: Small molecule disruption of G signaling inhibits the progression of heart failure. *Circ Res* 107: 532-539, 2010.
28. Daaka Y, Luttrell LM and Lefkowitz RJ: Switching of the coupling of the beta2-adrenergic receptor to different G proteins by protein kinase A. *Nature* 390: 88-91, 1997.
29. Senarath K, Kankanamge D, Samaradivakara S, Ratnayake K, Tennakoon M and Karunarathne A: Regulation of G protein $\beta\gamma$ signaling. *Int Rev Cell Mol Biol* 339: 133-191, 2018.
30. Tang X, Sun Z, Runne C, Madsen J, Domann F, Henry M, Lin F and Chen S: A critical role of Gbetagamma in tumorigenesis and metastasis of breast cancer. *J Biol Chem* 286: 13244-13254, 2011.
31. Xu S, Zhang Z, Ogawa O, Yoshikawa T, Sakamoto H, Shibasaki N, Goto T, Wang L and Terada N: An EP4 antagonist ONO-AE3-208 suppresses cell invasion, migration, and metastasis of prostate cancer. *Cell Biochem Biophys* 70: 521-527, 2014.
32. Liu Y, Pan YF, Xue YQ, Fang LK, Guo XH, Guo X, Liu M, Mo BY, Yang MR, Liu F, *et al*: uPAR promotes tumor-like biologic behaviors of fibroblast-like synoviocytes through PI3K/Akt signaling pathway in patients with rheumatoid arthritis. *Cell Mol Immunol* 15: 171-181, 2018.
33. Nygaard G and Firestein GS: Restoring synovial homeostasis in rheumatoid arthritis by targeting fibroblast-like synoviocytes. *Nat Rev Rheumatol* 16: 316-333, 2020.
34. Rajagopal S and Shenoy SK: GPCR desensitization: Acute and prolonged phases. *Cell Signal* 41: 9-16, 2018.
35. Smith JS and Rajagopal S: The β -Arrestins: Multifunctional regulators of G protein-coupled receptors. *J Biol Chem* 291: 8969-8977, 2016.
36. Kamal FA, Smrcka AV and Blaxall BC: Taking the heart failure battle inside the cell: Small molecule targeting of G $\beta\gamma$ subunits. *J Mol Cell Cardiol* 51: 462-467, 2011.