

Association between long non-coding RNA HOTAIR polymorphism and lung cancer risk: A systematic review and meta-analysis

CHUNLIN KE^{1*}, XUEGANG FENG^{2*}, JIE LI³, SIYU CHEN³ and XINYU HU³

¹Department of Radiotherapy, Cancer Center, The First Affiliated Hospital of Fujian Medical University, Fuzhou, Fujian 350005; Departments of ²Cardio-Thoracic Surgery and ³Oncology, 900 Hospital of The Joint Logistics Team, Fuzhou, Fujian 350025, P.R. China

Received April 12, 2022; Accepted June 7, 2022

DOI: 10.3892/etm.2022.11477

Abstract. Single nucleotide polymorphism (SNP) of long noncoding RNA (lncRNA) has been reported to be an important factor in cancer development. Recently, lncRNA homeobox transcript antisense intergenic RNA (HOTAIR) was indicated to induce tumorigenesis of several cancer types, but the association between the SNP of lncRNA HOTAIR and lung cancer susceptibility has remained undetermined. The present meta-analysis aimed to investigate the effect of HOTAIR polymorphism on susceptibility to lung cancer. The PubMed, Ovid Medline, Embase and Cochrane Library databases were thoroughly searched. Studies containing data on the incidence of lung cancer in patients with different HOTAIR SNPs were included. The Hardy-Weinberg equilibrium was analyzed to determine genotype distribution and allele frequencies. The odds ratio (OR) was pooled to evaluate the association of different SNPs with the susceptibility to lung cancer. A total of six studies comprising 1,715 patients with lung cancer and 2,745 healthy controls were finally included. A total of 4 SNPs (rs12826786, rs1899663, rs920778 and rs4759314) were reported. Analyses for all of these SNPs individually indicated that the lncRNA HOTAIR rs1899663 C>A polymorphism was a risk factor for lung cancer (dominant mode, AA+CA vs. CC: OR=0.816, 95% CI=0.707-0.942, P=0.005). The present study was the first meta-analysis investigating the association between lncRNA HOTAIR and lung cancer susceptibility. The results indicated that the lncRNA HOTAIR rs1899663 C>A polymorphism is a risk factor for lung cancer. lncRNA HOTAIR may be of value in lung

cancer screening, particularly for populations with high-risk factors, as well as prognosis prediction. Future investigations are required to further clarify the intrinsic mechanism of the role of HOTAIR in the oncogenesis of lung cancer.

Introduction

Lung cancer has brought a heavy burden to human society and families, with high morbidity and mortality rates among all types of malignant tumors (1). The statistical report on cancer incidence and mortality published by the International Cancer Center (GLOBOCAN 2018) suggests that lung cancer is a leading cause of death of patients with cancer in China and worldwide (2). Globally, it is estimated that there are 2.1 million new cases of lung cancer and 1.8 million lung cancer-related deaths each year (2). In China, the incidence of lung cancer is reported to be 5.7 per million and the total number of new cases is ~787,000, accounting for 20% of the total malignant tumor cases (3).

The occurrence and development of lung cancer are closely related to the tumor micro-environment, personal living habits and genetic differences (4,5). Smoking, history of lung-related diseases, air pollutants, family history of tumors, ethnic differences, obesity, occupational exposure and poor eating habits may increase the risk of lung cancer (6). Although smoking is a well-recognized risk factor for lung cancer, lung cancer also occurs in 10-25% of non-smokers. The influence of personal genetic factors on the pathogenesis of lung cancer should not be underestimated (7,8).

With the rise of molecular epidemiology, the ENCODE project indicated that up to 75% of human genome nucleotides may be transcribed into the original transcripts, producing a series of common RNAs including non-coding RNAs (ncRNA) (9). A large number of studies have indicated that most ncRNAs do not encode proteins but have specific functions (10). At the level of apparent modification, long ncRNA (lncRNA) affects numerous aspects of chromosome biology by remodeling chromatin structure and regulating a large number of genes. Based on its various characteristics and regulatory mechanisms, the abnormal expression of lncRNA may also be involved in the occurrence, development and metastasis of

Correspondence to: Dr Jie Li, Department of Oncology, 900 Hospital of The Joint Logistics Team, 156 Xierhuan Northern Road, Fuzhou, Fujian 350025, P.R. China
E-mail: arthur643@163.com

*Contributed equally

Key words: lncRNA, HOTAIR, SNP, lung cancer, susceptibility

tumors, thus affecting tumor staging and the occurrence of various cancers (11,12). The potential impact of lncRNA on the occurrence, development, treatment and prognosis of lung cancer is worthy of further investigation (13).

Single nucleotide polymorphism (SNP) refers to the polymorphism in the DNA sequence caused by the variation of a single nucleotide in the genome. It is a heritable variation that exists widely in the human genome (14). It is estimated that there is one SNP per 1,000 bases on average, which is recognized as the third-generation genetic marker after restriction fragment length polymorphism and microsatellites. SNP is involved in the transcription of coding genes by influencing the binding of transcription factors, thereby affecting gene expression. Researchers have discovered a large number of SNPs associated with diseases through genome-wide association studies and about one-third of the SNPs are located in non-protein coding regions, indicating the SNPs of lncRNA may also have an important role in disease (15). Emerging evidence suggested that the SNPs of lncRNAs are related to the susceptibility to and prognosis of various types of malignant tumor, such as breast cancer, gastric cancer and colorectal cancer (16).

In recent years, researchers have discovered that the homeobox (HOX) gene family participates in embryonic development (17). In their study from 2007, Rinn *et al* (18) used high-resolution microarray technology to identify and analyze hundreds of ncRNAs transcribed from the HOX gene cluster from 11 types of fibroblasts and discovered HOX transcript antisense intergenic RNA (HOTAIR) at the HOXC gene locus located on chromosome 12. HOTAIR was recently considered a novel oncogene, but it is controversial whether the SNP of lncRNA HOTAIR has a risk effect to influence lung cancer susceptibility (19). lncRNA HOTAIR is located at 12q13.13 and the length is 2.1 kb. There are four gene clusters in the genome: HOXA, HOXB, HOXC and HOXD (20). The transcription of these four gene clusters produces several long-chain ncRNAs and their expression has a temporal and spatial specificity (21). Although HOTAIR is produced from the HOXC sequence, it has the function of inhibiting the expression of the HOXD gene. The secondary structure of HOTAIR contains four independent folding units, two of which are evolutionarily conserved protein-binding domains (22). HOTAIR is produced and enriched in the nucleus and expressed in the cytoplasm. The HOTAIR-mediated epigenetic silencing function works by acting as a scaffold for the polycomb repressive complex 2 (PRC2) and lysine specific demethylase 1/RE1-silencing transcription factor (REST) co-repressor 1/REST complex. PRC2 binds to the 5' end of HOTAIR and regulates chromosome occupancy through the subunit EZH2 of the PRC2 complex, leading to the methylation of histone H3 lysine in the chromosome sequence region of the HOXD gene (23). Studies have indicated that at least 16 types of malignant tumor have increased expression of HOTAIR (24-26). Furthermore, it was reported that overexpressed HOTAIR combines with PRC2 to regulate gene expression patterns by changing the methylation status of H3K27, thus increasing the ability of cancer cells to metastasize and invade. Knockdown of HOTAIR or PRC2 is able to inhibit the invasion and metastasis ability of cancer cells (27). A number of studies focusing on lung cancer have

reported that HOTAIR has a proto-oncogene function in lung cancer. For instance, compared with adjacent tissues, the expression of HOTAIR in lung cancer tissue is significantly higher (28). The increased HOTAIR expression in non-small cell lung cancer is similar to that of lung adenocarcinoma and lung squamous cells (29). Another study determined that elevated HOTAIR expression was associated with the brain metastasis rate of lung cancer (30). The association between the SNP of lncRNA HOTAIR and lung cancer susceptibility remains to be further investigated.

Thus, the present systematic review and meta-analysis was performed to investigate the association between lncRNA HOTAIR SNP and lung cancer susceptibility.

Materials and methods

Literature search and search strategy. This systematic review and meta-analysis was performed following the preferred reporting items for systematic review and meta-analysis (PRISMA) guidelines (31). A computerized literature search was performed in the PubMed (<https://pubmed.ncbi.nlm.nih.gov/>), Ovid Medline (<https://www.wolterskluwer.com/en/solutions/ovid/ovid-medline-901>), Embase (<https://www.embase.com>) and Cochrane Library databases (<https://www.cochranelibrary.com/>) up to May 2nd 2021. Relevant studies were also searched in Chinese databases, including Chinese National Knowledge Infrastructure (CNKI; <https://www.cnki.net/>) and Wanfang (<https://www.wanfangdata.com.cn/index.html>). Google scholar (<https://scholar.google.com/>) was searched for gray literature that is not formally published in sources such as books or journal articles.

Regarding the search strategy, terms included the following items: ('lncRNA' or 'long non-coding RNA') and ('SNP' or 'polymorphism' or 'variants') and ('lung cancer' or 'lung tumor' or 'lung neoplasm' or 'adenocarcinoma' or 'lung carcinoma') and ('HOTAIR' or 'HOX transcript antisense intergenic RNA'). The items were searched for keywords and medical sub-headings. All results were downloaded and imported into Endnote X6 (Thomson Reuters) for further literature screening.

Inclusion and exclusion criteria. All of the studies evaluating and assessing the association between lncRNA HOTAIR polymorphism and the susceptibility to lung cancer were included in the present study. The inclusion criteria were as follows: i) Patients with lung cancer; ii) case-control studies on the polymorphism of lncRNA HOTAIR; iii) number of patients >50. The exclusion criteria were as follows: i) Reviews, observational studies, comments, case series and case reports; ii) lack of data on lncRNA HOTAIR; iii) lack of information on lung cancer risk; and iv) lack of available data that were able to be extracted from the text/table/graphs or obtained from the authors on request. There was no limitation in terms of the language of the studies included and the reference lists of reviews were also screened for the retrieval and further inclusion of studies.

Literature screening, data extraction and quality assessment. A total of two investigators (CK and XF) independently screened titles and abstracts according to the inclusion and




Figure 1. Flowchart of the literature screening. Records were searched via databases (n=46) and other sources (n=8). Finally, a total of 6 records were considered eligible and included in the present meta-analysis. SNP, single nucleotide polymorphism; lncRNA, long noncoding RNA; HOTAIR, homeobox transcript antisense intergenic RNA.

exclusion criteria. The full-text was further screened for data extraction or assessed for the inclusion criteria. In the case of any disagreements, a third investigator (XH) was consulted to resolve the divergence.

The data were extracted by two reviewers independently (CK and XF). All the data were imported into a standard form, including the name of the first author of the study, publication year, recruitment years and country, genotyping methods and the common polymorphism of HOTAIR. In addition, the sample size, as well as the age, sex, smoking history and alcohol use of healthy participants (control group) and patients with lung cancer (case group) were recorded. The raw data are provided as supplemental material (Table SI).

The study quality was evaluated according to Newcastle-Ottawa Scale (NOS), with a score of 7-9 considered to indicate high quality, whereas a score of 0-6 was indicative of low quality (32).

Statistical analysis. The meta-analysis was performed with Stata 15.0 software (StataCorp LP). The genotype frequencies of lncRNA HOTAIR polymorphisms for the Hardy-Weinberg equilibrium (HWE) were calculated for the controls using the χ^2 test and $P < 0.05$ was considered to indicate a significant disequilibrium. The risk was pooled by the odds ratio (OR) and 95% confidence intervals (CI). The

result of the pooled OR was visualized through the forest plot by using the *metan* module of Stata. The heterogeneity existing among studies was evaluated by the I^2 statistic and χ^2 test, with $I^2 \geq 50\%$ indicating the presence of heterogeneity. The random-effects model was used if heterogeneity was present among studies, while otherwise, the fixed-effects model was used. The risks of the heterozygote and variant homozygote compared with the wild-type homozygote were estimated. Furthermore, the dominant and recessive mode of the variant allele were evaluated. The sensitivity analysis was performed by reperforming the OR calculation while omitting one individual study at a time. Sensitivity analysis was performed for rs12826786, rs1899663, rs4759314 and rs920778 for all modes, including the dominant mode, recessive mode, additive mode, heterozygote mode, homozygote mode and allele mode. Egger's linear regression method was used to statistically assess the publication bias by Stata 15.0. The funnel plot for exploring the missed-reported articles was also analysed with the *metafunnel* module of Stata to visually display the outcomes of publication bias assessment. Subgroup analysis was performed using the *metan* module of Stata. The further heterogeneity analysis was performed based on the results of the subgroup analysis. $P < 0.05$ was considered to indicate statistical significance for all analyses.

Table I. Characteristics of the included studies.

Author (year)	Country	Recruitment year	Cancer type	Ethnicity	Genotyping method	Quality score	Control group					Case group					(Refs.)
							Sample, n	Age, years	Male sex, n (%)	Smoking, n (%)	Alcohol consumption, n (%)	Sample, n	Age, years	Male sex, n (%)	Smoking, n (%)	Alcohol consumption, n (%)	
Ren (2020)	China	2015-2017	All lung cancer	Asian	MassArray system	8	196	64.82±10.00	139 (70.9)	190 (96.9)	188 (95.9)	196	64.32±10.26	136 (69.4)	195 (99.4)	196 (100)	(33)
Minn (2020)	Japan	1995-2006	All lung cancer	Asian	TaqMan	8	1,241	NG	NG	NG	NG	129	NG	NG	NG	NG	(34)
Wang (2018)	China	2006-2010	All lung cancer	Asian	Assay-PCR	7	451	NG	280 (62.1)	159 (35.2)	NG	262	NG	188 (71.8)	169 (64.5)	NG	(35)
Li (2018)	China	NG	All lung cancer	Asian	TaqMan	7	551	56.69±15.65	165 (29.9)	111 (20.1)	NG	543	57.99±11.51	192 (35.4)	165 (30.4)	NG	(36)
Dadaş (2018)	Turkey	NG	NSCLC	Caucasian	Assay-PCR	6	93	61.77±12.00	67 (72.0)	61 (65.6)	NG	87	59.27±10.55	71 (81.6)	26 (29.9)	NG	(37)
Gong (2016)	China	2011-2013	All lung cancer	Asian	TaqMan	5	213	NG	80	NG	NG	498	NG	394 (79.1)	NG	NG	(38)

NG, not given; NSCLC, non-small cell lung cancer.

Results

Study selection and characteristics of included studies. A total of 46 studies were retrieved by searching the databases as described above, as well as 8 studies via other methods. Thus, a total of 54 studies focusing on the association of lncRNA HOTAIR with lung cancer were identified based on the search strategy. After excluding duplicates and other studies based on the inclusion and exclusion criteria, six studies were finally included in the present study (33-38). The literature screening process is presented in Fig. 1.

The characteristics of the included studies are described in Table I. The year of publication ranged from 2016 to 2020 and the recruitment period ranged from 1995 to 2017. Of the six studies, four were from China and the other two studies were from Turkey and Japan, respectively. Furthermore, two studies used MassArray to detect the SNPs of lncRNA, while the other four studies used the TaqMan assay instead. A total of 1,715 patients with lung cancer and 2,745 healthy participants were included in the analysis of the present study. The average age of the lung cancer group was 57.9-64.3 years, while it was 56.6-64.8 years for the healthy participants (control group). The proportion of male subjects ranged from 29.9 to 96.9% in the control group, as compared to 35.4-81.6% in the lung cancer group. A history of smoking was present in 20.1-96.6% of the healthy participants and in 29.9-99.4% of the patients with lung cancer. The results of the quality assessment are listed in Table I. Of the studies, four scored as >6 and were considered to be of high quality.

Genotype frequency distributions. Table II displays the genotype frequency distributions of the five SNPs of lncRNA HOTAIR. Regarding HOTAIR rs128261786 and HOTAIR rs1899663, $P_{HWE} < 0.05$ was calculated for the study by Wang *et al* (35), indicating unbalanced data in terms of the two SNPs.

Main findings. The association between different HOTAIR SNPs and lung cancer susceptibility is presented in Table III. No significant association with the susceptibility for lung cancer was obtained for SNPs rs12826786, rs4359714 and rs920778 ($P > 0.05$ for all modes). However, the lncRNA HOTAIR rs1899663 C>A polymorphism was a risk factor for lung cancer (dominant mode, AA+CA vs. CC: OR=0.816, 95% CI=0.707-0.942, $P=0.005$). However, for the recessive mode, additive mode, heterozygote mode, homozygote mode and allele mode, no significant association with lung cancer was observed.

The sensitivity analysis suggested that the omission of any single study did not significantly alter the overall pooled ORs in dominant mode, recessive mode, additive mode, heterozygote mode, homozygote mode and allele mode (Fig. 2).

Since the lncRNA HOTAIR rs1899663 C>A polymorphism was identified as a risk factor for lung cancer (OR=0.816; Fig. 3A), a further analysis was performed, including a publication bias analysis and subgroup analysis in terms of rs1899663 (dominant mode, AA+CA vs. CC). The P-value as calculated using Egger's test (Fig. 3B) was 0.466 and the funnel plot indicated a good symmetry (Fig. 3C). These results suggested that no significant publication bias was present. In the further subgroup analysis, the studies with higher quality

Table II. Common gene polymorphisms of homeobox transcript antisense intergenic RNA in the included studies.

A, rs920778

Author (year)	Controls				Cases				P _{HWE}	(Refs.)
	Total	AA	AG	GG	Total	AA	AG	GG		
Ren (2020)	183	104	69	10	184	114	60	10	0.7416	(33)
Minn (2020)	1,241	698	473	70	129	75	53	1	0.3844	(34)
Wang (2018)	451	233	192	26	262	110	132	20	0.0947	(35)
Li (2018)	551	326	203	22	543	324	197	22	0.1640	(36)

B, rs12826786

Author (year)	Controls				Cases				P _{HWE}	(Refs.)
	Total	AA	AG	GG	Total	AA	AG	GG		
Wang (2018)	451	355	85	11	262	200	49	13	0.0047	(35)
Li (2018)	551	381	156	14	543	340	185	18	0.6767	(36)
Dadaş (2018)	93	38	41	14	87	33	41	13	0.5934	(37)

C, rs1899663

Author (year)	Controls				Cases				P _{HWE}	(Refs.)
	Total	AA	AG	GG	Total	AA	AG	GG		
Ren (2020)	188	123	60	5	187	131	52	4	0.4656	(33)
Wang (2018)	452	254	189	9	262	178	66	18	0.0001	(35)
Dadaş (2018)	93	41	40	12	87	39	37	11	0.6495	(37)

D, rs4759314

Author (year)	Controls				Cases				P _{HWE}	(Refs.)
	Total	AA	AG	GG	Total	AA	AG	GG		
Wang (2018)	451	412	37	2	262	238	21	3	0.2463	(35)
Ren (2020)	184	168	16	0	175	158	17	0	0.5375	(33)
Li (2018)	551	489	61	1	543	526	16	1	0.5268	(36)

HWE, Hardy-Weinberg equilibrium.

(NOS score >6) also suggested a significant association of the rs1899663 C>A polymorphism with the risk of lung cancer (OR=0.77, 95% CI=0.65-0.92, P=0.003; Fig. 4A). Similar results were obtained in the subgroup of cancer type (lung cancer) (Fig. 4B). As for the studies with P_{HWE}≥0.05 (Fig. 4C), the trend for the risk of lung cancer was similar (OR=0.92, 95% CI=0.75-1.12, P=0.406). As for the study by Wang *et al* (35) with P_{HWE}<0.05, a significant association was observed (OR=0.73, 95% CI=0.60-0.90, P=0.003). Of note, in each of the subgroups, no heterogeneity was detected (I²=0 for all subgroups with the number of studies >1), indicating that

the NOS quality, type of cancer and gene frequency disequilibrium may be potential sources of heterogeneity. However, the overall heterogeneity for the rs1899663 C>A polymorphism was minor (I²=37.7%, P=0.201).

Discussion

The present study indicated that the lncRNA HOTAIR rs1899663 C>A polymorphism is a risk factor for lung cancer. In previous studies on the topic of HOTAIR-associated lung cancer susceptibility, two or three SNPs were usually

Table III. Association of homeobox transcript antisense intergenic RNA gene polymorphisms and lung cancer risk.

SNP/mode	Comparison	N	OR (95% CI)	P-value	I ² (%)	Model for meta-analysis
rs12826786						
Dominant	TT+CT vs. CC	3	0.964 (0.793-1.172)	0.716	56.30	Random-effects
Recessive	TT vs. CC+CT	3	1.101 (0.732-1.657)	0.643	42.50	Fixed-effects
Additive	TT+CC vs. TC	3	1.043 (0.987-1.102)	0.134	26.40	Fixed-effects
Heterozygote	CT vs. CC	3	0.899 (0.783-1.032)	0.131	29.60	Fixed-effects
Homozygote	TT vs. CC	3	1.085 (0.730-1.613)	0.686	49.90	Fixed-effects
Allele	T vs. C	3	0.990 (0.782-1.254)	0.934	66.50	Random-effects
rs1899663						
Dominant	AA+CA vs. CC	3	0.816 (0.707-0.942)	0.005	37.70	Fixed-effects
Recessive	AA vs. CC+CA	3	1.527 (0.552-4.225)	0.415	72.10	Random-effects
Additive	AA+CC vs. AC	3	1.135 (0.969-1.331)	0.117	69.90	Random-effects
Heterozygote	CA vs. CC	3	0.801 (0.607-1.056)	0.116	64.50	Random-effects
Homozygote	AA vs. CC	3	1.382 (0.588-3.247)	0.457	62.30	Random-effects
Allele	A vs. C	3	0.886 (0.762-1.030)	0.115	0.00	Fixed-effects
rs4759314						
Dominant	GG+AG vs. AA	3	1.630 (0.720-3.690)	0.241	85.00	Random-effects
Recessive	GG vs. AA+AG	2	1.933 (0.447-8.350)	0.377	0.00	Fixed-effects
Additive	AA+GG vs. AG	3	0.966 (0.906-1.030)	0.285	83.60	Random-effects
Heterozygote	AG vs. AA	3	1.612 (0.670-3.878)	0.286	86.20	Random-effects
Homozygote	GG vs. AA	2	1.983 (0.456-8.617)	0.361	0.00	Fixed-effects
Allele	G vs. A	3	1.649 (0.769-3.537)	0.198	82.90	Random-effects
rs920778						
Dominant	GG+AG vs. AA	4	1.029 (0.902-1.173)	0.670	53.60	Random-effects
Recessive	GG vs. AA+AG	4	0.914 (0.640-1.304)	0.620	43.80	Fixed-effects
Additive	AA+GG vs. AG	4	0.967 (0.908-1.029)	0.289	36.90	Fixed-effects
Heterozygote	AG vs. AA	4	1.050 (0.957-1.151)	0.305	42.90	Fixed-effects
Homozygote	GG vs. AA	4	0.984 (0.555-1.747)	0.957	53.60	Random-effects
Allele	G vs. A	4	1.008 (0.865-1.176)	0.917	57.70	Random-effects

SNP, single nucleotide polymorphism; OR, odds ratio.

investigated in an individual study. However, at least four SNPs (rs128261786, rs1899663, rs920778 and rs4759314) of HOTAIR have been reported in the lung cancer population. The present meta-analysis included all of these SNPs into the analysis and therefore arrived at a more comprehensive conclusion. The previous results remain controversial in that each SNP has been reported to be related to lung cancer in one or two of these previous studies. However, in the present meta-analysis, these data were integrated and according to the results, only rs1899663 was a significant risk factor. To the best of our knowledge, the present study was the first systematic review and meta-analysis focusing on the association between lncRNA HOTAIR and lung cancer susceptibility that has integrated all these four SNPs into its analysis, which is the novelty of this work.

Genetic differences in DNA sequences lead to phenotypic variation, which affects changes in phenotypic characteristics, disease risks and responses to the environment (39). At the same time, human genetic diversity is not limited to individual polymorphisms, but also to the specific combinations of tightly

linked alleles. The most common form of gene polymorphism is SNP, which refers to the polymorphism in the DNA sequence caused by the variation of a single nucleotide base in the genome. The types of SNP include conversion, transversion, insertion and deletion (40). SNP is associated with the function and expression level of the downstream protein by changing the protein-coding sequence and influencing the splicing changes of exons and introns, thereby affecting the individual's susceptibility to disease, prognosis and responsiveness to drugs (41).

To date, only a small number of studies have reported on the relationship between the SNP of the HOTAIR sequence and the susceptibility to malignancies. A study reported an enhancer-like sequence on the intron 1,719-2,353 bp downstream of the transcription start position of HOTAIR (42). There was a risk site for esophageal squamous cell carcinoma, rs920778, carrying the rs920778T allele. The expression level of the reported gene was higher than that of the rs920778C allele. This study also performed SNP typing and quantitative PCR using esophageal squamous




Figure 2. Sensitivity analysis for each single nucleotide polymorphism. The sensitivity was performed by omitting one study at a time. In each row of the forest plot, one study was omitted. Sensitivity analysis was performed for rs12826786, rs1899663, rs4759314 and rs920778 for all modes including (A) dominant mode, (B) recessive mode, (C) additive mode, (D) heterozygote mode, (E) homozygote mode and (F) allele mode.

cell carcinoma tissues and determined that individuals with the genotype rs920778TT had higher expression of HOTAIR than patients carrying rs920778CC (42). A study on papillary thyroid carcinoma indicated that in females, the rs920778TT and rs920778CT genotype increased the risk of developing papillary thyroid carcinoma by 0.75- and 0.46-fold, respectively (43). The study also verified the effect of the rs920778 locus polymorphism on HOTAIR expression in cells and tissues. It was indicated that cells carrying rs920778T alleles had higher HOTAIR expression than those with rs920778C alleles. In thyroid squamous cell carcinoma tissues and normal tissues, the expression of HOTAIR in cancer tissues and adjacent normal tissues of rs920778TT and rs920778CT genotype carriers was higher than that in rs920778CC genotype carriers (43). Previous results suggested that the expression of HOTAIR in individuals carrying the rs920778T allele was higher than that in individuals with rs920778C (44). Except for the proto-oncogene role in lung cancer, HOTAIR has also been indicated to be related to resistance to chemotherapy.

Liu *et al* (45) reported that HOTAIR expression was significantly upregulated in cisplatin-resistant lung cancer cells and that knockdown of HOTAIR by RNA interference was able to resensitize the responses to cisplatin both *in vitro* and *in vivo*. On the other hand, overexpression of HOTAIR decreased the sensitivity of lung cancer cells to cisplatin. The HOTAIR-associated chemosensitivity regulation was indicated to involve the inhibition of cell proliferation, induction of G0/G1 cell-cycle arrest and apoptosis enhancement through regulation of cyclin-dependent kinase inhibitor p21^{WAF1}/CIP1 (p21) expression (45). Accordingly, another clinical investigation by the same group (28) suggested that high levels of HOTAIR expression were correlated with a more advanced pathological stage of non-small cell lung cancer (NSCLC), lymph-node metastasis, as well as a relatively poor prognosis. The upregulated HOTAIR may regulate NSCLC cell invasion and metastasis, partially via downregulation of HOXA5 (28). In the present study, a critical literature review was performed to further understand the mechanisms of HOTAIR in cancer development.




Figure 3. Analysis for the rs1899663 C>A polymorphism. (A) The forest plot indicates the pooled OR for the effect of the rs1899663 C>A polymorphism on the risk of lung cancer. (B) Egger's test plot and (C) funnel plot for assessment of publication bias. OR, odds ratio; SND, standard normal distribution; se, standard error.

Apart from the p21^{WAF1}/CIP1 and HOXA5 pathways, the most commonly acknowledged mechanism of HOTAIR is through the axis of microRNA (miR) and downstream signaling molecules. These axes include but are not limited to the following: HOTAIR/miR-222-3p/Cyclin Dependent Kinase 19 axis (46), HOTAIR/specificity protein 1/miR-199a axis (47), HOTAIR/miR-29b/phosphate and tension homology deleted on chromosome 10/PI3K axis (48), HOTAIR/miR-203/zinc finger E-box binding homeobox 1 axis (49), HOTAIR/estrogen receptor 1/miR-130b-3p axis (50), HOTAIR/miR-34a/Janus kinase 2/STAT3 axis (51), HOTAIR/miR-129-5p/ribosomal protein L14 axis (52), HOTAIR/miR-129-5p/frizzled class receptor 7 axis (53) and HOTAIR/miR-149-5p/doublecortin-like kinase 1 axis (54). Furthermore, HOTAIR also interacts with several classic signaling pathways, which are commonly involved in oncogenesis, such as CCL22 signaling (55), the Wnt/ β -catenin signaling pathway (56), EZH2 and H3K27 methylation signaling (57). All of this evidence, as well as that provided by the present meta-analysis, suggests the importance of lncRNA HOTAIR SNP in the development of lung cancer.

Of note, there are certain limitations to the present study. First, due to the small sample of the included studies, it was




Figure 4. Subgroup analysis. Subgroup analysis was performed based on (A) Newcastle-Ottawa Scale score quality assessment, (B) type of cancer and (C) P-value of Hardy-Weinberg equilibrium. OR, odds ratio; NSCLC, non-small cell lung cancer.

not possible to analyze certain SNPs because only one study reported the respective relation. Furthermore, lung cancer susceptibility was indicated to be associated with the individual characteristics, such as lifestyle and family history, which was not possible to be extracted. In addition, the majority of the populations were from Asia and additional data are required for other populations. This work was not entered in registries such as the Cochrane Library or PROSPERO, yet the present meta-analysis was performed based on the PRISMA guidelines.

In conclusion, the present study was the first to systematically explore the association between lncRNA HOTAIR SNP and lung cancer susceptibility. It was indicated that populations carrying the lncRNA HOTAIR rs1899663 C>A polymorphism may have a high risk of developing lung cancer. lncRNA HOTAIR may be an important novel target in lung cancer prevention and prognosis prediction and future investigations are required to clarify the intrinsic mechanism of HOTAIR in lung cancer.

Acknowledgements

We thank our collaborator, Professor Jianjun Qiao (Department of Dermatology, the First Affiliated Hospital of Zhejiang University Medicine School, Hangzhou, China), for language and scientific editing support.

Funding

No funding was received.

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

Design of the meta-analysis: CK, XF and XH. Literature screening: CK and XF. Quality assessment: SC and XH. Check and approving the authenticity of the raw data: CK and XF. Statistical analysis: CK and XF and JL. Manuscript writing and revision: CK, XF, JL, SC and XH. All authors have read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Barta JA, Powell CA and Wisnivesky JP: Global epidemiology of lung cancer. *Ann Glob Health* 85: 8, 2019.
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68: 394-424, 2018.
- Cao M and Chen W: Epidemiology of lung cancer in China. *Thorac Cancer* 10: 3-7, 2019.
- Taghvaei S, Sowlat MH, Hassanvand MS, Yunesian M, Naddafi K and Sioutas C: Source-specific lung cancer risk assessment of ambient PM_{2.5}-bound polycyclic aromatic hydrocarbons (PAHs) in central Tehran. *Environ Int* 120: 321-332, 2018.
- Brasky TM, White E and Chen CL: Long-term, supplemental, one-carbon metabolism-related vitamin B use in relation to lung cancer risk in the vitamins and lifestyle (VITAL) cohort. *J Clin Oncol* 35: 3440-3448, 2017.
- Hicks BM, Filion KB, Yin H, Sakr L, Udell JA and Azoulay L: Angiotensin converting enzyme inhibitors and risk of lung cancer: Population based cohort study. *BMJ* 363: k4209, 2018.
- Lin KF, Wu HF, Huang WC, Tang PL, Wu MT and Wu FZ: Propensity score analysis of lung cancer risk in a population with high prevalence of non-smoking related lung cancer. *BMC Pulm Med* 17: 120, 2017.
- Fortunato O, Borzi C, Milione M, Centonze G, Conte D, Boeri M, Verri C, Moro M, Facchinetti F, Andriani F, *et al*: Circulating mir-320a promotes immunosuppressive macrophages M2 phenotype associated with lung cancer risk. *Int J Cancer* 144: 2746-2761, 2019.
- ENCODE Project Consortium: The ENCODE (ENCyclopedia of DNA elements) project. *Science* 306: 636-340, 2004.
- Huarte M: The emerging role of lncRNAs in cancer. *Nat Med* 21: 1253-1261, 2015.
- Wu Y, Shao A, Wang L, Hu K, Yu C, Pan C and Zhang S: The role of lncRNAs in the distant metastasis of breast cancer. *Front Oncol* 9: 407, 2019.
- Zhang Y and Tang L: The application of lncRNAs in cancer treatment and diagnosis. *Recent Pat Anticancer Drug Discov* 13: 292-301, 2018.
- Lu T, Wang Y, Di Chen JL and Jiao W: Potential clinical application of lncRNAs in non-small cell lung cancer. *Onco Targets Ther* 11: 8045-8052, 2018.
- Engle L, Simpson C and Landers J: Using high-throughput SNP technologies to study cancer. *Oncogene* 25: 1594-1601, 2006.
- Syvänen AC: Toward genome-wide SNP genotyping. *Nat Genet* 37 (Suppl): S5-S10, 2005.
- Andrew AS, Gui J, Sanderson AC, Mason RA, Morlock EV, Schned AR, Kelsey KT, Marsit CJ, Moore JH and Karagas MR: Bladder cancer SNP panel predicts susceptibility and survival. *Hum Genet* 125: 527-539, 2009.
- Duboule D: The rise and fall of Hox gene clusters. *Development* 134: 2549-2560, 2007.
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, Goodnough LH, Helms JA, Farnham PJ, Segal E and Chang HY: Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 129: 1311-1323, 2007.
- Hajjari M and Salavaty A: HOTAIR: an oncogenic long non-coding RNA in different cancers. *Cancer Biol Med* 12: 1-9, 2015.
- Loewen G, Jayawickramarajah J, Zhuo Y and Shan B: Functions of lncRNA HOTAIR in lung cancer. *J Hematol Oncol* 7: 90, 2014.
- Fatica A and Bozzoni I: Long non-coding RNAs: New players in cell differentiation and development. *Nat Rev Genet* 15: 7-21, 2014.
- Somarowthu S, Legiewicz M, Chillón I, Marcia M, Liu F and Pyle AM: HOTAIR forms an intricate and modular secondary structure. *Mol Cell* 58: 353-361, 2015.
- Bhan A and Mandal SS: LncRNA HOTAIR: A master regulator of chromatin dynamics and cancer. *Biochim Biophys Acta* 1856: 151-164, 2015.
- Zhuang Y, Wang X, Nguyen HT, Zhuo Y, Cui X, Fewell C, Flemington EK and Shan B: Induction of long intergenic non-coding RNA HOTAIR in lung cancer cells by type I collagen. *J Hematol Oncol* 6: 35, 2013.
- Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, Tsai MC, Hung T, Argani P, Rinn JL, *et al*: Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature* 464: 1071-1076, 2010.
- Wu Y, Zhang L, Wang Y, Li H, Ren X, Wei F, Yu W, Wang X, Zhang L, Yu J and Hao X: Long noncoding RNA HOTAIR involvement in cancer. *Tumour Biol* 35: 9531-9538, 2014.
- Xu ZY, Yu QM, Du YA, Yang LT, Dong RZ, Huang L, Yu PF and Cheng XD: Knockdown of long non-coding RNA HOTAIR suppresses tumor invasion and reverses epithelial-mesenchymal transition in gastric cancer. *Int J Biol Sci* 9: 587-597, 2013.
- Liu XH, Liu ZL, Sun M, Liu J, Wang ZX and De W: The long non-coding RNA HOTAIR indicates a poor prognosis and promotes metastasis in non-small cell lung cancer. *BMC Cancer* 13: 464, 2013.
- Zhao W, An Y, Liang Y and Xie XW: Role of HOTAIR long noncoding RNA in metastatic progression of lung cancer. *Eur Rev Med Pharmacol Sci* 18: 1930-1936, 2014.
- Nakagawa T, Endo H, Yokoyama M, Abe J, Tamai K, Tanaka N, Sato I, Takahashi S, Kondo T and Satoh K: Large noncoding RNA HOTAIR enhances aggressive biological behavior and is associated with short disease-free survival in human non-small cell lung cancer. *Biochem Biophys Res Commun* 436: 319-324, 2013.
- Moher D, Liberati A, Tetzlaff J and Altman DG; PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med* 6: e1000097, 2009.
- Stang A: Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 25: 603-605, 2010.
- Ren MM, Xu S, Wei YB, Yang JJ, Yang YN, Sun SS, Li YJ, Wang PY and Xie SY: Roles of HOTAIR in lung cancer susceptibility and prognosis. *Mol Genet Genomic Med* 8: e1299, 2020.
- Minn AKK, Sato N, Mieno MN, Arai T and Muramatsu M: Association study of long non-coding RNA HOTAIR rs920778 polymorphism with the risk of cancer in an elderly Japanese population. *Gene* 729: 144263, 2020.

35. Wang C, Li Y, Li YW, Zhang HB, Gong H, Yuan Y, Li WT, Liu HY and Chen J: HOTAIR lncRNA SNPs rs920778 and rs1899663 are associated with smoking, male gender, and squamous cell carcinoma in a Chinese lung cancer population. *Acta Pharmacol Sin* 39: 1797-1803, 2018.
36. Li H, Yang Z, Li J, Lv X, Gao M, Bi Y, Zhang Z, Wang S, Li S, Li N, *et al*: Genetic variants in lncRNA HOTAIR are associated with lung cancer susceptibility in a Chinese Han population in China: A case-control study. *Cancer Manag Res* 10: 5209-5218, 2018.
37. Dadaş E and Aydın M: Effect of HOTAIR rs12826786 and rs1899663 polymorphisms on lung cancer susceptibility and clinicopathological characteristics in a Turkish population: A hospital-based case-control study. *Cell Mol Biol (Noisy-le-grand)* 64: 97-102, 2018.
38. Gong WJ, Yin JY, Li XP, Fang C, Xiao D, Zhang W, Zhou HH, Li X and Liu ZQ: Association of well-characterized lung cancer lncRNA polymorphisms with lung cancer susceptibility and platinum-based chemotherapy response. *Tumour Biol* 37: 8349-8358, 2016.
39. Dor Y and Cedar H: Principles of DNA methylation and their implications for biology and medicine. *Lancet* 392: 777-786, 2018.
40. Shimizu Y, Sato S, Noguchi Y, Koyamatsu J, Yamanashi H, Higashi M, Nagayoshi M, Kadota K, Kawashiri SY, Nagata Y, *et al*: Impact of single nucleotide polymorphism on short stature and reduced tongue pressure among community-dwelling elderly Japanese participants: A cross-sectional study. *Environ Health Prev Med* 22: 62, 2017.
41. Leaché AD and Oaks JR: The utility of single nucleotide polymorphism (SNP) data in phylogenetics. *Annu Rev Ecol Syst* 48: 69-84, 2017.
42. Zhang X, Zhou L, Fu G, Sun F, Shi J, Wei J, Lu C, Zhou C, Yuan Q and Yang M: The identification of an ESCC susceptibility SNP rs920778 that regulates the expression of lncRNA HOTAIR via a novel intronic enhancer. *Carcinogenesis* 35: 2062-2067, 2014.
43. Zhu H, Lv Z, An C, Shi M, Pan W, Zhou L, Yang W and Yang M: Onco-lncRNA HOTAIR and its functional genetic variants in papillary thyroid carcinoma. *Sci Rep* 6: 31969, 2016.
44. Pan W, Liu L, Wei J, Ge Y, Zhang J, Chen H, Zhou L, Yuan Q, Zhou C and Yang M: A functional lncRNA HOTAIR genetic variant contributes to gastric cancer susceptibility. *Mol Carcinog* 55: 90-96, 2016.
45. Liu Z, Sun M, Lu K, Liu J, Zhang M, Wu W, De W, Wang Z and Wang R: The long noncoding RNA HOTAIR contributes to cisplatin resistance of human lung adenocarcinoma cells via downregulation of p21(WAF1/CIP1) expression. *PLoS One* 8: e77293, 2013.
46. Fan L, Lei H, Lin Y, Zhou Z, Li J, Wu A, Shu G, Ruger S and Yin G: HotaIR promotes the migration and proliferation in ovarian cancer by miR-222-3p/CDK19 axis. *Cell Mol Life Sci* 79: 254, 2022.
47. Chen J, Hou SF, Tang FJ, Liu DS, Chen ZZ, Zhang HL and Wang SH: HOTAIR/Spl/miR-199a critically regulates cancer stemness and malignant progression of cutaneous squamous cell carcinoma. *Oncogene* 41: 99-111, 2022.
48. Zhang W, Wu Q, Liu Y, Wang X, Ma C and Zhu W: lncRNA HOTAIR promotes chemoresistance by facilitating epithelial to mesenchymal transition through miR-29b/PTEN/PI3K signaling in cervical cancer. *Cells Tissues Organs* 211: 16-29, 2022.
49. Zhang W, Liu J, Wu Q, Liu Y and Ma C: HOTAIR contributes to stemness acquisition of cervical cancer through regulating miR-203 interaction with ZEB1 on epithelial-mesenchymal transition. *J Oncol* 2021: 4190764, 2021.
50. Zhang M, Wu K, Zhang P, Qiu Y, Bai F and Chen H: HOTAIR facilitates endocrine resistance in breast cancer through ESR1/miR-130b-3p axis: Comprehensive analysis of mRNA-miRNA-lncRNA network. *Int J Gen Med* 14: 4653-4663, 2021.
51. Deng S, Wang J, Zhang L, Li J and Jin Y: lncRNA HOTAIR promotes cancer stem-like cells properties by sponging miR-34a to activate the JAK2/STAT3 pathway in pancreatic ductal adenocarcinoma. *Onco Targets Ther* 14: 1883-1893, 2021.
52. Sun N, Zhang W, Liu J, Yang X and Chu Q: Propofol inhibits the progression of cervical cancer by regulating HOTAIR/miR-129-5p/RPL14 axis. *Onco Targets Ther* 14: 551-564, 2021.
53. Wu D, Zhu J, Fu Y, Li C and Wu B: lncRNA HOTAIR promotes breast cancer progression through regulating the miR-129-5p/FZD7 axis. *Cancer Biomark* 30: 203-212, 2021.
54. Zhan Y, Abuduwaili K, Wang X, Shen Y, Nuerlan S and Liu C: Knockdown of long non-coding RNA HOTAIR suppresses cisplatin resistance, cell proliferation, migration and invasion of DDP-resistant NSCLC cells by targeting miR-149-5p/doublecortin-like kinase 1 axis. *Cancer Manag Res* 12: 7725-7737, 2020.
55. Liang H and Peng J: lncRNA HOTAIR promotes proliferation, invasion and migration in NSCLC cells via the CCL22 signaling pathway. *PLoS One* 17: e0263997, 2022.
56. Tang Y, Song G, Liu H, Yang S, Yu X and Shi L: Silencing of long non-coding RNA HOTAIR alleviates epithelial-mesenchymal transition in pancreatic cancer via the Wnt/ β -catenin signaling pathway. *Cancer Manag Res* 13: 3247-3257, 2021.
57. Dai ZY, Jin SM, Luo HQ, Leng HL and Fang JD: lncRNA HOTAIR regulates anoikis-resistance capacity and spheroid formation of ovarian cancer cells by recruiting EZH2 and influencing H3K27 methylation. *Neoplasma* 68: 509-518, 2021.



This work is licensed under a Creative Commons Attribution 4.0 International (CC BY-NC 4.0) License