

Pioglitazone protects PC12 cells against oxidative stress injury: An *in vitro* study of its antiapoptotic effects via the PPAR γ pathway

YALI LI^{1,2}, JUN LONG¹, LIBO LI³, ZIYAO YU⁴, YANJING LIANG¹, BIN HOU¹, LI XIANG⁵ and XIAOLIN NIU⁶

¹Department of Rehabilitation, Weihai Municipal Hospital, Cheeloo College of Medicine, Shandong University, Weihai, Shandong 264200; ²Department of Rehabilitation Medicine, Affiliated Hospital of Weifang Medical University, Weifang, Shandong 261000; ³Department of Rehabilitation, The Second Affiliated Hospital, School of Medicine, Xi'an Jiaotong University, Xi'an, Shaanxi 710004; ⁴College of Sports and Health, Shandong Sport University, Jinan, Shandong 250000; Departments of ⁵Neurology and ⁶Cardiology, The Second Affiliated Hospital, School of Medicine, Xi'an Jiaotong University, Xi'an, Shaanxi 710004, P.R. China

Received July 22, 2022; Accepted January 20, 2023

DOI: 10.3892/etm.2023.12221

Abstract. To the best of our knowledge, the role of peroxisome proliferator-activated receptor γ (PPAR γ) in oxidative stress-induced PC12 cell damage is unknown. Using a PC12 cell model with H₂O₂ treatment, the present study investigated the expression levels of apoptosis-related genes and neuronal apoptosis after oxidative stress injury. The present study further investigated the protective effect and mechanism of pioglitazone, a PPAR γ agonist. PC12 cells treated with H₂O₂ were used as a model of oxidative stress injury. An MTT assay and flow cytometry were used to detect the effect of H₂O₂ on PC12 cell viability and the protective effect of pioglitazone. A TUNEL assay was used to detect neuronal apoptosis. The expression levels of PPAR γ , Bax, Bcl-2 and caspase-3 were examined by reverse transcription-quantitative PCR and western blotting. H₂O₂ reduced PC12 cell viability in a dose- and time-dependent manner. H₂O₂ significantly upregulated the protein expression levels of Bax and the cleaved caspase-3/caspase-3 ratio ($P<0.01$), decreased the protein expression levels of Bcl-2 ($P<0.01$), and increased the apoptosis rate of PC12 cells. Pioglitazone significantly reduced the protein expression levels of Bax and the cleaved caspase-3/caspase-3 ratio ($P<0.01$), increased the expression levels of Bcl-2 ($P<0.01$), decreased the Bax/Bcl-2 expression ratio ($P<0.01$) and increased the viability of H₂O₂-damaged PC12 cells in a dose-dependent manner. Treatment with the PPAR γ antagonist GW9662 or PPAR γ small interfering RNA counteracted the protective effect of pioglitazone on PC12

cells to different extents ($P<0.01$). Therefore, the present study reported the role of PPAR γ in protecting PC12 cells against oxidative stress injury, which may lead to novel therapeutic approaches for neurodegenerative diseases.

Introduction

Neurodegenerative diseases resulting from the progressive loss of structure and/or function of neurons contribute to different paralysis degrees and loss of cognition (1). Progressive neuronal loss is a prominent pathological feature of neurodegenerative diseases, in which oxidative stress serves a vital role in neuronal apoptosis, and is difficult to restore (2). Neurons contain polyunsaturated fatty acids, which are sensitive to free radicals. Polyunsaturated fatty acids are easily attacked by free radicals, and neurons have a low content of antioxidant enzymes. Therefore, the antioxidant capacity of neurons is reduced (3). Together, these factors promote the sensitivity of neurons to oxidative stress injury (3,4). Methods to effectively reduce oxidative stress injury in neurons have attracted increasing attention. PC12 cells, which have the properties of neurosecretory cells and neurons, along with high stability, homogeneity and a high degree of differentiation, are currently widely used in the study of nerve cell function, differentiation, development and death as a cell model (5).

It is well known that apoptosis is a tightly regulated process, which involves changes in the expression of a distinct set of genes (6). Bax and Bcl-2 are major genes responsible for regulating apoptosis. Bax is a member of the Bcl-2 family, and promotes apoptosis, while Bcl-2 blocks cell death (7). The Bax/Bcl-2 ratio is a widely used parameter to determine cell susceptibility to apoptosis. Caspase-3 is the most important executing protease in the process of apoptosis (8). Cleaved caspase-3 is the activated form of caspase-3 (9). Previous studies have confirmed that H₂O₂ can induce PC12 cell injury (10) and the expression and activation of the apoptosis-related gene caspase-3 (11).

Peroxisome proliferator-activated receptor γ (PPAR γ) is a ligand-activated nuclear receptor that regulates glucose and

Correspondence to: Dr Li Xiang, Department of Neurology, The Second Affiliated Hospital, School of Medicine, Xi'an Jiaotong University, 157 Xiwu Road, Xi'an, Shaanxi 710004, P.R. China
E-mail: lixiange@ sina.cn

Key words: peroxisome proliferator-activated receptor γ , PC12 cells, oxidative stress

lipid metabolism, endothelial function and inflammation (12). Thiazolidinediones (TZDs) are ligands that are known to bind to and activate nuclear PPAR γ and are currently used as insulin sensitizers in type 2 diabetes (13). Our previous study has demonstrated that PPAR γ agonists TZDs could protect the neuronal microenvironment and preserve nerve cells in the hippocampi of spontaneously hypertensive rats (SHRs) via antioxidative and antiapoptotic pathways (14). However, the underlying mechanism needs to be further studied. In the present study, a model of oxidative stress damage was generated in PC12 cells using H₂O₂ to observe whether pioglitazone had a neuroprotective effect and to determine the underlying mechanism.

Materials and methods

Cell culture and preconditioning protocols. PC12 cells were obtained from the Institute of Neurobiology, School of Medicine, Xi'an Jiaotong University (Xi'an, China). The cells were plated at a density of 3×10^5 cells/well in 6-well plates and maintained in DMEM/F12 supplemented with 10% heat-inactivated foetal bovine serum (both Gibco; Thermo Fisher Scientific, Inc.), 100 U/ml penicillin and 100 μ g/ml streptomycin (Thermo Fisher Scientific, Inc.) at 37°C under an atmosphere of 5% CO₂ and 95% air. The culture medium was changed three times per week. PC12 cells were preconditioned by 0, 6, 12, 24 or 36 h of exposure to different concentrations (0, 25, 50, 100, 200 and 400 μ mol/l) of H₂O₂ at 37°C. Experiments were performed at least three times.

Transfection experiments. A total of three alternative siRNA sequences targeting PPAR γ were used to knock down PPAR γ protein expression. The following groups were used: NC group, siRNA-NC (non-targeting) group, PPAR γ -siRNA 1 group, PPAR γ -siRNA 2 group and PPAR γ -siRNA 3 group. PPAR γ -siRNAs and negative control siRNA were designed and synthesized by Shanghai GenePharma Co., Ltd. First, PC12 cells were seeded in 24-well plates at an optimized concentration of 1×10^5 cells/well, 24 h before transfection. On the following day, when cell confluence had reached 60–70%, they were transfected with PPAR γ -siRNA 1 (10 μ M; 10 μ l/well), PPAR γ -siRNA 2 (10 μ M; 10 μ l/well), PPAR γ -siRNA 3 (10 μ M; 10 μ l/well) or siRNA-NC (10 μ M; 10 μ l/well) using Lipofectamine® 2000 (cat. no. 11668019; Invitrogen; Thermo Fisher Scientific, Inc.) at 37°C for 2 days according to the manufacturer's protocol. The sequence with the best inhibition rate, as determined by PPAR γ protein expression analysis 48 h after transfection, was selected. The sequences are shown in Table I. Following 48 h of transfection, the cells were collected for subsequent experiments.

MTT assay

Experiment 1: Effect of H₂O₂ on the viability of PC12 cells. The viability of cultured cells was measured using an MTT assay (15). First, PC12 cells were preconditioned by 0, 6, 12, 24 or 36 h of exposure to different concentrations (0, 25, 50, 100, 200 and 400 μ mol/l) of H₂O₂ at 37°C. Subsequently, cells were incubated with MTT solution (0.5 mg/ml in PBS) at 37°C for 4 h. The MTT solution was removed from the plate and the plate was dried. DMSO (100 μ l) was added to

each well to dissolve the formazan crystals before the optical density was measured at 570 nm. The results are presented as the percentage of MTT reduction, and the absorbance of the control cells was set as 100%. Experiments were performed at least three times.

Experiment 2: Protective effects of pioglitazone on PC12 cells with H₂O₂-induced injury. In this experiment, PC12 cells were divided into five groups: The control, H₂O₂ (100 μ mol/l H₂O₂), low-concentration pioglitazone (1×10^{-7} mol/l; cat. no. 111025-46-8; Alexis Biochemicals; Enzo Life Sciences), medium-concentration pioglitazone (1×10^{-6} mol/l) and high-concentration pioglitazone (1×10^{-5} mol/l) groups. Pioglitazone at different concentrations was utilized to precondition PC12 cells for a 1-h period at 37°C, whereas the H₂O₂ group cells were instead treated with 0.9% saline; all the PC12 cells of H₂O₂ and pioglitazone groups were then treated with 100 μ M H₂O₂ for a 24 h period at 37°C. For the control group, PC12 cells were treated with cell culture medium for 25 h at 37°C. The viability of the cultured cells was measured by an MTT assay as aforementioned. Experiments were performed at least three times.

Experiment 3: Role of PPAR γ in the protective effect of pioglitazone on PC12 cells with H₂O₂-induced injury. In this next experiment, PC12 cells were divided into five groups: The control, H₂O₂, pioglitazone + H₂O₂, pioglitazone + GW9662 + H₂O₂ and pioglitazone + PPAR γ -siRNA 3 + H₂O₂ groups. In the H₂O₂ group, PC12 cells were treated with 100 μ M H₂O₂ for 24 h at 37°C. PC12 cells in the control group were treated with cell culture medium at 37°C for 24 h. For the pioglitazone + H₂O₂ group, 1×10^{-5} mol/l pioglitazone was utilized to precondition PC12 cells for a 1-h period at 37°C, and then cells were treated with 100 μ M H₂O₂ for 24 h at 37°C. The pioglitazone + GW9662 + H₂O₂ group was treated with GW9662 (1×10^{-6} mol/l; cat. no. M6191; MilliporeSigma) for 1 h at 37°C, followed by 1×10^{-5} mol/l pioglitazone for 1 h at 37°C and H₂O₂ (100 μ mol/l) for 24 h at 37°C. For the pioglitazone + PPAR γ -siRNA 3 + H₂O₂ group, after PPAR γ -siRNA transfection, 1×10^{-5} mol/l pioglitazone was added to the cells for 1 h at 37°C prior to treatment with H₂O₂ (100 μ mol/l) for 24 h at 37°C. The viability of cultured cells was measured by an MTT assay as aforementioned. Experiments were performed at least three times.

Measurement of malondialdehyde (MDA) content, and superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) activities. Based on the MTT assay results in Experiment 1, the desired H₂O₂ concentration, action time and cell plating density for modelling oxidative stress injury in PC12 cells were 100 μ mol/l, 24 h and 1×10^5 cells/ml, respectively. To verify the successful modelling of PC12 cells, the MDA content and SOD and GSH-Px activity were measured in the control and H₂O₂ groups. In the H₂O₂ group, PC12 cells were treated with 100 μ M H₂O₂ for 24 h at 37°C. PC12 cells in the control group were treated with cell culture medium at 37°C for 24 h. Cells were collected and rinsed with PBS, and lysis buffer containing 20 mM Tris (pH 7.5), 150 mM NaCl, 1 mM PMSF and 1% Triton X-100 was then added for homogenization. The cell supernatants were collected. The content of MDA (cat. no. A003-1-2) and the activities of SOD (cat. no. A001-1-2) and GSH-Px (cat. no. A005-1-2) were all determined according to

Table I. PPAR γ -siRNA sequences.

siRNA	Sequence (5'-3')
PPAR γ -siRNA 1 (sense)	AGAUAAAGCUUCUGGAUUU
PPAR γ -siRNA 1 (antisense)	AAAUCCAGAAGCUUUAUCU
PPAR γ -siRNA 2 (sense)	AGGAAAGACAACAGACAAA
PPAR γ -siRNA 2 (antisense)	UUUGUCUGUUGUCUUUCCU
PPAR γ -siRNA 3 (sense)	CCUCCCUGAUGAAUAAAGATT
PPAR γ -siRNA 3 (antisense)	UCUUUAUUCaucagggaggtt
Negative control siRNA (sense)	UUCUCCGAACGUGUCACGUTT
Negative control siRNA (antisense)	ACGUGACACGUUCGGAGAATT

PPAR γ , peroxisome proliferator-activated receptor γ ; siRNA, small interfering RNA.

the instructions provided by Nanjing Jiancheng Bioengineering Institute as previously described (16). Briefly, MDA content was determined using the thiobarbituric acid method. The wavelength of colorimetry was 532 nm. SOD activity was determined using the hydroxylamine method and the wavelength of colorimetry was 550 nm. GSH-Px activity was determined using a colorimetric method and the wavelength of colorimetry was 412 nm. The optical density value was read using a microplate reader, and two parallel samples were used to ensure the accuracy of the experiment. Experiments were performed at least three times.

Detection of apoptotic cells by flow cytometry. Cells in the control, H₂O₂, pioglitazone + H₂O₂, pioglitazone + PPAR γ -siRNA 3 + H₂O₂ and pioglitazone + GW9662 + H₂O₂ groups were used to detect apoptosis by annexin V-FITC and PI staining followed by flow cytometry (Beckman Coulter, Inc.). The treatment of each group was performed as described for the MTT assay experiment 3. The procedure described in the documentation of the annexin V-FITC/PI detection test kit (cat. no. C1062; Beyotime Institute of Biotechnology) was followed. Cells were resuspended at a concentration of 1×10^6 cells/ml in 400 μ l 1X binding buffer solution. Then, 5 μ l annexin V-FITC and 10 μ l PI were added to stain the cells for 15 min at room temperature in the dark. Then, the cell apoptosis was detected by flow cytometric analysis (NL-CLC 1L-3L; Cytex NL-CLC Full Spectrum Flow Cytometer; Shanghai Xiatai Biotechnology Co., Ltd.). The excitation wavelength was 488 nm and the emission wavelength was 530 nm. Green fluorescence of FITC and red fluorescence of PI were observed after excitation. Data were analysed using FlowJo (v10; FlowJo LLC). Experiments were performed at least three times.

Reverse transcription-quantitative PCR (RT-qPCR) of PPAR γ in PC12 cells. RT-qPCR was performed for the NC, siRNA-NC, PPAR γ -siRNA 1, PPAR γ -siRNA 2 and PPAR γ -siRNA 3 groups after the transfection experiments. RT-qPCR was then performed to analyse cells in the control, H₂O₂, pioglitazone + H₂O₂, pioglitazone + PPAR γ -siRNA 3 + H₂O₂ and pioglitazone + GW9662 + H₂O₂ groups. The treatment of each group was performed as described for the MTT assay experiment 3. Total RNA was extracted

using TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.). RNA samples were transcribed into cDNA using the PrimeScript RT Master Mix Kit (Takara Bio, Inc.) according to the manufacturer's instructions. The SYBR ExScript RT-PCR Kit (Takara Bio, Inc.) was used for RT-qPCR on the IQ Multicolor Real-Time PCR Detection System (Bio-Rad Laboratories, Inc.). The primers used to amplify rat PPAR γ (forward, 5'-GGAGCCTAAGTTTGAGTTTGCTGTG-3' and reverse, 5'-TGCAGCAGGTTGTCTTGGATG-3') and β -actin (forward, 5'-GGAGATTACTGCCCTGGCTCCTA-3' and reverse, 5'-GACTCATCGTACTCCTGCTTGCTG-3') were designed and synthesized by Takara Biotechnology Co., Ltd. Amplification was carried out with the following thermal cycling program: 95°C for 30 sec, followed by 40 cycles of 95°C for 3 sec and 60°C for 30 sec. Cycle threshold values were obtained with Bio-Rad iQ5 2.0 Standard Edition Optical System software (Bio-Rad Laboratories, Inc.). β -actin was used as the internal control. Relative quantification was performed by the comparative cycle threshold ($2^{-\Delta\Delta C_q}$) method (17), and the data are presented as the mean \pm SD of three separate experiments performed with triplicate samples.

Western blot analysis. PPAR γ , Bax, Bcl-2, cleaved caspase-3 and caspase-3 protein expression was examined in the control, H₂O₂, pioglitazone + H₂O₂, pioglitazone + PPAR γ -siRNA 3 + H₂O₂ and pioglitazone + GW9662 + H₂O₂ groups. The treatment of each group was performed as described for the MTT assay experiment 3. PC12 cells were harvested by scraping into ice-cold PBS and centrifuged at 12,000 \times g for 8 min at 4°C. Afterwards, RIPA cell lysis buffer (P0013B; Beyotime Institute of Biotechnology) was used for extraction of cellular protein using 1 mM PMSF. A BCA kit (P0010; Beyotime Institute of Biotechnology) was used to determine protein concentrations. Equal amounts of protein (20 μ g/lane) from different samples were separated on 12% SDS polyacrylamide gels, transferred to PVDF membranes and blocked in 10% non-fat milk at room temperature for 2 h. The membranes were incubated at 4°C overnight with rabbit polyclonal antibodies against PPAR γ (1:400; cat. no. ab66343; Abcam), Bax (1:800; cat. no. bs2538; Bioworld Technology, Inc.), Bcl-2 (1:800; cat. no. bs1511; Bioworld Technology, Inc.), cleaved caspase-3 (1:1,000; cat. no. ab2302; Abcam), caspase-3

(1:800; cat. no. bs7004; Bioworld Technology, Inc.) and mouse monoclonal anti- β -actin (1:1,000; cat. no. sc-517582; Santa Cruz Biotechnology, Inc.). The membranes were washed with Tris-buffered saline containing Tween 20 (0.1%) three times and incubated with HRP-conjugated goat anti-rabbit IgG (1:5,000; cat. no. BS13278; Bioworld Technology, Inc.) and HRP-conjugated goat anti-mouse IgG (1:5,000; cat. no. BS12478; Bioworld Technology, Inc.) at room temperature for 2 h. After washing, protein bands were detected by incubation with chemiluminescent HRP substrate (SuperSignal West Pico; Thermo Fisher Scientific, Inc.) for 5 min at room temperature in the dark and exposure to X-ray film (FUJIFILM Wako Pure Chemical Corporation). Quantity One software 4.6.2 (Bio-Rad Laboratories, Inc.) was used to semi-quantify the band intensity, which was normalized to that of the loading control β -actin. The data are presented as the mean \pm SD of three separate experiments performed with triplicate samples.

TUNEL assay. A DeadEnd™ Colorimetric TUNEL System (cat. no. G7130; Promega Corporation) was used to detect PC12 apoptosis in the control, H_2O_2 , pioglitazone + H_2O_2 , pioglitazone + PPAR γ -siRNA 3 + H_2O_2 , pioglitazone + GW9662 + H_2O_2 groups. The treatment of each group was performed as described for the MTT assay experiment 3. The coverslips were placed at the bottom of a 24-well plate. PC12 cells were inoculated into the 24-well plate and treated according to the grouping. Subsequently, the coverslips were fixed with 4% paraformaldehyde at room temperature (15–25°C) for 30 min, incubated with protease K at room temperature for 20 min and washed three times with PBS. The coverslips were covered with equilibration buffer for 10 min and with rTdT incubation buffer (50 μ l; containing TUNEL reagent) at 37°C in the dark for 1 h. The coverslips were immersed in 2X saline-sodium citrate buffer for 15 min and rinsed with PBS. The coverslips were counterstained with DAPI (1:1,000; MilliporeSigma) for 5 min at room temperature, washed in PBS and mounted with 50% glycerin (IH0272; Beijing Legen Biotechnology Co., Ltd.). Fluorescence microscopy was carried out using an Olympus BX51 microscope (Olympus Corporation) with a mercury lamp power supply. Neurons with bright green nuclei were identified as TUNEL-positive neurons. The ratio of TUNEL-positive cells to DAPI-positive cells was used as the experimental indicator to calculate the apoptosis rate of PC12 cells. The number of TUNEL-positive cells was normalized to the number of DAPI-stained cells. A 20X objective lens was used to count apoptotic cells in nine random fields (3 samples in each group; three fields in each sample), and the observer was blinded to the treatment groups.

Statistical analysis. Statistical analysis was carried out using SPSS 16.0 software (SPSS, Inc.). Quantitative data are presented as the mean \pm SD and are representative of three independent experiments. Unpaired Student's t-test and one-way ANOVA followed by Tukey's post hoc test were used to assess the significance of differences among groups. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

PC12 cell apoptosis is induced by H_2O_2 . To find the appropriate concentration of H_2O_2 for the PC12 cell damage model, different concentrations of H_2O_2 (25, 50, 100, 200 and 400 μ mol/l) were used. PC12 cell viability was evaluated using an MTT assay after culture with H_2O_2 for 0, 6, 12, 24 and 36 h. The MTT assay results revealed that with the increase of the H_2O_2 concentration and incubation time, the viability of PC12 cells decreased gradually (Fig. 1A). To simulate the physiological conditions, the present study aimed to select a treatment method with slow action and resulting in a specific degree of damage. Conditions leading to a cell viability rate between 50 and 60% in the MTT assay were suitable for the apoptosis experiments. In the present experiment, treatment of PC12 cells with 100 μ mol/l H_2O_2 for 24 h inhibited PC12 cell viability to 50–60% of that in the control group. Thus, the desired H_2O_2 concentration and action time were 100 μ mol/l and 24 h, respectively.

Successful establishment of the PC12 cell model. The MDA content and the activities of SOD and GSH-Px were measured in the control and H_2O_2 groups to verify the successful establishment of the PC12 cell model (Fig. 2). After treatment with 100 μ mol/l H_2O_2 for 24 h, the content of the active peroxidation product MDA in the H_2O_2 group was increased but the activities of SOD and GSH-Px were decreased compared with those in the control group, indicating that oxidative stress injury in PC12 cells was successfully modelled.

Effects of preconditioning with different concentrations of pioglitazone on PC12 cell viability reduced by H_2O_2 . PC12 cells in the H_2O_2 group were treated with 100 μ mol/l H_2O_2 for 24 h, while cells in the control group were treated with cell culture medium. The MTT assay results showed that H_2O_2 significantly reduced the viability of PC12 cells ($P < 0.01$ vs. control group). Pioglitazone served a protective role in H_2O_2 -treated PC12 cells in a concentration-dependent manner. The concentration at which pioglitazone exhibited its maximum effects was 1×10^{-5} mol/l, and 1×10^{-4} mol/l pioglitazone did not increase PC12 viability compared with 1×10^{-5} mol/l pioglitazone (data not shown). Therefore, 1×10^{-5} mol/l was selected as the concentration for the next experiment.

Pioglitazone increases PPAR γ mRNA and protein expression. The RT-qPCR and western blotting results are shown in Fig. 3. PC12 cells were transfected with siRNA to knock down PPAR γ protein expression. The following three groups were used: NC group, siRNA-NC group and PPAR γ -siRNA group, divided into PPAR γ -siRNA 1, PPAR γ -siRNA 2 and PPAR γ -siRNA 3. The gene knockdown rate in all groups was detected by RT-qPCR (Fig. 3A) and western blotting (Fig. 3B and C). The results demonstrated that PPAR γ expression was decreased in PC12 cells transfected with PPAR γ -siRNA compared with siRNA-NC, with PPAR γ -siRNA 3 showing superior knockdown efficiency. Therefore, PPAR γ -siRNA 3 was selected for subsequent experiments.

Compared with the control group, H_2O_2 induced increases in PPAR γ mRNA and protein expression ($P < 0.01$). Pioglitazone pretreatment significantly increased PPAR γ




Figure 1. MTT assay results showing changes in the viability of PC12 cells. (A) PC12 cell viability decreased with increasing H₂O₂ concentration and in a time-dependent manner. Conditions leading to a cell viability rate between 50 and 60% in the MTT assay were suitable for the apoptosis experiments. In the present experiment, treatment of PC12 cells with 100 μmol/l H₂O₂ for 24 h inhibited PC12 cell viability to 50-60% of that in the control group. Thus, the appropriate H₂O₂ concentration and action time were 100 μmol/l and 24 h, respectively. (B) Changes in the viability of PC12 cells treated with H₂O₂ after preconditioning with pioglitazone at different concentrations. Control group, first group with no treatment; H₂O₂ group, second group with treatment with 100 μmol/l H₂O₂ for 24 h. *P<0.01, *P<0.05 vs. control group; #P<0.01, ΔP<0.05 vs. H₂O₂ group; ▼P<0.05 vs. pioglitazone group (1×10⁻⁷ mol/l).




Figure 2. MDA content and SOD and GSH-Px activities in the control and H₂O₂ groups. *P<0.01 vs. control group. GSH-Px, glutathione peroxidase; MDA, malondialdehyde; SOD, superoxide dismutase; pro, protein.

mRNA and protein expression (P<0.01 vs. control and H₂O₂ groups). Pioglitazone treatment increased PPARγ mRNA and protein expression 4.36- and 4.4-fold, respectively, compared with the control group. Compared with pioglitazone treatment, treatment with PPARγ-siRNA 3 and the PPARγ antagonist GW9662 significantly reduced PPARγ mRNA and protein expression (P<0.01).

Neuroprotective effect of pioglitazone on H₂O₂-treated PC12 cells. To investigate the protective effect of PPARγ on H₂O₂-treated PC12 cells, an MTT assay, a TUNEL assay

and flow cytometry were used to observe the neuroprotective effect of pioglitazone.

MTT assay results. As shown in Fig. 4, treatment with 100 μmol/l H₂O₂ significantly reduced PC12 cell viability to 56.8% of the value in the control group (P<0.01). Treatment with 1×10⁻⁵ mol/l pioglitazone increased PC12 cell viability to 80.2% of that in the control group, and there was a significant difference between the pioglitazone + H₂O₂ and H₂O₂ groups (P<0.01), suggesting pioglitazone may exert a protective effect on oxidative stress-injured PC12 cells. However, the viability of cells in the pioglitazone + H₂O₂ group also differed from that of cells in the control group (P<0.01), suggesting that the protective effect of pioglitazone is insufficient to completely protect PC12 cells from oxidative damage. Pretreatment with either PPARγ-siRNA 3 or the PPARγ antagonist GW9662 reversed the neuroprotective effects of pioglitazone to similar degrees. The viability of PC12 cells treated with PPARγ-siRNA and GW9662 decreased to 59.6 and 59.1% of that in the pioglitazone + H₂O₂ group, respectively, with both showing a significant difference compared with the pioglitazone + H₂O₂ group (P<0.01), suggesting that pioglitazone exerts its neuroprotective effect through PPARγ activation.

Flow cytometry results. As shown in Fig. 5, PC12 cell apoptosis in the control group was 6.76% (Fig. 5A). The percentages in Fig. 5F are the averages for all three experimental repeats. After H₂O₂ treatment, the mean apoptosis increased significantly to 25.48% (P<0.01 vs. control group; Fig. 5B). Pioglitazone decreased the apoptosis of PC12 cells to 12.93% (Fig. 5C), which was significantly different from that in the H₂O₂ group and the control group (P<0.01 and P<0.05, respectively), suggesting that the neuroprotective effect of pioglitazone was insufficient to completely reverse the oxidative damage caused by H₂O₂. Pretreatment with PPARγ-siRNA 3 (Fig. 5D) and the PPARγ antagonist GW9662 (Fig. 5E) increased the apoptosis rates of PC12 cells to 23.14 and 21.51%, respectively, which both showing a significant difference compared with the pioglitazone + H₂O₂ group (P<0.01). This




Figure 3. Effects of H₂O₂, pioglitazone, GW9662 and PPAR γ -siRNA on PPAR γ mRNA and protein expression. (A) PPAR γ expression in PC12 cells transfected with PPAR γ -siRNA detected by reverse transcription-quantitative PCR. (B) PPAR γ expression in PC12 cells transfected with PPAR γ -siRNA detected by western blotting and (C) semi-quantitative analysis of PPAR γ protein expression. *P<0.01 vs. siRNA-NC group; [#]P<0.05 vs. PPAR γ -siRNA 1 group; [#]P<0.01 vs. PPAR γ -siRNA 2 group. (D) Effects of H₂O₂, pioglitazone, GW9662 and PPAR γ -siRNA on PPAR γ mRNA expression. (E) Effects of H₂O₂, pioglitazone, GW9662 and PPAR γ -siRNA on PPAR γ protein expression. (F) Representative western blot image showing PPAR γ protein expression. *P<0.01 vs. control group; [#]P<0.01 vs. H₂O₂ group; [#]P<0.01 vs. pioglitazone + H₂O₂ group. NC, negative control; PPAR γ , peroxisome proliferator-activated receptor γ ; siRNA, small interfering RNA.




Figure 4. Effects of pioglitazone, PPAR γ -siRNA 3 and GW9662 on the viability of H₂O₂-treated PC12 cells, as determined by an MTT assay. *P<0.01 vs. control group; [#]P<0.01 vs. H₂O₂ group; [#]P<0.01 vs. Pioglitazone + H₂O₂ group. PPAR γ , peroxisome proliferator-activated receptor γ ; siRNA, small interfering RNA.

result suggested that all types of PPAR γ inhibition reversed the neuroprotective effect of pioglitazone to different degrees and that the neuroprotective effect of pioglitazone was mediated through the PPAR γ activation pathway.

TUNEL assay results. As Fig. 6 shows, the control group showed an intact cell morphology, a low fluorescence intensity, larger nuclei, more DAPI-positive cells and fewer TUNEL-positive cells. The TUNEL/DAPI ratio was 5.20% (Fig. 6A). After treatment with H₂O₂ for 24 h, the number of DAPI-positive nuclei decreased, and high-intensity concentrated fluorescence appeared in the nuclei. The nuclei decreased in size and became condensed and fragmented, showing the characteristics of apoptotic cells (Fig. 6B). The mean TUNEL/DAPI ratio increased to 26.77% (P<0.01 vs. control group; Fig. 6F), suggesting that H₂O₂ promoted apoptosis in PC12 cells. Pioglitazone decreased the mean TUNEL/DAPI ratio to 11.90% (Fig. 6C), showing differences compared with the H₂O₂ group and the control group (P<0.01 and P<0.05, respectively), suggesting that pioglitazone could protect PC12 cells from H₂O₂-induced oxidative damage. However, the protective effect was insufficient to completely reverse the oxidative damage. Pretreatment with the PPAR γ antagonist GW9662 and PPAR γ -siRNA increased the mean TUNEL/DAPI ratio to 22.34% (Fig. 6D) and 23.27% (Fig. 6E), respectively, with significant differences compared with the pioglitazone + H₂O₂ group (P<0.01). This result suggested that pioglitazone exerts a neuroprotective effect through PPAR γ activation.

Pioglitazone decreases the H₂O₂-induced increases in the Bax/Bcl-2 ratio and cleaved caspase-3/caspase-3 ratio. As shown in Fig. 7, after H₂O₂ treatment, the Bax and cleaved




Figure 5. Effect of pioglitazone on apoptosis in H₂O₂-treated PC12 cells, as determined by flow cytometry. (A) Control group. (B) H₂O₂ group. (C) Pioglitazone + H₂O₂ group. (D) Pioglitazone + PPAR γ -siRNA 3 + H₂O₂ group. (E) Pioglitazone + GW9662 + H₂O₂ group. (F) Analysis of the flow cytometry results. *P<0.01, *P<0.05 vs. control group; #P<0.01 vs. H₂O₂ group; &P<0.01 vs. Pioglitazone + H₂O₂ group. PPAR γ , peroxisome proliferator-activated receptor γ ; siRNA, small interfering RNA; UL, upper left; UR, upper right; LL, lower left; LR, lower right.




Figure 6. Effect of pioglitazone on apoptosis in H₂O₂-treated PC12 cells, as determined by a TUNEL assay (magnification, x400; scale bar, 50 μ m). (A) Control group. (B) H₂O₂ group. (C) Pioglitazone + H₂O₂ group. (D) Pioglitazone + GW9662 + H₂O₂ group. (E) Pioglitazone + PPAR γ -siRNA 3 + H₂O₂ group. (F) Analysis of the TUNEL assay results. *P<0.01, *P<0.05 vs. control group; #P<0.01 vs. H₂O₂ group; &P<0.01 vs. Pioglitazone + H₂O₂ group. PPAR γ , peroxisome proliferator-activated receptor γ ; siRNA, small interfering RNA.

caspase-3/caspase-3 ratio increased by 3.95- and 3.11-fold, respectively (both $P < 0.01$ vs. control group; Fig. 7A and D), while Bcl-2 protein expression was significantly reduced ($P < 0.01$ vs. control group; Fig. 7B). The Bax/Bcl-2 ratio in the H_2O_2 group was significantly higher than that in the control group ($P < 0.01$; Fig. 7C). Pioglitazone reduced the mean protein levels of Bax and cleaved caspase-3/caspase-3 ratio to 46.91 and 59.38% of the value in the H_2O_2 group, respectively ($P < 0.01$), and increased the mean expression levels of Bcl-2 by 1.53-fold ($P < 0.01$) compared with those in the H_2O_2 group, thus significantly reducing the Bax/Bcl-2 ratio ($P < 0.01$). Compared with the pioglitazone + H_2O_2 group, the pioglitazone + GW9662 + H_2O_2 and pioglitazone + PPAR γ -siRNA 3 + H_2O_2 groups showed increased Bax protein levels and cleaved caspase-3/caspase-3 ratio levels, reduced Bcl-2 expression and an increased Bax/Bcl-2 ratio ($P < 0.01$). However, no significant differences between the pioglitazone + GW9662 + H_2O_2 and pioglitazone + PPAR γ -siRNA 3 + H_2O_2 groups were observed. This result suggested that the PPAR γ agonist pioglitazone can increase the expression levels of the antiapoptotic protein Bcl-2 and decrease the expression levels of the proapoptotic proteins Bax and the cleaved caspase-3/caspase-3 ratio, thus protecting PC12 cells from H_2O_2 -induced oxidative damage. Pioglitazone mainly exerts its antiapoptotic effect through the PPAR γ pathway (18).

Discussion

Our previous study demonstrated that SHRs exhibited an age-dependent increase in TUNEL-positive cells in the CA1 subfield of the hippocampus, which was accompanied by increased expression of oxidative stress markers and reduced mRNA and protein expression levels of PPAR γ (19). PPAR γ agonist rosiglitazone can exert neuroprotective effects through antioxidative and antiapoptotic pathways independent of blood pressure control (14). However, the limitation of our previous study was that PPAR γ was not inhibited in animal experiments (14). In the present study, to verify the role of PPAR γ activation in protection against oxidative stress injury, PC12 cells were treated with H_2O_2 as a cellular model of oxidative stress injury. The PPAR γ antagonist GW9662 and PPAR γ -siRNA were used to block PPAR γ expression in PC12 cells. Pioglitazone exerted an antiapoptotic effect and promoted the survival of PC12 cells with oxidative stress injury. This effect was mediated through PPAR γ activation.

PC12 is a tumour cell line isolated from a rat adrenal pheochromocytoma. Differentiated PC12 cells have typical neuronal characteristics in terms of morphology and function and are widely used as a cell model to study the apoptosis and differentiation of nerve cells (20). H_2O_2 is considered to be the major precursor of reactive oxygen species (ROS) and is widely used to induce oxidative stress injury (5). Exogenous H_2O_2 easily enter cells through the cell membrane, resulting in the production of large amounts of ROS (21). Thus, H_2O_2 is commonly used to simulate cellular peroxidative damage *in vitro* (22). There have been a number of studies on PC12 cell apoptosis induced by H_2O_2 , and the conclusions were consistent (23,24). However, the effective concentrations and action times were different. In general, low and moderate concentrations (50–500 $\mu\text{mol/l}$) of H_2O_2 could induce oxidative stress,

while high concentrations could rapidly cause cell necrosis. To determine the appropriate concentration and working time of H_2O_2 , viability of PC12 cells was analysed using an MTT assay. The results showed that cell viability was decreased in PC12 cells following treatment with 100 $\mu\text{mol/l}$ H_2O_2 for 24 h.

ROS act on the unsaturated lipids in cell membranes, causing peroxidation of membrane lipids and leading to cell damage and the formation of lipid peroxidation products (25). MDA is an important ROS metabolite in cells and is a good indicator of the degree of tissue peroxidation (26). SOD and GSH-Px are two important antioxidant enzymes in living organisms (27). In the present study, H_2O_2 induced an increase in the content of the active peroxidation product MDA and decreases in the SOD and GSH-Px activities. This result indicated the successful establishment of the oxidative stress injury model in PC12 cells.

The pathways by which H_2O_2 induces apoptosis differ among cell types. Dumont *et al* (28) reported that H_2O_2 -induced apoptosis in T cells mainly depended on mitochondrial ROS and NF- κ B activation. H_2O_2 promoted apoptosis by activating caspase-3 in HL-60 cells (29). Kitamura *et al* (30) found that Bcl-2 and Bax expression did not increase significantly after treatment with H_2O_2 but mediated apoptosis by increasing P53 expression. H_2O_2 induced apoptosis in hepatoblastoma cells not only by upregulating the expression of P53 but also by decreasing the protein levels of Bcl-2 and Bax (31). In the present study, H_2O_2 upregulated the expression levels of Bax and caspase-3 and downregulated the expression levels of Bcl-2. It was suggested that the mechanism by which H_2O_2 induced apoptosis in PC12 cells may involve increasing the levels of proapoptotic proteins and decreasing those of antiapoptotic proteins, thus changing the apoptotic environment in PC12 cells. This is consistent with recent research showing that H_2O_2 could increase the Bax and cleaved caspase-3 protein levels in PC12 cells (32).

The process of apoptosis in neurons is similar to that in other cells. After recognition of death signals, apoptosis-promoting proteins such as Bax and BH3 interacting domain death agonist are translocated to the outer mitochondrial membrane and interact with antiapoptotic proteins such as Bcl-2, which abolishes the apoptosis-inhibiting effect of antiapoptotic proteins, increases the permeability of the mitochondrial membrane, releases cytochrome c into the cytoplasm and activates caspase-3, which eventually leads to apoptosis (33). The Bax/Bcl-2 ratio serves a decisive role in determining whether apoptosis is initiated in cells; thus, the Bax/Bcl-2 expression ratio is often used to evaluate the degree of apoptosis (34). The present results showed that H_2O_2 increased the protein levels of the proapoptotic factors Bax and cleaved caspase-3 and reduced the expression levels of the antiapoptotic protein Bcl-2, thus increasing the Bax/Bcl-2 ratio, which activated caspase-3 and induced apoptosis. The results of the TUNEL assay also confirmed these findings.

PPAR γ is a ligand-activated nuclear transcription factor. PPAR γ , when activated by its ligands, can bind to specific DNA response elements and regulate gene transcription and expression (35). Our previous study demonstrated that the PPAR γ agonist rosiglitazone could upregulate PPAR γ mRNA and protein expression in aged SHRs, while it reduced the expression of oxidative stress markers (inducible nitric oxide




Figure 7. Effect of pioglitazone on the protein expression levels of Bax, Bcl-2 and cleaved caspase-3/caspase-3 ratio in PC12 cells. (A) Effects of H_2O_2 , pioglitazone, GW9662 and PPAR γ -siRNA 3 on Bax protein expression. (B) Effects of H_2O_2 , pioglitazone, GW9662 and PPAR γ -siRNA 3 on Bcl-2 protein expression. (C) Effects of H_2O_2 , pioglitazone, GW9662 and PPAR γ -siRNA 3 on the Bax/Bcl-2 ratio in PC12 cells. (D) Effects of H_2O_2 , pioglitazone, GW9662 and PPAR γ -siRNA 3 on the ratio of cleaved caspase-3/caspase-3. (E) Bax, Bcl-2, cleaved caspase-3 and caspase-3 expression in different groups detected by western blotting * $P < 0.01$ vs. control group; # $P < 0.01$ vs. H_2O_2 group; & $P < 0.01$ vs. pioglitazone + H_2O_2 group. PPAR γ , peroxisome proliferator-activated receptor γ ; siRNA, small interfering RNA.

synthase and NADPH oxidase subunit gp47^{phox}) and proapoptotic markers (Bax and caspase-3) (14). Fuenzalida *et al.* (36) demonstrated that rosiglitazone upregulated Bcl-2 protein expression in neurons, induced mitochondrial stabilization, and prevented oxidative stress and apoptosis. These results indicated that the PPAR γ agonist rosiglitazone may exert neuroprotective effects through antioxidative and antiapoptotic mechanisms. To confirm the protective effect of PPAR γ agonists on PC12 cells, PC12 cells were incubated with different concentrations of pioglitazone before exposure to H₂O₂ for 1 h. The MTT assay results showed that pioglitazone concentration-dependently increased the survival rate of PC12 cells. The results of flow cytometric analysis and the TUNEL assay also confirmed the conclusion of the MTT assay. There were fewer early and late apoptotic cells in the pioglitazone + H₂O₂ group than in the H₂O₂ group, and the apoptosis rate was considerably lower in the pioglitazone + H₂O₂ group. RT-qPCR and western blot analyses confirmed that pioglitazone significantly increased PPAR γ expression in PC12 cells to a level 4.4-fold higher than that in the control group. This confirmed a previous report that neuroprotective concentrations of pioglitazone can induce a 5-fold increase in PPAR γ expression, thereby maintaining responsiveness of cortical neurons by increasing the expression of its receptors (37).

In addition, H₂O₂ induced an increase in PPAR γ expression, which may be a compensatory protective mechanism for the cells against oxidative stress injury (38). Western blotting was used to evaluate the protein expression levels of Bax, Bcl-2 and cleaved caspase-3/caspase-3 ratio. The results demonstrated that 100 μ mol/l H₂O₂ increased the protein expression levels of Bax and cleaved caspase-3/caspase-3 ratio, decreased Bcl-2 protein expression and increased the ratio of Bax to Bcl-2. Pioglitazone downregulated the protein expression levels of Bax and cleaved caspase-3/caspase-3 ratio, upregulated Bcl-2 protein expression, thus reducing the ratio of Bax to Bcl-2. These results suggested that pioglitazone could attenuate the H₂O₂-induced proapoptotic environment in PC12 cells. To further explore whether rosiglitazone can serve a role in the activation of PPAR γ , the PPAR γ antagonist GW9662 and PPAR γ -siRNA were used to block PPAR γ expression in PC12 cells. The results demonstrated that pretreatment with GW9662 significantly increased the Bax/Bcl-2 ratio and cleaved caspase-3/caspase-3 ratio in PC12 cells compared with those in the pioglitazone + H₂O₂ group. PPAR γ -siRNA had the same effects. Therefore, GW9662 and PPAR γ -siRNA could offset the protective effect of pioglitazone on PC12 cells with H₂O₂-induced injury.

In conclusion, in the present study, a model of neuronal apoptosis induced by oxidative stress was established *in vitro*, and the neuroprotective effect of pioglitazone was studied. The results showed that pioglitazone increased antioxidant activity in PC12 cells with H₂O₂-induced injury, increased the expression levels of Bcl-2, and decreased the protein levels of Bax and cleaved caspase-3, thus ameliorating the proapoptotic environment and reducing the apoptosis rate of PC12 cells. Treatment with a PPAR γ antagonist or PPAR γ -siRNA inhibited the protective effect of pioglitazone on PC12 cells with H₂O₂-induced injury, suggesting that pioglitazone could protect PC12 cells against oxidative stress injury through PPAR γ activation. In conclusion, pioglitazone

exerted an antiapoptotic effect and promoted the survival of PC12 cells in the presence of oxidative stress injury. This effect occurred through PPAR γ activation. Therefore, the present study suggested that PPAR γ activation might have intervention potential in neurodegenerative disorders. The limitation of the present study was that primary neuronal cultures, which represent neuronal properties better than PC12 cells, were not used. In addition, ROS were not directly detected before and after H₂O₂ treatment, which would be an improved approach to support the successful establishment of the PC12 cell model.

Acknowledgements

The authors would like to thank Professor Yi Zhu (Department of Physiology and Pathophysiology, Tian Jin Medical University, Tianjin, China), Professor Dengfeng Gao (Department of Cardiovascular Medicine, Xi'an Jiaotong University, Xi'an, China), Dr Jianbo Yu (Department of Rehabilitation, Weihai Municipal Hospital, Weihai, China) and Dr Yinchun Jiao (Department of Rehabilitation, Rushan Hospital of Traditional Chinese Medicine, Weihai, China) for their excellent technical assistance.

Funding

The current study was supported by the Doctoral Research Fund of the Affiliated Hospital of Weifang Medical University (grant no. 2022BSQD11), the Natural Science Foundation of Shandong Province (grant no. ZR202102240754), and the Shandong Medicine and Health Science and Technology Development Plan Project (grant no. 202003070101).

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YalL, LX and XN conceived and designed the experiments. LL, BH, LX and JL performed all experiments. YanL, BH, JL and ZY collected experimental data and performed the statistical analysis. YalL and LX wrote the paper, which was revised and polished by XN. JL and YanL confirmed the authenticity of all the raw data. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Gitler AD, Dhillon P and Shorter J: Neurodegenerative disease: Models, mechanisms, and a new hope. *Dis Model Mech* 10: 499-502, 2017.
- Huang H, Chen L, Mao G, Bach J, Xue Q, Han F, Guo X, Otom A, Chernykh E, Alvarez E, *et al*: The 2019 yearbook of neurorestoratology. *J Neurorestoratology* 8: 1-11, 2020.
- Tang XQ, Feng JQ, Chen J, Chen PX, Zhi JL, Cui Y, Guo RX and Yu HM: Protection of oxidative preconditioning against apoptosis induced by H₂O₂ in PC12 cells: Mechanisms via MMP, ROS, and Bcl-2. *Brain Res* 1057: 57-64, 2005.
- Shao A, Lin D, Wang L, Tu S, Lenahan C and Zhang J: Oxidative stress at the crossroads of aging, stroke and depression. *Aging Dis* 11: 1537-1566, 2020.
- Wiatrak B, Kubis-Kubiak A, Piwowar A and Barg E: PC12 cell line: Cell types, coating of culture vessels, differentiation and other culture conditions. *Cells* 9: 958, 2020.
- D'Arcy MS: Cell death: A review of the major forms of apoptosis, necrosis and autophagy. *Cell Biol Int* 43: 582-592, 2019.
- Jensen K, WuWong DJ, Wong S, Matsuyama M and Matsuyama S: Pharmacological inhibition of Bax-induced cell death: Bax-inhibiting peptides and small compounds inhibiting Bax. *Exp Biol Med*(Maywood) 244: 621-629, 2019.
- Zhou Z, Meng M and Ni H: Chemosensitizing effect of astragalus polysaccharides on nasopharyngeal carcinoma cells by inducing apoptosis and modulating expression of Bax/Bcl-2 ratio and caspases. *Med Sci Monit* 23: 462-469, 2017.
- Jiang H, Zhao PJ, Su D, Feng J and Ma SL: Paris saponin I induces apoptosis via increasing the Bax/Bcl-2 ratio and caspase-3 expression in gefitinib-resistant non-small cell lung cancer *in vitro* and *in vivo*. *Mol Med Rep* 9: 2265-2272, 2014.
- Jang JH and Surh YJ: Protective effects of resveratrol on hydrogen peroxide-induced apoptosis in rat pheochromocytoma (PC12) cells. *Mutat Res* 496: 181-190, 2001.
- Del Rio MJ and Velez-Pardo C: Monoamine neurotoxins-induced apoptosis in lymphocytes by a common oxidative stress mechanism: Involvement of hydrogen peroxide (H₂O₂), caspase-3, and nuclear factor kappa-B (NF-kappaB), p53, c-Jun transcription factors. *Biochem Pharmacol* 63: 677-688, 2002.
- Stump M, Mukohda M, Hu C and Sigmund CD: PPAR γ regulation in hypertension and metabolic syndrome. *Curr Hypertens Rep* 17: 89, 2015.
- Wang SB, Dougherty EJ and Danner RL: PPAR γ signaling and emerging opportunities for improved therapeutics. *Pharmacol Res* 111: 76-85, 2016.
- Li Y, Yu G, Liu L, Long J, Su S, Zhao T, Liu W, Shen S and Niu X: Rosiglitazone attenuates cell apoptosis through antioxidative and anti-apoptotic pathways in the hippocampi of spontaneously hypertensive rats. *Int J Mol Med* 43: 693-700, 2019.
- Kumar P, Nagarajan A and Uchil PD: Analysis of cell viability by the MTT assay. *Cold Spring Harb Protoc* 2018, 2018.
- Xiao Z, Yu X, Zhang S and Liang A: The expression levels and significance of GSH, MDA, SOD, and 8-OHdG in osteochondral defects of rabbit knee joints. *Biomed Res Int* 2022: 6916179, 2022.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Refaie MMM and El-Hussieny M: Protective effect of pioglitazone on ovarian ischemia reperfusion injury of female rats via modulation of peroxisome proliferator activated receptor gamma and heme-oxygenase 1. *Int Immunopharmacol* 62: 7-14, 2018.
- Li Y, Liu J, Gao D, Wei J, Yuan H, Niu X and Zhang Q: Age-related changes in hypertensive brain damage in the hippocampi of spontaneously hypertensive rats. *Mol Med Rep* 13: 2552-2560, 2016.
- Lahiani A, Brand-Yavin A, Yavin E and Lazarovici P: Neuroprotective effects of bioactive compounds and MAPK pathway modulation in 'ischemia'-stressed PC12 pheochromocytoma cells. *Brain Sci* 8: 32, 2018.
- Satoh T, Sakai N, Enokido Y, Uchiyama Y and Hatanaka H: Free radical-independent protection by nerve growth factor and Bcl-2 of PC12 cells from hydrogen peroxide-triggered apoptosis. *J Biochem* 120: 540-546, 1996.
- Guo Z, Huang M, Yuan Y, Guo Y, Song C, Wang H and Dang X: Nischarin downregulation attenuates cell injury induced by oxidative stress via Wnt signaling. *Neuroreport* 31: 1199-1207, 2020.
- Nakayama H, Nakahara M, Matsugi E, Soda M, Hattori T, Hara K, Usami A, Kusumoto C, Higashiyama S and Kitaichi K: Protective effect of ferulic acid against hydrogen peroxide induced apoptosis in PC12 cells. *Molecules* 26: 90, 2020.
- Xiao L, Liao F, Ide R, Horie T, Fan Y, Saiki C and Miwa N: Enzyme-digested Colla Corii Asini (E'jiao) prevents hydrogen peroxide-induced cell death and accelerates amyloid beta clearance in neuronal-like PC12 cells. *Neural Regen Res* 15: 2270-2272, 2020.
- Islam MT: Oxidative stress and mitochondrial dysfunction-linked neurodegenerative disorders. *Neurol Res* 39: 73-82, 2017.
- de Oliveira Ulbrecht MO, Gonçalves DA, Zanoni LZG and do Nascimento VA: Association between selenium and malondialdehyde as an efficient biomarker of oxidative stress in infantile cardiac surgery. *Biol Trace Elem Res* 187: 74-79, 2019.
- Wang XF, Huang YF, Wang L, Xu LQ, Yu XT, Liu YH, Li CL, Zhan JY, Su ZR, Chen JN and Zeng HF: Photo-protective activity of pogostone against UV-induced skin premature aging in mice. *Exp Gerontol* 77: 76-86, 2016.
- Dumont A, Hehner SP, Hofmann TG, Ueffing M, Dröge W and Schmitz ML: Hydrogen peroxide-induced apoptosis is CD95-independent, requires the release of mitochondria-derived reactive oxygen species and the activation of NF-kappaB. *Oncogene* 18: 747-757, 1999.
- Matsura T, Kai M, Fujii Y, Ito H and Yamada K: Hydrogen peroxide-induced apoptosis in HL-60 cells requires caspase-3 activation. *Free Radic Res* 30: 73-83, 1999.
- Kitamura Y, Ota T, Matsuoka Y, Tooyama I, Kimura H, Shimohama S, Nomura Y, Gebicke-Haerter PJ and Taniguchi T: Hydrogen peroxide-induced apoptosis mediated by p53 protein in glial cells. *Glia* 25: 154-164, 1999.
- Jiang MC, Liang HJ, Liao CF and Lu FJ: Methyl methanesulfonate and hydrogen peroxide differentially regulate p53 accumulation in hepatoblastoma cells. *Toxicol Lett* 106: 201-208, 1999.
- Li RL, Zhang Q, Liu J, Sun JY, He LY, Duan HX, Peng W and Wu CJ: Hydroxy- α -sanshool possesses protective potentials on H₂O₂-stimulated PC12 cells by suppression of oxidative stress-induced apoptosis through regulation of PI3K/Akt signal pathway. *Oxid Med Cell Longev* 2020: 3481758, 2020.
- Senichkin VV, Pervushin NV, Zuev AP, Zhivotovsky B and Kopeina GS: Targeting Bcl-2 family proteins: What, where, when? *Biochemistry (Mosc)* 85: 1210-1226, 2020.
- Oltvai ZN, Millman CL and Korsmeyer SJ: Bcl-2 heterodimerizes *in vivo* with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* 74: 609-619, 1993.
- Ding YP, Kang J, Liu S, Xu Y and Shao B: The protective effects of peroxisome proliferator-activated receptor gamma in cerebral ischemia-reperfusion injury. *Front Neurol* 11: 588516, 2020.
- Fuenzalida K, Quintanilla R, Ramos P, Piderit D, Fuentealba RA, Martinez G, Inestrosa NC and Bronfman M: Peroxisome proliferator-activated receptor gamma up-regulates the Bcl-2 anti-apoptotic protein in neurons and induces mitochondrial stabilization and protection against oxidative stress and apoptosis. *J Biol Chem* 282: 37006-37015, 2007.
- Cimini A, Benedetti E, Cristiano L, Sebastiani P, D'Amico MA, D'Angelo B and Di Loreto S: Expression of peroxisome proliferator-activated receptors (PPARs) and retinoic acid receptors (RXRs) in rat cortical neurons. *Neuroscience* 130: 325-337, 2005.
- Wang N, Han J, Xu Y, Liang H, Cheng Y and Sun L: Hydrogen peroxide-induced changes of PPAR γ in primary cultured cortical neurons. *Zhonghua Yi Xue Za Zhi* 95: 1686-1690, 2015 (In Chinese).



Copyright © 2023 Li et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.