

Associations of recurrent lumbar disc herniation after percutaneous endoscopic lumbar discectomy with age, body mass index, modic change, disc degeneration and sacral slope: A quantitative review

JINLONG ZHAO¹⁻³, LINGFENG ZENG¹⁻³, SHUAI ZHAO², GUIHONG LIANG¹⁻³,
BANGXIN SHA⁴, HAOYANG FU², WEIYI YANG², JUN LIU¹⁻⁴ and YUPING ZENG^{1,2}

¹Department of Orthopedics, The Second Clinical College of Guangzhou University of Chinese Medicine/Guangdong Provincial Hospital of Chinese Medicine, Guangzhou, Guangdong 510405; ²Department of Orthopedics, The Second Affiliated Hospital of Guangzhou University of Chinese Medicine; ³The Research Team on Bone and Joint Degeneration and Injury, Guangdong Provincial Academy of Chinese Medical Sciences, Guangzhou, Guangdong 510120; ⁴Department of Orthopedics, Guangdong Second Chinese Medicine Hospital (Guangdong Province Engineering Technology Research Institute of Traditional Chinese Medicine), Guangzhou, Guangdong 510095, P.R. China

Received August 13, 2023; Accepted February 20, 2024

DOI: 10.3892/etm.2024.12483

Abstract. Recurrent lumbar disc herniation (rLDH) seriously affects the quality of life of patients and increases the medical burden. The purpose of the present study was to determine the risk factors for rLDH after percutaneous endoscopic lumbar discectomy (PELD). The PubMed, Cochrane Library and Embase databases were searched for studies on the factors associated with rLDH after PELD. The databases were searched from inception to March 30, 2023. The combined effects of categorical variables and continuous variables were measured using odds ratios (ORs) and weighted mean differences (WMDs), respectively, and their corresponding 95% confidence intervals (CIs) were calculated. RevMan 5.3 software

was used for data analysis. A total of 9 case-control studies were included in this meta-analysis, comprising 5,446 patients. This study explored a total of 18 potential risk factors for rLDH after PELD; ultimately, 5 factors were associated with the risk of rLDH. Meta-analysis showed that older age (WMD=6.49, 95% CI: 2.52 to 10.46), greater body mass index (WMD=1.16, 95% CI: 0.69 to 1.62), modic change (OR=2.48, 95% CI: 1.54 to 3.99), Pfirrmann grade ≥ 4 (OR=2.84, 95% CI: 1.3 to 6.16) and greater sacral slope angle (WMD=3.48, 95% CI: 0.53 to 6.42) were risk factors for rLDH after PELD. The risk factors identified in the present study may enable clinicians to identify high-risk populations early and to select appropriate surgical procedures to reduce the risk of rLDH. Perioperative interventions targeting the modifiable factors identified in this study may be beneficial for reducing the risk of rLDH.

Correspondence to: Professor Jun Liu, Department of Orthopedics, The Second Clinical College of Guangzhou University of Chinese Medicine/Guangdong Provincial Hospital of Chinese Medicine, 111 Dade Road, Yuexiu, Guangzhou, Guangdong 510405, P.R. China
E-mail: gzucmljujun@foxmail.com

Professor Yuping Zeng, Department of Orthopedics, The Second Affiliated Hospital of Guangzhou University of Chinese Medicine, 111 Dade Road, Yuexiu, Guangzhou, Guangdong 510120, P.R. China
E-mail: zyp8365@163.com

Abbreviations: BMI, body mass index; CI, confidence interval; DHI, disc height index; DM, diabetes mellitus; IV, inverse variance; M-H, Mantel-Haenszel test; NOS, Newcastle-Ottawa scale; NR, not reported; OR, odds ratio; PELD, percutaneous endoscopic lumbar discectomy; rLDH, recurrent lumbar disc herniation; sROM, sagittal range of motion

Key words: risk factors, recurrent lumbar disc herniation, percutaneous endoscopic lumbar discectomy, meta-analysis

Introduction

Lumbar disc herniation (LDH) refers to a series of syndromes characterized by low back pain and lower limb radiation pain induced by the stimulation or compression of nerves due to degeneration or injury of the lumbar intervertebral disc, leading to rupture of the fibrous ring and protrusion of the nucleus pulposus (1,2). LDH mainly occurs at L4-5 and L5-S1 and is mainly observed in middle-aged and elderly individuals, with a male predominance (3). LDH can limit mobility and the ability to perform physical activities, and cause disability (4), thereby reducing the quality of life of patients and increasing medical costs. Although conservative treatment is the first choice for treating LDH, surgery is necessary for severe LDH to achieve good clinical outcomes (5,6).

With the continuous development of minimally invasive spinal technology, percutaneous endoscopic lumbar discectomy (PELD) is the main method of surgical treatment for LDH (7). PELD has numerous advantages, such as minimal trauma,

less bleeding, a clear surgical field, relatively low surgical costs and high patient satisfaction (8). In addition, PELD can be performed under local anaesthesia, which may lead to the acquisition of knowledge regarding a patient's neurological and vascular injuries during surgery (9). Whether from a medical or patient perspective, the use of PELD to treat LDH has numerous technical advantages. However, the widespread application of endoscopic technology has led to post-PELD complications, including recurrent LDH (rLDH), receiving increased attention (10). According to the literature, 2-25% of patients who undergo PELD experience rLDH after surgery (10-12). Surgical intervention is necessary for patients with rLDH who have been clearly diagnosed and have not yet achieved remission after conservative treatment. However, scar tissue after the initial surgery increases the difficulty of repeated discectomy and increases the risk of dural tears or nerve damage (13,14). In addition, removing the posterior structure during reoperation may increase the risk of lumbar segmental instability (15,16). There is no doubt that both reoperation and the clinical symptoms associated with rLDH have a significant negative impact on the psychological burden and economic pressure of patients.

PELD is a common surgical method for the treatment of LDH, and rLDH is a possible serious complication. In this context, identifying the risk factors for rLDH after PELD has important clinical value for formulating appropriate surgical protocols and evaluating postoperative efficacy. However, there is a lack of high-level evidence regarding the risk factors for rLDH after PELD. To fill this gap, a meta-analysis was performed to evaluate and identify the risk factors for rLDH after PELD. The conclusions of this study are intended to provide a theoretical basis for the clinical prevention and reduction of postoperative rLDH after PELD.

Materials and methods

Data sources and retrieval strategies. The design and implementation of the present study followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (17). Three databases, including PubMed (www.pubmed.ncbi.nlm.nih.gov), Cochrane Library (www.cochranelibrary.com) and Embase (www.embase.com), were searched to identify studies that examined risk factors for rLDH after PELD. The databases were searched from inception to March 30th, 2023. Furthermore, the reference lists of the included studies were manually searched to obtain additional eligible studies. The search strategy included a combination of MeSH terms and free words. The key words included 'recurrent lumbar disc herniation', 'rLDH', 'reoperation', 'repeat discectomy', 'PELD', 'percutaneous endoscopic lumbar discectomy', 'percutaneous transforaminal endoscopic discectomy', and 'Yeung endoscopic spine system'. The details of the search strategies for each database are provided in Appendix S1.

Inclusion and exclusion criteria. The inclusion criteria were as follows: i) Study participants of any age, sex, ethnicity or region, including patients in need to be diagnosed with LDH and treated with PELD; ii) the research topic was risk factors for rLDH and complete data on grouping comparison of rLDH and non-rLDH were available in the literature; iii) due to the fact that the present study is a secondary literature analysis,

the study type was limited to cohort and case-control studies based on the existing literature. Ostensibly, rLDH typically occurs after surgery. Due to ethical considerations and implementation difficulties, randomized controlled trials may not be the most appropriate method for studying risk factors, and thus, the present quantitative review only included cohort studies and case-control studies; iv) the included literature was required to contain reports of at least one risk factor; v) the outcome measurement data were required to be expressed as the frequency (%) or mean \pm standard deviation in order to help reduce statistical heterogeneity. There were no restrictions regarding the language of the publications.

The exclusion criteria were as follows: i) Duplicate literature, case reports and animal experiments; ii) obvious statistical errors in the literature; iii) repeated publications of the same study population; and iv) the full text cannot be obtained.

Data extraction. Two researchers (GL, BS) independently extracted data from the included literature and cross-checked them. Disagreements were resolved through discussion or consultation with the corresponding author. The process of literature selection included selecting the literature that met the research purpose by reading the titles and abstracts, then reading the full texts of the literature according to the inclusion and exclusion criteria, and finally selecting the literature that met the research purpose. The following data were extracted: First author, publication time, research design type, sample size, follow-up time and risk factors.

Literature quality evaluation. The quality of the included studies was evaluated by two researchers using the Newcastle-Ottawa scale (NOS) (18). The NOS consists of three components, including study subject selection (4 points), intergroup comparability (2 points) and outcome or exposure factor evaluation (3 points), with a total score of 9 points. Scores of 0 to 3, 4 to 6 and 7 to 9 were considered to indicate low-, moderate- and high-quality research, respectively.

Data analysis. RevMan 5.3 software (The Cochrane Collaboration) was used for data analysis. For categorical and continuous variables, odds ratios (ORs) and weighted mean differences (WMDs) were used as effect size measures, respectively, and point estimates and 95% confidence intervals (CIs) for each effect size measure were calculated. The random-effects model was used for all meta-analyses. The test level for meta-analysis was set to $\alpha=0.05$. Sensitivity analysis was performed by comparing the result consistency between the fixed-effects and random-effects models and eliminating studies with greater impact. And the method exclusion of one study at a time and re-performing the meta-analysis was also performed. A funnel plot was constructed to assess whether there was evidence of publication bias. In addition, each outcome was required to be examined by at least 3 studies to be considered for pooled data analysis.

Results

Literature search results. Based on the search strategy of the present study, a total of 391 relevant documents were retrieved. After removing duplicate publications and reading the titles




Figure 1. Flow diagram of systematic literature search. PELD, percutaneous endoscopic lumbar discectomy; rLDH, recurrent lumbar disc herniation.

and abstracts of the studies, 27 articles remained for full-text screening. Referring to the inclusion criteria of the present study and after reading the full texts, 9 case-control studies were ultimately included in the present meta-analysis (19-27). The specific details of the study selection process are presented in Fig. 1.

Basic characteristics of the included studies. A total of 9 case-control studies were included in the present study and they were published between 2020 and 2023, indicating that research on rLDH after PELD has received extensive attention from researchers in recent years. A total of 5,446 patients who underwent PELD were included in this meta-analysis, with 560 rLDH patients (343 males and 217 females) and 4,886 non-rLDH patients (2,811 males and 2,075 females). The weighted mean age was 50.75 in the rLDH group and 44.53 in the non-rLDH group. The follow-up period ranged from 6-48 months. The operative segments of LDH were mainly L4/5 and L5/S1. The basic characteristics of the nine included studies are presented in Table I.

Results of the literature quality evaluation. The NOS scores of the nine included studies (19-27) are shown in Table II. The NOS scores ranged from 7-8, which indicates that the methodological quality of all studies included in the present study is high.

Meta-analysis

Sociodemographic and anthropometry factors. A total of 9 related factors were analysed (Table III) and the corresponding forest maps of sociodemographic and anthropometry factors

are shown in Figs. S1-S9. The meta-analysis found that age and body mass index (BMI) are both significant risk factors for rLDH. Sex, symptom duration, smoking, drinking, DM and hypertension were not risk factors for rLDH (all $P > 0.05$). A total of 8 studies (19,21-27) reported an association between age and rLDH, and there was heterogeneity among the studies ($I^2 = 93\%$). The random-effects model was used to perform meta-analysis and the results showed that higher patient age was associated with a greater risk of rLDH after PELD (WMD=6.49, 95% CI: 2.52 to 10.46, $P = 0.001$).

A total of 8 studies (19,21-27) reported the association between BMI and rLDH. There was heterogeneity among the eight studies, and thus, a random-effects model was used for meta-analysis. It was found that a higher BMI significantly increased the risk of rLDH (WMD=1.16, 95% CI: 0.69 to 1.62, $P < 0.001$).

Clinical and imaging factors. Similarly, this study also analysed the association of 9 clinical and imaging factors with rLDH (Table III) the corresponding forest maps are provided in Figs. S10-S18. A total of 7 studies (19,20,22-25,27) reported the effect of modic change on rLDH, and there was heterogeneity among these studies ($I^2 = 73\%$). The meta-analysis results showed that there was an association between modic change and the risk of rLDH after PELD (OR=2.48, 95% CI: 1.54 to 3.99, $P = 0.0002$).

A total of 3 studies (19,25,27) reported the association of the Pfirrmann grade with rLDH. The meta-analysis results showed that a Pfirrmann grade ≥ 4 was a risk factor for rLDH after PELD (OR=2.84, 95% CI: 1.30 to 6.16) and this association was statistically significant ($P = 0.008$).

Table I. Characteristics of the included studies.

First author, year	Design	Sample size (M/F), n		Mean age, years		Level of herniated disk, n (L2/L3, L3/L4, L4/5, L5/S1)		Type of herniation, n (central/paracentral/ lateral or extreme lateral)		Follow-up (months)	(Refs.)
		RLDH	Non-RLDH	RLDH	Non-RLDH	RLDH	Non-RLDH	RLDH	Non-RLDH		
He <i>et al</i> , 2023	Case-control study	63 (45/18)	627 (410/217)	51.49	42.87	0/0/37/26	0/0/370/257	NR	NR	24	(19)
Jia <i>et al</i> , 2021	Case-control study	32 (18/14)	320 (212/108)	NR	NR	0/2/18/12	0/12/165/143	1/31/0	3/177/140	6	(20)
Kong <i>et al</i> , 2020	Case-control study	46 (22/24)	608 (333/275)	54.4	44.9	NR	NR	31/13/20	165/212/231	28	(21)
Li <i>et al</i> , 2023	Case-control study	56 (30/26)	589 (296/293)	51.82	48.32	4/6/29/17	6/22/412/149	4/26/11	50/360/179	24	(22)
Ren <i>et al</i> , 2023	Case-control study	130 (80/50)	1059 (640/419)	46.62	44.2	NR	NR	NR	NR	38	(23)
Shi <i>et al</i> , 2021	Case-control study	68 (47/21)	136 (94/42)	46.21	47.36	0/0/41/27	0/0/82/54	NR	NR	24	(24)
Wang <i>et al</i> , 2022	Case-control study	57 (39/18)	885 (462/423)	56.7	41.2	0/0/30/27	3/18/405/459	6/42/9	69/732/84	24	(25)
Yu <i>et al</i> , 2020	Case-control study	46 (25/21)	438 (229/209)	52.98	47.74	NR	NR	13/27/6	100/237/101	12 to 48	(26)
Zhao <i>et al</i> , 2021	Case-control study	62 (37/25)	224 (135/89)	52.1	44.9	NR	NR	NR	NR	24	(27)

rLDH, recurrent lumbar disc herniation; M, male; F, female; NR, not reported.

Table II. Newcastle-Ottawa scale for risk of bias assessment of studies included in the meta-analysis.

First author, year	Selection			Outcome				Overall	(Refs.)	
	Representativeness of exposed cohort	Selection of nonexposed	Ascertainment of exposure	Outcome not present at start	Comparability	Assessment of outcome	Adequate follow-up length			Adequacy of follow-up
He <i>et al</i> , 2023	-	*	*	*	**	*	*	*	8	(19)
Jia <i>et al</i> , 2021	-	*	*	*	**	*	*	-	7	(20)
Kong <i>et al</i> , 2020	-	*	*	*	**	-	*	*	7	(21)
Li <i>et al</i> , 2023	-	*	*	*	**	*	*	*	8	(22)
Ren <i>et al</i> , 2023	-	*	*	*	**	-	*	*	7	(23)
Shi <i>et al</i> , 2021	-	*	*	*	**	*	*	*	7	(24)
Wang <i>et al</i> , 2022	-	*	*	*	**	*	*	*	8	(25)
Yu <i>et al</i> , 2020	-	*	*	*	**	*	*	*	8	(26)
Zhao <i>et al</i> , 2021	-	*	*	*	**	-	*	*	7	(27)

*, score of 1; **, score of 2; -, score of 0.

Table III. Meta-analysis of the risk factors.

A, Sociodemographic and anthropometry factors						
Factor	Studies, n	OR or WMD	95% CI	P-value	I^2 , %	Analysis model
Age, years	8	6.49	2.52-10.46	0.001	93	IV, random
Male sex	9	1.08	0.90-1.30	0.410	2	M-H, random
Female sex	9	0.92	0.77-1.11	0.410	2	M-H, random
BMI, kg/m ²	8	1.16	0.69-1.62	<0.001	69	IV, random
Symptom duration, months	4	1.71	-0.18-3.59	0.080	97	IV, random
Smoking (yes vs. no)	8	1.43	0.95-2.14	0.090	69	M-H, random
Drinking (yes vs. no)	5	1.13	0.83-1.53	0.450	0	M-H, random
DM (yes vs. no)	7	1.22	0.85-1.74	0.280	5	M-H, random
Hypertension (yes vs. no)	4	1.00	0.69-1.47	0.990	1	M-H, random
B, Clinical and imaging factors						
Factor	Studies, n	OR or WMD	95% CI	P-value	I^2 , %	Analysis model
Operation time, min	3	-0.19	-5.54-5.15	0.940	59	IV, random
Modic change (yes vs. no)	7	2.48	1.54-3.99	<0.001	73	M-H, random
Pfirsman grade ≥ 4	3	2.84	1.30-6.16	0.008	81	M-H, random
DHI	5	-0.01	-0.04-0.02	0.640	92	IV, random
sROM, °	5	1.50	-0.59-3.58	0.160	96	IV, random
Facet orientation, °	3	-1.82	-3.72-0.07	0.060	76	IV, random
Facet tropism, °	3	-0.36	-0.80-0.08	0.110	10	IV, random
Lumbar lordosis angle, °	4	-1.18	-6.43-4.07	0.660	95	IV, random
Sacral slope angle, °	3	3.48	0.53-6.42	0.020	86	IV, random

The forest maps of all risk factors are shown in Figs. S1-S18. BMI, body mass index; DM, diabetes mellitus; DHI, disc height index; sROM, sagittal range of motion; OR, odds ratio; WMD, weighted mean difference; M-H, Mantel Haenszel test; IV, inverse variance; random, random-effects model.

A total of 3 studies (23,26,27) reported the association between sacral slope angle and rLDH, and there was heterogeneity among these three studies ($I^2=86\%$). Meta-analysis showed that a larger sacral slope angle was a risk factor for rLDH after PELD (WMD=3.48, 95% CI: 0.53 to 6.42), with a statistically significant difference ($P=0.02$).

Publication bias. Four funnel plots (age, male, female, BMI) were constructed to evaluate publication bias (Fig. 2). The scattered points in the funnel diagram are basically symmetrical, suggesting that there is no evidence of publication bias in the evaluation of age, male, female and BMI.

Discussion

Due to population ageing and changes in modern lifestyles, the occurrence of LDH exhibits a trend of growth (28). Epidemiological studies have shown that the overall incidence rate of LDH ranges from 2-3%, but the incidence rates among men and women aged >35 years are 4.8 and 2.5%, respectively (28). PELD is often used to treat LDH that is not effectively managed by conservative treatment (6). Identifying

the risk factors for rLDH after PELD surgery and formulating the best treatment strategy or informing patients of the risks before surgery is an important task. In the past 20 years, there have been 2 systematic reviews evaluating the risk factors for rLDH (29,30), but these two studies have certain limitations that may prevent us from applying their conclusions to the identification of rLDH after PELD. The two systematic evaluations (29,30) comprised patients who underwent surgery, including microdiscectomy, laminotomy, PELD, decompression, lumbar fusion and open discectomy, indicating significant clinical heterogeneity among the included studies. In addition, both systematic evaluations are concerned with a small number of potential risk factors, providing less evidence for clinical application. Since PELD is the most commonly used surgical method, only the risk factors for rLDH after PELD were examined; the present results may allow clinicians to evaluate risk of rLDH following PELD. In addition, there were 18 potential risk factors observed in the present meta-analysis, but only 5 factors were significant, which provide more comprehensive information. The present study found that higher age, greater BMI, modic change, Pfirsman grade ≥ 4 and greater sacral slope angle are all significant risk




Figure 2. Funnel plots of (A) age, (B) male, (C) female, and (D) BMI. BMI, body mass index.

factors for rLDH after PELD. Therefore, PELD may not be the best option at the stage of developing medical strategies for patients with one or more of the above risk factors. The findings of the present study will help spine surgeons develop appropriate surgical protocols and better inform patients of surgical risks.

Regarding sociodemographic and anthropometry factors, consistent with previously published research findings (31-33), the present study also found a significant association between higher age and the occurrence of rLDH. A retrospective cohort study showed that the reoperation rate of patients aged ≥ 57 years who underwent PELD was higher than that of patients aged < 57 years (34), and this study found that rLDH after PELD was the main reason for reoperation. As ageing progresses, the degradation of spinal tissue becomes increasingly severe. Ehrendorfer *et al* (35) found that the severity of preoperative disc degeneration in patients with rLDH is significantly higher than that in non-rLDH patients. A study indicated that obese patients are more prone to LDH (36) and are prone to adverse clinical outcomes after surgery. A Cox regression analysis showed that a BMI ≥ 25 kg/m² is an important risk factor for rLDH after PELD (37). The study by Yaman *et al* (38) indicated that the reoperation rate of overweight or obese patients after PELD is significantly higher than that of patients with LDH with a normal BMI, which may be related to increased load on the intervertebral disc. Therefore, it may be indicated that both older age and a greater BMI are highly associated with the risk of rLDH.

Clinical and imaging factors were also examined in the present meta-analysis. Modic changes typically include reactive vertebral changes associated with inflammation,

an unstable microenvironment or degenerative disc disease (39,40). As the intervertebral disc itself does not contain vascular tissue, the micropores in the endplate are exchange channels for nutrients, water and other metabolic products. When the endplate changes, its nutritional effect on the intervertebral disc decreases (39,40). Therefore, it is not difficult to understand that modic changes are associated with a higher risk of rLDH. The present study found that a Pfirrmann grade ≥ 4 is associated with a higher risk of rLDH, which is consistent with the findings of Kim *et al* (41). This latter study concluded that a higher degree of intervertebral disc degeneration is associated with a higher risk of rLDH (41). During the process of intervertebral disc degeneration, type I collagen increases, while type II collagen decreases, and the content of proteoglycans and elastin decreases (42), which causes the nucleus pulposus to lose elasticity and the annulus fibrosus to appear cracked. Therefore, more severe degeneration of the intervertebral disc is associated with poorer self-repairing ability of the fibrous ring, ultimately leading to the protrusion of the nucleus pulposus (42). The present study found that the sacral slope angle is associated with the risk of rLDH after PELD. Studies have shown that a large sacral slope angle can lead to L5-S1 vertebral body slippage and its mechanism is increased stress in the L5-S1 vertebral body (43-45), which may indicate a potential association between the susceptibility to rLDH after PELD. A biomechanical study found that a larger sacral slope angle can lead to lumbar lordosis (45) and increase stress in the lumbar intervertebral disc. Therefore, correcting the sacral slope angle can reduce the risk of rLDH after PELD. A larger sacral slope angle can be considered a risk factor for rLDH.

The present study has several limitations, and resolving these limitations will further increase the reliability of its findings. Due to these limitations, readers need to consider not only the conclusions of this study but also real clinical scenarios to interpret them. First, the present meta-analysis involved only association analyses, so the determination of causal relationships requires further prospective cohort studies or Mendelian randomization studies in the future. Furthermore, although strict inclusion and exclusion criteria were established to ensure comparability among studies, unclear definitions of rLDH, inconsistent sample sizes and differences in follow-up time between certain studies may reduce the reliability of the conclusions of the present study. Finally, there was high statistical heterogeneity for most risk factors in the present study, but our attempt to find the source of heterogeneity was not successful. Therefore, the application of the conclusions of the present study requires clinical workers to consider their clinical experiences.

In conclusion, as significant risk factors for rLDH after PELD surgery, older age, higher BMI, modic change, intervertebral disc degeneration and larger sacral slope angle were identified in the present study. These findings will enable medical workers to identify high-risk populations early and to choose appropriate surgical procedures to reduce the risk of rLDH. Future research should further validate the modifiable risk factors identified in the present study to improve outcomes for patients with rLDH.

Acknowledgements

Not applicable.

Funding

This work was supported by the National Key Research and Development Program (grant no. 2021YFC1712804), Guangdong Basic and Applied Basic Research Foundation (grant no. 2022A1515220131) and Research Fund for Bajan Talents of Guangdong Provincial Hospital of Chinese Medicine (grant no. BJ2022KY01).

Availability of data and materials

The data generated in the present study may be requested from the corresponding author.

Authors' contributions

JZ, JL and YZ conceived the study and are responsible for the overall content. Literature screening and data processing were performed by GL and BS. SZ, LZ, GL, BS, HF, WY and JZ analyzed and interpreted the data. JZ, LZ and GL prepared the manuscript. JL and LZ edited the manuscript. JZ, LZ and SZ contributed equally to this work. JL and YZ confirm the authenticity of all the raw data. All authors have read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Amin RM, Andrade NS and Neuman BJ: Lumbar disc herniation. *Curr Rev Musculoskelet Med* 10: 507-516, 2017.
2. Benzakour T, Igoumenou V, Mavrogenis AF and Benzakour A: Current concepts for lumbar disc herniation. *Int Orthop* 43: 841-851, 2019.
3. Gadjradj PS, Smeele NVR, de Jong M, Depauw PRAM, van Tulder MW, de Bekker-Grob EW and Harhangi BS: Patient preferences for treatment of lumbar disc herniation: A discrete choice experiment. *J Neurosurg Spine* 36: 704-712, 2022.
4. Protzer LA, Glassman SD, Mummaneni PV, Bydon M, Bisson EF, Djurasovic M and Carreon LY: Return to work in patients with lumbar disc herniation undergoing fusion. *J Orthop Surg Res* 16: 534, 2021.
5. Cho S, Lim YC, Kim EJ, Park Y, Ha IH, Lee YS and Lee YJ: Analysis of conservative treatment trends for lumbar disc herniation with radiculopathy in Korea: A population-based cross-sectional study. *Healthcare (Basel)* 11: 2353, 2023.
6. Saravi B, Zink A, Ülkümen S, Couillard-Despres S, Wollborn J, Lang G and Hassel F: Clinical and radiomics feature-based outcome analysis in lumbar disc herniation surgery. *BMC Musculoskelet Disord* 24: 791, 2023.
7. Sun DD, Lv D, Wu WZ, Ren HF, Bao BH, Liu Q and Sun ML: Estimation and influence of blood loss under endoscope for percutaneous endoscopic lumbar discectomy (PELD): A clinical observational study combined with in vitro experiment. *J Orthop Surg Res* 15: 281, 2020.
8. Jarebi M, Awaf A, Lefranc M and Peltier J: A matched comparison of outcomes between percutaneous endoscopic lumbar discectomy and open lumbar microdiscectomy for the treatment of lumbar disc herniation: A 2-year retrospective cohort study. *Spine J* 21: 114-121, 2021.
9. Mooney J, Laskay N, Erickson N, Salehani A, Mahavadi A, Ilyas A, Mainali B, Nowak B and Godzik J: General vs local anesthesia for percutaneous endoscopic lumbar discectomy (PELD): A systematic review and meta-analysis. *Global Spine J* 13: 1671-1688, 2023.
10. Shepard N and Cho W: Recurrent lumbar disc herniation: A review. *Global Spine J* 9: 202-209, 2019.
11. Dower A, Chatterji R, Swart A and Winder MJ: Surgical management of recurrent lumbar disc herniation and the role of fusion. *J Clin Neurosci* 23: 44-50, 2016.
12. Hlubek RJ and Mundis GM Jr: Treatment for recurrent lumbar disc herniation. *Curr Rev Musculoskelet Med* 10: 517-520, 2017.
13. Luo L, Zhao C, Li P, Liu L, Zhou Q, Luo F and Liang L: Posterior dynamic stabilization with limited redisection for recurrent lumbar disc herniation. *Pain Res Manag* 2021: 1288246, 2021.
14. Le H, Sandhu FA and Fessler RG: Clinical outcomes after minimal-access surgery for recurrent lumbar disc herniation. *Neurosurg Focus* 15: E12, 2003.
15. Arif S, Brady Z, Enchev Y and Peev N: Is fusion the most suitable treatment option for recurrent lumbar disc herniation? A systematic review. *Neurol Res* 42: 1034-1042, 2020.
16. Chen Z, Zhao J, Liu A, Yuan J and Li Z: Surgical treatment of recurrent lumbar disc herniation by transforaminal lumbar interbody fusion. *Int Orthop* 33: 197-201, 2009.
17. O'Dea RE, Lagisz M, Jennions MD, Koricheva J, Noble DWA, Parker TH, Gurevitch J, Page MJ, Stewart G, Moher D and Nakagawa S: Preferred reporting items for systematic reviews and meta-analyses in ecology and evolutionary biology: Aprisma extension. *Biol Rev Camb Philos Soc* 96: 1695-1722, 2021.
18. Stang A: Critical evaluation of the newcastle-ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 25: 603-605, 2010.
19. He H, Ma J, Xiong C, Wei T, Tang A, Chen Y and Xu F: Development and validation of a nomogram to predict the risk of lumbar disk reherniation within 2 years after percutaneous endoscopic lumbar discectomy. *World Neurosurg* 172: e349-e356, 2023.

20. Jia M, Sheng Y, Chen G, Zhang W, Lin J, Lu S, Li F, Ying J and Teng H: Development and validation of a nomogram predicting the risk of recurrent lumbar disc herniation within 6 months after percutaneous endoscopic lumbar discectomy. *J Orthop Surg Res* 16: 274, 2021.
21. Kong M, Xu D, Gao C, Zhu K, Han S, Zhang H, Zhou C and Ma X: Risk factors for recurrent L4-5 disc herniation after percutaneous endoscopic transforaminal discectomy: A retrospective analysis of 654 cases. *Risk Manag Healthc Policy* 13: 3051-3065, 2020.
22. Li X, Pan B, Cheng L, Li G, Liu J and Yuan F: Development and validation of a prognostic model for the risk of recurrent lumbar disc herniation after percutaneous endoscopic transforaminal discectomy. *Pain Physician* 26: 81-90, 2023.
23. Ren G, Liu L, Jiang P, Xie Z, Wang P, Zhang W, Wang H, Shen M, Deng L, Tao Y, *et al*: Machine learning predicts recurrent lumbar disc herniation following percutaneous endoscopic lumbar discectomy. *Global Spine J* 14: 146-152, 2024.
24. Shi H, Zhu L, Jiang ZL and Wu XT: Radiological risk factors for recurrent lumbar disc herniation after percutaneous transforaminal endoscopic discectomy: A retrospective matched case-control study. *Eur Spine J* 30: 886-892, 2021.
25. Wang F, Chen K, Lin Q, Ma Y, Huang H, Wang C and Zhou P: Earlier or heavier spinal loading is more likely to lead to recurrent lumbar disc herniation after percutaneous endoscopic lumbar discectomy. *J Orthop Surg Res* 17: 356, 2022.
26. Yu C, Zhan X, Liu C, Liao S, Xu J, Liang T, Zhang Z and Chen J: Risk factors for recurrence of L5-S1 disc herniation after percutaneous endoscopic transforaminal discectomy: A retrospective study. *Med Sci Monit* 26: e919888-1-e919888-12, 2020.
27. Zhao C, Zhang H, Wang Y, Xu D, Han S, Meng S, Han J, Liu H, Zhou C and Ma X: Nomograms for predicting recurrent herniation in PETD with preoperative radiological factors. *J Pain Res* 14: 2095-2109, 2021.
28. Vialle LR, Vialle EN, Suárez Henao JE and Giraldo G: LUMBAR DISC HERNIATION. *Rev Bras Ortop* 45: 17-22, 2010.
29. Huang W, Han Z, Liu J, Yu L and Yu X: Risk factors for recurrent lumbar disc herniation: A systematic review and meta-analysis. *Medicine (Baltimore)* 95: e2378, 2016.
30. Yang J, Liu R, Miao Y, Nian L and Meng X: Risk factors for recurrence after percutaneous endoscopic lumbar discectomy: A meta-analysis. *World Neurosurg* 172: 88-93, 2023.
31. Shin EH, Cho KJ, Kim YT and Park MH: Risk factors for recurrent lumbar disc herniation after discectomy. *Int Orthop* 43: 963-967, 2019.
32. Kang MS, Hwang JH, Choi DJ, Chung HJ, Lee JH, Kim HN and Park HJ: Clinical outcome of biportal endoscopic revisional lumbar discectomy for recurrent lumbar disc herniation. *J Orthop Surg Res* 15: 557, 2020.
33. Rossi V, Maalouly J and Choi JYS: Lumbar arthroplasty for treatment of primary or recurrent lumbar disc herniation. *Int Orthop* 47: 1071-1077, 2023.
34. Kim CH, Chung CK, Choi Y, Shin S, Kim MJ, Lee J and Park BJ: The selection of open or percutaneous endoscopic lumbar discectomy according to an age cut-off point: Nationwide cohort study. *Spine (Phila Pa 1976)* 40: E1063-E1070, 2015.
35. Ehrendorfer S: Ipsilateral recurrent lumbar disc herniation. *J Bone Joint Surg Br* 81: 368, 1999.
36. Patgaonkar P, Goyal V, Agrawal U, Marathe N and Patel V: Impact of body weight, height, and obesity on selection of skin entry point for transforaminal endoscopic lumbar discectomy. *Asian J Neurosurg* 17: 262-267, 2022.
37. Yao Y, Liu H, Zhang H, Wang H, Zhang C, Zhang Z, Wu J, Tang Y and Zhou Y: Risk factors for recurrent herniation after percutaneous endoscopic lumbar discectomy. *World Neurosurg* 100: 1-6, 2017.
38. Yaman ME, Kazancı A, Yaman ND, Baş F and Ayberk G: Factors that influence recurrent lumbar disc herniation. *Hong Kong Med J* 23: 258-263, 2017.
39. Udby PM, Modic M, Elmoose S, Carreon LY, Andersen MØ, Karppinen J and Samartzis D: The clinical significance of the modic changes grading score. *Global Spine J* 2022: 21925682221123012, 2022 (Epub ahead of print).
40. Li Z, Gao X, Ding W, Li R and Yang S: Asymmetric distribution of modic changes in patients with lumbar disc herniation. *Eur Spine J* 32: 1741-1750, 2023.
41. Kim KT, Park SW and Kim YB: Disc height and segmental motion as risk factors for recurrent lumbar disc herniation. *Spine (Phila Pa 1976)* 34: 2674-2678, 2009.
42. Moriguchi Y, Borde B, Berlin C, Wipplinger C, Sloan SR, Kirnaz S, Pennicooke B, Navarro-Ramirez R, Khair T, Grunert P, *et al*: In vivo annular repair using high-density collagen gel seeded with annulus fibrosus cells. *Acta Biomater* 79: 230-238, 2018.
43. Labelle H, Mac-Thiong JM and Roussouly P: Spino-pelvic sagittal balance of spondylolisthesis: A review and classification. *Eur Spine J* 20(Suppl 5): 641-646, 2011.
44. Roussouly P, Gollogly S, Berthonnaud E, Labelle H and Weidenbaum M: Sagittal alignment of the spine and pelvis in the presence of L5-s1 isthmus lysis and low-grade spondylolisthesis. *Spine (Phila Pa 1976)* 31: 2484-2490, 2006.
45. Kumaran Y, Nishida N, Tripathi S, Mumtaz M, Sakai T, Elgafy H and Goel VK: Effects of sacral slope changes on the intervertebral disc and hip joint: A finite element analysis. *World Neurosurg* 176: e32-e39, 2023.



Copyright © 2024 Zhao et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.