

Fenofibrate prevents myocardial inflammation and fibrosis via PPAR α /I κ B ζ signaling pathway in rat autoimmune myocarditis

HE CHANG^{1,2}, YUE WANG², YANG WU², SONG-LIN SHI² and ZHI QI²

¹Department of Geriatrics, Xiang'an Hospital of Xiamen University, Xiamen, Fujian 361100, P.R. China;

²Department of Basic Medical Sciences, School of Medicine, Xiamen University, Xiamen, Fujian 361102, P.R. China

Received September 19, 2024; Accepted March 21, 2025

DOI: 10.3892/etm.2025.12879

Abstract. Fenofibrate is a peroxisome proliferator-activated receptor α (PPAR α) agonist currently utilized clinically to reduce lipid levels. Experimental autoimmune myocarditis (EAM) is a T cell-mediated autoimmune myocarditis with histopathological evidence of extensive inflammatory cells infiltrated into the myocardium, resulting in myocardial fibrosis. Previously, the authors demonstrated that long-term fenofibrate could alleviate rat EAM; however, the underlying mechanism remains unclear. In the present study, EAM was induced in rats and fenofibrate was administered to evaluate its short-term effects. The CD4(+) T cells were purified from rats with EAM and PPAR α (-/-) mice and determined the levels of I κ B ζ , pNF- κ Bp65 and IL-6. The findings of the present study revealed that fenofibrate treatment improved the progression of EAM in rats, as evidenced by a decreased ratio of the heart weight to body weight, reduced inflammatory cell infiltration, and relieved cardiac function. It also inhibited the transcription levels of T helper 17 (Th17)-related inflammatory cytokines, fibrosis-associated factors, and the expression of I κ B ζ , and dose-dependently downregulated I κ B ζ expression in differentiated Th17 cells from rats with EAM. The PPAR α antagonist MK886 could reverse the effects in a dose-dependent manner. PPAR α deficiency significantly upregulated I κ B ζ and pNF- κ Bp65 expression in the CD4(+) T cells of PPAR α (-/-) mice. PPAR α activation by three different PPAR α agonists significantly inhibited I κ B ζ and IL-6 levels. The results suggested that I κ B ζ plays an important role in the pathogenesis of autoimmune myocarditis, and fenofibrate treatment

ameliorates EAM by preventing myocardial inflammation and fibrosis, possibly through the PPAR α /I κ B ζ signaling pathway.

Introduction

Peroxisome proliferator-activated receptor α (PPAR α) is a critical regulator of cardiac lipid metabolism and has a significant effect on various functions, such as glucose and lipid homeostasis, cardiac metabolism substrate conversion, and inflammation and autoimmune disease development (1). PPAR α can also reduce the release of pro-inflammatory factors, inhibit the production of chemokines, and promote T cell differentiation (2). Fenofibrate lowers cholesterol and triglyceride levels and is a widely used as a PPAR α agonist in clinical practice. It has multiple effects on the heart, including prevention of myocardial inflammation, attenuation of isoproterenol-induced acute myocardial ischemic injury, and inhibition of macrophage and T lymphocyte infiltration into the left ventricle (3). Experimental autoimmune myocarditis (EAM) is an autoimmune disease induced by CD4(+) T cells. Histopathology has shown that CD4(+) T cells infiltrate the myocardium in the acute phase, leading to severe myocardial damage and subsequent cardiac fibrosis (4-6). Fenofibrate treatment can alleviate EAM (7,8). It has been previously shown that PPAR α plays a crucial role in T helper 17 (Th17) cell differentiation, and fenofibrate can alleviate EAM (9). Methyl- β -cyclodextrin, a specific cholesterol-depleting agent, also ameliorates EAM by suppressing myocarditis-induced apoptosis (10).

I κ B ζ is the most recently identified member of the I κ B family and interacts with the NF- κ Bp50 subunit to positively regulate the expression of numerous inflammatory factors (11,12). I κ B ζ plays a central role in inflammatory diseases and various autoimmune diseases, making it a key Th17-associated factor (13,14). I κ B ζ mediates the inflammation response to TNF- α and IL-17, and its inhibition mediates toll-like receptor transcriptional responses (15,16). However, the role of I κ B ζ in EAM and the underlying mechanisms remain unclear.

In the present study, the authors investigated whether fenofibrate could regulate the I κ B ζ signaling pathway in rat EAM and explore its possible mechanisms. The results showed that fenofibrate treatment ameliorated EAM by preventing myocardial inflammation and fibrosis. It was also found that I κ B ζ is

Correspondence to: Dr He Chang, Department of Geriatrics, Xiang'an Hospital of Xiamen University, 2000 Xiang'an East Road, Xiamen, Fujian 361100, P.R. China
E-mail: changhe@xmu.edu.cn

Professor Zhi Qi, Department of Basic Medical Sciences, School of Medicine, Xiamen University, 4221 Xiang'an South Road, Xiamen, Fujian 361102, P.R. China
E-mail: qizhi@xmu.edu.cn

Key words: fenofibrate, peroxisome proliferator-activated receptor α , I κ B ζ , NF- κ B, autoimmune myocarditis

a key factor in EAM, and the PPAR α /IkB ζ signaling pathway is involved in the pathogenesis of EAM. This suggested that IkB ζ may be a new molecular target of fenofibrate for treating autoimmune myocarditis.

Materials and methods

Animals. Male Lewis rat aged 6-8 weeks (180-200 g) and PPAR α (-/-) mice (129S4/SvJae) aged 8-10 weeks (20-25 g) were acquired from Beijing Vital River Lab Animal Technology Co., Ltd. and Jackson ImmunoResearch Laboratories, Inc., respectively. Male C57BL/6J mice as PPAR α (+/+) wild-type (WT) mice were purchased from Shanghai SLAC Laboratory Animal Company, Ltd. Animal care and experiments were conducted in accordance with the procedures approved by the Animal Care and Use Committee of Xiamen University (approval no. XMULAC 20220111; Xiamen, China). Animals were provided free access to standard rodent chow and water and were housed in a SPF conditions at 23 $\pm$ 2°C and a relative humidity of 60 $\pm$ 5%. Throughout the studies, all animals were treated in accordance with the guidelines for animal experiments of our institution.

Rat EAM model. EAM model was established as previously described (5,6). Throughout the studies, all the experiment operations were performed under the anesthesia environment to animal welfare consideration. The duration of the experiment was 21 days. On day 0 the rats were anesthetized with 2% isoflurane inhalation and immunized once by subcutaneous injection with a 0.2-ml emulsion containing 1 mg of cardiac myosin and an equal volume of complete Freund's adjuvant in both footpads; the morbidity and survival rate were 100% in rats immunized by this method (4). A total of 24 rats were divided into three groups (n=8 for each group). The control group rats received only complete Freund's adjuvant for immunization. The immunized EAM rats underwent daily oral gavage administration with fenofibrate or a solvent alone from day 14 to day 21 for 7 consecutive days. All the rats' health and behavior were monitored and body weight were measured every 3 days. It was previously reported that rat EAM cardiac inflammation occurred in the acute phase and peaked on day 14 to day 21, characterized by severe heart failure (5,6). On day 21 all the experimental rats were anesthetized with 2% isoflurane inhalation and euthanized through cervical dislocation, the area of the chest were sterilized with 75% alcohol, and an aseptic surgical knife was subsequently used to fully expose the heart. A total of 5 ml blood from the inferior vena cava were drawn out. After dissecting the heart, heart tissues were carefully harvested, washed, and weighted. Body weight (BW) and heart weight (HW) were measured to calculate the ratio of HW/BW.

Histopathological examination. Heart tissues were fixed in 10% formalin at room temperature for 3 days and 4% paraformaldehyde in PBS and embedded in paraffin wax. Sections were cut at 5- μ m thickness for hematoxylin & eosin (H&E) staining and scored macroscopically as follows: i) 0, no inflammation; ii) 1, presence of a small discolored focus; iii) 2, presence of multiple small discolored foci; iv) 3, diffuse discolored areas not exceeding a total of 1/3 of the cardiac

surface; and v) 4, diffuse discolored areas totaling >1/3 of the cardiac surface (5,9,10). Inflammatory cell infiltration was examined under a light microscope, the ratio of the area of inflammatory cell infiltration in each field to the area of the whole field was calculated, and its mean value was used for microscopic scoring as follows: i) 0, no inflammation; ii) 1, <25% of the heart section involved; iii) 2, 25-50%; iv) 3, 50-75%; and v) 4, >75% (5).

Cardiac function assessment. Echocardiography was performed on day 21 for all the experimental rats using a 14-MHz probe (Vivid 7; GE Healthcare). Wall thickness and left ventricular (LV) dimensions (including LV internal dimensions in systole, LV internal dimensions in diastole, and LV posterior wall of diastole), interventricular septal thickness at diastolic (IVS), and heart rate were measured. LV ejection fraction (LVEF) and LV fractional shortening (FS) were assessed as previously described (9,10).

CD4(+) T cell isolation. CD4(+) T cells were isolated and purified from rats with EAM and PPAR α (-/-) mice spleens using microbeads [CD4(+) T cell isolation kit; MiltenyiBiotec, Inc.] as previously described (9). CD4(+) T cells from rats with EAM were induced Th17 cell differentiation and incubated with fenofibrate 20 μ M (Abcam) and PPAR α antagonist MK886 20 μ M (Abcam) for 24 h for reverse transcription-quantitative PCR (RT-qPCR) analyses. CD4(+) T cells from PPAR α (-/-) mice were activated using 1 μ g/ml anti-CD3 (BD Biosciences) bound to plates and 2 μ g/ml anti-CD28 (BD Biosciences) in solution for 3, 6, 12 and 24 h. CD4(+) T cells from PPAR α (+/+) mice were added and incubated with three PPAR α agonists, fenofibrate 100 μ mol/l (Abcam), Wy14643 50 μ mol/l (Abcam) or GW7646 1 μ mol/l (Abcam) for 24 h. Cells were collected for western blotting and RT-qPCR analyses.

Western blot analysis. Cardiac ventricles from the EAM rats and cells were homogenized in a lysis buffer composed of 8 M urea, 1 mM dithiothreitol, 1 mM ethylenediaminetetraacetic acid (EDTA), and 50 mM Tris-HCl at pH 8.0. Following precipitation with trichloro-acetate and sodium deoxycholate, the protein samples were quantified using Lowry's method. SDS-PAGE (12.5%) was used to isolate 10 μ g of proteins, which were subsequently transferred to a PVDF membrane, the membrane was blocked with 5% non-fat milk or 5% BSA (Beijing Solarbio Science & Technology Co., Ltd.) for 1 h at room temperature and incubated overnight at 4°C with antibodies against anti-vimentin (1:1,000; Thermo Fisher Scientific, Inc.), anti-collagen I (1:1,000; Abcam), anti-RAR-related orphan receptor gamma (ROR γ t; 1:1,000; BD Biosciences), anti-IkB ζ (1:1,000; Cell Signaling Technology, Inc.), anti-pNF- κ Bp65 (1:1,000; Thermo Fisher Scientific, Inc.), anti- β -actin (1:2,000; Thermo Fisher Scientific, Inc.), anti-GAPDH (1:2,000; Thermo Fisher Scientific, Inc.) followed by incubation with goat anti-mouse IgG antibody (1:2,000; Cell Signaling Technology, Inc.) or goat anti-rabbit IgG antibody (1:2,000; Cell Signaling Technology, Inc.) at room temperature for 2 h. Membranes were eventually visualized with the ChemiDoc Touch Imaging System (Bio-Rad Laboratories, Inc.). Quantification of the resulting bands was achieved using densitometry software ImageJ (version 1.5.4; National Institutes of Health).

Table I. Echocardiographic parameters.

Parameters	Control	EAM	EAM + Feno
LVEDs (mm)	3.055±0.22	6.291±0.884 ^a	3.904±0.604 ^c
LVEDd (mm)	5.816±0.186	8.591±0.768 ^a	5.753±0.931 ^c
LVPW (mm)	1.347±0.18	1.732±0.475	1.513±0.185
IVS (mm)	1.208±0.137	2.520±0.285 ^a	1.399±0.168
LVEF (%)	77.87±3.823	52.014±6.184 ^b	54.319±8.652 ^c
LVFS (%)	47.327±3.972	27.999±3.807 ^b	30.297±5.603 ^c
Heart rate (time/min)	294±13	458±24 ^b	342±28 ^d

All rats were sacrificed on day 21 (n=8 for each group). EAM vs. control, ^aP<0.01; ^bP<0.001. EAM + Feno vs. EAM, ^cP<0.05; ^dP<0.001. EAM, experimental autoimmune myocarditis; EAM + Feno, EAM and fenofibrate; LVEDs, left ventricular end-systolic internal diameter; LVEDd, left ventricular end-diastolic internal diameter; LVPW, left ventricular posterior wall at diastolic; IVS, interventricular septal thickness at diastolic; LVEF, left ventricular ejection fraction; LVFS, left ventricular fractional shortening.




Figure 1. Feno alleviates rat EAM. (A) Representative hematoxylin & eosin staining of ventricular sections from control, EAM and feno-treated rats with EAM (upper images, 10x10; scale bar, 2 mm and lower images, 10x40; scale bar, 20 μ m). (B) Ratio of heart HW/BW. (C) EAM macroscopic scores. (D) Microscopic scores (inflammatory cell infiltration) (n=8 for each group). EAM vs. control, ***P<0.001 and ****P<0.0001. EAM + Feno vs. EAM, #P<0.05 and ###P<0.001. EAM, experimental autoimmune myocarditis; EAM + Feno, EAM and fenofibrate; HW/BW, heart weight to body weight.

RT-qPCR analysis. Total RNA was extracted from the rat hearts, spleen, and isolated CD4(+) T cells using TRIzol[®] reagent (Invitrogen; Thermo Fisher Scientific, Inc.) in accordance with the manufacturer's protocol. Random primers and reverse transcriptase were employed to synthesize cDNA from 2 μ g of the total RNA using the PrimeScript RT reagent Kit

(Takara Bio, Inc.) according to the manufacturer's protocol. The reaction temperature and time were set according to the same protocol. qPCR was performed using SuperReal PreMix Plus (SYBR Green; Tiangen Biotech Co., Ltd.) on an ABI 7500 Fast Real-Time PCR Detection system (Applied Biosystems; Thermo Fisher Scientific, Inc.). Gene amplification was




Figure 2. Feno inhibits IL-6, TGF- β , IL-23 and ROR γ t expression in the heart, splenocyte and CD4(+) T cells. (A) Heart, (B) splenocytes and (C) CD4(+) T cells were purified from the spleens of rats. Reverse transcription-quantitative PCR was employed to measure the relative transcript levels, with GAPDH serving as the internal reference for normalization (n=5 for each group). EAM vs. control, ***P<0.001. EAM + Feno vs. EAM, *P<0.05; **P<0.01 and ###P<0.001. EAM, experimental autoimmune myocarditis; EAM + Feno, EAM and fenofibrate; IL, interleukin; TGF- β , transforming growth factor beta; ROR γ t, RAR-related orphan receptor gamma t.

performed using specific primer pairs (Table SI). The reaction program was determined as 3 min at 95°C, followed by 40 cycles of denaturation at 95°C for 30 sec, and annealing at 60°C for 60 sec, and extension at 75°C for 60 sec. Comparative analysis of qPCR results by utilizing the $2^{-\Delta\Delta C_q}$ method (17) was conducted to assess the relative levels of these molecules. This analysis was carried out after normalizing the results to GAPDH or β -actin expression.

Cytokine ELISA. The serum protein levels of IL-17 (eBioscience; Thermo Fisher Scientific, Inc.), IFN- γ (R&D Systems), and IL-4 (R&D Systems) in the control, EAM, and fenofibrate-treated EAM rats were measured using rat ELISA kit (Table SII) at day 21.

Immunohistochemistry staining (IHC). Deparaffinized cardiac tissue sections (5 μ m) were heated in EDTA buffer, treated

with 3% H₂O₂ in methanol for 10 min and blocked with 5% normal serum at room temperature for 30 min. These sections were incubated with the primary antibodies anti-ROR γ t (1:100; Abcam), anti-vimentin (1:100; Thermo Fisher Scientific, Inc.), and anti-I κ B ζ (1:100; Cell Signaling Technology, Inc.) at 4°C overnight, followed by secondary antibodies rabbit anti-mouse IgG-HRP (1:5,000; Abcam) or goat anti-rabbit IgG-HRP (1:5,000; Abcam) (Table SII) for 1 h at room temperature. The sections were then visualized with DAB chromogen, counterstained with hematoxylin for 10 sec at room temperature, mounted with an antifade mounting medium (Applygen Technologies, Inc.) and analyzed using an Olympus fluorescent Microscope (IX51; Olympus Corporation) for the immunohistochemical examinations of vimentin, I κ B ζ , and ROR γ t.

Statistical analysis. The results are presented as the mean values with standard deviations. Statistical analysis was




Figure 3. Feno inhibits the expression of Th17-related factors in the heart of rats with EAM. (A-E) TNF- α , IL-1 β , IFN- γ , IL-4 and IL-17. Rat hearts were isolated and examined by reverse transcription-quantitative PCR analysis, with GAPDH serving as the internal reference for normalization (n=5 for each group). (F) Serum levels of IFN γ , IL-4 and IL-17 on day 21 of EAM rats were tested by ELISA (n=5 for each group). EAM vs. control, *P<0.05; **P<0.01 and ***P<0.001. EAM + Feno vs. EAM, #P<0.05 and ###P<0.001. EAM, experimental autoimmune myocarditis; EAM + Feno, EAM and fenofibrate; Th17, T helper 17; TNF- α , tumor necrosis factor alpha; IL, interleukin; IFN- γ , interferon gamma.

conducted using one-way ANOVA, followed by Bonferroni's test for multiple comparisons. Statistical significance was established at P<0.05. All statistical analyses were performed using GraphPad Prism 7.0 software (GraphPad; Dotmatics).

Results

Fenofibrate treatment ameliorates rat EAM. Preliminary studies about the long-term effects of fenofibrate on EAM were conducted using different dosages from day 0 to day 21. The results showed that both 100 mg/kg and 200 mg/kg of fenofibrate ameliorated EAM while the dosage of 200 mg/kg had improved treatment effect compared with 100 mg/kg (Fig. S1). Therefore, fenofibrate was administered at a dosage of 200 mg/kg to evaluate its long-term effects in a previous study conducted by the authors (9). In the present study, fenofibrate (200 mg/kg) was used to evaluate its short-term effect. H&E staining of transverse cardiac ventricle sections revealed severe myocarditis in the rats with EAM, characterized by extensive inflammatory cells infiltration. Fenofibrate treatment significantly reduced inflammatory cells infiltration and mitigated EAM-induced myocardial inflammation (Fig. 1A). EAM in rats caused macroscopic and microscopic alterations with marked inflammatory cells infiltration and necrosis, and fenofibrate treatment ameliorated EAM as evidenced by a decrease in HW/BW, myocardial damage scores and inflammatory cell infiltrate scores (Fig. 1B-D). According to echocardiographic parameters, the rats with EAM showed enlarged LV, thick IVS and decreased LVEF leading to heart failure. Meanwhile, fenofibrate treatment improved the cardiac function as revealed by a reduction in LV end diameter at systole and diastole and an increase in LVEF and FS (Table I).

Fenofibrate inhibits the expression of Th17-related inflammatory cytokines. The experimental rats were euthanized after 21 days. Hearts, splenocytes and CD4(+) T cells were isolated. RT-qPCR quantified IL-6, TGF- β , IL-23 and ROR γ t expression levels, revealing that fenofibrate treatment suppressed the expression of these inflammatory cytokines in the hearts, splenocytes and CD4⁺ T cells of the rats with EAM (Fig. 2A-C). Fenofibrate treatment also significantly reduced the expression of Th17-related factors TNF- α , IL-1 β , IFN γ , IL-4 and IL-17 (Fig. 3A-E). ELISA confirmed that fenofibrate treatment suppressed the serum protein levels of IFN γ , IL-4 and IL-17 in the rats with EAM (Fig. 3F).

Fenofibrate inhibits the expression of fibrosis-associated factors. PPAR α can suppress TGF- β -induced myocardial fibrosis (18). Therefore, the expression of fibrosis-associated factors TGF- β , tissue inhibitors of metalloproteinase (TIMP), fibronectin (FN) and galectin-3 (GAL3), and fibrosis markers vimentin and collagen type I (collagen I) were assessed in the hearts of rats with EAM. Fenofibrate treatment significantly reduced their expression (Fig. 4A-D). IHC and western blot analysis revealed that fenofibrate treatment significantly inhibited the expression of vimentin and collagen I (Fig. 4E and F).

Fenofibrate inhibits IkB ζ expression in the heart of rats with EAM. IkB ζ is critical in inflammatory and autoimmune diseases, and ROR γ t is essential for Th17 differentiation in EAM (19). IHC staining of transverse cardiac sections revealed the extensive infiltration of IkB ζ -positive and ROR γ t-positive cells in the myocardium at EAM lesions, which were significantly reduced by fenofibrate treatment (Fig. 5A). Western




Figure 4. Fenofibrate inhibits the expression of fibrosis-associated factors in the heart of rats with EAM. mRNA levels of (A) TGF- β , (B) TIMPs, (C) FN and (D) GAL3 were determined using reverse transcription-quantitative PCR analysis with GAPDH serving as the internal reference for normalization ($n=5$ for each group). (E) Vimentin, representative immunohistochemistry image of ventricular sections. Scale bar, 20 μm . (F) Vimentin and collagen I protein expression in the heart examined by western blot analysis. β -actin serving as the internal reference ($n=3$). EAM vs. control, * $P<0.05$, ** $P<0.01$ and *** $P<0.001$. EAM + Feno vs. EAM, # $P<0.05$, ## $P<0.01$ and ### $P<0.001$. EAM, experimental autoimmune myocarditis; EAM + Feno, EAM and fenofibrate; TGF- β , transforming growth factor beta; TIMPs, tissue inhibitors of metalloproteinase; FN, fibronectin; GAL3, galectin 3.

blot analysis confirmed that fenofibrate suppressed I κ B ζ and ROR γ t expression (Fig. 5B).

Fenofibrate inhibits I κ B ζ expression in the CD4(+) T cells from rats with EAM. The CD4(+) T cells were purified from the spleen of rats with EAM and induced Th17 cell differentiation using recombinant (r) IL-6 and rTGF- β . RT-qPCR analysis demonstrated that fenofibrate, at concentrations ranging from 0 to 20 μM , dose-dependently decreased I κ B ζ (Fig. 6A). MK886, which acts as a PPAR α antagonist, displayed a dose-dependent reversal of these effects (Fig. 6B). These findings indicated that I κ B ζ is involved in Th17 differentiation during EAM development.

PPAR α deficiency upregulates I κ B ζ and IL-6 expression. Spleen-derived CD4(+) T cells from PPAR α (-/-) mice were exposed to anti-CD3 (1 $\mu\text{g}/\text{ml}$) and anti-CD28 (2 $\mu\text{g}/\text{ml}$) monoclonal antibodies and incubated for periods of 3, 6, 12 and 24 h. Western blot analysis demonstrated that I κ B ζ and pNF- κ Bp65 levels were significantly increased in the CD4(+) T cells of PPAR α (-/-) mice compared with that in PPAR α (+/+) mice (Fig. 7A and B). IL-6 is regulated by I κ B- ζ and NF- κ B activation. I κ B ζ controls Th17 differentiation by ROR γ t and IL-23 activation (20). It was observed that the levels of IL-6 and ROR γ t were higher at all tested time

points in the CD4(+) T cells of PPAR α (-/-) mice. However, the IL-23 expression did not change significantly (Fig. 7C-E).

Activation of PPAR α inhibits I κ B ζ and IL-6 expression. CD4(+) T cells from PPAR α (+/+) mice were added and incubated with three PPAR α agonists (fenofibrate 100 $\mu\text{mol}/\text{l}$, Wy14643 50 $\mu\text{mol}/\text{l}$, GW7646 1 $\mu\text{mol}/\text{l}$) to study whether PPAR α affects IL-6 expression. Western blot analysis revealed that these PPAR α agonists suppressed I κ B ζ expression (Fig. 8A). ELISA results also indicated that these PPAR α agonists significantly reduced IL-6 secretion (Fig. 8B). All these findings indicated that IL-6 mediates Th17 differentiation via the PPAR α /I κ B ζ pathway.

Discussion

Myocarditis is an inflammatory cardiomyopathy that can lead to acute heart failure and dilated cardiomyopathy and currently has no specific treatment. EAM in rats is similar to human giant cell myocarditis, and recurrent forms can lead to dilated cardiomyopathy (4-6). In a previous study conducted by the authors, it was found that the peak of EAM cardiac inflammation occurred on days 14 to 21 and was characterized by infiltration of the CD4(+) T cells from the spleen into the myocardium. It was also revealed that Th17




Figure 5. Feno inhibits IκBζ and RORγt expression in the heart of rats with EAM. (A) Representative immunohistochemistry image of IκBζ and RORγt in the ventricular sections. Scale bar, 20 μm. (B) Western blot and statistical analysis. GAPDH serving as the internal reference (n=3). EAM vs. control, ***P<0.001. EAM + Feno vs. EAM, #P<0.05 and ###P<0.001. EAM, experimental autoimmune myocarditis; EAM + Feno, EAM and fenofibrate; RORγt, RAR-related orphan receptor gamma t.




Figure 6. Feno inhibits IκBζ expression in the CD4(+) T cells from rats with EAM. (A and B) IκBζ expression in the CD4(+) T cells extracted from rats with EAM examined by reverse transcription-quantitative PCR. β-actin serving as the internal reference (n=5 for each group). CD4(+) T cells were obtained from EAM rat spleens and incubated with rIL-6 (40 ng/ml) and rTGF-β (10 ng/ml), and treated with either fenofibrate alone or a combination of fenofibrate and the PPARα antagonist MK886 for 72 h. **P<0.01, #P<0.05, ##P<0.01 and ###P<0.001. EAM, experimental autoimmune myocarditis; Feno, fenofibrate; r-, recombinant.

cells play an important role in the development of EAM and that fenofibrate can improve EAM by inhibiting Th17 differentiation (9).

In the present study, it was demonstrated that short-term administration of fenofibrate alleviated EAM. The expression levels of typical Th17-related factors including IL-6, TGF-β




Figure 7. Time course expression of IκBζ-related factors in the CD4(+) T cells from PPARα(-/-) mice. (A and B) IκBζ and pNF-κBp65 protein levels determined by western blotting and statistical analysis. CD4(+) T cells were purified from the spleens of PPARα(+/-) and PPARα(-/-) mice, and incubated with anti-CD3 (1 μg/ml) and anti-CD28 (2 μg/ml) antibodies for 12 and 24 h. GAPDH serving as the internal reference (n=5). *P<0.05 and **P<0.01. (C-E) Reverse transcription-quantitative PCR analysis was employed to measure the relative transcript levels of IL-6, RORγt and IL-23 in the CD4(+) T cells. PPARα, peroxisome proliferator-activated receptor α; RORγt, RAR-related orphan receptor gamma t; IL, interleukin.




Figure 8. Activation of PPARα by three different PPARα agonists inhibits IκBζ and IL-6 expression. (A) IκBζ expression in the CD4(+) T cells determined by western blotting and statistical analysis; GAPDH was used as a control for internal comparison (n=5 for each group). (B) Production of IL-6 in the culture supernatants. Splenic CD4(+) T cells were isolated from PPARα wild type mice and incubated with three different PPARα agonists for 24 h (Feno 100 μmol/l, Wy14643 50 μmol/l and GW7646 1 μmol/l), and IL-6 levels in the culture supernatants were tested by ELISA analysis. (n=5 for each group). **P<0.01 and ***P<0.001. IL-6, interleukin-6; Feno, fenofibrate.

and IL-23 in the heart and splenic CD4(+) T cells from rats with EAM were significantly increased. The proinflammatory cytokines TNF-α, IL-1β, IFNγ and IL-17 were also significantly upregulated. The levels of these factors were reduced significantly by fenofibrate treatment. These findings suggested that fenofibrate attenuates cardiac inflammation and exerts anti-inflammatory effects by suppressing Th17-related inflammatory cytokines secretion. Thus, this provides a more comprehensive understanding of fenofibrate's therapeutic potential on autoimmune myocarditis. This result is consistent with the effect of fenofibrate on other inflammatory diseases (21,22).

In the pathological process of EAM, the infiltration of inflammatory cells into myocardium leads to myocardial

fibrosis. Although cardiac fibrosis is beneficial for enhancing the structural stability of the heart, it can also lead to heart structure remodeling and impaired cardiac function. Therefore, improving cardiac fibrosis may help avoid further deterioration caused by EAM. The expression levels of types I, III and IV collagen, FN, matrix metalloproteinases, and TIMP are increased during the progression of myocardial fibrosis (23). PPARα activation can inhibit the TGF-β-induced cardiac fibrosis pathway. Fenofibrate can also reduce myocardial inflammation and collagen deposition by modulating the PPARα pathway and therefore, fenofibrate can relieve cardiac fibrosis and reverse cardiac dysfunction (24,25). In the present study, the expression levels of fibrosis-related factors TIMP, FN and GAL3 as well as fibrosis markers vimentin and collagen I, were detected by western blotting and RT-qPCR. Consistent with previous studies, the results also showed that these fibrosis-related factors were significantly upregulated in the hearts of EAM rats, and that fenofibrate treatment inhibited the levels of these factors and improved cardiac fibrosis.

IκBζ is a key Th17-related factor that plays an important role in the development of autoimmune diseases such as psoriasis (26,27). In mice with IκBζ-deficiency, the development of psoriasis induced by IL-17, IL-23 and imiquimod was significantly inhibited (28,29). IL-17 and its family members, produced by CD4(+) T cells and various innate immune cells, are implicated in the pathogenesis of EAM (30); however, its role in EAM remains to be investigated. The results of the present study indicated that fenofibrate significantly inhibited the expression of IκBζ in the heart of rats with EAM. IL-6 plays an important role in EAM initiation through RORγt-mediated Th17 differentiation. The present study also revealed that fenofibrate treatment significantly inhibited the upregulation of RORγt expression in the hearts of rats with EAM. These findings suggested that IκBζ is a molecular target involved in Th17 differentiation in autoimmune myocarditis.

By stimulating CD4(+) T cells isolated from the spleen of EAM rats to induce Th17 cell differentiation, it was found that PPAR α agonist fenofibrate upregulated I κ B ζ expression in a dose-dependent manner. This effect was reversed by PPAR α antagonist MK886 in a dose-dependent manner. These results suggested that PPAR α promotes Th17 cell differentiation through the I κ B ζ signaling pathway.

To further explore the potential mechanism of fenofibrate in treating EAM, CD4(+) T cells isolated from the spleen of PPAR α -/- mice were activated and it was revealed that the mRNA and protein levels of I κ B ζ were upregulated in activated PPAR α (-/-) mice CD4(+) T cells compared with those in PPAR α (+/+) mice. I κ B ζ interacts with the NF- κ Bp50 subunit to positively regulate the expression of pro-inflammatory cytokines such as IL-6, IL-12 and CCL2 (31). The activation of NF- κ B and stimulation of Toll-like receptor ligands and IL-1 β are required to induce I κ B ζ expression (32). Previously, several studies have reported that I κ B ζ deficiency in LPS-induced macrophages prevents the production of the key pro-inflammatory cytokine IL-6 (33). In the absence of I κ B ζ , T cells exhibit serious defects in the development of Th17 cells (34). I κ B ζ is induced by IL-17R and collaboratively regulates IL-17 expression with ROR γ t. IL-6, which is a key factor in inducing Th17 cell differentiation, activates STAT3 and increases the expression of ROR γ t (35). Th17 cell differentiation and IL-6 secretion can be inhibited by PPAR α agonists (36).

In the present study, the correlation between PPAR α and I κ B ζ was confirmed. *In vivo*, it was shown that PPAR α activation inhibited I κ B ζ expression on EAM. *In vitro*, it was revealed that PPAR α deficiency upregulated I κ B ζ and IL-6 expression in the CD4(+) T cells from PPAR α (-/-) mice. By conducting Chromatin immunoprecipitation (ChIP) assays, Muromoto *et al* (34,37) found that two different I κ B- ζ promoter regions and STAT3 constitutively binds to the genomic promoter region of I κ B- ζ TSS1. Luciferase reporter assays of the I κ B- ζ promoter activity, revealed that catalytic activity of TYK2 and its substrate transcription factor STAT3, is required for I κ B- ζ promoter activity. The limitation of the present study is the lack of direct molecular evidence of PPAR α binding or transcriptional modulation with I κ B- ζ . In a future study, the authors will provide further molecular evidences and focus on specific mechanisms regarding how PPAR α and fenofibrate interacts with I κ B ζ and whether PPAR α directly interacts with or binds the promoter regions of I κ B ζ gene. Luciferase reporter assays will be performed to evaluate the effects of fenofibrate on the PPAR α /I κ B ζ pathway activation, and the promoter activity of I κ B ζ and mRNA stability will be examined using ChIP assay. Further mechanistic studies will be continued.

In summary, it was demonstrated that I κ B ζ contributes to the pathogenesis of autoimmune myocarditis, and fenofibrate treatment ameliorates EAM by preventing myocardial inflammation and fibrosis possibly through the PPAR α /I κ B ζ signaling pathway. Thus, I κ B ζ may be a new molecular target for fenofibrate treatment in autoimmune myocarditis.

Acknowledgements

Not applicable.

Funding

The present study was supported by the Natural Science Foundation of Fujian Province of China (grant no. 2021J01014), the Scientific Research Foundation for Advanced Talents, Xiang'an Hospital of Xiamen University, Fujian, China (grant no. PM201809170018).

Availability of data and materials

The data generated in this study may be requested from the corresponding author.

Authors' contributions

HC and ZQ designed the study, wrote and revised the manuscript. YWa and YWu conducted the experiments to collected data. SLS analyzed the data. HC and ZQ confirm the authenticity of all the raw data. All authors read and approved the final version of the manuscript.

Ethics approval and consent to participate

Animal care and experiments were conducted in accordance with the procedures approved by the Ethics Committee of Animal Care and Use of Xiamen University (approval no. XMULAC 20220111; Xiamen, China). All animals were treated in accordance with the principles of The Declaration of Helsinki and welfare considerations were taken to minimize the number of animals used and their suffering.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Manoharan I, Suryawanshi A, Hong Y, Ranganathan P, Shanmugam A, Ahmad S, Swafford D, Manicassamy B, Ramesh G, Koni PA, *et al*: Homeostatic PPAR α signaling limits inflammatory responses to commensal microbiota in the intestine. *J Immunol* 196: 4739-4749, 2016.
2. Riaz F, Wei P and Pan F: PPARs at the crossroads of T cell differentiation and type 1 diabetes. *Front Immunol* 14: 1292238, 2023.
3. Balakumar P, Rohilla A and Mahadevan N: Pleiotropic actions of fenofibrate on the heart. *Pharmacol Res* 63: 8-12, 2011.
4. Kodama M, Matsumoto Y, Fujiwara M, Masani F, Izumi T and Shibata A: A novel experimental model of giant cell myocarditis induced in rats by immunization with cardiac myosin fraction. *Clin Immunol Immunopathol* 57: 250-262, 1990.
5. Zhong C, Wu Y, Chang H, Liu C, Zhou L, Zou J and Qi Z: Effect of PKC inhibitor on experimental autoimmune myocarditis in Lewis rats. *Oncotarget* 8: 54187-54198, 2017.
6. Zhong C, Chang H, Wu Y, Zhou L, Wang Y, Wang M, Wu P, Qi Z and Zou J: Up-regulated Cx43 phosphorylation at Ser368 prolongs QRS duration in myocarditis. *J Cell Mol Med* 22: 3537-3547, 2018.
7. Cheng H, Xi Y, Chi X, Wu Y and Liu G: Fenofibrate treatment of rats with experimental autoimmune myocarditis by alleviating Treg/Th17 disorder. *Cent Eur J Immunol* 41: 64-70, 2016.

8. Maruyama S, Kato K, Kodama M, Hirano S, Fuse K, Nakagawa O, Nakazawa M, Miida T, Yamamoto T, Watanabe K and Aizawa Y: Fenofibrate, a peroxisome proliferator-activated receptor α activator, suppresses experimental autoimmune myocarditis by stimulating the interleukin-10 pathway in rats. *J Atheroscler Thromb* 9: 87-92, 2002.
9. Chang H, Zhao F, Xie X, Liao Y, Song Y, Liu C, Wu Y, Wang Y, Liu D, Wang Y, *et al*: PPAR α suppresses Th17 cell differentiation through IL-6/STAT3/ROR γ t pathway in experimental autoimmune myocarditis. *Exp Cell Res* 375: 22-30, 2019.
10. Chang H, Wang Y, Wu Y, Ma P, Song Y, Liu C, Ye Y, Qi JH and Qi Z: Cardiac apoptosis caused by elevated cholesterol level in experimental autoimmune myocarditis. *Exp Cell Res* 395: 112169, 2020.
11. Feng Y, Chen Z, Xu Y, Han Y, Jia X, Wang Z, Zhang N and Lv W: The central inflammatory regulator I κ B ζ : Induction, regulation and physiological functions. *Front Immunol* 14: 1188253, 2023.
12. Hövelmeyer N, Schmidt-Supprian M and Ohnmacht C: NF- κ B in control of regulatory T cell development, identity, and function. *J Mol Med (Berl)* 100: 985-995, 2022.
13. Kakiuchi N, Yoshida K, Uchino M, Kihara T, Akaki K, Inoue Y, Kawada K, Nagayama S, Yokoyama A, Yamamoto S, *et al*: Frequent mutations that converge on the NFKBIZ pathway in ulcerative colitis. *Nature* 577: 260-265, 2020.
14. Choi MC, Maruyama T, Chun CH and Park Y: Alleviation of murine osteoarthritis by cartilage-specific deletion of I κ B ζ . *Arthritis Rheumatol* 70: 1440-1449, 2018.
15. Bambouskova M, Gorvel L, Lampropoulou V, Sergushichev A, Loginicheva E, Johnson K, Korenfeld D, Mathyer ME, Kim H, Huang LH, *et al*: Electrophilic properties of itaconate and derivatives regulate the I κ B ζ -ATF3 inflammatory axis. *Nature* 556: 501-504, 2018.
16. Slowikowski K, Nguyen HN, Noss EH, Simmons DP, Mizoguchi F, Watts GFM, Gurish MF, Brenner MB and Raychaudhuri S: CUX1 and I κ B ζ (NFKBIZ) mediate the synergistic inflammatory response to TNF and IL-17A in stromal fibroblasts. *Proc Natl Acad Sci USA* 117: 5532-5541, 2020.
17. Rao X, Huang X, Zhou Z and Lin X: An improvement of the 2⁻(-delta delta CT) method for quantitative real-time polymerase chain reaction data analysis. *Biostat Bioinforma Biomath* 3: 71-85, 2013.
18. Bansal T, Chatterjee E, Singh J, Ray A, Kundu B, Thankamani V, Sengupta S and Sarkar S: Arjunolic acid, a peroxisome proliferator-activated receptor α agonist, regresses cardiac fibrosis by inhibiting non-canonical TGF- β signaling. *J Biol Chem* 292: 16440-16462, 2017.
19. Yamazaki S: The nuclear NF- κ B regulator I κ B ζ : Updates on its molecular functions and pathophysiological roles. *Cells* 13: 1467, 2024.
20. Hawkes JE, Yan BY, Chan TC and Krueger JG: Discovery of the IL-23/IL-17 signaling pathway and the treatment of psoriasis. *J Immunol* 201: 1605-1613, 2018.
21. Park A and Heo TH: IL-17A-targeting fenofibrate attenuates inflammation in psoriasis by inducing autophagy. *Life Sci* 326: 121755, 2023.
22. Elaidy SM, Essawy SS, Hussain MA, El-Kherbetawy MK and Hamed ER: Modulation of the IL-23/IL-17 axis by fenofibrate ameliorates the ovalbumin/lipopolysaccharide-induced airway inflammation and bronchial asthma in rats. *Naunyn Schmiedeberg Arch Pharmacol* 391: 309-321, 2018.
23. Cao JW, Duan SY, Zhang HX, Chen Y and Guo M: Zinc deficiency promoted fibrosis via ROS and TIMP/MMPs in the myocardium of mice. *Biol Trace Elem Res* 196: 145-152, 2020.
24. Zhang Y, Ji H, Qiao O, Li Z, Pecoraro L, Zhang X, Han X, Wang W, Zhang X, Man S, *et al*: Nanoparticle conjugation of ginsenoside Rb3 inhibits myocardial fibrosis by regulating PPAR α pathway. *Biomed Pharmacother* 139: 111630, 2021.
25. Qiu Z, Zhao Y, Tao T, Guo W, Liu R, Huang J and Xu G: Activation of PPAR α ameliorates cardiac fibrosis in Dsg2-deficient arrhythmogenic cardiomyopathy. *Cells* 11: 3184, 2022.
26. Bertelsen T, Ljungberg C, Litman T, Huppertz C, Hennze R, Rønholdt K, Iversen L and Johansen C: I κ B ζ is a key player in the antipsoriatic effects of secukinumab. *J Allergy Clin Immunol* 145: 379-90, 2020.
27. Gautam P, Maenner S, Cailotto F, Reboul P, Labialle S, Jouzeau JY, Bourgaud F and Moulin D: Emerging role of I κ B ζ in inflammation: Emphasis on psoriasis. *Clin Transl Med* 12: e1032, 2022.
28. Bertelsen T, Iversen L and Johansen C: The human IL-17A/F heterodimer regulates psoriasis-associated genes through I κ B ζ . *Exp Dermatol* 27: 1048-1052, 2018.
29. Mok BR, Kim AR, Baek SH, Ahn JH, Seok SH, Shin JU and Kim DH: PFN1 prevents psoriasis pathogenesis through I κ B ζ regulation. *J Invest Dermatol* 142: 2455-2463.e9, 2022.
30. Mills KHG: IL-17 and IL-17-producing cells in protection versus pathology. *Nat Rev Immunol* 23: 38-54, 2023.
31. Ohto-Ozaki H, Hayakawa M, Kamoshita N, Maruyama T, Tominaga SI and Ohmori T: Induction of I κ B ζ augments cytokine and chemokine production by IL-33 in mast cells. *J Immunol* 204: 2033-2042, 2020.
32. Yamamoto M, Yamazaki S, Uematsu S, Sato S, Hemmi H, Hoshino K, Kaisho T, Kuwata H, Takeuchi O, Takeshige K, *et al*: Regulation of Toll/IL-1-receptor-mediated gene expression by the inducible nuclear protein IkappaBzeta. *Nature* 430: 218-222, 2004.
33. Lorscheid S, Müller A, Löffler J, Resch C, Bucher P, Kurschus FC, Waismann A, Schäkel K, Hailfinger S, Schulze-Osthoff K and Kramer D: Keratinocyte-derived I κ B ζ drives psoriasis and associated systemic inflammation. *JCI Insight* 4: e130835, 2019.
34. Muromoto R, Sato A, Komori Y, Nariya K, Kitai Y, Kashiwakura JI and Matsuda T: Regulation of NFKBIZ gene promoter activity by STAT3, C/EBP β , and STAT1. *Biochem Biophys Res Commun* 613: 61-66, 2022.
35. Wang J, Liu T, Chen X, Jin Q, Chen Y, Zhang L, Han Z, Chen D, Li Y, Lv Q and Xie M: Bazedoxifene regulates Th17 immune response to ameliorate experimental autoimmune myocarditis via inhibition of STAT3 activation. *Front Pharmacol* 11: 613160, 2021.
36. Zhou Z, Sun W, Liang Y, Gao Y, Kong W, Guan Y, Feng J and Wang X: Fenofibrate inhibited the differentiation of T helper 17 cells in vitro. *PPAR Res* 2012: 145654, 2012.
37. Muromoto R, Tawa K, Ohgakiuchi Y, Sato A, Saino Y, Hirashima K, Minoguchi H, Kitai Y, Kashiwakura JI, Shimoda K, *et al*: I κ B- ζ expression requires both TYK2/STAT3 activity and IL-17-regulated mRNA stabilization. *Immunohorizons* 3: 172-185, 2019.



Copyright © 2025 Chang et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.