

Genetic and epigenetic alterations of *RIZ1* and the correlation to clinicopathological parameters in liver fluke-related cholangiocarcinoma

PRASONG KHAENAM^{1,2,5}, PATCHAREE JEARANA KOON^{2,5}, CHAWALIT PAIROJKUL^{3,5},
VAJARABHONGSA BHUDHISAWASDI^{4,5} and TEMDUANG LIMPAIBOON^{2,5}

¹Department of Biomedical Sciences, Graduate School, ²Centre for Research and Development of Medical Diagnostic Laboratories, Faculty of Associated Medical Sciences, Departments of ³Pathology, ⁴Surgery, and ⁵Liver Fluke and Cholangiocarcinoma Research Center, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand

Received October 15, 2009; Accepted December 22, 2009

DOI: 10.3892/etm_00000060

Abstract. The retinoblastoma interacting zinc finger (*RIZ1*) gene is adjacent to DIS228 where microsatellite instability has been associated with poor patient survival in liver fluke-associated cholangiocarcinoma (CCA). An understanding of the molecular mechanisms underlying the carcinogenesis and pathogenesis of CCA is necessary to improve patient survival. Therefore, we determined the genetic and epigenetic alterations of *RIZ1* in 81 CCA samples and 69 matched non-tumor tissues. Methylation was found in 31 of 81 (38%) tumor samples and in 5 of 69 (7%) matched non-tumor tissues. Frameshift mutations (2 of 81) and loss of heterozygosity (LOH) (14 of 81) were not common. Statistical analysis found no significant correlation between *RIZ1* alterations and clinicopathological features, but RIZPro704 LOH was associated with patient survival in the multivariate analysis. *RIZ1* hypermethylation may be one of the crucial molecular events contributing to cholangiocarcinogenesis, and RIZPro704 LOH may adversely impact patient survival. The biological function of *RIZ1* in CCA should be further investigated in order to verify its potential role in regulating this cancer.

Introduction

Cholangiocarcinoma (CCA), a malignancy of the biliary epithelium, is the major type of liver cancer found in north-east Thailand (1). The high incidence of CCA in this region is strongly associated with a high prevalence of liver fluke (*Opisthorchis viverrini*) infection. Chronic irritation and inflammation caused by liver fluke infection are major factors contributing to the carcinogenesis and pathogenesis of CCA (2). Surgical resection is currently the most successful and accessible therapeutic method for CCA patients but is associated with poor survival. Hence, insights into the molecular mechanisms of carcinogenesis and pathogenesis are necessary for coping with this disease.

Our previous study on fine mapping at 1p36-pter revealed a significant association of microsatellite instability (MSI) at DIS228 with poor survival in CCA patients (3). DIS228 is adjacent to the gene, *retinoblastoma interacting zinc finger (RIZ)* (4). There are two isoforms of *RIZ*, *RIZ1* and *RIZ2*, which are encoded by different promoters (5). Their amino acid sequences are almost identical except for the presence of an N-terminal PR (PRDI-BF1 and *RIZ*) domain in *RIZ1* resulting in a difference in biological function. An important function of the PR domain is histone methyltransferase activity which catalyzes methylation at lysine 9 of histone H3 leading to repression of transcription (6). In previous studies, expression of *RIZ1* was found to be decreased in several types of human cancers (7,8), whereas *RIZ2* was uniformly expressed in all of the examined cases (7-10), suggesting a tumor-suppressive activity of *RIZ1* that harbors the PR domain and an oncogenic activity of *RIZ2* that lacks the PR domain. Moreover, it was demonstrated that *RIZ1*-knockout mice are tumor-prone (11), while adenovirus-mediated *RIZ1* expression caused G2-M cell cycle arrest and/or apoptosis in breast, liver and MSI⁺ colon cancer cells (8-10). *RIZ1* was also found to regulate the expression of IGF-1 resulting in a reduction in cell proliferation and an induction of apoptosis (12).

Several studies have demonstrated that *RIZ1* is a downstream effector of the estrogen receptor (ER) pathway (13,14),

Correspondence to: Dr Temduang Limpai boon, Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen 40002, Thailand
E-mail: temduang@kku.ac.th

Abbreviations: *RIZ*, the retinoblastoma interacting zinc finger gene; LOH, loss of heterozygosity; CCA, cholangiocarcinoma; Pro704, proline residue 704 of *RIZ1*; ER, estrogen receptor

Key words: retinoblastoma interacting zinc finger, cholangiocarcinoma, promoter hypermethylation, frameshift mutation, loss of heterozygosity

Table I. Primer sequences, annealing temperature (T_m) and product size.

Methods	Sequences	T_m ($^{\circ}$ C)	Product size (bp)	Reference no.
MSP				
Methylation	F: GTGGTGGTTATTGGGCGACGGC R: GCTATTTTCGCCGACCCCGACG	68	177	32
Unmethylation	F: TGGTGGTTATTGGGTGATGGT R: ACTATTTACCAACCCCAACA	64	175	32
LOH				
RIZCA	F: GGTGAAAACACTGAAATTCGAAAACACTG R: CAGAGCATAGTTGTCAATTTGTCT	58	~207	22
RIZPro704	F: CCCAAGATAAACTAACTCCT R: ACTCCATGCTGGTGAGTC	58	~266	22
Frameshift mutation				
RIZA8	F: GAGCTCAGCAAAATGTCGTC R: CAAGTCGGCCTTCTGCTTTG	62	116	23
RIZA9	F: TCTCACATCTGCCCTTACTG R: GTGATGAGTGCCACCTTTC	62	144	23

and its expression is decreased after estradiol treatment (14,15). In the absence of estradiol (E2), biological active estrogen, *RIZ1* was found to bind directly to the DNA adjacent to the promoter region of ER target genes and to inhibit the transcription of these genes by methylating lysine 9 of histone H3 (14). The presence of E2 changes the role of *RIZ1* from being a histone methyltransferase to an ER coactivator thus enhancing the maximum response to E2 (14). In addition, the ER signaling pathway can be activated by either estrogen or the growth factor signaling pathway such as IGF-1 (16).

Alterations of *RIZ1* through both genetic and epigenetic mechanisms have been reported (17,18). Epigenetic inactivation by promoter hypermethylation is the most common mechanism leading to decreased expression of this gene in many types of cancers (19,20). As for genetic alterations, the majority are frameshift mutations at polyadenosine tracts, A8 and A9, located at the PR binding domain (21). The second most common genetic defect of *RIZ1* in many types of cancer is loss of heterozygosity (LOH) (17,18,22). Other types of mutations are rare (21,23). The purpose of this study was to investigate the genetic and epigenetic defects of *RIZ1* in CCA samples. Associations between *RIZ1* alterations and clinicopathological data were analyzed. Univariate and multivariate Cox regression were used for survival analysis.

Materials and methods

Patients. Informed consent was obtained from each patient according to the guidelines of the Ethics Committee of Khon Kaen University (HE500634). Blood and liver resection samples were obtained from 81 intrahepatic CCA patients undergoing surgery at Srinagarind Hospital, Faculty of Medicine, Khon Kaen University. DNA was extracted from leukocytes, frozen tissues and microdissected tissues as described previously (3,24). DNA samples obtained from

frozen liver tissues were used for methylation analysis, and leukocyte and microdissected DNA samples were used for genetic studies including intragenic allelic alteration and frameshift mutation.

Primers. Primer sequences and annealing temperatures used for the analysis of the methylation status, intragenic allelic loss, MSI and frameshift mutations are listed in Table I. Forward strands of primer sets for frameshift mutation and intragenic allelic alteration analyses were labeled at the 5'-end with fluorescein dye 4,7,2',4',5',7'-hexachloro-6-carboxyfluorescein (HEX) (Bio Basic Inc., Canada).

Methylation-specific PCR (MSP). After bisulfite modification, DNA derived from 81 tumor and 69 matched non-tumor tissues of CCA patients were analyzed for *RIZ1* promoter methylation using MSP as described previously (18,24,25). The concentration of $MgCl_2$ used was 5 mM and the PCR reaction was hot-started at 95 $^{\circ}$ C for 5 min before addition of 1.5 units of Taq polymerase. Human placental DNA treated with *SssI* methylase (New England Biolabs, Ipswich, MA, USA) and human leukocyte DNA served as positive controls for the methylated and unmethylated reactions, respectively.

Intragenic allelic alteration and frameshift mutation analysis. LOH and MSI were determined as described previously (22,26). Markers included RIZCA and RIZPro704 located at the intron preceding exon 5 and amino acid residue 704 (Pro704) in exon 8, respectively. LOH was determined for both RIZCA and RIZPro704, and MSI was determined for RIZCA. Frameshift mutations were analyzed by PCR amplification of the repeated sequences in the coding regions (27). Primer sequences of A8 and A9 tracts were obtained from a previous report (23). Genetic alterations were analyzed using the GS-3000 gel scan fragment auto analyzer (Corbett Research, Australia).




Figure 1. Methylation status of *RIZ1* determined by MSP in representative CCA cases. Case no. 87 was methylated strongly in tumor (T) but weakly in non-tumor (N) tissue, whereas case no. 82 was unmethylated. M and U lanes indicate methylated and unmethylated products, respectively.




Figure 2. Intragenic allelic losses at the RIZCA and RIZPro704 markers. Arrowheads indicate LOH. RIZCA is a microsatellite marker. The lower band found with the RIZPro704 marker resulted from a deletion polymorphism at amino acid proline residue 704 (Pro704). L, leukocyte DNA; T, tumor DNA.

Statistical analysis. Clinicopathological features of the CCA patients including age, gender, tumor stages, histological types, blood vessel invasion, nerve invasion and lymphatic invasion were analyzed for correlations with *RIZ1* alterations using the Chi-square test. Survival was assessed using the Kaplan-Meier log-rank method and Cox regression. All variables shown to be significant ($P < 0.150$) in the univariate analyses were entered into a multivariate model using Cox's proportional hazards model in a backward stepwise manner and the log-likelihood ratio approach. Statistical analyses were performed using SPSS for Windows, version 15 (SPSS, Inc., Chicago, IL, USA). Two-sided values of $P < 0.05$ were considered statistically significant.

Results

***RIZ1* promoter hypermethylation and intragenic alteration in CCA patients.** The frequency of *RIZ1* promoter hypermethylation determined using MSP in 81 tumors and 69 matched non-tumor specimens from CCA patients was 38 (31 of 81) and 7% (5 of 69), respectively ($P = 0.006$). DNA methylation was found in non-tumor samples only when its matched tumor sample also showed methylation. Representative results concerning the determination of *RIZ1* methylation are shown in Fig. 1.

LOH was observed in 14 of 81 (17%) CCA cases, comprising 4 of 56 (7%) at RIZCA and 10 of 52 (19%) at RIZPro704 (representative results in Fig. 2). LOH at RIZPro704 and LOH at RIZCA were significantly independent ($P = 0.029$). Frameshift mutations were found only at the A9 tract in 2 (2.5%) cases (Fig. 3). MSI at RIZCA was found in 8 (10%) cases.




Figure 3. Frameshift mutations at A8 and A9 tracts. None of the cases showed a frameshift mutation at A8 tract but a mutation was detected in two cases at A9 tract (arrowheads).




Figure 4. Venn-Euler diagram of the distribution of *RIZ1* alterations in 81 CCA cases. There were 31 cases with methylation-positivity (Met) which included 7 cases with methylation-positivity alone, 2 cases with a frameshift mutation (FS), 3 cases with LOH and 19 non-informative (NI) cases in the LOH analysis. There were 50 unmethylated cases which included 11 cases with LOH, 24 NI cases and 15 cases without any alteration.

As shown in the Venn-Euler diagram (Fig. 4), a *RIZ1* alteration was found in 42 (52%) cases. A simultaneous alteration was found in 5 (6%) cases, 3 of which were methylated with LOH and 2 methylated with a frameshift mutation. *RIZ1* methylation alone was found in 26 (32%) cases. LOH alone was found in 11 (14%) cases; 39 (48%) cases had no LOH or DNA methylation.

Statistical analysis. *RIZ1* alterations were not correlated to patient clinicopathological features. Correlations between post-operative survival and *RIZ1* alterations were evaluated using the univariate Kaplan-Meier log-rank test and multivariate Cox regression (Table II). Histological type was the only factor found to be correlated with patient survival in the Kaplan-Meier analysis ($P = 0.042$). Adenosquamous and squamous carcinomas, defined as 'others', were poor prognostic factors, while papillary adenocarcinoma was associated with a better patient survival. Other variables were not correlated with patient survival. However, only variables presenting $P < 0.150$ in the univariate analysis were included in the multivariate Cox regression analysis. LOH at RIZPro704 was an independent prognostic factor with a hazard ratio 2.77 (95% CI, 1.12-2.84; $P = 0.027$).

Discussion

DNA methylation was detected in matched non-tumor samples (7%) suggesting that methylation occurs early in carcinogenesis. This finding corroborates that of a previous study which found *RIZ1* methylation in precancerous lesions (17). Since one study involving prostate cancer showed that *RIZ1* methylation

Table II. Univariate and multivariate survival analyses of *RIZ1* alterations and patient clinicopathological features.

Features	No.	Univariate ^a		Multivariate ^b	
		HR (95% CI)	P-value	HR (95% CI)	P-value
Gender				NS	NS
Male	57	Reference			
Female	24	0.63 (0.37-1.08)	0.092		
Age				NS	NS
≤54 years	39	Reference			
>54 years	42	0.71 (0.44-1.13)	0.146		
Stage			0.222	-	-
II	2	Reference	1.000		
III	12	0.40 (0.08-1.92)	0.254		
IV	60	0.74 (0.176-3.09)	0.677		
Histological types			0.063	NS	NS
Papillary adenocarcinoma	17	Reference			
Well differentiated	23	1.62 (0.80-3.28)	0.177		
Moderately differentiated	9	1.45 (0.61-3.44)	0.401		
Poorly differentiated	22	1.39 (0.68-2.80)	0.359		
Other ^c	7	4.44 (1.64-12.00)	0.003		
Blood vessel invasion				-	-
Absent	24	Reference			
Present	48	1.44 (0.83-2.45)	0.188		
Nerve invasion				NS	NS
Absent	39	Reference			
Present	33	1.66 (0.99-2.76)	0.053		
Lymphatic invasion				-	-
Absent	16	Reference			
Present	56	1.39 (0.74-2.62)	0.309		
RIZ1 methylation				-	-
Absent	50	Reference			
Present	31	0.78 (0.48-1.26)	0.306		
RIZ LOH					
LOH ⁻	24	Reference			
LOH ⁺	14	1.93 (0.93-4.01)	0.078	NS	NS
RIZCA					
LOH ⁻	52	Reference			
LOH ⁺	4	1.39 (0.49-3.89)	0.534	NS	NS
RIZPro704					
LOH ⁻	42	Reference			
LOH ⁺	10	1.72 (0.83-3.59)	0.145	2.77 (1.12-6.84)	0.027

^aVariables presenting P<0.150 in univariate analysis were selected for multivariate analysis (bold). ^bMultivariate analysis using Cox regression, backward stepwise method. ^cIncludes adenosquamous and squamous carcinomas. HR, hazard ratio; 95% CI, 95% confidence interval; NS, not significant; -, not included in multivariate analysis

is not associated with patient clinicopathological features but may be associated with carcinogenesis (28), it is likely that inactivation of *RIZ1* by promoter hypermethylation may play a similar role in CCA. Moreover, the non-tumor cells used in our study, although having a normal appearance under gross and microscopic examination, may have already undergone genetic

and/or epigenetic alterations. Nevertheless, the methylated bands found in most of the non-tumor samples were much less intense than those observed in the tumor specimens.

LOH at RIZPro704 was a significant independent predictor for postoperative survival (Cox regression, P=0.027). This finding corroborated previous studies involving colorectal

cancer (7) and parathyroid tumors (18) where RIZPro704 LOH was higher than and mostly independent of RIZCA LOH. Almost all RIZPro704 LOH⁺ samples (8 of 10) lost the smaller allele (Pro704⁻) which resulted from a deletion polymorphism. RIZPro704 is located in the *RIZ1* coding region; however, its contribution to RIZ1 function in cancer is not much understood. Since RIZPro704 is close to the ER binding motif (amino acids 864-1,046 of RIZ1 protein) (13), this residue may be important for maintaining RIZ1 conformation. For this reason, interaction between ER and RIZ1 may occur only with the wild-type RIZ1 (Pro704⁺), which does not harbor a deletion polymorphism at Pro704. Loss of Pro704⁻ with remaining Pro704⁺ might be favorable for interaction between RIZ1 and ER. In a previous study, ER was up-regulated in 80% of CCA cases, while it was rarely expressed in normal liver tissues (29). *IGF-1* and *IGF-1R* expression was found to be repressed by RIZ1 (12) while expression increased to approximately 60% in human intrahepatic CCA cases, whereas their expression was not detected in normal human liver tissues (29). Taken together, we postulated that the up-regulation of ER in CCA inhibits the tumor suppressive activity of RIZ1 and activates the expression of some target genes involved in cell proliferation such as *IGF-1* resulting in poor prognosis of the patient. However, its response to estrogen and its association with bone mineral density in women remains controversial (30,31). Therefore, the biological role of *RIZ1* and its response to ER signaling in CCA require further investigation.

The percentages of MSI at RIZCA (9.9%) and at D1S228 (11.2%) (3) are similar, indicating the defect of mismatch repair genes. Thus, we expected that the frequency of *RIZ1* frameshift mutation in these samples might be similar to the MSI frequency found in both loci. Surprisingly, the frequency of frameshift mutations in *RIZ* was very low (2.5%) indicating that a frameshift mutation is not a common mechanism for *RIZ* inactivation in CCA, although its frequency is higher in other types of tumors (23).

In conclusion, the present study showed that, in CCA, genetic alterations of *RIZ1* such as LOH and frameshift mutations are not common compared to epigenetic alterations such as promoter hypermethylation. Epigenetic inactivation in *RIZ1* may occur at an early step in the process of carcinogenesis. Pro704 LOH was correlated to poor patient survival; however, further study is needed to elucidate the mechanisms involved in CCA.

Acknowledgements

This study was supported by the Thailand Research Fund through The Royal Golden Jubilee PhD Program, grant no. PHD/0084/2546 to P. Khaenam and T. Limpaboon, and by the Centre for Research and Development of the Medical Diagnostic Laboratories, Faculty of Associated Medical Sciences, Khon Kaen University, Thailand.

References

- Sripa B and Pairojkul C: Cholangiocarcinoma: lessons from Thailand. *Curr Opin Gastroenterol* 24: 349-356, 2008.
- Sripa B, Kaewkes S, Sithithaworn P, *et al*: Liver fluke induces cholangiocarcinoma. *PLoS Med* 4: e201, 2007.
- Limpaboon T, Tapdara S, Jearanaikoon P, Sripa B and Bhudhisawasdi V: Prognostic significance of microsatellite alterations at 1p36 in cholangiocarcinoma. *World J Gastroenterol* 12: 4377-4382, 2006.
- Buyse IM, Takahashi EI and Huang S: Physical mapping of the retinoblastoma interacting zinc finger gene RIZ to D1S228 on chromosome 1p36. *Genomics* 34: 119-121, 1996.
- Liu L, Shao G, Steele-Perkins G and Huang S: The retinoblastoma interacting zinc finger gene RIZ produces a PR domain-lacking product through an internal promoter. *J Biol Chem* 272: 2984-2991, 1997.
- Xie M, Shao G, Buyse IM and Huang S: Transcriptional repression mediated by the PR domain zinc finger gene RIZ. *J Biol Chem* 272: 26360-26366, 1997.
- Chadwick RB, Jiang GL, Bennington GA, *et al*: Candidate tumor suppressor RIZ is frequently involved in colorectal carcinogenesis. *Proc Natl Acad Sci USA* 97: 2662-2667, 2000.
- Jiang G, Liu L, Buyse IM, Simon D and Huang S: Decreased RIZ1 expression but not RIZ2 in hepatoma and suppression of hepatoma tumorigenicity by RIZ1. *Int J Cancer* 83: 541-546, 1999.
- He L, Yu JX, Liu L, *et al*: RIZ1, but not the alternative RIZ2 product of the same gene, is underexpressed in breast cancer, and forced RIZ1 expression causes G2-M cell cycle arrest and/or apoptosis. *Cancer Res* 58: 4238-4244, 1998.
- Jiang GL and Huang S: Adenovirus expressing RIZ1 in tumor suppressor gene therapy of microsatellite-unstable colorectal cancers. *Cancer Res* 61: 1796-1798, 2001.
- Steele-Perkins G, Fang W, Yang XH, *et al*: Tumor formation and inactivation of RIZ1, an Rb-binding member of a nuclear protein-methyltransferase superfamily. *Genes Dev* 15: 2250-2262, 2001.
- Pastural E, Takahashi N, Dong WF, *et al*: RIZ1 repression is associated with insulin-like growth factor-1 signaling activation in chronic myeloid leukemia cell lines. *Oncogene* 26: 1586-1594, 2007.
- Abbondanza C, Medici N, Nigro V, *et al*: The retinoblastoma-interacting zinc-finger protein RIZ is a downstream effector of estrogen action. *Proc Natl Acad Sci USA* 97: 3130-3135, 2000.
- Carling T, Kim KC, Yang XH, Gu J, Zhang XK and Huang S: A histone methyltransferase is required for maximal response to female sex hormones. *Mol Cell Biol* 24: 7032-7042, 2004.
- Gazzerro P, Abbondanza C, D'Arcangelo A, *et al*: Modulation of RIZ gene expression is associated to estradiol control of MCF-7 breast cancer cell proliferation. *Exp Cell Res* 312: 340-349, 2006.
- Kahlert S, Nuedling S, van Eickels M, Vetter H, Meyer R and Grohe C: Estrogen receptor alpha rapidly activates the IGF-1 receptor pathway. *J Biol Chem* 275: 18447-18453, 2000.
- Tokumaru Y, Nomoto S, Jeronimo C, *et al*: Biallelic inactivation of the RIZ1 gene in human gastric cancer. *Oncogene* 22: 6954-6958, 2003.
- Carling T, Du Y, Fang W, Correa P and Huang S: Intragenic allelic loss and promoter hypermethylation of the RIZ1 tumor suppressor gene in parathyroid tumors and pheochromocytomas. *Surgery* 134: 932-940, 2003.
- Oshimo Y, Oue N, Mitani Y, *et al*: Frequent epigenetic inactivation of RIZ1 by promoter hypermethylation in human gastric carcinoma. *Int J Cancer* 110: 212-218, 2004.
- Akahira J, Suzuki F, Suzuki T, *et al*: Decreased expression of RIZ1 and its clinicopathological significance in epithelial ovarian carcinoma: correlation with epigenetic inactivation by aberrant DNA methylation. *Pathol Int* 57: 725-733, 2007.
- Poetsch M, Dittberner T and Woenckhaus C: Frameshift mutations of RIZ, but no point mutations in RIZ1 exons in malignant melanomas with deletions in 1p36. *Oncogene* 21: 3038-3042, 2002.
- Fang W, Piao Z, Simon D, Sheu JC and Huang S: Mapping of a minimal deleted region in human hepatocellular carcinoma to 1p36.13-p36.23 and mutational analysis of the RIZ (PRDM2) gene localized to the region. *Genes Chromosomes Cancer* 28: 269-275, 2000.
- Piao Z, Fang W, Malkhosyan S, *et al*: Frequent frameshift mutations of RIZ in sporadic gastrointestinal and endometrial carcinomas with microsatellite instability. *Cancer Res* 60: 4701-4704, 2000.
- Limpaboon T, Khaenam P, Chinnasri P, *et al*: Promoter hypermethylation is a major event in hMLH1 gene inactivation in liver fluke-related cholangiocarcinoma. *Cancer Lett* 217: 213-219, 2005.

25. Herman JG, Graff JR, Myohanen S, Nelkin BD and Baylin SB: Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci USA* 93: 9821-9826, 1996.
26. Limpaboon T, Krissadarak K, Sripa B, *et al*: Microsatellite alterations in liver fluke-related cholangiocarcinoma are associated with poor prognosis. *Cancer Lett* 181: 215-222, 2002.
27. Yamamoto H, Sawai H and Perucho M: Frameshift somatic mutations in gastrointestinal cancer of the microsatellite mutator phenotype. *Cancer Res* 57: 4420-4426, 1997.
28. Hasegawa Y, Matsubara A, Teishima J, *et al*: DNA methylation of the RIZ1 gene is associated with nuclear accumulation of p53 in prostate cancer. *Cancer Sci* 98: 32-36, 2007.
29. Alvaro D, Barbaro B, Franchitto A, *et al*: Estrogens and insulin-like growth factor 1 modulate neoplastic cell growth in human cholangiocarcinoma. *Am J Pathol* 169: 877-888, 2006.
30. Grundberg E, Carling T, Brandstrom H, *et al*: A deletion polymorphism in the RIZ gene, a female sex steroid hormone receptor coactivator, exhibits decreased response to estrogen in vitro and associates with low bone mineral density in young Swedish women. *J Clin Endocrinol Metab* 89: 6173-6178, 2004.
31. Stolk L, van Meurs JB, Arp PP, Hofman A, Pols HA and Uitterlinden AG: The RIZ Pro704 insertion-deletion polymorphism, bone mineral density and fracture risk: the Rotterdam study. *Bone* 42: 286-293, 2008.
32. Du Y, Carling T, Fang W, Piao Z, Sheu JC and Huang S: Hypermethylation in human cancers of the RIZ1 tumor suppressor gene, a member of a histone/protein methyltransferase superfamily. *Cancer Res* 61: 8094-8099, 2001.