

Comano's (Trentino) thermal water interferes with interleukin-6 production and secretion and with cytokeratin-16 expression by cultured human psoriatic keratinocytes: Further potential mechanisms of its anti-psoriatic action

ANNA CHIARINI¹, ILARIA DAL PRA¹, RAFFAELLA PACCHIANA¹,
GIUSEPPE ZUMIANI², MAURO ZANONI³ and UBALDO ARMATO¹

¹Histology and Embryology Unit, Department of Biomedical and Surgical Sciences,
University of Verona Medical School, Verona I-37134; ²Dermatology Operative Unit,
Saint Chiara Hospital, Trento I-38100; ³Research Centre, Terme di Comano, Trentino, Italy

Received June 13, 2006; Accepted September 5, 2006

Abstract. Thermal balneotherapy with Comano's spa water (CW; Trentino, Italy) is used for psoriasis and other skin disorders but its mechanisms of action are mostly unknown. Previously, we showed that CW can interfere with the expression and secretion of various VEGF-A isoforms by cultured human psoriatic epidermal keratinocytes. In this study, confluent cultures of IL-6-hypersecreting keratinocytes isolated from 6 psoriatic patients were exposed for 11-15 days to DMEM, the chemicals of which had been dissolved in either deionised water (DW-DMEM, controls) or CW (CW-DMEM, treated cells). As detected by means of immunocytochemistry, Western immunoblotting, and ELISA assays, the intracellular levels and secretion rates of IL-6 were drastically curtailed in the CW-DMEM-incubated keratinocytes and in their cell-conditioned media. A nearly maximal inhibition of IL-6 release had already been induced by a CW fraction in the DMEM as low as 25%. CW exposure also promptly, intensely, and persistently down-regulated the expression of cytokeratin-16 (CK-16), a marker associated with keratinocyte psoriatic phenotype. Hence, CW balneotherapy may beneficially affect the clinical manifestations of psoriasis via an attenuation of the local deregulation of

several cytokines/chemokines, including IL-6 and VEGF-A isoforms, and of a concurrent, abnormal cell differentiation program entailing the expression, amongst other proteins, of CK-16.

Introduction

Psoriasis, a chronic inflammatory dermatosis affecting approximately 2% of the Western population, is clinically marked by relapsing-remitting manifestations of well-defined, symmetrical erythematous plaques covered by scales. Albeit genetically founded, the pathogenesis of psoriasis remains unclear (1). Currently, psoriasis is believed to be a T lymphocyte-driven disorder (2). The formation of tortuous, dilated, inflamed, and hyper-permeable venous limbs of capillary plexuses in the upper dermal papillae precedes the plaque's epidermal hyperplasia and dermal infiltration by inflammatory cells (i.e. neutrophils, T lymphocytes, monocytes) (3-6). It has been suggested that psoriasis is an angioproliferative ailment due to the local release of angiogenic molecules by the epidermis (7-12). Local fibroblast activation and increased keratinocyte production and release of several cytokines/chemokines, such as interleukin-1 (IL-1), IL-6, IL-8, IL-20, vascular endothelial growth factor-A (VEGF-A) isoforms, endothelial cell stimulating angiogenesis factor (ESAF), tumour necrosis factor- α (TNF- α), amphiregulin, transforming growth factor- α (TGF- α), and platelet-derived endothelial cell growth factor/thymidine phosphorylase (TP), are also included in the typical features of the disease (10,13-20).

IL-6 is a multifunctional cytokine of the haemopoietins family that also comprises leukemia inhibitory factor (LIF), oncostatin M (OSM), IL-11, granulocyte-colony stimulating factor (G-CSF), ciliary neurotrophic factor (CNTF), and cardiotrophin (referenced in refs. 17,21). All haemopoietins share the gp130 signal-transducing subunit (21). IL-6 acts as a growth factor for keratinocytes either directly via its specific receptor signalling (22-24) and/or indirectly by inducing the production and release of keratinocyte growth factor (KGF) by dermal fibroblasts (25). Non-activated normal keratinocytes

Correspondence to: Professor Ubaldo Armato, Histology and Embryology Unit, Department of Biomedical and Surgical Sciences, University of Verona Medical School, Strada Le Grazie 8, Verona I-37134, Italy
E-mail: ubaldo.armato@univr.it

Abbreviations: CK-16, cytokeratin-16; CW, Comano's thermal water; DW, deionised water; IL-6, interleukin-6; WB, Western immunoblotting

Key words: chemokines, Comano's thermal water, cytokeratin-16, cytokines, human keratinocytes, interleukin-6, psoriasis

Table I. Components of Comano's water.^a

Ions	mM
Sodium	0.182
Potassium	0.026
Magnesium	1.010
Calcium	2.440
Bicarbonate	6.340
Chloride	0.047
Sulfuric acid	0.144
Silicon	0.163
Fluorine	0.048
Lithium	0.0002
Aluminum	0.00246
Manganese	0.00064
Iron	0.0038
Copper	0.0017
Zinc	0.00143
Strontium	0.00605

^aThis water is hypotonic as its dry residue amounts to only 190 mg/l.

express low levels of IL-6-specific mRNA and release only tiny amounts of IL-6 (26). IL-6 expression is significantly heightened at psoriatic lesional skin sites (27-33), chiefly in transitional zones in which moderate epidermal cell hyper-proliferation occurs (31). IL-6 levels are increased even in the supernatants of lesional psoriatic skin (34). However, actual levels of IL-6 production and secretion may vary in psoriatic skin samples and supernatants from different patients according to disease stage and genetic background (1,35,36). In fact, IL-6 concentrations in lesional skin go up with the worsening or go down with the improvement of the clinical signs of psoriasis (29,31,36). Moreover, IL-6 production by keratinocytes is induced by IGF-II via the activation of NF- κ B (37), augmented by LIF (38) and by several other cytokines that are overproduced in psoriasis (i.e. IL-1 α , IL-1 β , IL-2, IL-4, GM-CSF, TNF- α) (31,39), and conversely inhibited by treatment with (-)-epigallocatechin-3-gallate, a major green tea polyphenol (40). By means of paracrine and autocrine loops mediated via its specific high-affinity functional receptors (24), IL-6 also may enhance the proliferation of human adult dermal fibroblasts and/or upregulate the production of collagen, glycosaminoglycans (GAGs), interstitial collagenase, and stromelysin-1 by such cells (24,41,42), thereby favouring the inflammatory processes related not only to psoriasis but even to wound healing and hypertrophic burn scarring (26,31,34,41,43-45). The increased circulating levels of IL-6 detected in psoriasis are thought to mediate, via the induction of other cytokines (e.g. IL-2) and adhesion molecules (e.g. ICAM-1), both the proliferative and functional activities of B, T, and natural killer (NK) cells, thereby modulating the systemic immune responses of the host in psoriasis and other pathologies (26,31,34,46,47).

Amongst the crucial features of a hyper-proliferative skin condition such as psoriasis is an abnormal keratinocyte differentiation program entailing the expression of proteins that are otherwise absent from normal skin, such as cytokeratin-16 (CK-16), CK-6, and CK-17, and the antimicrobial/elastase inhibitor SKALP/elafin (48). Hence, these proteins or their mRNAs have been used as stand-in markers in drug-screening procedures (48), and CK-16 is usually referred to as a strongly expressed constituent of the psoriasis-associated keratinocyte phenotype (49).

Comano (Trentino, Italy) spa's water (CW) is a thermal hypotonic water containing various electrolytes (Table I). The major dermatological diseases so far treated via CW balneotherapy are psoriasis and atopic dermatitis (50). Other dermatoses also cared for with CW include contact dermatitis, seborrhoeic dermatitis, *lichen planus*, and palmoplantar keratosis (50). Previous *in vivo* studies showed the effectiveness of CW balneotherapy in the treatment of psoriasis, since it both significantly lessened hyperkeratosis, acanthosis, and dermal papillomatosis and improved skin hydration (50). It must be recalled here that the permeability barrier of normal epidermis is severely disturbed in psoriatic skin (51,52), and that bathing in hypotonic salt solutions triggers anti-inflammatory effects in lesional skin sites (53). However, most of the mechanisms through which the clinical signs of psoriasis (and of the other above mentioned skin disorders) are improved by means of CW balneotherapy have not as yet been clarified. In a previous study, we showed that exposure to CW interferes with VEGF-A isoform expression and secretion by the human psoriatic keratinocytes (54). To further clarify the mechanism(s) possibly involved in the therapeutic effectiveness of CW balneotherapy in psoriasis, we investigated CW's effects on IL-6 hyper-production and hyper-secretion and on CK-16 expression by epidermal keratinocytes isolated from lesional skin biopsies. We show that the addition of CW (in total or partial stead of DW) to the growth medium thwarts the heightened production and release of IL-6 and the expression of CK-16 on the part of the psoriatic keratinocytes. These findings are consistent with CW being endowed with a complex phenotype- and cytokine/chemokine-regulating potential that translates into valuable anti-psoriatic therapeutic benefits.

Materials and methods

In vitro cell culture. For this work human epidermal keratinocytes were isolated from skin biopsies taken, after informed consent, from 6 psoriatic patients. After rapid transfer to the laboratory, the biopsies were incubated at 4°C overnight in a dispase II solution (0.25% w/v; Roche, Milan, Italy). Weak enzymatic digestion allowed the epidermis (as a single lamina) to easily detach from the underlying dermis and subcutaneous tissue. By incubating the isolated epidermal sheet in trypsin solutions (0.25% w/v), suspensions of keratinocytes were obtained. Trypsin action was next inhibited by adding an excess of serum, and the cell suspensions were soon spun down at 600 rpm for 10 min at 4°C. The supernatants were decanted, the pellets resuspended, and the living cells counted in a Neubauer chamber. Keratinocytes were next seeded into plastic flasks pre-coated with a feeder-layer of preirradiated 3T3-J2 cells. To expand the keratinocyte


IL-6

Figure 1. Exposure to a CW-DMEM medium elicited an early, deep down-regulation and a late, modest rise of the intracellular IL-6 levels in cultured otherwise IL-6-hypersecreting psoriatic epidermal keratinocytes. (A) Psoriatic keratinocytes incubated in 100% CW-DMEM exhibited a decrease at day 3 and an increase at day 11 of the fluorescent signal specifically related to IL-6 with respect to untreated (DW-DMEM-kept) counterparts. The confocal pictures shown here were deconvolved as indicated in the Materials and methods and are representative of 6 distinct experiments. (B and C) WB observations and densitometric analyses of the proteins extracted from keratinocytes exposed to a DW- or a CW-DMEM for 3, 5, 7, and 11 days showed an early fall followed by a late moderate rise of the intracellular IL-6 levels in CW-treated keratinocytes. The immunoblot shown in B is representative, and the points on the curves in C are means $\pm$ SEM of 6 experiments. ** $p < 0.001$ and * $p < 0.01$ between time-corresponding DW and CW points.

population, MCDB153:1 medium [consisting of three parts of Dulbecco's modified Eagle's medium (DMEM) and one part of F12 medium; Sigma-Aldrich, Milan, Italy] was used, to which foetal bovine serum (FBS; 10% v/v; BioWhittaker Europe, Belgium), antibiotics (solution of penicillin-streptomycin 1% w/v; BioWhittaker), epidermal growth factor (EGF; $0.1 \mu\text{g ml}^{-1}$; PeproTech, UK), insulin (20 ng ml^{-1} ; PeproTech), and hydrocortisone ($0.5 \mu\text{g ml}^{-1}$; PeproTech) were added. This medium was replaced every two days with fresh samples of the same medium. Human psoriatic keratinocytes proliferated rapidly starting from minute clusters and formed a single layer of small and highly adherent epithelial cells. They had a mitotic doubling time of approximately 48 h. Once cultured *in vitro*, such keratinocytes kept steadily secreting into the medium, as determined by ELISA assays, amounts of IL-6 several fold greater than did normal keratinocytes (55).

Experimental protocol. IL-6-hypersecreting psoriatic keratinocytes were detached from the culture flasks by a mild trypsin treatment and then seeded at 1.0×10^6 cells into wells containing 2.0 ml of either DMEM medium, whose chemical constituents had been dissolved in DW (controls in DW-DMEM), or in one of three different CW-DMEM media, in

which DW had been totally (100%) or in part (50% or 25%) substituted with CW. Between days 3 and 15 of experimental treatment, the cultured cells and/or the cell-conditioned media were sampled and their respective contents of IL-6 and/or CK-16 were determined.

Immunocytochemistry. At chosen time points, psoriatic keratinocytes exposed to either DW- or 100% CW-DMEM were fixed with absolute methanol at -20°C for 10 min, washed twice with PBS, and permeabilised in 0.1% Triton X-100 at room temperature for 15 min. Then the cells were washed with PBS-FBS (1%) (Cambrex BioScience, Milan, Italy) at room temperature for 1 h and incubated for 1 h at 37°C with an anti-CK-16 IgG mouse monoclonal antibody (final dilution $10 \mu\text{g ml}^{-1}$; Chemicon International, Inc.) or with an anti-IL-6 rabbit polyclonal antibody (final dilution $10 \mu\text{g ml}^{-1}$; Santa Cruz Biotechnology, Inc., Santa Cruz, CA). Next, keratinocytes were washed three times with PBS-BSA (1%) and incubated for 1 h at room temperature in the dark with specific secondary antibodies (1:100 dilution) conjugated with Alexa Fluor-488 or -555 (Molecular Probes, Invitrogen Corporation, Carlsbad, CA). Control cells not exposed to the primary antibody were always run in parallel. The cells were finally examined under

an LSM 510 confocal microscope (Carl Zeiss S.p.A., Milan, Italy). Deconvolved fluorescence images were obtained with Huygens Professional Software for Windows (Scientific Volume Imaging b.v., Hilversum, The Netherlands).

Western immunoblotting (WB). After 3, 7, and 11 days *in vitro*, psoriatic keratinocytes kept in 100% CW- or in DW-DMEM were scraped into cold PBS and sedimented at 200 x g for 10 min. The sedimented cells were homogenized in T-PER™ tissue protein extraction reagent (Pierce Chemical Co., Rockford, IL) containing a complete EDTA-free protease inhibitor cocktail (Roche Diagnostics, Monza, Italy). The protein contents of the samples were assayed by Bradford's method (56) using bovine serum albumin as a standard. Equal amounts (10 or 20 µg) of proteins from each cell lysate or cell-conditioned DW- or CW-DMEM (25 µl) were boiled in sample buffer (0.0625 M Tris-HCl, pH 6.8, 2% w/v SDS, 5% w/v β-mercaptoethanol, 10% v/v glycerol, 0.002% w/v bromophenol blue) and electrophoresed in 10% w/v SDS-polyacrylamide gel. The separated proteins were blotted onto a nitrocellulose membrane (0.45 µm; Bio-Rad Laboratories, Hercules, CA). To immunodetect IL-6 and CK-16, the blots were probed with the same specific primary antibodies as used for immunocytochemistry at a final dilution of 1.0 µg ml⁻¹. Blots were next incubated with alkaline phosphatase-conjugated anti-mouse or anti-rabbit IgG (Santa Cruz), and stained with BCIP/NBT liquid substrate reagent (Sigma). Developed blots were photographed with an Olympus 3300™ digital camera, and the determination of the M_r and the densitometric analysis of each specific protein band were carried out using Sigmagel™ software (Jandel Corp., Erkrath, Germany).

ELISA assay of IL-6. Human psoriatic keratinocytes were cultured for 15 days in four different growth media containing DMEM components dissolved in the following percent fluid fractions: i) CW 100%/DW 0%; ii) CW 50%/DW 50%; iii) CW 25%/DW 75%; and iv) CW 0%/DW 100% (control medium). Cell-conditioned samples of the four kinds of growth media were taken at days 3, 5, 7, 10, 12, and 15 of culture and stored at -80°C to be subsequently assayed for their IL-6 content. To this aim, a specific commercial ELISA kit was used (CLB, Amsterdam, The Netherlands). The tests were performed according to the instructions of the manufacturer. The sensitivity of the assays for IL-6 was 0.5 pg ml⁻¹. The results were expressed as daily secretion values per duplicate cultures.

Statistical analysis. One-way analysis of variance (ANOVA) with post hoc Bonferroni's test was used to compare mean values and a significance level of ≤0.05 was chosen.

Results

Effects of CW on intracellular levels of IL-6. WB analyses revealed that human psoriatic keratinocytes produced IL-6 species endowed with molecular masses close to 45 kDa (Fig. 1B), suggesting that the original 23- to 25-kDa IL-6 protein moiety underwent significant post-translational modifications (N- and O-glycosylations and phosphorylations) within the epidermal epithelial cells (57). Both confocal

sIL-6




Figure 2. Fractions of CW ranging from 25% to 100% in the DMEM medium deeply and progressively curtailed the hyper-secretion of IL-6 on the part of cultured psoriatic keratinocytes bringing it down to within normal range values (i.e. 6- to 8-ng ml⁻¹ per 10⁶ cells) (55). Points on the curves are means from 6 experiments carried out in duplicate. SEMs, not shown, were within ± 15% of each mean value. Between day 5 and 15 all values pertaining to CM-treated keratinocytes exhibited a statistical significance of at least $p < 0.01$ vs. the values of time-corresponding DW-DMEM-incubated (control) cells. sIL-6, IL-6 secreted into the growth media.

microscopy (Fig. 1A), WB observations (Fig. 1B), and densitometric determinations (Fig. 1C) showed that by day 3 after the onset of the experiments the intracellular levels of IL-6 had significantly fallen (-73%, $p < 0.001$ in WB specimens) in the 100% CW-DMEM-incubated keratinocytes with respect to the DW-DMEM-kept (untreated) cells. Next, between days 5 and 7, intracellular IL-6 levels became close in both untreated and CW-treated keratinocytes (Fig. 1A-C). Finally, by day 11, intracellular IL-6 levels were found to have risen (+59%, $p < 0.01$ in WB specimens) in CW-DMEM-kept keratinocytes with respect to untreated ones (Fig. 1A-C). Thus, an early deep cutback of intracellular IL-6 levels was induced by incubating keratinocytes in CW-DMEM medium, whereas a tardy discrete intracellular IL-6 surge was the likely upshot of a quite strongly hindered secretion of IL-6, the production of which had meanwhile been down-regulated.

Effects of CW on IL-6 secretion. When untreated (i.e. incubated in DW-DMEM), the psoriatic keratinocytes released massive amounts (up to >50 ng ml⁻¹ per 10⁶ cells) of IL-6 into the medium from the third day onwards (Fig. 2). However, when exposing keratinocytes from day 5 onwards to DMEM containing various percentages (i.e. from 25% to 100%) of CW fractions, the IL-6 hyper-secretory activity of the same cells was strikingly and progressively cut down (e.g., at day 5, from -69% to -76%, $p < 0.001$ vs. parallel untreated keratinocytes; at day 15, from -88% to -96%, $p < 0.001$ vs. parallel untreated cells) (Fig. 2). Most interestingly, a nearly maximum inhibitory effect on IL-6 secretion had already been achieved by exposing keratinocytes to CW 25%/DW 75%-DMEM

CK-16




Figure 3. Exposure to CW persistently down-regulated the expression of CK-16, a marker of the psoriatic phenotype, by human lesional keratinocytes cultured *in vitro*. (A) Deconvolved confocal pictures representative of 6 experiments shows that after 3 days a sharp reduction in the intracellular CK-16 content of keratinocytes exposed to 100% CW-DMEM had occurred with respect to that of DW-DMEM-incubated (control) cells. (B and C) WB observations and densitometric analyses showed that a strong and persistent down-regulation of CK-16 took place in CW-DMEM-kept keratinocytes with respect to DW-DMEM-incubated keratinocytes. The immunoblot shown in B is typical, and the points on the curves in C are means $\pm$ SEM of 6 experiments. * $p < 0.001$ between the values pertaining to time-corresponding keratinocyte samples kept in either DW-DMEM or CW-DMEM.

(Fig. 2). Hence, a persisting exposure to CW fractions ranging from 25% to 100% in the DMEM similarly brought IL-6 secretion rates down to within the range of normal values (i.e. ~ 6 - to 8 -ng ml^{-1} per 10^6 keratinocytes) (55).

Effects of CW on CK-16 expression. Observations under the confocal microscope revealed that after 3 days of exposure to 100% CW-DMEM the intensity of the fluorescent signal specifically related to CK-16 had remarkably weakened in the cytoplasm of psoriatic keratinocytes with respect to parallel controls kept in DW-DMEM (Fig. 3A). The results of WB observations and of corresponding densitometric assessments showed that after a 3-day exposure to 100% CW-DMEM, the density of the CK-16-specific 48-kDa protein band had diminished by $\sim 76\%$ ($p < 0.001$) vs. that of

parallel DW-DMEM-incubated keratinocytes (Fig. 3B and C). Moreover, after an 11-day exposure to 100% CW-DMEM, the density of the CK-16-specific protein band had been reduced merely to one-tenth ($p < 0.001$) that proper of parallel DW-DMEM-kept keratinocytes (Fig. 3B and C). Thus, a lasting exposure to CW severely hindered the expression of CK-16, a marker of the psoriatic phenotype (48,49), by the human epidermal keratinocytes.

Discussion

In this work we tested CW's effects on IL-6 hyper-production and hyper-secretion by psoriatic keratinocytes kept in pure *in vitro* cultures, i.e. in the complete absence of T cells (58). Our results show that the addition of CW (instead of DW) to

the DMEM significantly curtailed both the heightened intracellular levels and secretion rates of IL-6 by these psoriatic keratinocytes. Notably, IL-6 hyper-secretion was cut down to within the range of normal values (55) by incubating the keratinocytes in DMEM whose CW fraction was as little as 25%. This strong inhibition of IL-6 release went so far as to elicit a late discrete intracellular accumulation of IL-6 even though IL-6 production had also been down-regulated. The operative mechanisms underlying these IL-6-interfering effects elicited by CW components in keratinocytes remain to be elucidated. Collectively, our findings suggest that, by interfering with IL-6 hyper-production and hyper-secretion by the psoriatic keratinocytes, the exposure to CW significantly hinders the mitogenic, proinflammatory, and proangiogenic actions sustained by the occurrence of an IL-6 surplus within psoriatic skin lesions (10,17,22,30,31,59).

The programmed expression of cytokeratins (CKs), which is determined by the location and functioning of the keratinocytes, is commonly taken as a set of phenotypic markers related to the stages of development and differentiation of the epidermal cells (60). The proliferating keratinocytes residing in the basal layer produce CK-5, CK-14, and low amounts of CK-15; the differentiating keratinocytes placed in the suprabasal layers express CK-1, CK-2, and CK-10; conversely, like CK-6 and CK-17, CK-16 is uniquely expressed in activated, hyper-proliferating keratinocytes such as those harboured in psoriatic lesions (48,60). Therefore, CK-16 is usually referred to as a marker associated with the psoriatic phenotype (49,61). The present results show that exposure to CW elicits a massive and persistent down-regulation of CK-16 expression in cultured human adult psoriatic keratinocytes. Thus, the whole of the interferences brought about by CW exposure on VEGF-A isoforms (54), IL-6, and CK-16 expression suggest that CW has the ability to shift the keratinocytes' psoriatic phenotype towards a somewhat more normal pattern. Further studies should establish whether CW down-regulates other cytokines or chemokines, such as IL-1, IL-8 and TNF- α (17), and markers, such as CK-6, CK-17, and SKALP/elafin (49,62,63), all of which are strongly expressed by the psoriatic keratinocytes. However, our previous (54) and present findings clearly demonstrate that CW acts via identifiable and measurable biological mechanisms, thereby excluding that CW actions fall within the compass of placebo effects.

In conclusion, our previous (54) and present observations support the view that CW balneotherapy may elicit beneficial effects by interfering with an improper local production and secretion of several chemokines and cytokines, including IL-6 and VEGF-A isoforms, and by attenuating some facets, such as CK-16 expression, of the psoriatic phenotype, which altogether underlie the epidermal hyperplasia and dermal neo-angiogenesis, inflammation, and leukocyte infiltration phenomena proper of local psoriatic illness.

References

- Bowcock AM and Krueger JG: Getting under the skin: the immunogenetics of psoriasis. *Nat Rev Immunol* 5: 699-711, 2005.
- Bos JD and De Rie MA: The pathogenesis of psoriasis. *Immunol Today* 20: 40-46, 1999.
- Telner P and Fekete Z: The capillary responses in psoriatic skin. *J Invest Dermatol* 36: 225-230, 1961.
- Creamer D, Allen MH, Sousa A, Poston R and Barker JN: Localization of endothelial proliferation and microvascular expansion in active plaque psoriasis. *Br J Dermatol* 136: 859-865, 1997.
- Braverman IM and Sibley J: Role of the microcirculation in the treatment and pathogenesis of psoriasis. *J Invest Dermatol* 78: 12-17, 1982.
- Schubert C and Christophers E: Mast cells and macrophages in early relapsing psoriasis. *Arch Dermatol Res* 277: 352-358, 1985.
- Folkman J: Angiogenesis in psoriasis: therapeutic implications. *J Invest Dermatol* 59: 40-43, 1972.
- Barker JN: The pathophysiology of psoriasis. *Lancet* 338: 227-230, 1991.
- Goodfield M, Hull SM, Holland D, Roberts G, Wood E, Reid S and Cunliffe W: Investigations of the 'active' edge of plaque psoriasis: vascular proliferation precedes changes in epidermal keratin. *Br J Dermatol* 131: 808-813, 1994.
- Creamer D, Sullivan D, Bicknell R and Barker J: Angiogenesis in psoriasis. *Angiogenesis* 5: 231-236, 2002.
- Malhotra R, Stenn KS, Fernandez LA and Braverman IM: Angiogenic properties of normal and psoriatic skin associate with epidermis, not dermis. *Lab Invest* 61: 162-165, 1989.
- Barnhill RL, Parkinson EK and Ryan TJ: Supernatants from cultured human epidermal keratinocytes stimulate angiogenesis. *Br J Dermatol* 110: 273-281, 1984.
- Nickoloff BJ, Mitra RS, Varani J, Dixit VM and Polverini PJ: Aberrant production of interleukin-8 and thrombospondin-1 by psoriatic keratinocytes mediates angiogenesis. *Am J Pathol* 144: 820-828, 1994.
- Ettahadi P, Greaves MW, Wallach D, Aderka D and Camp RD: Elevated tumour necrosis factor- α (TNF- α) biological activity in psoriatic skin lesions. *Clin Exp Immunol* 96: 146-151, 1994.
- Elder JT, Fisher GJ, Lindquist PB, Bennett GL, Pittelkow MR, Coffey RJ Jr, Ellingsworth L, Derynck R and Voorhees JJ: Overexpression of transforming growth factor α in psoriatic epidermis. *Science* 243: 811-814, 1989.
- Creamer D, Jaggar R, Allen M, Bicknell R and Barker J: Overexpression of the angiogenic factor platelet-derived endothelial cell growth factor/thymidine phosphorylase in psoriatic epidermis. *Br J Dermatol* 137: 851-855, 1997.
- Bonifati C and Ameglio F: Cytokines in psoriasis. *Int J Dermatol* 38: 241-251, 1999.
- Blumberg H, Conklin D, Xu WF, *et al*: Interleukin 20: discovery, receptor identification, and role in epidermal function. *Cell* 104: 9-19, 2001.
- Detmar M, Brown LF, Claffey KP, Yeo KT, Kocher O, Jackman RW, Berse B and Dvorak HF: Overexpression of vascular permeability factor/vascular endothelial growth factor and its receptors in psoriasis. *J Exp Med* 180: 1141-1146, 1994.
- Bhushan M, McLaughlin B, Weiss JB and Griffiths CE: Levels of endothelial cell stimulating angiogenesis factor and vascular endothelial growth factor are elevated in psoriasis. *Br J Dermatol* 141: 1054-1060, 1999.
- Gearing DP: The leukemia inhibitory factor and its receptor. *Adv Immunol* 53: 31-58, 1993.
- Hernandez-Quintero M, Kuri-Harcuch W, Gonzalez Robles A and Castro-Munozledo F: Interleukin-6 promotes human epidermal keratinocyte proliferation and keratin cytoskeleton reorganization in culture. *Cell Tissue Res*, Mar 21, 2006 (Epub ahead of print).
- Waelte ER, Inaebit SP, Rast HP, Hunziker T, Limat A, Braathen LR and Wiesmann U: Co-culture of human keratinocytes on post-mitotic human dermal fibroblast feeder cells: production of large amounts of interleukin-6. *J Invest Dermatol* 98: 805-808, 1992.
- Hibi M, Murakami M, Saito M, Hirano T, Taga T and Kishimoto T: Molecular cloning and expression of an IL-6 signal transducer. *Cell* 63: 1149-1157, 1990.
- Brauchle M, Angermeyer K, Hubner G and Werner S: Large induction of keratinocyte growth factor expression by serum growth factors and pro-inflammatory cytokines in cultured fibroblasts. *Oncogene* 9: 3199-3204, 1994.
- Luger TA and Schwartz T: Epidermal cytokines. In: *Skin Immune System*. Bos J (ed). Boca Raton, CRC Press, pp257-292, 1990.
- Bonifati C, Ameglio F, Carducci M, Sacerdoti G, Pietravalle M and Fazio M: Interleukin-1-beta, interleukin-6, and interferon-gamma in suction blister fluids of involved and uninvolved skin and in sera of psoriatic patients. *Acta Derm Venereol Suppl* 186: 23-24, 1994.



SPANDIDOS ti C, Carducci M, Cordiali Fei P, Trento E, Sacerdoti G, PUBLICATIONS M and Ameglio F: Correlated increases of tumour necrosis

- factor- α , interleukin-6 and granulocyte monocyte-colony stimulating factor levels in suction blister fluids and sera of psoriatic patients - relationships with disease severity. *Clin Exp Dermatol* 19: 383-387, 1994.
29. Ameglio F, Bonifati C, Pietravalle M and Fazio M: Interleukin-6 and tumour necrosis factor levels decrease in the suction blister fluids of psoriatic patients during effective therapy. *Dermatology* 189: 359-363, 1994.
30. Grossman RM, Krueger J, Yourish D, Granelli-Piperno A, Murphy DP, May LT, Kupper TS, Sehgal PB and Gottlieb AB: Interleukin 6 is expressed in high levels in psoriatic skin and stimulates proliferation of cultured human keratinocytes. *Proc Natl Acad Sci USA* 86: 6367-6371, 1989.
31. Paquet P and Pierard GE: Interleukin-6 and the skin. *Int Arch Allergy Immunol* 109: 308-317, 1996.
32. Ameglio F, Bonifati C, Fazio M, Mussi A, Trento E, Cordiali Fei P, Donati P, Pimpinelli F, D'Auria L and Carducci M: Interleukin-11 production is increased in organ cultures of lesional skin of patients with active plaque-type psoriasis as compared with nonlesional and normal skin. Similarity to interleukin-1 beta, interleukin-6 and interleukin-8. *Arch Dermatol Res* 289: 399-403, 1997.
33. Ohta Y, Nishiyama S and Nishioka K: *In situ* expression of interleukin-6 in psoriatic epidermis during treatment. *J Dermatol* 21: 301-307, 1994.
34. Bonifati C, Mussi A, D'Auria L, Carducci M, Trento E, Cordiali-Fei P and Ameglio F: Spontaneous release of leukemia inhibitory factor and oncostatin-M is increased in supernatants of short-term organ cultures from lesional psoriatic skin. *Arch Dermatol Res* 290: 9-13, 1998.
35. Takematsu H and Tagami H: Lack of correlation between interleukin 6 and interleukin 1 levels in psoriatic lesional skin. *Tohoku J Exp Med* 172: 243-252, 1994.
36. Olaniran AK, Baker BS, Paige DG, Garioch JJ, Powles AV and Fry L: Cytokine expression in psoriatic skin lesions during PUVA therapy. *Arch Dermatol Res* 288: 421-425, 1996.
37. Kwon YW, Jang ER, Lee YM, Kim YS, Kwon KS, Jang HS, Oh CK and Kim KW: Insulin-like growth factor II induces interleukin-6 expression via NF- κ B activation in psoriasis. *Biochem Biophys Res Commun* 278: 312-317, 2000.
38. Villiger PM, Geng Y and Lotz M: Induction of cytokine expression by leukemia inhibitory factor. *J Clin Invest* 91: 1575-1581, 1993.
39. Hirano T, Akira S, Taga T and Kishimoto T: Biological and clinical aspects of interleukin 6. *Immunol Today* 11: 443-449, 1990.
40. Xia J, Song X, Bi Z, Chu W and Wan Y: UV-induced NF- κ B activation and expression of IL-6 is attenuated by (-)-epigallocatechin-3-gallate in cultured human keratinocytes *in vitro*. *Int J Mol Med* 16: 943-950, 2005.
41. Mateo RB, Reichner JS and Albina JE: Interleukin-6 activity in wounds. *Am J Physiol* 266: R1840-R1844, 1994.
42. Fehgali CA, Bost KL, Boulware DW and Levy LS: Human recombinant interleukin-4 induces proliferation and interleukin-6 production by cultured human skin fibroblasts. *Clin Immunol Immunopathol* 63: 182-187, 1992.
43. Elias JA, Freundlich B, Adams S and Rosenbloom J: Regulation of human lung fibroblast collagen production by recombinant interleukin-1, tumor necrosis factor, and interferon- γ . *Ann NY Acad Sci* 580: 233-244, 1990.
44. Brenneisen P, Wlaschek M, Wenk J, Baudschun R, Hinrichs R, Dissemond J, Krieg T and Scharffetter-Kochanek K: Ultraviolet-B induction of interstitial collagenase and stromelysin-1 occurs in human dermal fibroblasts via an autocrine interleukin-6-dependent loop. *FEBS Lett* 449: 36-40, 1999.
45. Duncan MR and Berman B: Stimulation of collagen and glycosaminoglycan production in cultured human adult dermal fibroblasts by recombinant human interleukin-6. *J Invest Dermatol* 97: 686-692, 1991.
46. Billiau A: Interferon β 2 as a promotor of growth and differentiation of B cells. *Immunol Today* 8: 84-87, 1987.
47. Luger TA, Schwarz T, Krutmann J, Kirnbauer R, Neuner P, Kock A, Urbanski A, Borth W and Schauer E: Interleukin-6 is produced by epidermal cells and plays an important role in the activation of human T-lymphocytes and natural killer cells. *Ann NY Acad Sci* 557: 405-414, 1989.
48. Pol A, van Ruissen F and Schalkwijk J: Development of a keratinocyte-based screening model for anti-psoriatic drugs using green fluorescent protein under the control of an endogenous promoter. *J Biomol Screen* 7: 325-332, 2002.
49. Leigh IM, Navsaria H, Purkis PE, McKay IA, Bowden PE and Riddle PN: Keratins (K16 and K17) as markers of keratinocyte hyperproliferation in psoriasis *in vivo* and *in vitro*. *Br J Dermatol* 135: 501-511, 1995.
50. Zumiani G, Zanoni M and Agostini G: Valutazione dell'efficacia dell'acqua della fonte termale di Comano versus acqua di acquedotto nella cura della psoriasi. *G Ital Dermatol Venereol* 135: 1-5, 2000.
51. Grice KA: Transepidermal water loss in pathological skin. In: *Physiology and Pathophysiology of the Skin*. Jarret A (ed). Academic Press, London, pp2147-2155, 1980.
52. Harding CR: The stratum corneum: structure and function in health and disease. *Dermatol Ther* 17 (suppl 1): 6-15, 2004.
53. Tsourelis-Nikita E, Menchini G, Gheretich I and Hercogova J: Alternative treatment of psoriasis with balneotherapy using Leopoldine spa water. *J Eur Acad Dermatol Venereol* 16: 260-262, 2002.
54. Chiarini A, Dal Pra I, Pacchiana R, Menapace L, Zumiani G, Zanoni M and Armato U: Comano's (Trentino) thermal water interferes with the expression and secretion of vascular endothelial growth factor-A (VEGF-A) protein isoforms by cultured human psoriatic keratinocytes: A potential mechanism of its anti-psoriatic action. *Int J Mol Med* 18: 17-25, 2006.
55. McKenzie RC, Venner TJ, Sauder DN and Farkas-Himsley H: Augmentation of interleukin-6 (IL-6) expression in squamous carcinoma cells and normal human keratinocytes treated with recombinant anti-neoplastic protein (ACP). *Anticancer Res* 14: 1165-1168, 1994.
56. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254, 1976.
57. Santhanam U, Ghayeb J, Sehgal PB and May LT: Post-translational modifications of human interleukin-6. *Arch Biochem Biophys* 274: 161-170, 1989.
58. Barker CL, McHale MT, Gillies AK, Waller J, Pearce DM, Osborne J, Hutchinson PE, Smith GM and Pringle JH: The development and characterization of an *in vitro* model of psoriasis. *J Invest Dermatol* 123: 892-901, 2004.
59. Penkowa M, Camats J, Hadberg H, Quintana A, Rojas S, Giral M, Molinero A, Campbell IL and Hidalgo J: Astrocyte-targeted expression of interleukin-6 protects the central nervous system during neuroglial degeneration induced by 6-aminocotinamide. *J Neurosci Res* 73: 481-496, 2003.
60. Freedberg IM, Tomic-Canic M, Komine M and Blumenberg M: Keratins and the keratinocyte activation cycle. *J Invest Dermatol* 116: 633-640, 2001.
61. Machesney M, Tidman N, Waseem A, Kirby L and Leigh I: Activated keratinocytes in the epidermis of hypertrophic scars. *Am J Pathol* 152: 1133-1141, 1998.
62. Pfundt R, Wiggins M, Bergers M, Zweers M, Frenken M and Schalkwijk J: TNF- α and serum induce SKALP/elafin gene expression in human keratinocytes by a p38 MAP kinase-dependent pathway. *Arch Dermatol Res* 292: 180-187, 2000.
63. Van Ruissen F, de Jongh GJ, Zeeuwen PL, Van Erp PE, Madsen P and Schalkwijk J: Induction of normal and psoriatic phenotypes in submerged keratinocyte cultures. *J Cell Physiol* 168: 442-452, 1996.