

# Interleukin-18 regulates T helper 1 or 2 immune responses of human cord blood CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> natural killer T cells

YUKA FUJIBAYASHI<sup>1</sup>, YOSHIHIRO FUJIMORI<sup>1</sup>, IKUYO KASUMOTO<sup>1</sup>, SHUNRO KAI<sup>3</sup>, HIROSHI HARA<sup>1</sup>, HARUKI OKAMURA<sup>2</sup>, HIROKO TSUTSUI<sup>4</sup>, HIROYASU OGAWA<sup>5</sup> and KENJI NAKANISHI<sup>6</sup>

Laboratories of <sup>1</sup>Cell Transplantation, and <sup>2</sup>Host Defenses, Institute for Advanced Medical Sciences,

<sup>3</sup>Division of Blood Transfusion, <sup>4</sup>Department of Microbiology, <sup>5</sup>Division of Hematology, Department of Medicine, and

<sup>6</sup>Department of Immunology and Medical Zoology, Hyogo College of Medicine, Nishinomiya, Hyogo 663-8501, Japan

Received March 14, 2007; Accepted April 19, 2007

**Abstract.** Natural killer T (NKT) cells, exhibiting both T-cell and NK-cell markers, are known to regulate immune responses by secreting T-helper (Th) 1 and Th2 cytokines. We analyzed NKT cells in cord blood (CB) for phenotypical and functional characteristics and regulatory mechanisms that control Th1 and Th2 determination. Human CB V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells were predominantly the CD4<sup>+</sup> single positive (SP) phenotype (approximately 96%), in contrast to adult peripheral blood V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells which are composed of a dominant population of the CD4<sup>+</sup>CD8<sup>-</sup> double negative (DN) phenotype and a minor population of the CD4<sup>+</sup> SP phenotype. The CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells, following stimulation with the primary culture, gained the capacity to secrete interferon (IFN)- $\gamma$ , a Th1 cytokine, and interleukin (IL)-13, a Th2 cytokine. The combination of IL-18 and IL-12 induced IFN- $\gamma$  production in CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells, while IL-18 in combination with IL-2 induced IL-13 production in these cells. Thus, IL-18 regulates the determination of the Th1 or Th2 immune response by human CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells through different cytokine combinations.

## Introduction

Natural killer T (NKT) cells have been implicated in various immune responses, including bacterial infection, tumor rejection, immune tolerance and the regulation of autoimmune diseases (1,2). NKT cells recognize glycolipids [ $\alpha$ -galactosylceramide ( $\alpha$ -GalCer)] presented by the MHC-like molecule CD1d. Although murine CD1d-restricted NKT cells express

an invariant T-cell receptor (TCR) consisting of V $\alpha$ 14/J $\alpha$ 18 paired with V $\beta$ 8, V $\beta$ 7 or V $\beta$ 2 (V $\alpha$ 14<sup>+</sup> NKT cells), human CD1d-restricted NKT cells express a TCR consisting of a V $\alpha$ 24/J $\alpha$ Q chain paired with a V $\beta$ 11 chain (V $\alpha$ 24<sup>+</sup> NKT cells) (1,2). In human adult peripheral blood (PB), CD4<sup>+</sup>CD8<sup>-</sup> double negative (DN) cells consist predominantly of V $\alpha$ 24<sup>+</sup> NKT cells (2-4) which have been reported to be decreased in patients with autoimmune diseases (5) and type 1 diabetes (6). NKT cells potentially have the ability to produce both T helper (Th) 1 cytokines, such as interferon (IFN)- $\gamma$ , and Th2 cytokines, such as interleukin (IL)-4 and IL-13 (1). The former is responsible for inducing the inflammatory process while the latter is responsible for the anti-inflammatory or allergic process (7).

Human umbilical cord blood (CB) is known to contain NKT cells with an activated/memory phenotype, usually observed in adult peripheral blood NKT cells (8,9), although the cellular immune system of the newborn infant is generally known to be immature and hypo-responsive in comparison to the adult system (10). Functionally, cord blood NKT cells do not produce cytokines upon primary stimulation (8), but produce Th1 or Th2 cytokines upon secondary activation (11).

IL-18 is a multifunctional cytokine which augments both innate and acquired immunity and potentiates the Th1 and Th2 reaction (12). IL-18 in combination with IL-12 strongly augments the IFN- $\gamma$  production by T cells (12,13). A high level of IL-18R $\alpha$  expression correlates with the ability of IL-18 to induce IFN- $\gamma$  production (14). The synergistic effect of IL-12 and IL-18 occurs in the absence of TCR ligation and this may represent an alternative pathway for the production of IFN- $\gamma$  during the inflammatory response (12). IL-18 also has the paradoxical capacity to promote the Th2 response under the different cytokine milieu. IL-18 in combination with IL-2 induces the production of Th2 cytokines, such as IL-4 and IL-13, in the T cells (12). In mice, the administration of IL-18 caused the production of Th2 cytokines by V $\alpha$ 14<sup>+</sup> NKT cells (15). However, the effect of IL-18 on human V $\alpha$ 24<sup>+</sup> NKT cells has not yet been investigated.

In the present study, we analyzed phenotypical and functional characteristics of CB V $\alpha$ 24<sup>+</sup> NKT cells and the regulatory mechanism that determines the Th1 and Th2 profile in these cells. We showed that IL-18 in combination with IL-12

---

*Correspondence to:* Dr Yoshihiro Fujimori, Laboratory of Cell Transplantation, Institute for Advanced Medical Sciences, Hyogo College of Medicine, 1-1 Mukogawa-cho, Nishinomiya, Hyogo 663-8501, Japan  
E-mail: fuji-y@hyo-med.ac.jp

**Key words:** natural killer T cells, cord blood, interleukin-18, T helper 1, T helper 2




Figure 1. Analysis of surface phenotype of  $V\alpha 24^+V\beta 11^+$  NKT cells in cord blood and adult blood.  $V\alpha 24^+$  cells were enriched from fresh cord blood and adult peripheral blood by immunomagnetic bead separation, and further subjected to flow cytometrical analysis. (A)  $V\alpha 24^+V\beta 11^+$  cells were gated in the lymphocyte fraction focused on forward scatter and side scatter. (B) CD3 and CD161 expression of  $V\alpha 24^+V\beta 11^+$  gated NKT cells. (C) CD4 and CD8 expression of the CB  $V\alpha 24^+V\beta 11^+$  NKT cells. The data are representative of 10 independent experiments (A-C). Proportion of CD4<sup>+</sup> (D), DN (E) or CD8<sup>+</sup> (F) NKT cells in cord blood (CB) (n=10) and adult peripheral blood (PB) (n=8)  $V\alpha 24^+V\beta 11^+$  NKT cells. The numbers indicate the mean percentages of positive cells of each phenotype in  $V\alpha 24^+V\beta 11^+$  NKT cells.

was able to induce the CB CD4<sup>+</sup>  $V\alpha 24^+$  NKT cells to produce Th1 cytokine, while IL-18 in combination with IL-2 was able to induce Th2 cytokine production in these cells. Therefore, IL-18 regulated the Th1 or Th2 determination by the CB CD4<sup>+</sup>  $V\alpha 24^+$  NKT cells.

## Materials and methods

**Cell preparation and culture.** Umbilical cord blood (CB) was obtained from normal, full-term deliveries and adult peripheral blood was obtained by venous puncture after receiving informed consent. Mononuclear cells (MC) were separated by density gradient centrifugation using Ficoll-Paque (Amersham Pharmacia, Uppsala, Sweden).  $V\alpha 24^+$  cells were obtained by treatment with a phycoerythrin (PE)-conjugated-anti- $V\alpha 24$  antibody (Immunotech, Marseille Cedex, France) and anti-PE-magnetic beads (Miltenyi Biotec, CA, USA) and by sorting with the automated magnetic cell sorter (autoMACS) system (Miltenyi Biotec).  $V\alpha 24^+$  cells were cultured for 14 days in RPMI-1640 medium (Sigma Chemical Co., St. Louis, MO, USA) containing 10% fetal calf serum (Gibco BRL, Grand Island, NY, USA) in the presence of 10 ng/ml human recombinant IL-2 (Pepro Tech Inc., Rocky Hill, NJ) and 10 ng/ml  $\alpha$ -galactosylceramide [ $\alpha$ -GalCer] KRN 7000; kindly provided by Kirin Brewery Co. Ltd., Tokyo, Japan] (16).

**Flow cytometry.**  $V\alpha 24^+$  cells freshly prepared or cultured in the presence of  $\alpha$ -GalCer and IL-2 were stained with various combinations of monoclonal antibodies (Ab), and subjected to a four-color immunofluorescence analysis using an EPICS XL flow cytometer (Beckman Coulter, Miami, FL, USA).

Fluorescein isothiocyanate (FITC)-conjugated anti- $V\beta 11$  Ab (Immunotech), PE- or biotin-conjugated anti- $V\alpha 24$  Ab, phycoerythrin-Texas Red (ECD)-conjugated-anti-CD8 Ab (Beckman Coulter), PC5 conjugated anti-CD3 or CD4 Ab and PE-conjugated-anti-CD161 (Immunotech), and PE-conjugated anti-IL-12R $\beta 1$  and -IL-18R $\alpha$  Ab (R&D Systems, Minneapolis, MN) were used. The intracellular cytokines were stained with PE-conjugated anti-IL-4 Ab and FITC-conjugated anti-IFN- $\gamma$  Ab (Immunotech), after the cells were fixed and permeabilized using an Intra Prep permeabilization kit (Beckman Coulter). Ionomycin (Sigma) and Phorbol-12-myristate-13-acetate (PMA) (Calbiochem, Germany) were used for the augmentation of cytokine production. Brefeldin A (Sigma) was added for 4 h before the end of incubation. Appropriate isotype-matched control Abs were used in all experiments.

**Assay for cytokines produced by NKT cells.** CD4<sup>+</sup>  $V\alpha 24^+$  NKT cells cultured in the presence of IL-2 and  $\alpha$ -GalCer were washed twice and incubated in RPMI-1640 medium in the presence of different combinations of IL-2 (10 ng/ml), IL-12 (10 ng/ml) (Pepro Tech) and IL-18 (10 ng/ml) (Hayashibara Biochemical Laboratories, Inc., Okayama, Japan) for 7 days in 96-well flat-bottom tissue culture plates. The culture supernatants were harvested and assayed for IFN- $\gamma$  and IL-4 using commercially available ELISA kits (Biosource, Camarillo, CA, USA).

**Statistical analysis.** The data are presented as the mean  $\pm$  standard deviation (SD). Student's t-test was used to compare two groups. P values < 0.05 were considered to be statistically significant.




Figure 2. Cytokine production and IL-12R $\beta$ 1/IL-18R $\alpha$  expression in CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells. Immunomagnetic bead-separated V $\alpha$ 24<sup>+</sup> cells were cultured with  $\alpha$ -GalCer and IL-2 for 14 days, which resulted in the preferential expansion of CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells with >95% purity. Intracellular staining for IL-13 and IFN- $\gamma$  in fresh (A) and cultured CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells (B) after stimulation by ionomycin/PMA for 4 h. Expression of IL-12R $\beta$ 1 (C) and IL-18R $\alpha$  (D) on cultured CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells. The data are representative of 4 independent experiments (A-D).

## Results

**Phenotypical analysis of cord blood NKT cells.** We first phenotypically analyzed the V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells that were freshly prepared from the cord blood (CB) and the adult peripheral blood (PB). We enriched the V $\alpha$ 24<sup>+</sup> cells using immunomagnetic beads, and identified the V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> cells by flow cytometry (Fig. 1A). Seventy percent of the CB CD3<sup>+</sup> V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> cells expressed a natural killer marker, CD161 (Fig. 1B). Freshly prepared CB V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells were mostly the CD4<sup>+</sup> single positive (SP) phenotype (Fig. 1C). As shown in Fig. 1D, 96.5 $\pm$ 1.6% of the V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells were positive for CD4, in sharp contrast to the finding that only 21.5 $\pm$ 10.1% of the adult PB NKT cells were positive for CD4 ( $P$ <0.001). In addition, only 3.2 $\pm$ 1.3% of the CB V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells were the DN phenotype (Fig. 1E). In contrast, 76.3 $\pm$ 10.7% of the adult PB V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells were the DN phenotype ( $P$ <0.001). We observed that 1.3 $\pm$ 1.0% of the CB NKT cells and 5.0 $\pm$ 0.9% of the adult PB NKT cells were positive for CD8 (Fig. 1F).

**Cytokine production and receptor expression by CB NKT cells.** We examined the cytokine production in CB NKT cells by intracellular staining. Freshly prepared CB NKT cells did not produce IFN- $\gamma$  or IL-13 upon primary stimulation (Fig. 2A) (8). Culturing CB V $\alpha$ 24<sup>+</sup> cells with IL-2 and  $\alpha$ -GalCer for 14 days yielded >95% CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> V $\beta$ 11<sup>+</sup> NKT cells. The CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells produced both IFN- $\gamma$  and IL-13 upon treatment with PMA/ionomycin (Fig. 2B), indicating that CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells have the capacity to produce both Th1 and Th2 cytokines. We next examined the IL-18 receptor




Figure 3. IFN- $\gamma$ /IL-13 secretion by CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells. Cultured CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells were washed and incubated in the presence of IL-2 or IL-12 with or without IL-18. The culture supernatants were harvested 7 days later and analyzed by ELISA. The data are shown as the mean  $\pm$  SD from 5 separate experiments.

expression on CB NKT cells. Freshly prepared CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells expressed negligible levels of the IL-12 receptor (R)  $\beta$ 1 and low levels of IL-18R $\alpha$  (data not shown). Culturing CB CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells with IL-2 and  $\alpha$ -GalCer for 14 days up-regulated the IL-12R $\beta$ 1 and IL-18R $\alpha$  expression (Fig. 2C and D).

**IL-18 regulates Th1 and Th2 responses of NKT cells.** We examined the effect of IL-18 on the cytokine production in human CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells in the presence of different combinations of IL-12, IL-2 and IL-18, as shown in Fig. 3. When CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells were incubated with IL-12 alone, low levels of IFN- $\gamma$  were produced. Combined stimulation with IL-18 and IL-12 induced markedly high levels of IFN- $\gamma$ . However, IL-13 was not produced by the stimulation with IL-12 alone or IL-12+IL-18. In contrast, IL-18 or IL-2 alone produced low levels of IL-13, and a combination of IL-18 and IL-2 induced markedly high levels of IL-13. These results show that IL-18 differentially regulates human CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells to produce IFN- $\gamma$  and IL-13 in collaboration with IL-12 and IL-2, respectively.

## Discussion

The secretion of Th1 and Th2 cytokines by NKT cells underlies the immuno-regulatory properties of these cells. NKT cells can suppress type 1 diabetes in non-obese diabetic (NOD) mice through the secretion of IL-4 and IL-10 (17). Defects of NKT cells may contribute to the pathogenesis of type 1 diabetes in both NOD mice and humans (6). NKT cells are reported to suppress carcinogenesis through IFN- $\gamma$  (18). Furthermore, the secretion of the Th2 cytokine IL-13 was found to inhibit the immune rejection of tumor grafts (19). These findings suggest that the regulated expression of Th1 and Th2 cytokines by NKT cells may control the outcome of some disease conditions. It is thus important to understand how the production of Th1 and Th2 cytokines by human NKT cells is regulated. In this study, we addressed this issue using human CB V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells.

Although both murine and human NKT cells are restricted by CD1d, there are several phenotypical and functional

differences between murine and human NKT cells and even between human neonatal and adult NKT cells (1,20,21). Murine V $\alpha$ 14<sup>+</sup> NKT cells primarily appear in the thymus as a result of TCR- $\alpha$  rearrangement in CD4 CD8 double-positive (DP) thymocytes. These cells are selected by CD1d in a thymic environment, and matured into CD4<sup>+</sup> single positive (SP) and CD4 CD8 double-negative (DN) V $\alpha$ 14<sup>+</sup> T cells (21). The ratio of mature CD4<sup>+</sup> to DN NKT thymocytes is approximately 6:4 in the murine thymus, which is close to that in the peripheral blood and other tissues. Murine CD4<sup>+</sup> and DN V $\alpha$ 14<sup>+</sup> T cells are functionally identical and depend mainly on IL-15 for their survival and homeostatic expansion in the periphery (21). In humans, in contrast to mice, NKT cells are produced in the thymus with CD4<sup>+</sup> dominance (>90%), and peripheral blood CD4<sup>+</sup> NKT cells are maintained mainly by the thymic output with limited peripheral expansion (20). On the other hand, DN NKT cells undergo extensive peripheral expansion to form a major population in adult peripheral blood (20). In this study, we confirmed the essential phenotypical differences between NKT cells in cord blood and adult blood. DN NKT cells constituted the major population in adult PB V $\alpha$ 24<sup>+</sup>V $\beta$ 11<sup>+</sup> NKT cells, whereas CD4<sup>+</sup> NKT cells were the majority in CB V $\alpha$ 24<sup>+</sup> NKT cells (Fig. 1).

There have been a limited number of reports documenting the mechanism underlying the Th2 versus Th1 cytokine secretion by NKT cells. In mice, altered  $\alpha$ GalCer ligands with shorter sphingosine chains activate the Th2 function (22), while NK1.1 signaling favors the Th1 response (23). Although both murine CD4<sup>+</sup> and DN V $\alpha$ 14<sup>+</sup> T cells are functionally identical, human V $\alpha$ 24<sup>+</sup> NKT cells in the adult peripheral blood consist of two populations with distinct functions: CD4<sup>+</sup> NKT cells that produce both IFN- $\gamma$  and IL-4 and DN NKT cells that produce IFN- $\gamma$  but not IL-4 (24). We showed that cord blood NKT cells were mainly composed of CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells (Fig. 1) and able to produce both Th1 and Th2 cytokines through secondary but not primary stimulation (Fig. 2A and B). Using cord blood NKT cells, Kadowaki *et al* have demonstrated that type 1 myeloid dendritic cells (DC1) preferentially induce CB NKT cells to produce IFN- $\gamma$ , while type 2 lymphoid DC (DC2) preferentially induce IL-4 production (11). In the present study, we showed that the determination toward Th1 or Th2 cytokine production by human CD4<sup>+</sup> NKT cells was regulated by cytokine stimulation. We found that IL-18 was a key cytokine for Th1 or Th2 determination of human NKT cells. In human CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells (Fig. 3), the combined stimulation of IL-18 with IL-12 promoted IFN- $\gamma$  production, while the combined stimulation of IL-18 with IL-2 promoted IL-13 production. This regulation of the Th1/Th2 determination occurred in the absence of TCR engagement. Since the dendritic cells produce IL-12 and IL-18, and DC1 is a high producer of IL-12, our observation that IL-12 plus IL-18 promotes IFN- $\gamma$  production may be consistent with the observation by Kadowaki *et al* that DC1 preferentially induces IFN- $\gamma$  production in CB NKT cells (11).

In conclusion, the present study showed that IL-18 controls the Th1 and Th2 immune response of human CD4<sup>+</sup> V $\alpha$ 24<sup>+</sup> NKT cells through different cytokine combinations. Present results may provide an understanding of the regulation of the

Th1 or Th2 determination of NKT cells which is a critical event for immuno-regulation by these cells.

### Acknowledgements

We are grateful to Ms. Hiromi Takeda for her excellent technical assistance. This study was supported by a research grant for the 'High-Tech Research Center' Project for Private Universities from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and a grant from Research on Human Genome, Tissue Engineering Food Biotechnology of the Ministry of Health, Labor and Welfare of Japan.

### References

1. Bendelac A, Savage PB and Teyton L: The biology of NKT cells. *Annu Rev Immunol* 25: 297-336, 2007.
2. Kim CH, Butcher EC and Johnston B: Distinct subsets of human V $\alpha$ 24-invariant NKT cells: cytokine responses and chemokine receptor expression. *Trends Immunol* 23: 516-519, 2002.
3. Takahashi T, Chiba S, Nieda M, Azuma T, Ishihara S, Shibata Y, Juji T and Hirai H: Analysis of human V  $\alpha$  24+CD8+ NKT cells activated by alpha-galactosylceramide-pulsed monocyte-derived dendritic cells. *J Immunol* 168: 3140-3144, 2002.
4. Takahashi T, Nieda M, Koezuka Y, Nicol A, Porcelli SA, Ishikawa Y, Tadokoro K, Hirai H and Juji T: Analysis of human V  $\alpha$  24+ CD4+ NKT cells activated by alpha-glycosylceramide-pulsed monocyte-derived dendritic cells. *J Immunol* 164: 4458-4464, 2000.
5. Kojo S, Adachi Y, Keino H, Taniguchi M and Sumida T: Dysfunction of T cell receptor AV24AJ18+, BV11+ double-negative regulatory natural killer T cells in autoimmune diseases. *Arthritis Rheum* 44: 1127-1138, 2001.
6. Wilson SB, Kent SC, Patton KT, Orban T, Jackson RA, Exley M, Porcelli S, Schatz DA, Atkinson MA, Balk SP, Strominger JL and Hafler DA: Extreme Th1 bias of invariant V $\alpha$ 24J $\alpha$ Q T cells in type 1 diabetes. *Nature* 391: 177-181, 1998.
7. Crane IJ and Forrester JV: Th1 and Th2 lymphocytes in autoimmune disease. *Crit Rev Immunol* 25: 75-102, 2005.
8. D'Andrea A, Goux D, De Lalla C, Koezuka Y, Montagna D, Moretta A, Dellabona P, Casorati G and Abrignani S: Neonatal invariant V $\alpha$ 24+ NKT lymphocytes are activated memory cells. *Eur J Immunol* 30: 1544-1550, 2000.
9. van Der Vliet HJ, Nishi N, de Gruijl TD, von Blomberg BM, van den Eertwegh AJ, Pinedo HM, Giaccone G and Scheper RJ: Human natural killer T cells acquire a memory-activated phenotype before birth. *Blood* 95: 2440-2442, 2000.
10. Harris DT, Schumacher MJ, Locascio J, Besencon FJ, Olson GB, DeLuca D, Shenker L, Bard J and Boyse EA: Phenotypic and functional immaturity of human umbilical cord blood T lymphocytes. *Proc Natl Acad Sci USA* 89: 10006-10010, 1992.
11. Kadowaki N, Antonenko S, Ho S, Risoan MC, Soumelis V, Porcelli SA, Lanier LL and Liu YJ: Distinct cytokine profiles of neonatal natural killer T cells after expansion with subsets of dendritic cells. *J Exp Med* 193: 1221-1226, 2001.
12. Nakanishi K, Yoshimoto T, Tsutsui H and Okamura H: Interleukin-18 regulates both Th1 and Th2 responses. *Annu Rev Immunol* 19: 423-474, 2001.
13. Tominaga K, Yoshimoto T, Torigoe K, Kurimoto M, Matsui K, Hada T, Okamura H and Nakanishi K: IL-12 synergizes with IL-18 or IL-1 $\beta$  for IFN- $\gamma$  production from human T cells. *Int Immunol* 12: 151-160, 2000.
14. Yoshimoto T, Takeda K, Tanaka T, Ohkusu K, Kashiwamura S, Okamura H, Akira S and Nakanishi K: IL-12 up-regulates IL-18 receptor expression on T cells, Th1 cells, and B cells: synergism with IL-18 for IFN- $\gamma$  production. *J Immunol* 161: 3400-3407, 1998.
15. Yoshimoto T, Min B, Sugimoto T, Hayashi N, Ishikawa Y, Sasaki Y, Hata H, Takeda K, Okumura K, Van Kaer L, Paul WE and Nakanishi K: Nonredundant roles for CD1d-restricted natural killer T cells and conventional CD4<sup>+</sup> T cells in the induction of immunoglobulin E antibodies in response to interleukin 18 treatment of mice. *J Exp Med* 197: 997-1005, 2003.



SPANDIDOSashi E, Motoki K, Uchida T, Fukushima H and Koezuka Y:  
PUBLICATIONS)00, a novel immunomodulator, and its antitumor activities.  
ONCOL Res 7: 529-534, 1995.

17. Hammond KJ, Poulton LD, Palmisano LJ, Silveira PA, Godfrey DI and Baxter AG: alpha/beta-T cell receptor (TCR)+CD4-CD8-(NKT) thymocytes prevent insulin-dependent diabetes mellitus in nonobese diabetic (NOD)/Lt mice by the influence of interleukin (IL)-4 and/or IL-10. *J Exp Med* 187: 1047-1056, 1998.
18. Smyth MJ, Thia KY, Street SE, Cretney E, Trapani JA, Taniguchi M, Kawano T, Pelikan SB, Crowe NY and Godfrey DI: Differential tumor surveillance by natural killer (NK) and NKT cells. *J Exp Med* 191: 661-668, 2000.
19. Terabe M, Matsui S, Noben-Trauth N, Chen H, Watson C, Donaldson DD, Carbone DP, Paul WE and Berzofsky JA: NKT cell-mediated repression of tumor immunosurveillance by IL-13 and the IL-4R-STAT6 pathway. *Nat Immunol* 1: 515-520, 2000.
20. Baev DV, Peng XH, Song L, Barnhart JR, Crooks GM, Weinberg KI and Metelitsa LS: Distinct homeostatic requirements of CD4+ and CD4- subsets of Valpha24-invariant natural killer T cells in humans. *Blood* 104: 4150-4156, 2004.
21. Gapin L, Matsuda JL, Surh CD and Kronenberg M: NKT cells derive from double-positive thymocytes that are positively selected by CD1d. *Nat Immunol* 2: 971-978, 2001.
22. Miyamoto K, Miyake S and Yamamura T: A synthetic glycolipid prevents autoimmune encephalomyelitis by inducing TH2 bias of natural killer T cells. *Nature* 413: 531-534, 2001.
23. Exley M, Porcelli S, Furman M, Garcia J and Balk S: CD161 (NKR-P1A) costimulation of CD1d-dependent activation of human T cells expressing invariant V alpha 24 J alpha Q T cell receptor alpha chains. *J Exp Med* 188: 867-876, 1998.
24. Lee PT, Benlagha K, Teyton L and Bendelac A: Distinct functional lineages of human V(alpha)24 natural killer T cells. *J Exp Med* 195: 637-641, 2002.