

Cultured rat calvarial osteoblast-like cells are provided with orexin type 1 receptors

AGNIESZKA ZIOLKOWSKA¹, MARCIN RUCINSKI¹, CINZIA TORTORELLA²,
MARIANNA TYCZEWSKA¹, GASTONE G. NUSSDORFER² and LUDWIK K. MALENDOWICZ¹

¹Department of Histology and Embryology, School of Medicine, Karol Marcinkowski University of Medical Sciences, PL-60781 Poznan, Poland; ²Department of Human Anatomy and Physiology, Section of Anatomy, School of Medicine, University of Padua, Via Gabelli 65, I-35121 Padua, Italy

Received July 19, 2007; Accepted August 24, 2007

Abstract. Orexins A and B are hypothalamic peptides which are derived from the proteolytic cleavage of prepro-orexin and act via two subtypes of receptors, named OX1-R (that almost exclusively binds orexin-A) and OX2-R (nonselective for both orexins). Several lines of evidence show that other neuropeptides, which like orexins are involved in the central control of energy homeostasis (e.g. leptin and ghrelin), may play a role in the regulation of bone metabolism, acting via autocrine-paracrine or endocrine routes. Therefore, we studied by reverse transcription-polymerase chain reaction (RT-PCR) the expression of the orexin system in rat calvarial osteoblast-like (ROB) cells, whose osteoblastic lineage was immunocytochemically demonstrated by their osteonectin and collagen-1 α content at day 14 of culture. Conventional PCR detected the mRNA expression of OX1-R, but not OX2-R and prepro-orexin in ROB cells at days 2, 7 and 21 of culture. Semiquantitative real time-PCR evidenced a gradual down-regulation of OX1-R mRNA in relation to the duration of culture. This novel finding suggests that rat osteoblasts could be a target for circulating orexin-A, especially during their early stages of differentiation into mature osteoblasts.

Introduction

Orexins A and B are hypothalamic neuropeptides that stimulate both food intake and energy expenditure, and play a role in sleep-wakefulness regulation and modulation of hypothalamic-pituitary-adrenal axis activity. Orexins originate from the proteolytic cleavage of prepro-orexin and act through two subtypes of receptors, named OX1-R and OX2-R.

OX1-R almost exclusively binds orexin-A, while OX2-R is nonselective for both orexins (1-5).

Evidence has been provided that hypothalamic orexinergic neurons are regulated by hormonal signals, including ghrelin and leptin (6-8). Available findings indicate that certain neuropeptides involved in energy homeostasis, such as leptin and ghrelin, may have a role in the regulation of bone metabolism, their receptors being expressed in osteoblasts (see Discussion). However, there are no data available concerning the expression of orexin system components in bone cells. Hence, we investigated the expression of prepro-orexin and OX1-R and OX2-R mRNAs in cultured rat calvarial osteoblast-like (ROB) cells.

Materials and methods

Animals and reagents. Female Wistar rats (final body weight 100-110 g) were kept under a 14-h light:10-h dark cycle (illumination onset at 6.00 a.m.) at 23°C, and maintained on a standard diet and tap water *ad libitum*. The study protocol was approved by the local Ethics Committee for Animal Studies. Rabbit anti-onectin and mouse anti-collagen-1 α primary polyclonal antibodies were purchased from Acris Antibodies (Herford, Germany), and goat anti-rabbit MFP488 (emission wavelength, 524 nm) and anti-mouse MFP590 (emission wavelength, 624 nm) secondary antibodies from MoBi Tech (Göttingen, Germany). 4',6'-diamine-2'-phenylindole (DAPI) was obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Dulbecco's modified Eagle's medium (DMEM), fetal calf serum (FCS), bovine serum albumin (BSA), phosphate-buffered saline (PBS), and all other laboratory reagents were provided by Sigma-Aldrich Corp. (St. Louis, MO).

ROB cell culture. The technique used was described by Boden *et al* (9), with few modifications (10). Briefly, calvarias of 8 rats were placed in DMEM, and the connective tissue was removed. Calvarias were then cut into small fragments, which were dissociated to cell suspensions by enzymatic digestion with 0.1% collagenase-I for 30 min at 37°C. ROB cells were harvested by centrifugation and re-suspended in DMEM supplemented with NaHCO₃, 6% FCS and antibiotic-

Correspondence to: Professor G.G. Nussdorfer, Department of Human Anatomy and Physiology, Section of Anatomy, School of Medicine, University of Padua, Via Gabelli 65, I-35121 Padua, Italy
E-mail: gastone.nussdorfer@unipd.it

Key words: orexins, orexin receptors, calvarial osteoblast-like cells, rat




Figure 1. Osteonectin (A) and collagen-1 α immunoreactivity (B) in ROB cells cultured for 14 days, as demonstrated by immunofluorescence. Nuclei were stained with DAPI. Magnification as defined by bars, 10 μ m.

antimycotic solution. Cells were then plated in culture dishes (10^4 cells/dish), and cultured for up to 21 days at 37°C in a humidified atmosphere of 95% air-5% CO₂; the medium being changed every 24 h.

Immunocytochemistry. ROB cells cultured for 14 days were fixed in 4% paraformaldehyde for 10 min at room temperature, washed in PBS and then preincubated with 5% BSA for 30 min. Cells were incubated with the primary anti-osteonectin and anti-collagen-1 α antibodies (1:400 dilution) for 18 h at 4°C, and then, after washing in PBS, with the secondary fluorescent antibodies (1:1,000 dilution) for 60 min at room temperature. Control specimens were obtained using primary antibodies preabsorbed with antigen excess (11). Slides were washed in PBS, nuclei were stained with DAPI, and specimens were observed and photographed using a Nikon Eclipse E600 microscope.

Reverse transcription (RT)-polymerase chain reaction (PCR). Total RNA was extracted from ROB cells cultured for 2, 7 and 21 days, as well as from the hypothalamus and anterior pituitary of two rats, and reverse transcribed to cDNA (12-14). Conventional PCR was carried out in a Perkin-Elmer 480 DNA Thermal Cycler (Perkin-Elmer Life Sciences, Milan, Italy), using the amplification program previously described (15) and the primers indicated in the legend of Fig. 2. The specificity of the PCR was verified by sequencing analysis (16). Semiquantitative real time-PCR of OX1-R mRNA was performed in a Roche Light Cycler 2.0 with software version 4.0, as detailed previously (17-19). The following program was used: a predenaturation step to activate Taq DNA polymerase (95°C for 10 min), followed by a three-step amplification (denaturation at 95°C for 10 sec; annealing at 56°C for 5 sec; and extension at 72°C for 2 sec). Subsequently, a melting curve (60-90°C with a heating rate of 0.1°C/sec) was carried out to check the specificity of amplification and the presence of by-products. All samples (n=6) were amplified in duplicate, and data were normalized in relation to the housekeeping gene hypoxanthine phosphoribosyl-transferase 1 (HPRT) and expressed as the means \pm SEM.




Figure 2. Ethidium bromide-stained 2% agarose gel showing cDNA amplified with rat specific primers from RNA of exemplary ROB cells cultured for 2 days (lanes 2 and 3), 7 days (lanes 4 and 5) and 21 days (lanes 6 and 7). Positive controls for prepro-orexin (hypothalamus) and other genes (anterior pituitary) are shown in lanes 8 and 9. Primer sequences were: prepro-orexin sense (83-103) CAGACACCATGAACCTTCCTT, antisense (191-208) CAGAGCAGTCGGGCAGAG and probe (148-155) CTGCTGCC (amplicon 126 bp; accession no. NM013179); OX1-R sense (1476-1495) TGCTGGTTTTTGCACCTCTGT, antisense (1530-1549) GCGAAACATCCCAAACACTC and probe (1498-1505) TCTGCCCA (amplicon, 74 bp; accession no. NM013064); OX2-R sense (782-806) GGCTTATCTCCAAA TATTCCGTAAA, antisense (828-850) CTCTGAACCACAGAAGAAGT TCC and probe (814-821) GCAGACAG (amplicon, 69 bp; accession no. NM013074); and HPRT sense (391-412) CAGTCAACGGGGGACATAAA AG, antisense (515-536) ATTTTGGGGCTGTACTGCTTGA and probe (417-424) TGGTGGAG (amplicon, 146 bp; accession no. NM012583). Lane 1 was loaded with 200 ng of a DNA size marker (O'Range Ruler 50 bp DNA Ladder, MBI Fermentas, Vilnius, Lithuania).

Results

Immunocytochemistry revealed the expression of both osteonectin and collagen-1 α in all ROB cells at 14 days of culture (Fig. 1).

Conventional PCR did not detect prepro-orexin and OX2-R mRNA expression in ROB cells, but detected it in the rat hypothalamus and anterior pituitary, respectively. Conversely, OX1-R mRNA was observed in ROB cells at days 2, 7 and 21 of culture (Fig. 2). Real time-PCR evidenced a gradual decrease in OX1-R mRNA expression in ROB cells in relation to the duration of culture (Fig. 3).




Figure 3. Relative OX1-R mRNA expression (normalized to HPRT) in ROB cells cultured for 2, 7 and 21 days. Bars are the means \pm SEM (n=6).

Discussion

ROB cell primary culture is one of the most common *in vitro* models used to investigate specific gene events associated with osteoblast proliferation, differentiation and mineralization of extracellular matrix. Cultured ROB cells spontaneously differentiate into osteoblasts, and this process is coupled to the increased expression of both collagen and non-collagenous bone proteins, including osteonectin, osteocalcin and osteopontin (9,10,14,20-24). Our immunofluorescence findings showed that all cultured ROB cells expressed both osteonectin and collagen-1 α , suggesting their osteoblastic lineage.

Consistent findings indicate that both bone formation and remodeling are regulated by several biologically active peptides, that act via endocrine, paracrine and autocrine mechanisms. Such peptides include substance-P, VIP, CGRP and neuropeptide-Y (25,26), as well as leptin and the endogenous ligand of growth hormone secretagogue (GHS) receptor ghrelin, which, like orexins, play a major role in the central control of energy homeostasis. Evidence has been provided that leptin is expressed in and secreted from human osteoblasts in primary culture (27-29), and that functionally active leptin-receptor isoforms are present in human osteoblast-derived cell lines and rat osteoblasts (30,31). Leptin was found to enhance human marrow stromal cell differentiation into osteoblasts (32), to stimulate proliferation of cultured osteoblasts, and to promote matrix production and mineralization (27-29). In contrast, *in vivo* studies evidenced an inhibitory effect of leptin on bone formation (33). Ghrelin and its GHS-R1a have been shown to be expressed in human and rat osteoblasts, and ghrelin has been reported to stimulate *in vitro* their proliferation and differentiation (34-36).

Our study demonstrated that ROB cells express OX1-R mRNA, but not prepro-orexin and OX2-R mRNAs. This novel finding may suggest that rat osteoblasts could be a target for circulating orexin A, for which OX1-R is selective. Moreover, real time-PCR reveals a time-dependent down-regulation of OX1-R expression in cultured ROB cells, which could be related to the stage of their differentiation into mature osteoblasts. Further studies are underway to elucidate the physiological relevance of these findings and to

ascertain whether orexins may be included in that group of neuropeptides regulating bone metabolism.

References

- De Lecea L, Kilduff T, Peyron C, Gao X, Foye P, Danielson P, Fukuhara C, Battenberg E, Gautvik V, Bartlett F, Frankel W, Van den Pol A, Bloom F, Gautvik K and Sutcliffe J: The hypocretins: hypothalamus specific peptides with neuro-excitatory activity. *Proc Natl Acad Sci USA* 95: 322-327, 1998.
- Beuckmann CT and Yanagisawa M: Orexins: from neuropeptides to energy homeostasis and sleep/wake regulation. *J Mol Med* 80: 329-342, 2002.
- Kukkonen JP, Holmqvist T, Ammoun S and Akerman KEO: Functions of the orexinergic/hypocretinergic system. *Am J Physiol* 283: C1567-C1591, 2002.
- Sakurai T: Reverse pharmacology of orexin: from an orphan GPCR to integrative physiology. *Regul Pept* 126: 3-10, 2005.
- Spinazzi R, Andreis PG, Rossi GP and Nussdorfer GG: Orexins in the regulation of the hypothalamic-pituitary-adrenal axis. *Pharmacol Rev* 58: 46-57, 2006.
- Cai XJ, Evans ML, Lister CA, Lesie RA, Arch JR, Wilson S and Williams G: Hypoglycemia activates orexin neurons and selectively increases hypothalamic orexin-B levels: responses inhibited by feeding and possibly mediated by the nucleus of the solitary tract. *Diabetes* 50: 105-112, 2000.
- Funahashi H, Hori T, Shimoda Y, Mizushima H, Ryushi T, Katoh S and Shioda S: Morphological evidence for neural interactions between leptin and orexin in the hypothalamus. *Regul Pept* 92: 31-35, 2000.
- Toshinai K, Date Y, Murakami N, Shimada M, Mondal MS, Shimbara T, Guan JL, Wang QP, Funahashi H, Sakurai T, Shioda S, Matsukura S, Kangawa K and Nakazato M: Ghrelin-induced food intake is mediated via the orexin pathway. *Endocrinology* 144: 1506-1512, 2003.
- Boden SC, McCuaig K, Hair G, Racine M, Titus L, Wozney JM and Nanes MS: Differential effects and glucocorticoid potentiation of bone morphogenetic protein action during rat osteoblast differentiation *in vitro*. *Endocrinology* 137: 3401-3407, 1996.
- Ziolkowska A, Rucinski M, Pucher A, Tortorella C, Nussdorfer GG and Malendowicz LK: Expression of osteoblast marker genes in rat calvarial osteoblast-like cells, and effects of endocrine disruptors diphenylolpropane, benzophenone-3, resveratrol and silymarin. *Chem Biol Interact* 164: 147-156, 2006.
- Belloni AS, Petrelli L, Guidolin D, De Toni R, Bova S, Spinazzi R, Gerosa G, Rossi GP and Nussdorfer GG: Identification and localization and adrenomedullin-storing cardiac mast cells. *Int J Mol Med* 17: 709-713, 2006.
- Albertin G, Carraro G and Nussdorfer GG: Human adrenomedullin gene silencing by short interfering RNAs: A preliminary study. *Int J Mol Med* 15: 579-583, 2005.
- Albertin G, Carraro G, Petrelli L, Guidolin D, Neri G and Nussdorfer GG: Endothelin-1 and adrenomedullin enhance the growth of human adrenocortical carcinoma-derived SW-13 cell line by stimulating proliferation and inhibiting apoptosis. *Int J Mol Med* 15: 469-474, 2005.
- Rucinski M, Ziolkowska A, Hochól A, Pucher A, Macchi C, Belloni AS, Nussdorfer GG and Malendowicz LK: Estradiol and resveratrol stimulating effect on osteocalcin, but not osteonectin and collagen-1 α gene expression in primary culture of rat calvarial osteoblast-like cells. *Int J Mol Med* 18: 565-570, 2006.
- Spinazzi R, Ziolkowska A, Neri G, Nowak M, Rebuffat P, Nussdorfer GG, Andreis PG and Malendowicz LK: Orexins modulate the growth of cultured rat adrenocortical cells, acting through type 1 and type 2 receptors coupled to the MAPK p42/p44- and p38-dependent cascades. *Int J Mol Med* 15: 847-852, 2005.
- Rossi GP, Sticchi D, Giuliani L, Bernante P, Zavattero S, Pessina AC and Nussdorfer GG: Adiponectin receptor expression in the human adrenal cortex and aldosterone-producing adenomas. *Int J Mol Med* 17: 975-980, 2006.
- Albertin G, Ruggero M, Guidolin D and Nussdorfer GG: Gene silencing of human RAMP2 mediated by short-interfering RNA. *Int J Mol Med* 18: 531-535, 2006.
- Carraro G, Albertin G, Aragona F, Forneris M, Casale V, Spinazzi R and Nussdorfer GG: Age-dependent decrease in the ghrelin gene expression in the human adrenal cortex: A real-time PCR study. *Int J Mol Med* 17: 319-321, 2006.

19. Ziolkowska A, Tortorella C, Nussdorfer GG, Rucinski M, Majchrzak M and Malendowicz LK: Accumulation of steroidogenic acute regulatory protein mRNA and decrease in the secretory and proliferative activity of rat adrenocortical cells in the presence of proteasome inhibitors. *Int J Mol Med* 17: 865-868, 2006.
20. Aubin JE, Liu F, Malaval L and Gupta AK: Osteoblast and chondroblast differentiation. *Bone* 17: 77S-83S, 1995.
21. Pockwinse SM, Stein JL, Lian JG and Stein GS: Developmental stage-specific cellular responses to vitamin-D and glucorticoids during differentiation of the osteoblast phenotype: inter-relationship of morphology and gene expression by *in situ* hybridization. *Exp Cell Res* 216: 244-260, 1995.
22. Plant A and Tobias JH: Characterization of the temporal sequence of osteoblast gene expression during estrogen-induced osteogenesis in female mice. *J Cell Biochem* 82: 683-691, 2001.
23. Hinoi E, Fujimori S, Takemori A and Yoneda Y: Cell death by pyruvate deficiency in proliferative cultured calvarial osteoblasts. *Biochem Biophys Res Commun* 294: 1177-1183, 2002.
24. Tsukamoto Y, Ishihara Y, Miyagawa-Tomita S and Hagiwara H: Inhibition of ossification *in vivo* and differentiation of osteoblasts *in vitro* by tributyltin. *Biochem Pharmacol* 68: 739-746, 2004.
25. Manolagas SC: Birth and death of bone cells: basic regulatory mechanisms and implications for the pathogenesis and treatment of osteoporosis. *Endocr Rev* 21: 115-137, 2000.
26. Elefteriou F: Neuronal signaling and the regulation of bone remodeling. *Cell Mol Life Sci* 62: 2339-2349, 2005.
27. Reseland JE, Syversen U, Bakke I, Qvigstad G, Eide LG, Hjertner O, Gordeladze JO and Drevon CA: Leptin is expressed in and secreted from primary cultures of human osteoblasts and promotes bone mineralization. *J Bone Miner Res* 16: 1426-1433, 2001.
28. Gordeladze JO, Drevon CA, Syversen U and Reseland JE: Leptin stimulates human osteoblastic cell proliferation, *de novo* collagen synthesis, and mineralization: impact on differentiation markers, apoptosis, and osteocalcin signaling. *J Cell Biochem* 85: 825-836, 2002.
29. Iwamoto I, Fujino T and Douchi T: The leptin receptor in human osteoblasts and the direct effect of leptin on bone metabolism. *Gynecol Endocrinol* 19: 97-104, 2004.
30. Cornish J, Callon KE, Bava U, Lin C, Naot D, Hill BL, Grey AB, Broom N, Myers DE, Nicholson GC and Reid IR: Leptin directly regulates bone cell function *in vitro* and reduces bone fragility *in vivo*. *J Endocrinol* 175: 405-415, 2002.
31. Lee YJ, Park JH, Ju SK, You KH, Ko JS and Kim HM: Leptin receptor isoform expression in rat osteoblasts and their functional analysis. *FEBS Lett* 528: 43-47, 2002.
32. Thomas T, Gori F, Khosla S, Jensen MD, Burguera B and Riggs BL: Leptin acts on human marrow stromal cells to enhance differentiation to osteoblasts and to inhibit differentiation to adipocytes. *Endocrinology* 140: 1630-1638, 1999.
33. Takeda S: Central control of bone remodeling. *Biochem Biophys Res Commun* 328: 697-699, 2005.
34. Fukushima N, Hanada R, Teranishi H, Fukue Y, Tachibana T, Ishikawa H, Takeda S, Takeuchi Y, Fukumoto S, Kangawa K, Nagata K and Kojima M: Ghrelin directly regulates bone formation. *J Bone Miner Res* 20: 790-798, 2005.
35. Maccarinelli G, Sibilia V, Torsello A, Raimondo F, Pitto M, Giustina A, Netti C and Cocchi D: Ghrelin regulates proliferation and differentiation of osteoblastic cells. *J Endocrinol* 184: 249-256, 2005.
36. Delhanty PJ, Van der Eerden BC, Van der Velde M, Gauna C, Pols HA, Jahr H, Chiba H, Van der Lely AJ and Van Leeuwen JP: Ghrelin and unacylated ghrelin stimulate human osteoblast growth via mitogen-activated protein kinase (MAPK)/ phosphoinositide 3-kinase (PI3K) pathways in the absence of GHS-R1a. *J Endocrinol* 188: 37-47, 2006.