

Effect of the release from mechanical stress on osteoclastogenesis in RAW264.7 cells

KENJIRO SHIBATA^{1,2}, YOSHITAKA YOSHIMURA², TAKASHI KIKUIRI³, TOMOKAZU HASEGAWA⁴,
YUMI TANIGUCHI³, YOSHIKI DEYAMA², KUNIAKI SUZUKI² and JUNICHIRO IIDA¹

Departments of ¹Orthodontics, ²Molecular Cell Pharmacology and ³Pediatric Dentistry, Hokkaido University Graduate School of Dental Medicine, Kita-ku, Sapporo 060-8586; ⁴Department of Pediatric Dentistry, School of Dentistry, Iwate Medical University, Morioka 020-8505, Japan

Received November 12, 2010; Accepted December 30, 2010

DOI: 10.3892/ijmm.2011.675

Abstract. The effects of mechanical stress release on osteoclastogenesis may be as important as those of mechanical stress application. However, the direct effects of mechanical stress on the behavior of osteoclasts has not been thoroughly investigated and there is limited information on the results of the release from mechanical stress. In this study, the effects of mechanical stress application and its release on osteoclast differentiation were examined. The number of tartrate-resistant acid phosphatase (TRAP)-positive multinucleated osteoclasts derived from RAW264.7 cells were measured and the expression of osteoclast differentiation genes, which was altered in response to the release from mechanical stress according to the Flexercell tension system was evaluated by real-time PCR. Osteoclast differentiation and fusion were suppressed by mechanical stress application and were rapidly induced after mechanical stress release. The mRNA expression of the osteoclast specific genes, TRAP, matrix metalloproteinase-9 (MMP-9), cathepsin-K (cath-k), calcitonin receptor (CTR), ATPase H⁺ transporting vacuolar proton pump member I (ATP6i), chloride channel-7 (ClC7) and dendritic cell-specific transmembrane protein (DC-STAMP) was decreased with mechanical stress application, and increased up to 48 h after the release from it. These alterations in gene mRNA expression were associated with the number of osteoclasts and large osteoclasts. Inducible nitric oxide synthetase (iNOS) mRNA was increased with mechanical stress and decreased after its release. Nitric oxide (NO) production was increased with mechanical stress. Nuclear factor of activated T cells cytoplasmic (NFATc) family mRNAs were not altered with mechanical stress, but were up-regulated up to 48 h after the release from it. These findings indicate that the suppression of osteoclast differentia-

tion and fusion induced by mechanical stress is the result of NO increase via iNOS, and that the promotion of osteoclast differentiation and fusion after the release from mechanical stress is related to the NFATc family genes, whose expression remained constant during mechanical stress but was up-regulated after the release from mechanical stress.

Introduction

Bone homeostasis is maintained by the balance between bone formation and bone resorption (1). Tipping the balance in favor of osteoclasts leads to diseases with a low bone mass, including osteoporosis, whereas impaired osteoclastic bone resorption results in diseases with high bone mass, including osteopetrosis (2). The relationship between the immune and skeletal systems has gradually become clearer (3,4). Accumulating data on the molecular mechanisms of osteoclast differentiation have demonstrated that the receptor activator of nuclear factor- κ B (RANK)-ligand (RANKL) pathway and the macrophage colony-stimulating factor (M-CSF) are necessary and sufficient for the differentiation of osteoclasts (5,6). In addition, the role of the nuclear factor of activated T cells cytoplasmic (NFATc) 1 gene, a master switch for the regulation of the terminal differentiation of osteoclasts which functions downstream of RANKL, has been demonstrated both *in vitro* and *in vivo* (7-9). NFATc1 expression is dependent on both the TRAF6/NF- κ B and the c-Fos pathways. The NFAT family of transcription factors was originally discovered in T cells of the immune system, but they are involved in the regulation of various biological systems including the Ca²⁺-regulated pathway (9).

Mechanical stress can largely affect this balanced bone metabolism. Bone adjusts to load by changing its mass, shape or microarchitecture (10). It has been reported that removing the load results in bone loss (11), whereas application of a load causes bone apposition (12). However, the precise mechanism by which loading is transduced into cellular control of bone remodeling is unknown. A number of studies *in vitro* and *in vivo* have described the effects of mechanical stress on the bone cells. To investigate the influence of mechanical stress on the cells, several *in vitro* models have been established with varying methods, including sheer stress

Correspondence to: Dr Yoshitaka Yoshimura, Hokkaido University Graduate School of Dental Medicine, Kita 13 Nishi 7, Kita-ku, Sapporo 060-8586, Japan
E-mail: yoshi@den.hokudai.ac.jp

Key words: mechanical stress, osteoclast, nitric oxide

(fluid flow) (13,14) compressive (15,16) and tensile force (17), hydrostatic pressure (18), microgravity (19), and so on (20,21). Furthermore, in addition to an *in vivo* experimental system including transgenic mice, many studies have indicated the effect of mechanical stress on the bone's adaptive remodeling behavior in osteocytes (22), osteoblasts (23) and in a co-culture between osteoblasts and osteoclasts (24). Several studies have shown that nitric oxide (NO) acts as a mediator of mechanically induced bone remodeling (10,25,26). However, the direct effect of mechanical stress on the behavior of osteoclasts (27) has only been investigated in a few studies and there is limited information on the effects of the release from mechanical stress, which thus became the focus of the present study.

The aim of this study was to elucidate which osteoclast differentiation genes are altered in response to the release from mechanical stress and whether such an alteration is mediated by loading-induced NO production. To validate that NO may modulate the differentiation of osteoclasts in response to mechanical stress, we inhibited the release of NO by an NO synthetase (NOS) inhibitor.

Materials and methods

Cell culture and the Flexercell tension system. We used the murine monocyte/macrophage cell line RAW264.7 cells (RAW cells) (ATCC no. TIB-71™, USA) as osteoclast precursors. The cells were grown to subconfluence in 100 mm standard dishes (Falcon™, Becton-Dickinson, Franklin Lakes, NJ, USA) with Dulbecco's modified Eagle's medium (DMEM) (Wako Pure Chemical, Japan) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Invitrogen, MD, USA) and kanamycin sulfate (Meiji Seika, Japan) at 37°C in a humidified atmosphere of 95% air and 5% CO₂. Subsequently, RAW cells (3x10⁴ cells/well) were transferred to Flexcell 6-well culture plates with a type I collagen-coated and flexible-bottomed well (Flexcell Corp., Hillsborough, NC, USA), and cultured with α -MEM (Wako Pure Chemical) supplemented with 10% heat-inactivated FBS, 50 ng/ml RANKL (Oriental Yeast, Japan), 2 mM L-alanyl-L-glutamine (Sigma-Aldrich, St. Louis, MO, USA), 284 μ M L-ascorbic acid 2-phosphate (Sigma-Aldrich) and 66.7 μ g/ml kanamycin sulfate at 37°C in a humidified atmosphere of 95% air and 5% CO₂. RAW cells differentiated into TRAP-positive multinucleated osteoclastic cells in the presence of RANKL (27). Mechanical stress was applied to RAW cells with a Flexercell tension system (28). In this device, a flexible silicon membrane is stretched across a loading post by the application of a vacuum pressure. The experimental regimens used in this study delivered 10% elongation at 30 cycles per min (0.5 Hz) (29). Mechanical stress was applied for a period of 48 h over 3-4 days (3-4 MS). After the termination of the mechanical stress application, culturing of the RAW cells continued for the indicated periods. The culture medium was replaced on days 2, 3, 5 and 7 of culturing.

Tartrate-resistant acid phosphatase (TRAP) staining. The cells cultured for a given period were fixed with 10% neutral formalin. They were then washed with distilled water and stained in the TRAP staining solution (pH 5.0) with Fast Red Violet LB salt (Sigma-Aldrich) (30). TRAP-positive

Table I. The assay ID of the primers used.

Gene	Assay ID
RANK	Mm00475698_m1
<i>c-fms</i>	Mm00432689_m1
OSCAR	Mm00558663_m1
NFATc1	Mm00479445_m1
NFATc2	Mm00477776_m1
NFATc3	Mm01249194_m1
IFN- β	Mm00439546_s1
iNOS	Mm00440485_m1
nNOS	Mm00435175_m1
TRAP	Mm00475698_m1
CTR	Mm00432271_m1
MMP-9	Mm00432271_m1
cath-k	Mm00484036_m1
CIC7	Mm00442400_m1
ATP6i	Mm00469395_g1
DC-STAMP	Mm01168058_m1
GAPDH	Mm99999915_g1

cells with >2 nuclei under a light microscope were counted as osteoclasts. Osteoclasts with >2 nuclei were distinguished from large osteoclasts with ≥ 8 nuclei (27). The control culture was grown under the same condition without mechanical stress. The counted range was 1 cm² dividing the circle in a fan-shape.

Reverse transcription-polymerase chain reaction and real time PCR. Total RNA was extracted using the TRIzol reagent isolation kit (Invitrogen) (31). Reverse transcription was performed with 1 μ g of total RNA in a total volume of 20 μ l per reaction. cDNA was amplified by Rever Tra Ace- α FSK-101 (Toyobo, Japan). The following primers (Applied Biosystems, USA) were used: RANK, M-CSF receptor (*c-fms*), osteoclast-associated receptor (OSCAR), TRAP, matrix metalloproteinase-9 (MMP-9), cathepsin-K (cath-k), calcitonin receptor (CTR), dendritic cell-specific transmembrane protein (DC-STAMP) ATPase H⁺ transporting vacuolar proton pump member I (ATP6i), chloride channel-7 (CIC7), NFATc1, NFATc2, NFATc3, inducible NOS (iNOS) neuronal NOS (nNOS) and interferon- β (IFN- β) as shown in Table I. As an internal control for RNA quantity, the same cDNA was amplified using primers specific for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA. All amplification reactions were performed in triplicate. The ABI Prism 7300 sequence detection system (Applied Biosystems) (32) was used for real-time PCR analysis. At the end of each PCR run, the data were automatically analyzed by the system and amplification plots were obtained. The $\Delta\Delta$ Ct method was used to calculate the real-time PCR results (amount of target = 2^{- $\Delta\Delta$ Ct}).

Inhibition of nitric oxide production. NO release was inhibited by the addition of N^G-Nitro-L-arginine methyl ester, hydrochloride (L-NAME), (Dojindo, Japan). L-NAME was




Figure 1. The number of TRAP-positive multinucleated osteoclasts was suppressed with mechanical stress, and then was increased after mechanical stress release through osteoclast differentiation. Mechanical stress was applied for 48 h over 3-4 days, but not to control. After the release from mechanical stress, TRAP-positive multinucleated osteoclasts were counted until 72 h every 24 h. (A) The number of TRAP-positive multinucleated cells. (B) The number of large osteoclasts with ≥ 8 nuclei. Magnification $\times 100$. Results are shown as mean \pm SD (n=12).

dissolved in PBS and added to the culture medium (final concentration, 100 μ M) (33).

Nitrite and nitrate measurements. The NO metabolite (NO₂ and NO₃) levels were analyzed in the conditioned medium by means of a colorimetric assay kit: NO₂/NO₃ Assay kit-C II (Dojindo). Samples were read at 540 nm with a Spectra Microplate Auto reader (Bio-Rad Model 680, Hercules, CA, USA).

Statistics. All results are given as mean \pm SD. Comparisons between 2 groups were analyzed using the 2-tailed unpaired Student's t-test. P-values of <0.05 were considered statistically significant.

Results

The number of osteoclasts increased rapidly after the release from mechanical stress. The effect of mechanical stress and the release from mechanical stress on osteoclast differentiation was examined. Mechanical stress was applied for 48 h over 3-4 days. The number of TRAP-positive multinucleated osteoclasts was assessed every 24 h after the release from mechanical stress up to 72 h. The control group was grown under the same conditions without mechanical stress. Mechanical stress caused suppression in the number of osteoclasts to less than one third of that in the control group (Fig. 1A). The mechanical stress-induced decrease in the number of large osteoclasts with ≥ 8 nuclei (Fig. 1B) was more pronounced than that in the osteoclasts with >2 nuclei. On the other hand, osteoclast differentiation and fusion were rapidly promoted after the release from mechanical stress and exceeded the control group 48 h after the release from it (Fig. 1).

The effects of mechanical stress application and its release on the expression of osteoclast differentiation genes. We investigated how the expression of osteoclast differentiation genes was altered by mechanical stress application and its release, using a real-time PCR analysis (Fig. 2) for osteoclast

specific genes (TRAP, CTR, MMP-9, cath-k, CIC7, ATP6i), which are differentiation markers detected after the formation of mature osteoclasts, and for DC-STAMP. The expression of these genes was decreased in the mechanical stress group, and increased after the release from mechanical stress up to 48 h compared to the control group. The expression of iNOS mRNA was increased in the mechanical stress group, and then decreased after the release from mechanical stress compared to the control group. On the other hand, the expression of nNOS mRNA was slightly decreased in the mechanical stress group, and remained fairly constant after the release from mechanical stress compared to the control group. We could not detect eNOS mRNA expression in these osteoclastic RAW cells (data not shown). The expression of IFN- β mRNA was significantly decreased in the mechanical stress group, and then continued decreasing after the release from mechanical stress compared to the control group. The mRNA expression of NFATc family gene members was not altered during mechanical stress application, and was up-regulated after the release from mechanical stress and up to 48 h compared to the control group. The expression of RANK mRNA was unchanged and that of OSCAR mRNA was increased 48 h after the release from mechanical stress, although they were slightly decreased in the mechanical stress group compared to the control group. The expression of *c-fms* mRNA was increased 24 h after the release from mechanical stress, although it was not significantly difference in the mechanical stress group compared to the control group.

Mechanical stress suppressed osteoclastogenesis and increased NO production and the NOS inhibitor, L-NAME, reduced the effect of mechanical stress. A significant increase in NO production was observed in the mechanical stress group. The concentration of NO₂/NO₃ in the conditioned medium with the mechanical stress showed an approximately 2-fold increase compared to the control group (Fig. 3). The concentration of NO₂/NO₃ in the conditioned medium with the mechanical stress and L-NAME showed a decrease compared with the mechanical stress only group (Fig. 3).




Figure 2. Effects of mechanical stress application and release on the expressions of RANK, *c-fms*, OSCAR, the NFATc family, iNOS, nNOS, IFN- β , TRAP, CTR, MMP-9, cath-k, CIC7, ATP6i and DC-STAMP. Real-time PCR analysis was performed with indicated primers in the mechanical stress group (A), and at 24 h (B), 48 h (C) and 72 h (D) after the release of mechanical stress. Results are shown as fold-changes or relative quantitation of target expression ($2^{-\Delta\Delta Ct}$ method) relative to control after normalization against GAPDH expression and as mean \pm SD (n=6). The control group which was cultured for the same days without mechanical stress was defined as the standard (n=1).

L-NAME had no effect on differentiation under basal conditions in osteoclasts (Fig. 4A). Addition of L-NAME to the culture medium with mechanical stress partially blocked the effect of mechanical stress on osteoclast differentiation and




Figure 3. Mechanical stress (MS) increased nitric oxide (NO) production and L-NAME reduced the effect of mechanical stress on osteoclastogenesis in RAW cells. Mechanical stress was applied to RAW cells with a Flexercell tension system for a period of 48 h over 3-4 days (grey bar), but not to the control (white bar) in the absence (-) or in the presence (+) of L-NAME. Total $\text{NO}_2^- + \text{NO}_3^-$ accumulated in the conditioned medium were shown as NO production. The data shown are representative of the results obtained in three experiments. Results are shown as the mean \pm SD (n=3).

fusion. The number of both osteoclasts and large osteoclasts induced by mechanical stress application and L-NAME treatment was increased approximately 2-fold compared to the mechanical stress only group (Fig. 4).

The effect of mechanical stress and L-NAME on the expressions of osteoclast differentiation genes. In order to confirm the affect of nitrite accumulation induced by mechanical stress and L-NAME treatment, we employed real-time PCR (Fig. 5). The expression of osteoclast specific gene mRNAs (TRAP, CTR, MMP-9, cath-k, CIC7, ATP6i) and DC-STAMP mRNA was increased with L-NAME treatment in the mechanical stress group compared to the mechanical stress only group. The expression of iNOS mRNA showed a tendency to decrease, however, the difference was not statistically significant. The expression IFN- β mRNA significantly increased with L-NAME treatment in the mechanical stress group compared with the mechanical stress only group. The expression of RANK, *c-fms* and NFATc1 mRNA was unchanged between the two groups.

Discussion

Application of mechanical stress on bone tissue generates extracellular matrix deformation, and contributes to the regulation of bone modeling and remodeling. Thereby, mechanical stress is translated into mechanical signals. Several studies have shown that these mechanical signals could adaptively change the functions of osteoblasts, osteocytes and bone stromal cells (10,34,35). In contrast, very few studies have focused on the direct effects of mechanical stress on osteoclast RAW cells. Furthermore, many studies have shown that loading is necessary to generate an anabolic effect in the bone (36), and that the anabolic response to loading declines soon after loading is initiated and mechanosensitivity is increased by the insertion of rest periods (37). However, the precise consequences of the removal of mechanical signals from bone tissue are currently difficult to predict *in vivo*, because they are dependent on the interactions between genetics, gender and the specific anatomical site. The effects




Figure 4. Suppression of the number of TRAP-positive multinucleated osteoclasts induced by mechanical stress (MS) was partially recovered by treatment with the NOS inhibitor, L-NAME. RAW cells were cultured in Flexcell 6-well plates with RANKL in the absence (-) or in the presence (+) of L-NAME. (A) The number of TRAP-positive multinucleated cells. (B) The number of large osteoclasts with ≥ 8 nuclei. Magnification $\times 100$. Results are shown as the mean \pm SD (n=3).




Figure 5. Effect of mechanical stress together with L-NAME on the expression of TRAP, CTR, MMP-9, cath-k, CIC7, ATP6i, DC-STAMP, IFN- β , iNOS, RANK, *c-fms* and NFATc1. RAW cells were cultured in Flexcell 6-well plates with mechanical stress in the absence (grey bar) or in the presence (black bar) of L-NAME. Results are presented as the mean \pm SD (n=3) of fold-changes or relative quantitation of target expression ($2^{-\Delta\Delta Ct}$ method) relative to the mechanical stress group after normalization against GAPDH expression. The mechanical stress group were defined as the standard (=1). MS, mechanical stress group; MS + L-NAME, treatment of L-NAME with MS.

of release from mechanical stress may be as important as the effects of mechanical stress. For this reason, in the present study we have evaluated the effects of mechanical stress applied with the Flexercell tension system and of the release from mechanical stress on osteoclastogenesis and further investigated the molecular mechanisms involved.

Previously, we have shown that mechanical stress directly suppresses osteoclast differentiation suggesting delayed differentiation in RAW cells (27). In the present study, we have demonstrated that the suppression of osteoclast differentiation and fusion with mechanical stress is related to the increase of NO via iNOS, and that the promotion of osteoclast differentiation and fusion after the release from mechanical stress is related to the NFATc family genes, whose expression remained constant during mechanical stress but was up-regulated after the release from mechanical stress. NO is a small, diffusible, diatomic free radical involved in several biological processes, including the regulation of bone formation, resorption, remodeling, mechanotransduction and repair in physiological or pathophysiological conditions. NO is an important regulator of the response of the bone to mechanical stress, and is produced through the activities of constitutive eNOS or iNOS (38). Several *in vitro* studies have shown that NO rapidly increases in response to mechanical stress in bone cells (39). NO mediates adaptive bone formation, protects osteocytes against apoptosis and mediates osteoclast activity (26). High levels of NO reduce osteoclast activity, while the inhibition of NO production increases osteoclastogenesis and osteoclast activity (40). iNOS has been detected in osteocytes, osteoclasts and chondrocytes and is expressed in response to inflammatory cytokines. Once induced, iNOS produces large amounts of NO (38). Thus, our results together with the previous findings may further contribute to the understanding of the relationship between mechanical stress and NO in osteoclasts. Indeed, in the present study we have shown that NO via iNOS is increased with mechanical stress application and suppresses osteoclast differentiation and fusion, although we could not detect eNOS mRNA expression in these osteoclastic RAW cells. IFN- β has been identified as a RANKL-stimulated autocrine negative feedback inhibitor that limits osteoclastogenesis and triggers iNOS/NO as an important negative feedback signal during osteoclastogenesis (41). In this study, the expression of IFN- β mRNA is decreased during mechanical stress and after the release from mechanical stress and is increased with L-NAME treatment. This result might contribute, at least in part, to the above-mentioned negative feedback signal.

NFATc1-deficient embryonic stem cells fail to differentiate into osteoclasts in response to RANKL stimulation and ectopic expression of NFATc1 causes precursor cells to undergo efficient differentiation without RANKL signaling, indicating that NFATc1 is a master switch for regulating the terminal differentiation of osteoclasts, functioning downstream of RANKL (7). Transcriptional activation of NFATc1 in osteoclasts is mediated by the RANKL/TRAF6/c-Fos signaling pathway because NFATc1 rescues osteoclastogenesis in precursors lacking c-Fos (3). In the present study, the mRNA expression of NFATc family genes was constant in spite of the suppression of osteoclast-specific genes, and was up-regulated after the release from mechanical stress. This result suggests that

homeostasis was maintained under conditions of mechanical stress through various courses of action.

With respect to the osteoclast specific genes (TRAP, CTR, MMP-9, cath-k, CIC7, ATP6i) and DC-STAMP which are essential for cell to cell fusion in osteoclasts (42), the expression of these mRNA was decreased with mechanical stress, and then increased after the release from mechanical stress up to 48 h compared to the control group, correlating with the number of both osteoclasts and large osteoclasts. However, 72 h after the release from mechanical stress, CTR whose expression inhibits osteoclast activity both *in vitro* and *in vivo* (43) was increased. This result might stand opposite to the osteoclasts which increased rapidly after the release from mechanical stress. Expression of the *c-fms* receptor is necessary for osteoclastogenesis and increased 24 h after the release from mechanical stress, although it was not significantly different in the mechanical stress group compared to the control group. RAW cells did not require exogenous M-CSF for RANKL-induced osteoclast formation. The capability of producing M-CSF in response to RANKL may enable RAW cells to differentiate into osteoclasts in the presence of RANKL alone (44). This capability may be different between physiological osteoclast precursors and RAW cells. OSCAR is critical for osteoclast differentiation. OSCAR mRNA expression was increased 48 h after the release from mechanical stress, although it was slightly decreased in the mechanical stress group compared to the control group. FcR γ -mediated signal transduction by OSCAR is involved in osteoclast differentiation (45). FcR γ -associated receptors are only activated by osteoblasts. In the RANKL/M-CSF system DAP12-associated receptors may be activated by endogenous ligands provided by osteoclast precursor cells themselves. DAP12 and FcR γ compensated for each other in the activation of the ITAM signal, and the cooperation of the RANKL and ITAM signals is essential for NFATc1 induction and osteoclastogenesis (9). Hence *c-fms* and OSCAR mRNA expression may be associated only to a small extent with the effects of mechanical stress on the osteoclastogenesis of RAW cells.

In summary, we have shown that the effects of the release from mechanical stress may be as important as the effects of mechanical stress on osteoclastogenesis. Our aim was to investigate the osteoclast differentiation genes which are altered in response to the release from mechanical stress applied with the Flexercell tension system. We have found that the suppression of osteoclast differentiation and fusion induced by mechanical stress resulted from the increase of NO via iNOS, and that the promotion of osteoclast differentiation and fusion after the release from mechanical stress is related to the NFATc family genes, whose expression remained constant during mechanical stress but was up-regulated after the release from mechanical stress.

Acknowledgements

We are grateful to Dr N. Suzuki and Dr T. Nakai for technical advice and support. This study was supported in part by the Japan Society for the Promotion of Science Grant-in-Aid for Scientific Research (C) (grant no. 22592274 to Y.Y. and no. 21592584 to T.K.) and a Grant-in-Aid for Young Scientists (B) (grant no. 22792032 to Y.T.).

References

1. Karsenty G and Wagner EF: Reaching a genetic and molecular understanding of skeletal development. *Dev Cell* 2: 389-406, 2002.
2. Teitelbaum SL and Ross FP: Genetic regulation of osteoclast development and function. *Nat Rev Genet* 4: 638-649, 2003.
3. Nakashima T and Takayanagi H: Osteoclasts and the immune system. *J Bone Miner Metab* 27: 519-529, 2009.
4. Nakashima T and Takayanagi H: Osteoimmunology: crosstalk between the immune and bone systems. *J Clin Immunol* 29: 555-567, 2009.
5. Yasuda H, Shima N, Nakagawa N, *et al*: Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci USA* 95: 3597-3602, 1998.
6. Yoshida H, Hayashi S, Kunisada T, *et al*: The murine mutation osteopetrosis is in the coding region of the macrophage colony stimulating factor gene. *Nature* 345: 442-444, 1990.
7. Takayanagi H, Kim S, Koga T, *et al*: Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev Cell* 3: 889-901, 2002.
8. Asagiri M, Sato K, Usami T, *et al*: Autoamplification of NFATc1 expression determines its essential role in bone homeostasis. *J Exp Med* 9: 1261-1269, 2005.
9. Takayanagi H: Mechanistic insight into osteoclast differentiation in osteoimmunology. *J Mol Med* 83: 170-179, 2005.
10. Ehrlich PJ and Lanyon LE: Mechanical strain and bone cell function: a review. *Osteoporos Int* 13: 688-700, 2002.
11. Schneider V, Oganov V, LeBlanc A, *et al*: Bone and body mass changes during space flight. *Acta Astronaut* 36: 463-466, 1995.
12. Srinivasan S, Weimer DA, Agans SC, Bain SD and Gross TS: Low-magnitude mechanical loading becomes osteogenic when rest is inserted between each load cycle. *J Bone Miner Res* 17: 1613-1620, 2002.
13. Mehrotra M, Saegusa M, Wadhwa S, Voznesensky O, Peterson D and Pilbeam C: Fluid flow induces RANKL expression in primary murine calvarial osteoclasts. *J Cell Biochem* 98: 1271-1283, 2006.
14. Tan SD, de Vries TJ, Kuijpers-Jagtman AM, Semeins CM, Everts V and Klein-Nulend J: Osteocytes subjected to fluid flow inhibit osteoclast formation and bone resorption. *Bone* 41: 745-751, 2007.
15. Ichimiya H, Takahashi T, Ariyoshi W, *et al*: Compressive mechanical stress promotes osteoclast formation through RANKL expression on synovial cells. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 103: 334-341, 2007.
16. Liu J, Zhao Z, Zou L, *et al*: Pressure-loaded MSCs during early osteodifferentiation promote osteoclastogenesis by increase of RANKL/OPG ratio. *Ann Biomed Eng* 37: 794-802, 2009.
17. Kanzaki H, Chiba M, Sato A, Miyagawa A, Arai K, Nukatsuka S and Mitani H: Cyclical tensile force on periodontal ligament cells inhibits osteoclastogenesis through OPG induction. *J Dent Res* 85: 457-462, 2006.
18. Rubin J, Biskobing D, Fan X, Rubin C, McLeod K and Taylor WR: Pressure regulates osteoclast formation and MCSF expression in marrow culture. *J Cell Physiol* 170: 81-87, 1997.
19. Makihira S, Kawahara Y, Yuge L, Mine Y and Nikawa H: Impact of the microgravity environment in a 3-dimensional clinostat on osteoblast- and osteoclast-like cells. *Cell Biol Int* 32: 1176-1181, 2008.
20. Kadow-Romacker A, Hoffmann JE, Duda G, Wildemann B and Schmidmaier G: Effect of mechanical stimulation on osteoblast- and osteoclast-like cells *in vitro*. *Cells Tissues Organs* 190: 61-68, 2009.
21. Kurata K, Uemura T, Nemoto A, *et al*: Mechanical strain effect on bone-resorbing activity and messenger RNA expressions of marker enzymes in isolated osteoclast culture. *J Bone Miner Res* 16: 722-730, 2001.
22. Robling AG, Niziolek PJ, Baldrige LA, *et al*: Mechanical stimulation of bone *in vivo* reduces osteocyte expression of Sost/sclerostin. *J Biol Chem* 283: 5866-5875, 2008.
23. Inoue D, Kido S and Matsumoto T: Transcriptional induction of FosB/DeltaFosB gene by mechanical stress in osteoblasts. *J Biol Chem* 279: 49795-49803, 2004.
24. Rubin J, Murphy TC, Fan X, Goldschmidt M and Taylor WR: Activation of extracellular signal-regulated kinase is involved in mechanical strain inhibition of RANKL expression in bone stromal cells. *J Bone Miner Res* 17: 1452-1460, 2002.
25. Rahnert J, Fan X, Case N, Murphy TC, Grassi F, Sen B and Rubin J: The role of nitric oxide in the mechanical repression of RANKL in bone stromal cells. *Bone* 43: 48-54, 2008.
26. Tan SD, Kuijpers-Jagtman AM, Semeins CM, *et al*: Fluid shear stress inhibits TNFalpha-induced osteocyte apoptosis. *J Dent Res* 85: 905-909, 2006.
27. Suzuki N, Yoshimura Y, Deyama Y and Suzuki K: Mechanical stress directly suppresses osteoclast differentiation in RAW264.7 cells. *Int J Mol Med* 21: 291-296, 2008.
28. Vande Geest JP, Di Martino ES and Vorp DA: An analysis of the complete strain field within Flexercell membranes. *J Biomech* 37: 1923-1928, 2004.
29. Nakai T, Yoshimura Y, Deyama Y and Suzuki K: Mechanical stress up-regulates RANKL expression via the VEGF autocrine pathway in osteoblastic MC3T3-E1 cells. *Mol Med Rep* 2: 229-234, 2009.
30. Takeyama S, Yoshimura Y, Deyama Y, Sugawara Y, Fukuda H, and Matsumoto A: Phosphate decreases osteoclastogenesis in coculture of osteoblast and bone marrow. *Biochem Biophys Res Commun* 282: 798-802, 2001.
31. Nakamura K, Deyama Y, Yoshimura Y, Suzuki K and Morita M: Toll-like receptor 3 ligand-induced antiviral response in mouse osteoblastic cells. *Int J Mol Med* 19: 771-775, 2007.
32. Aguirre JJ, Plotkin LL, Gortazar AR, *et al*: A novel ligand-independent function of the estrogen receptor is essential for osteocyte and osteoblast mechanotransduction. *J Biol Chem* 282: 25501-25508, 2007.
33. Pitsillides AA, Rawlinson SC, Suswillo RF, Bourrin S, Zaman G and Lanyon LE: Mechanical strain-induced NO production by bone cells: a possible role in adaptive bone (re)modeling? *FASEB J* 9: 1614-1622, 1995.
34. Rubin J, Rubin C and Jacobs CR: Molecular pathways mediating mechanical signaling in bone. *Gene* 367: 1-16, 2006.
35. Galli C, Passeri G and Macaluso GM: Osteocytes and WNT: the mechanical control of bone formation. *J Dent Res* 89: 331-343, 2010.
36. Judex S, Gupta S and Rubin C: Regulation of mechanical signals in bone. *Orthod Craniofac Res* 12: 94-104, 2009.
37. Robling AG, Burr DB and Turner CH: Recovery periods restore mechanosensitivity to dynamically loaded bone. *J Exp Biol* 204: 3389-3399, 2001.
38. Van't Hof RJ and Ralston SH: Nitric oxide and bone. *Immunology* 103: 255-261, 2001.
39. Bacabac RG, Smit TH, Mullender MG, Dijcks SJ, Van Loon JJ and Klein-Nulend J: Nitric oxide production by bone cells is fluid shear stress rate dependent. *Biochem Biophys Res Commun* 315: 823-829, 2004.
40. Wimalawansa SJ: Nitric oxide and bone. *Ann NY Acad Sci* 1192: 391-403, 2010.
41. Zheng H, Yu X, Osdoby PC and Osdoby P: RANKL stimulates iNOS expression and NO production in developing osteoclasts: an autocrine negative feedback mechanism triggered by RANKL-induced IFN- β via NF- κ B that restrains osteoclastogenesis and bone. *J Biol Chem* 281: 15809-15820, 2006.
42. Yagi M, Miyamoto T, Sawatani Y, *et al*: DC-STAMP is essential for cell-cell fusion in osteoclasts and foreign body giant cells. *J Exp Med* 202: 345-351, 2005.
43. Mancini L, Moradi-Bidhendi N, Brandi ML, Perretti M and MacIntyre I: Modulation of the effects of osteoprotegerin (OPG) ligand in a human leukemic cell line by OPG and calcitonin. *Biochem Biophys Res Commun* 279: 391-397, 2000.
44. Islam S, Hassan F, Tumurkhuu G, *et al*: Receptor activator of nuclear factor-kappa B ligand induces osteoclast formation in RAW 264.7 macrophage cells via augmented production of macrophage-colony-stimulating factor. *Microbiol Immunol* 52: 585-590, 2008.
45. Ishikawa S, Arase N, Suenaga T, *et al*: Involvement of Fc γ in signal transduction of osteoclast-associated receptor (OSCAR). *Int Immunol* 16: 1019-1025, 2004.