

Effect of fibroblasts on breast cancer cell mammosphere formation and regulation of stem cell-related gene expression

FENGCHUN ZHANG¹, CAILI SONG¹, YUE MA², LEI TANG², YINGCHUN XU² and HONGXIA WANG²

¹Department of Oncology, Suzhou Kowloon Hospital, Shanghai Jiaotong University School of Medicine, Suzhou 215021; ²Department of Oncology, Shanghai Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200127, P.R. China

Received February 24, 2011; Accepted April 20, 2011

DOI: 10.3892/ijmm.2011.700

Abstract. The purpose of this study was to investigate the regulatory effects of breast cancer fibroblasts (BCFs) vs. normal mammary fibroblasts (NMFs) on mammosphere formation and stem cell-related gene expression in breast cancer cells. Breast cancer cells (MCF-7) were cultured in suspension to generate primary and secondary mammospheres. The proportion of CD44⁺/CD24^{low/-} cells was assessed by flow cytometry (FCM), and *Wnt1*, *Notch1*, β -catenin, *CXCR4*, *SOX2* and *ALDH3A1* gene expression was detected by quantitative real-time PCR. The fibroblasts from either breast cancer tissue or normal mammary tissue were purified from tissue specimens and co-cultured with breast cancer cells. The mammosphere formation efficacy was approximately 180/10,000 MCF-7 cells. FCM analysis showed that, compared to the 2.1% positive expression in the MCF-7 monolayer culture cells, the expression of CD44⁺/CD24^{low/-} in MCF-7 mammosphere cells was significantly elevated to 10.4% ($P < 0.01$). The proportion of the CD44⁺/CD24^{low/-} subpopulation of the cells in mammospheres was nearly 5-fold higher than that of general MCF-7 cells. Compared with MCF-7 monolayer culture cells, mammosphere cells showed significantly ($P < 0.01$) enhanced expression of *Wnt1* [fold-change (FC), 2.25], *Notch1* (FC, 2.45), β -catenin (FC, 1.72), *CXCR4* (FC, 4.68), *SOX2* (FC, 4.25) and *ALDH3A1* (FC, 5.38). When BCFs were co-cultured with MCF-7 cells under mammosphere culture conditions, the length of time of mammosphere formation decreased, the volume of the mammospheres increased and the mammosphere-forming efficiency (MFE) was higher than that of NMFs and the control group. Both the BCF and NMF groups showed enhanced gene expression for the following genes: *Wnt1* (FC, 3.18 and 1.27, respectively), β -catenin (FC, 1.75 and 1.22, respectively), *Notch1* (FC, 2.09 and 1.31, respectively), *CXCR4* (FC, 2.77 and

1.33, respectively), *SOX2* (FC, 2.77 and 1.80, respectively) and *ALDH3A1* (FC, 5.23 and 1.85, respectively). Cancer fibroblast cells can promote the MFE and up-regulate stem cell-related gene expression in breast cancer cells.

Introduction

According to the cancer stem cell (CSC) theory, a small population of tumor cells is capable of self-renewal, which gives rise to the heterogeneous nature of tumors. CSCs have been discovered in multiple cancers, including those of the hematopoietic system, brain, breast, prostate, and gastrointestinal tract (1-4). These cells can exclude dyes such as Hoechst or rhodamine due to the increased expression of membrane transporter proteins, including p-glycoproteins or breast cancer resistance proteins (BCRPs) (5). Therefore, it is difficult for conventional therapies to eradicate CSCs, which largely contributes to patient relapse, years after chemotherapy or radiotherapy.

In 2003, Clarke and his colleagues identified human breast cancer stem cells by their CD44⁺/CD24⁻ phenotype. These rare cell populations possessed an increased ability to form tumors when they were injected into etoposide-treated NOD/SCID mice; as few as 100 CD44⁺/CD24⁻ human breast cancer cells could recapitulate the human mammary tumors, whereas the injection of 10,000 cells with other phenotypes failed to give rise to tumors (2). Since then, several techniques have been established to isolate or enrich for tumorigenic breast cancer stem cells (BCSCs), including side-population (SP) separation and mammosphere culture (6,7). In breast cancer, the mammosphere culture system has been widely used to identify and enrich for putative stem cells using breast cancer cell lines (8). In a previous study, we described an *in vitro* culture system to propagate primary human mammary epithelial stem cells and progenitor cells in an undifferentiated state based on their ability to proliferate in suspension as spherical structures when cultured on non-adherent surfaces in the presence of growth factors. We found that cultured mammospheres exhibited the characteristics of BCSCs using this method (9).

In some reports, mammary stem/progenitor cells have been described as small, light cells that lie in an intermediate location, termed a stem cell niche (10) that is composed of stromal cells (myofibroblasts, endothelial cells and inflammatory

Correspondence to: Dr Hongxia Wang or Dr Yingchun Xu, Department of Oncology, Shanghai Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200127, P.R. China
E-mail: whx365@126.com
E-mail: xiaoxu2384@163.com

Key words: breast cancer, stem cell, mammosphere, fibroblast cells

cells) and extracellular matrix (ECM) components (e.g., laminin, fibronectin, collagen, and proteoglycans), which may influence mammary development. BCSCs also have a functional niche; stromal cells, together with ECM components, provide the microenvironment that is pivotal for BCSC self-renewal, proliferation, and maintenance in an undifferentiated state. Based on this finding, recent studies have mainly focused on how to regulate mammospheres. Fibroblasts are considered to play a central role in the complex process of tumor-stromal interactions and consequently also in tumorigenesis. Premalignant mammary epithelial cells exposed to senescent human fibroblasts irreversibly become invasive and undergo full malignant transformation. In invasive breast cancer, fibroblasts were found in a much higher proportion than *in situ* carcinomas and were predominantly at the invasive front.

However, the regulatory effects of fibroblasts on mammosphere formation remain unclear. We performed this study to further investigate the interactive influence of fibroblasts on BCSCs. The aim of the study was to compare the effect of breast cancer fibroblasts (BCFs) and normal mammary fibroblasts (NMFs) on the MCF-7 breast cancer cell mammosphere formation and signaling pathway regulation, which will help us to understand the interaction between the microenvironment and breast cancer stem cells.

Materials and methods

Cell culture and reagents. MCF-7 human breast epithelial adenocarcinoma cells were obtained from the American Type Culture Collection (ATCC, Manassas, VA). Cells were cultured routinely in Dulbecco's modified Eagle's medium/F12 (DMEM/F12, Hyclone, Logan, UT) supplemented with 10% fetal bovine serum (FBS), 100 U/ml penicillin and 100 µg/ml streptomycin (Life Technologies, Inc., Grand Island, NY) at 37°C in a humidified atmosphere with 5% CO₂.

Epidermal growth factor (EGF), insulin, basic fibroblast growth factor (bFGF), B27, TRIzol and primers for PCR were purchased from Invitrogen (Carlsbad, CA). Moloney murine leukemia virus (MMLV) reverse transcriptase was purchased from Promega (Madison, WI). Heparin, type I collagenase and hyaluronidase were obtained from Sigma (St. Louis, MO). Fibroblast-specific protein (FSP) antibody was purchased from NeoMarkers (Pittsburgh, PA).

Mammosphere culture. To obtain BCSCs and propagate them as mammospheres, cells floating in the supernatant of 2-day-old cultures were collected by centrifugation, washed in Hank's-buffered salt solution, and plated in ultralow attachment plates (Corning, NY) at a density of 2×10^4 viable cells/ml. The cells were then grown in serum-free mammary epithelial growth medium (BioWhittaker, Walkersville, MD) supplemented with B27, 10 ng/ml EGF, 5 µg/ml bovine insulin, 20 ng/ml bFGF2 and 4 µg/ml heparin. Bovine pituitary extract was excluded from the growth medium. Growth factors were added to the mammosphere cultures every 3 days, and mammospheres (>60 µm in size) were counted on Day 7.

By Day 7, non-adherent spherical clusters of cells were harvested by trypsinization and centrifugation and dispersed by passage through a 40-µm pore filter. The single cells were then collected for further analysis.

Flow cytometry. Mammospheres were dispersed to obtain single-cell populations as described above. Cells were washed in phosphate-buffer saline (PBS) with 2.5% bovine serum albumin and stained with FITC-anti-mouse CD24 and APC-anti-mouse CD44 antibodies (BD Pharmingen, San Jose, CA). Cells were incubated on ice for 30 min, washed twice with PBS and then fixed in PBS containing paraformaldehyde. Flow cytometric analysis was performed on a BD FACSCalibur system (BD Biosciences, San Jose, CA), and acquisition was performed with the BD CellQuest software (BD Biosciences). The FlowJo software (Tree Star, Ashland, OR) was used for data analysis.

Isolation and culture of fibroblasts. Tumor specimens were obtained after obtaining informed consent from the patients and receiving approval from the Ethics Committee. Malignant (n=5) and normal mammary tissue samples (from the same patient but from a site remote of the tumor area and free from tumor cells, as confirmed by subsequent histological analysis; n=5), without neoadjuvant chemotherapy were collected immediately after surgery and mechanically disaggregated under sterile conditions within 30 min.

The procedure was performed as previously described, with slight modifications. In brief, fresh tumor tissues were dissected to a volume of 1 mm³, with the elimination of fat, and subjected to 100 U/ml type 1 collagenase and 150 U/ml hyaluronidase treatment under constant stirring at 37°C for 18 h. The resulting cell suspension was passed through an 80-µm filter, and the single cell suspension was centrifuged at 80 x g for 6 min and seeded in 25- or 75-cm² tissue culture flask. Cultures were fed twice a week, and after 7 days, when 60-80% confluence had been reached, stromal fibroblasts were obtained by differential trypsinization. The resulting second-passage fibroblasts were then seeded into a separate flask and maintained in DMEM/F12 supplemented with 10% FBS, 100 U/ml penicillin, and 100 U/ml streptomycin in a 37°C incubator containing 5% CO₂. Cells were maintained in culture for up to 2-4 passages.

Immunocytochemistry. The suspension of fibroblasts obtained was cultured on sterile glass coverslips for 24 h, after which the coverslips were rinsed with PBS and the cells were fixed with 4% (vol/vol) acetic acid in methanol. The sections were treated with 3% hydrogen peroxide in methanol for 30 min and then 10% BSA for 30 min at room temperature. Primary antibodies against vimentin, α-SMA, and FSP were added separately, and the samples were incubated at 37°C for 24 h. After washing with PBS, a secondary antibody against the rabbit antibody (diluted 1:200 in PBS) was added for a 30 min incubation at 37°C, and the immunoreactivity was visualized by immersing the sample in a diaminobenzidine (DAB)-H₂O₂ solution (0.05 M Tris-HCl buffer, pH 7.6, with 0.05% DAB and 0.01% H₂O₂) for 5 min at room temperature. The sections were finally stained with hematoxylin and mounted for light microscopy analysis.

Transwell assay. Mammospheres were co-cultured with either NMFs or BCFs using a transwell system. Fibroblasts (2×10^5 cells in 2 ml/well) were seeded into the upper chamber of the transwell, and mammospheres (2×10^5 cells/ml) were cultured in

Table I. Primers used in real-time PCR analysis.

Gene	Primer sequence	Product size (bp)
<i>GAPDH</i>	(F) 5'-ACCCACTCCTCCACCTTTGA-3' (R) 5'-CTGTTGCTGTAGCCAAATTCGT-3'	101
<i>Wnt1</i>	(F) 5'-GAACCTGCTTACAGACTCCAAGAGT-3' (R) 5'-CCGGATTTTGGCGTATCAGA-3'	98
<i>Notch1</i>	(F) 5'-CCGCAGTTGTGCTCCTGAA-3' (R) 5'-ACCTTGGCGGTCTCGTAGCT-3'	109
<i>β-catenin</i>	(F) 5'-CCTTTGTCCCGCAAATCATG-3' (R) 5'-ACGTACGGCGCTGGGTATC-3'	101
<i>CXCR4</i>	(F) 5'-CAGTGGCCGACCTCCTCTT-3' (R) 5'-ACATGGACTGCCTTGCATAGG-3'	100
<i>SOX2</i>	(F) 5'-TGCAGAGCGCTGCACAT-3' (R) 5'-CGGGCAGCGTGTACTTATCC-3'	96
<i>ALDH3A1</i>	(F) 5'-TCCAGCAACGACAAGGTGATT-3' (R) 5'-GGCAGAGAGTGCAAGGTGATG-3'	101

the lower chamber in DMEM/F12 medium containing 10 ng/ml EGF, 5 ng/ml bFGF, 2.5 µg/ml insulin, and 2 µl/ml 50X B27. By Day 4, mammospheres were observed by microscopy, the mammosphere-forming efficiency (MFE) was counted (diameter of mammospheres >60 µm) and gene expression was detected by quantitative real-time PCR (qRT-PCR).

qRT-PCR. Total-RNA from the single-cell suspension was extracted using TRIzol Reagent according to the manufacturer's instructions, and the RNA was then stored at -80°C until use. The RNA from each sample was reverse transcribed using random primers and MMLV. Real-time PCR was performed on cDNA samples using a SYBR-Green Master Mix (Applied Biosystems, Carlsbad, CA). The sequences of the gene-specific primers and the lengths of the PCR products are listed in Table I; PCR primers were designed using Primer Express Software, version 2. The reactions were conducted using the ABI PRISM 7900HT (Applied Biosystems) according the following conditions: 50°C for 2 min, 95°C for 10 min and 40 cycles of 95°C for 15 sec and 60°C for 60 sec. The data were normalized to the reference gene GAPDH, and gene-specific amplification was confirmed by determining the melting curves of the PCR products.

Statistical analysis. The results are presented as the mean ± SD for at least three individual experiments for each group. The statistical analysis was performed using SPSS 16.0 statistical Software and GraphPad Prism 4.0 software (San Diego, CA). Statistical differences were determined using either the two-tailed Student's t-test or the one-way analysis of variance (ANOVA). The level of statistical significance was defined as P≤0.05.

Results

Mammospheres enrich breast CSCs. Single MCF-7 cells were cultured in suspension (2×10⁴ cells/ml), as described previously,

to produce mammospheres. In general, apparent mammospheres could be observed by microscopy by Day 6 (Fig. 1A and B). The shape of the mammospheres was round, with relatively regular morphology and an average diameter of 160 µm. The mammosphere formation efficacy (MFE) was about 180/10,000 MCF-7 cells, and the cells in the mammospheres showed a higher colony formation ability *in vitro* and higher tumorigenicity *in vivo* than whole MCF-7 cell populations, suggesting that the cell population was enriched with breast cancer stem/progenitor cells in agreement with a previous report (9).

Flow cytometry was then performed to assess the percentage of CD44⁺/CD24^{low/-} cells in mammosphere cultures and MCF-7 monolayer cultures. The cells in the upper left quadrant were classified as CD44⁺/CD24^{low/-} cells (Fig. 1C). Our analysis indicated that, compared with a 2.1% positive expression in the MCF-7 monolayer culture cells, the expression of CD44⁺/CD24^{low/-} in MCF-7 mammosphere cells was significantly elevated to 10.4% (P<0.01). These results are consistent with previous reports, in which the percentage of CD44⁺/CD24^{low/-} cells in mammosphere populations ranged from 10-12.1%, while in MCF-7 monolayer culture cells the percentage was from 1.6-2.1% (11,12).

We performed qRT-PCR analysis to further determine the expression of *Wnt1*, *Notch1*, *β-catenin*, *CXCR4*, *SOX2* and *ALDH3A1* in mammosphere cells and MCF-7 monolayer culture cells. The results indicated that mammosphere cells showed significantly (P<0.01) increased expression of *Wnt1* [fold change (FC), 2.25], *Notch1* (FC, 2.45), *β-catenin* (FC, 1.72), *CXCR4* (FC, 4.68), *SOX2* (FC, 4.25) and *ALDH3A1* (FC, 5.38) compared with MCF-7 monolayer culture cells (Fig. 1D).

Identification of normal mammary and breast cancer fibroblasts. Fibroblasts were isolated and cultured as described previously. After culturing for 1-2 weeks, a mixture of both fibroblast and epithelial cells was observed under inverted microscopy (Fig. 2A). In light of the lower tolerance to trypsin




Figure 1. Mammosphere culture of MCF-7 cells. (A) Mammosphere formation of MCF-7 cells (x4). (B) Mammosphere formation of MCF-7 cells (x20). (C) CD44 and CD24 expression in MCF-7 monolayer cells (left) and mammospheres (right) as detected by fluorescence activated cell-sorting (FACS). (D) Real-time PCR analysis of the gene expression of *Wnt1*, *Notch1*, *β-catenin*, *CXCR4*, *SOX2* and *ALDH3A1*.

that fibroblasts have compared to epithelial cells (stromal cells are more easily detached by trypsin), we cultured and repeatedly digested these cells to separate and purify fibroblasts. After two or three subcultured generations, pure fibroblasts populations were achieved; the cells had a uniform appearance of a long fusiform shape typical of fibroblastic morphology when viewed by inverted microscopy (Fig. 2B).

Fibroblasts are normally defined by the concurrent expression of the smooth muscle marker α -smooth muscle actin (α -SMA), the mesenchymal marker vimentin and FSP; on the other hand, CK (cytokeratin) was confirmed as an endothelial cell marker. To identify whether cells obtained by the above method were fibroblasts, the cells were stained with antibodies against α -SMA, vimentin, FSP and CK by immunohistochemistry. Vimentin, α -SMA, and FSP but not CK, were expressed in these cells, indicating that the cells we collected through this method were fibroblasts (Fig. 2C).

Effect of normal mammary and breast cancer fibroblasts on MCF-7 mammosphere formation. To determine the influence of fibroblasts on MCF-7 cell migration and mammosphere formation, MCF-7 cells were co-cultured with NMFs or BCFs in a transwell system (Fig. 3A). Mammospheres were observed




Figure 2. Identification of fibroblasts. (A) Admixture of fibroblasts and epithelial cells (x100). (B) Fibroblast cells are morphologically characterized as large spindle-shaped cells with indented nuclei (x100). (C) Immunohistochemistry for CK(-), Vimentin(+), α -SMA(+), FSP(+) (x100).

by microscopy daily, and the MFE in each group was calculated on Day 7.

We found that BCFs promoted the migration of MCF-7 cells in the transwell culture system. When BCFs were co-cultured with MCF-7 under mammosphere culture conditions, the time of mammosphere formation was shorter, the volumes of the mammospheres were greater and the MFE was higher than both NMFs and the control group. Mammosphere formation in the BCFs group was enhanced 3.2-fold over control cultures. NMFs showed an approximately 1.3-fold greater MFE compared with the control group, whereas BCFs showed an approximately 2.5-fold increase over the NMFs group. Our study demonstrates that fibroblasts, especially those in cancer tissues, increase the mammosphere formation efficiency of MCF-7 cells in our co-culture system, reflecting their ability to increase the clonogenic efficiency of breast cancer cells (Fig. 3B-D).

Regulation of mammosphere signaling proteins by breast cancer and normal mammary fibroblasts. To further determine the




Figure 3. Effect of breast cancer (BCFs) and normal mammary fibroblasts (NMFs) on MCF-7 cell migration and mammosphere formation. (A) Comparison of mammosphere-forming efficiency, NMFs, BCFs. (B) Migration of MCF-7 cells; the arrow points to the migrating MCF-7 cells. (C) MCF-7 cells co-cultured with NMFs, Day 2. (D) MCF-7 cells co-cultured with BCFs, Day 7.




Figure 4. Effect of breast cancer (BCFs) and normal mammary fibroblasts (NMFs) on stem cell-related gene expression. NMFs and BCFs were co-cultured with mammospheres.

impact of breast cancer and normal mammary fibroblasts on several major mammosphere signaling proteins, we examined the differential gene expression by qRT-PCR. The analysis of gene expression profiling showed that the expression levels of *Wnt1* in MCF-7 mammospheres were significantly up-regulated by 3.18-fold in the BCFs co-culture group and by 1.27-fold in the NMFs co-culture group over the control group (Fig. 4).

Similar trends were also observed for the expression of other signaling proteins, such as β -catenin, *Notch1*, *CXCR4*, *SOX2* and *ALDH3A1*, with either NMFs or BCFs, as follows: *Wnt1*, FC of 1.27 and 3.18, respectively; β -catenin, FC of 1.22 and 1.75, respectively; *Notch1*, FC of 1.31 and 2.09, respectively; *CXCR4*, FC of 1.73 and 2.77, respectively; *SOX2*, FC of 1.8 and 2.77, respectively; and *ALDH3A1*, FC of 1.85 and 5.23, respectively.

Discussion

Stem cells are characterized by their ability to self-renew as well as generate differentiated cells within each organ. Numerous studies have provided strong evidence for the existence of mammary stem cells capable of self-renewal and differentiation into the basal and luminal lineages comprising the functional mammary epithelium (13). Recently, Sheridan *et al* (14) reported the existence of a CD44⁺/CD24^{low/-} subpopulation in breast cancer cell lines that possess more invasive and proliferative properties than other breast cancer cell populations. Patrawala *et al* (6) confirmed that the CD44⁺/CD24^{low/-} subpopulation of cancer cells have BCSC properties, and as compared to unsorted cells, only a low number of CD44⁺/CD24^{low/-} cells is sufficient for tumor initiation.

During breast carcinogenesis, concurrent morphological and molecular changes occur in stromal and epithelial compartments, which form a microenvironment suitable for tumor growth. The tumor microenvironment is composed of mesenchymal cells, extracellular matrix components, vascular endothelial cells and inflammatory cells; fibroblasts (also termed myofibroblasts or cancer-associated fibroblasts) represent the majority of the stromal cells. The cells are able to modify the epithelial cell phenotype by direct cell-to-cell contacts, through soluble factors or by modification of ECM components. At the same time, breast cancer cells promote MMP-2 and MMP-9 expression in cancer-associated fibroblasts by stromal-epithelial interactions, which form a microenvironment that contributes to the invasion and metastasis of malignant tumors (15). Premalignant mammary epithelial cells exposed to senescent human fibroblasts were reported to irreversibly become invasive and undergo a full malignant transformation (16). Cancer cells were shown to be suppressed by young and quickly proliferating non-cancer cells, such as embryonic stem cells (17). These reports demonstrate that stromal alterations accompany or even precede the malignant conversion of epithelial cells.

In a previous study, we described an *in vitro* culture system to propagate primary mammary epithelial stem cells in an undifferentiated state based on their ability to proliferate as spherical structures (mammospheres) in the presence of growth factors, and that these cultured mammospheres exhibited BCSC properties (9). In this study, the mammosphere formation efficacy was approximately 180/10,000 MCF-7 cells. Flow cytometry (FCM) analysis showed that, compared with 2.1% expression in the MCF-7 monolayer culture cells, the expression of CD44⁺/CD24^{low/-} in MCF-7 mammosphere cells was significantly elevated (10.4%) ($P < 0.01$). The proportion of cells with the CD44⁺/CD24^{low/-} phenotype in the mammosphere subpopulation was nearly 5-fold higher than that in the general MCF-7 population, suggesting that, consistent with several reports, mammosphere cultivation enriches CSCs (18,19).

Studies on normal breast stem cells and BCSCs have revealed that several key signaling pathways are involved in the self-renewal and maintenance of the stem cell pool, including the Wnt/ β -catenin, Notch, Hedgehog (Hh), transforming growth factor (TGF)- β , PTEN and Bmi signaling pathways (20-22). The Wnt pathway is involved in the regulation of self-renewal in normal breast stem cells and causes an increase in mammary stem cells (23), and the expression of

Wnt1 and β -catenin increases in BCSCs compared to non-BCSCs (24). Notch signaling promotes the self-renewal of mammary stem cells as well as the proliferation of early progenitor cells, and unregulated Notch signaling prevents the terminal differentiation of mammary epithelial cells (25). Hh signaling components, such as PTCH1, Gli1 and Gli2 are highly expressed in normal human mammary stem/progenitor cells (26). Consistent with previous results, we observed an enhanced expression of *Wnt1*, *Notch1*, β -catenin, *CXCR4*, *SOX2* and *ALDH3A1* in mammosphere cells as compared with MCF-7 monolayer culture cells.

Furthermore, Kucia *et al* (27) found that CSCs exhibit the same metastasis mechanisms as normal stem cells, and that transplantation and the SDF-1/CXCR4 axis play an important part in these events. In our study, cancer mammospheres strongly expressed CXCR4, which suggests that CXCR4 may be involved in regulating breast cancer stem cell metastasis and invasion. SOX2 is a transcription factor essential for the self-renewal and differentiation of embryonic (28,29) and neural stem cells (30). Eriksson *et al* (31) found that CD44⁺CD24^{low/-} cells isolated from breast tumor patient pleural effusions expressed SOX2. ALDH3A1 is a NAD-dependent enzyme responsible for the oxidation of intracellular aldehydes. A recent study demonstrated that ALDH3A1 not only inhibits cell proliferation by affecting the MAPK pathway but also converts the activated drug to the inactive metabolite (32), and Ginestier *et al* (33) discovered that ALDH1 expression in breast carcinomas correlated with poor prognosis. Here, we found that the expression of CXCR4, SOX2 and ALDH3A1 were 4.68-, 4.25- and 5.38-fold higher, respectively, in mammosphere cells than in monolayer cells.

We further examined whether cancer-associated fibroblasts could influence MCF-7 cell mammosphere formation. Using the transwell co-culture system, we observed that cancer-associated fibroblasts significantly enhanced the formation of MCF-7 cell mammospheres, as mammosphere formation in the breast cancer fibroblast group was enhanced 3.2-fold over control cultures. Our study shows that fibroblasts, especially cancer fibroblasts, increase the mammosphere formation efficiency of MCF-7 cells in a co-culture system, reflecting their ability to increase the clonogenic efficiency of this cell line. NMFs increased the MCF-7 MFE approximately 1.3-fold compared with the control group, while BCFs caused an approximately 2.5-fold increase compared with the NMFs group. Our data indicates that tumor fibroblasts may provide a functional microenvironment to play an important role in mammosphere formation.

The study also revealed that fibroblasts have a broad effect on signaling protein expression; Wnt1 expression levels were significantly up-regulated 3.18-fold in the BCFs co-culture group over the control group, whereas the NMFs co-culture group caused a 1.27-fold increase over the control group. Similar trends were also observed for the expression of other signaling proteins.

In general, these studies show that the microenvironment provides a suitable condition for cancer cells by secreting cytokines, which are associated with CSC growth and differentiation. These results indicate that the role of cancer microenvironments cannot be ignored, and that they also provide a reference for clinical and fundamental research.

Cells exist in a complicated ecological system and make interactions by juxtacrine and paracrine signaling, and any form of inappropriate signaling can be carcinogenic. The approach of suppressing tumor growth by blocking specific signal transmission will be a breakthrough for breast cancer therapy. Our study suggests that cancer-associated fibroblasts may represent a new target for cancer therapy by blocking specific signaling molecules and downstream effectors to sacrifice malignant cells. These results may also be useful in finding new therapies for CSC eradication.

In summary, cancer fibroblast cells can promote the MFE and up-regulate stem cell-related gene expression in breast cancer cells, indicating that the cancer microenvironment provides a suitable condition for CSC maintenance.

Acknowledgements

The study was supported by the Leading Academic Discipline Project of the Shanghai Municipal Education Committee (J50208), the National Natural Science Foundation of China (30670798) and the Suzhou Planning Project of Science and Technology (SYS201062).

References

- Singh SK, Hawkins C, Clarke ID, *et al*: Identification of human brain tumour initiating cells. *Nature* 432: 396-401, 2004.
- Al-Hajj M, Wicha MS, Benito-Hernandez A, *et al*: Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 100: 3983-3988, 2003.
- Haraguchi N, Utsunomiya T, Inoue H, *et al*: Characterization of a side population of cancer cells from human gastrointestinal system. *Stem Cells* 24: 506-513, 2006.
- Bonnet D and Dick JE: Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 3: 730-737, 1997.
- Goodell MA, McKinney-Freeman S and Camargo FD: Isolation and characterization of side population cells. *Methods Mol Biol* 290: 343-352, 2005.
- Patrawala L, Calhoun T, Schneider-Broussard R, *et al*: Side population is enriched in tumorigenic, stem-like cancer cells, whereas ABCG2⁺ and ABCG2⁻ cancer cells are similarly tumorigenic. *Cancer Res* 65: 6207-6219, 2005.
- Xu JX, Morii E, Liu Y, *et al*: High tolerance to apoptotic stimuli induced by serum depletion and ceramide in side-population cells: high expression of CD55 as a novel character for side-population. *Exp Cell Res* 313: 1877-1885, 2007.
- Cariati M, Naderi A, Brown JP, *et al*: Alpha-6 integrin is necessary for the tumorigenicity of a stem cell-like subpopulation within the MCF7 breast cancer cell line. *Int J Cancer* 122: 298-304, 2008.
- Lin SC, Huang MZ, Zhang FC and Zhang YY: Enrichment of breast cancer stem cells and correlative gene expression. *J Shanghai Jiaotong Univ (Medical Sci)* 28: 2008.
- Chepko G and Dickson RB: Ultrastructure of the putative stem cell niche in rat mammary epithelium. *Tissue Cell* 35: 83-93, 2003.
- Phillips TM, McBride WH, Pajonk F: The response of CD24^{low}/CD44⁺ breast cancer-initiating cells to radiation. *J Natl Cancer Inst* 98: 1777-1785, 2006.
- Huang MZ, Zhang FC and Zhang YY: Influence factors on the formation of mammospheres from breast cancer stem cells. *Beijing Da Xue Xue Bao* 40: 500-504, 2008 (In Chinese).
- Visvader JE and Lindeman GJ: Mammary stem cells and mammo-poiesis. *Cancer Res* 66: 9798-9801, 2006.
- Sheridan C, Kishimoto H, Fuchs RK, *et al*: CD44⁺/CD24⁻ breast cancer cells exhibit enhanced invasive properties: an early step necessary for metastasis. *Breast Cancer Res* 8: R59, 2006.
- Singer CF, Kronsteiner N, Marton E, *et al*: MMP-2 and MMP-9 expression in breast cancer-derived human fibroblasts is differentially regulated by stromal-epithelial interactions. *Breast Cancer Res Treat* 72: 69-77, 2002.
- Parrinello S, Coppe JP, Krtolica A and Campisi J: Stromal-epithelial interactions in aging and cancer: senescent fibroblasts alter epithelial cell differentiation. *J Cell Sci* 118: 485-496, 2005.
- Ukrainitseva SV and Yashin AI: Treating cancer with embryonic stem cells: rationale comes from aging studies. *Front Biosci* 10: 588-595, 2005.
- Dontu G, Abdallah WM, Foley JM, *et al*: In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. *Genes Dev* 17: 1253-1270, 2003.
- Yuan X, Curtin J, Xiong Y, *et al*: Isolation of cancer stem cells from adult glioblastoma multiforme. *Oncogene* 23: 9392-9400, 2004.
- Zhou J, Wulfschlegel J, Zhang H, *et al*: Activation of the PTEN/mTOR/STAT3 pathway in breast cancer stem-like cells is required for viability and maintenance. *Proc Natl Acad Sci USA* 104: 16158-16163, 2007.
- Brennan KR and Brown AM: Wnt proteins in mammary development and cancer. *J Mammary Gland Biol Neoplasia* 9: 119-131, 2004.
- Boulanger CA, Wagner KU and Smith GH: Parity-induced mouse mammary epithelial cells are pluripotent, self-renewing and sensitive to TGF-beta1 expression. *Oncogene* 24: 552-560, 2005.
- Lindvall C, Evans NC, Zylstra CR, *et al*: The Wnt signaling receptor Lrp5 is required for mammary ductal stem cell activity and Wnt1-induced tumorigenesis. *J Biol Chem* 281: 35081-35087, 2006.
- Kruger JA, Kaplan CD, Luo Y, *et al*: Characterization of stem cell-like cancer cells in immune-competent mice. *Blood* 108: 3906-3912, 2006.
- Dontu G, Jackson KW, McNicholas E, *et al*: Role of Notch signaling in cell-fate determination of human mammary stem/progenitor cells. *Breast Cancer Res* 6: R605-R615, 2004.
- Liu S, Dontu G, Mantle ID, *et al*: Hedgehog signaling and Bmi-1 regulate self-renewal of normal and malignant human mammary stem cells. *Cancer Res* 66: 6063-6071, 2006.
- Kucia M, Reka R, Miekus K, *et al*: Trafficking of normal stem cells and metastasis of cancer stem cells involve similar mechanisms: pivotal role of the SDF-1-CXCR4 axis. *Stem Cells* 23: 879-894, 2005.
- Park IH, Zhao R, West JA, *et al*: Reprogramming of human somatic cells to pluripotency with defined factors. *Nature* 451: 141-146, 2008.
- Kopp JL, Ormsbee BD, Desler M and Rizzino A: Small increases in the level of Sox2 trigger the differentiation of mouse embryonic stem cells. *Stem Cells* 26: 903-911, 2008.
- Cavallaro M, Mariani J, Lancini C, *et al*: Impaired generation of mature neurons by neural stem cells from hypomorphic Sox2 mutants. *Development* 135: 541-557, 2008.
- Eriksson M, Guse K, Bauerschmitz G, *et al*: Oncolytic adenoviruses kill breast cancer initiating CD44⁺CD24^{low} cells. *Mol Ther* 15: 2088-2093, 2007.
- Muzio G, Trombetta A, Martinasso G, *et al*: Antisense oligonucleotides against aldehyde dehydrogenase 3 inhibit hepatoma cell proliferation by affecting MAP kinases. *Chem Biol Interact* 143-144: 37-43, 2003.
- Ginestier C, Hur MH, Charafe-Jauffret E, *et al*: ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell* 1: 555-567, 2007.