

Hydrogen sulfide protects PC12 cells against reactive oxygen species and extracellular signal-regulated kinase 1/2-mediated downregulation of glutamate transporter-1 expression induced by chemical hypoxia

LIANGCAN XIAO^{1*}, AIPING LAN^{2*}, LIQIU MO¹, WENMING XU³, NAN JIANG¹,
FEN HU², JIANQIANG FENG² and CHANGGRAN ZHANG³

¹Department of Anesthesiology, The First Affiliated Hospital, Sun Yat-Sen University, Guangzhou; ²Department of Physiology, Zhongshan School of Medicine, Sun Yat-Sen University, Guangzhou; ³Department of Internal Medicine, Region of Huangpu, The First Affiliated Hospital, Sun Yat-Sen University, Guangzhou, Guangdong, P.R. China

Received May 9, 2012; Accepted June 28, 2012

DOI: 10.3892/ijmm.2012.1090

Abstract. Hypoxia and/or ischemia are implicated in neurodegenerative disorders. In these diseases, hypoxia/ischemia may induce oxidative stress, including production of reactive oxygen species (ROS), which result in a decrease in glutamate transporter expression. Hydrogen sulfide (H₂S), as the third gasotransmitter, has neuroprotective effects and potent antioxidant properties. In the present study, we investigated the role of glutamate transporter-1 (GLT-1) in the protection of H₂S against chemical hypoxia-induced injury in PC12 cells. We found that cobalt chloride (CoCl₂), a chemical hypoxia agent, reduced the expression of GLT-1 in a time-dependent manner. Pretreatment with NaHS (a donor of H₂S) reversed the CoCl₂-induced downregulation of GLT-1 expression. Pretreatment with DHK (a selective inhibitor of GLT-1) for 30 min prior to NaHS preconditioning significantly inhibited the cytoprotection of H₂S against CoCl₂-induced injuries, leading to an increase in cytotoxicity and apoptosis as well as to a loss of mitochondrial membrane potential (MMP). In addition, we found that similar to the effect of NaHS, pretreatment with NAC (a ROS scavenger) or U0126 (a MEK1/2 inhibitor)

blocked the downregulation of GLT-1 expression induced by CoCl₂. Collectively, we demonstrated for the first time that ROS and extracellular signal-regulated kinase 1/2 (ERK1/2)-mediated reduction of GLT-1 expression may be involved in chemical hypoxia-induced neural injury and that H₂S attenuates this injury partly by upregulating GLT-1 expression in PC12 cells.

Introduction

It is well documented that hypoxia and/or ischemia can elicit the release of several neurotransmitters (1,2), including glutamate (3). Such elevated levels of glutamate, and the subsequent activation of ionotropic NMDA receptors, can trigger the neuronal damage during hypoxia and/or ischemia (4,5). Glutamate homeostasis is therefore crucial to prevent neuronal death after a hypoxic/ischemic episode. Glutamate transport is the only mechanism for the removal of glutamate from the extracellular fluid in the brain (6,7), and it is essential for maintaining extracellular glutamate below neurotoxic levels in the normal brain (8). Therefore, glutamate transporters are considered to play a key role in the process of increase in extracellular glutamate during hypoxia/ischemia.

To date, 5 distinguishing high-affinity, Na⁺-dependent glutamate transporters have been identified: excitatory amino acid transporter (EAAT)1, glutamate-aspartate transporter (GLAST), EAAT2, glutamate transporter-1 (GLT-1), EAAT3, excitatory amino acid carrier 1 (EAAC1), EAAT4 and EAAT5. These transporters are present throughout the central nervous system (CNS), with GLT-1 being highly abundant in astroglial cells, whereas GLAST exists at higher levels in Bergmann glia in the cerebellum (9,10). GLT-1 plays a critical role in CNS homeostasis, accounting for up to 70% of glutamate clearance (10,11).

The roles of GLT-1 in hypoxia/ischemia-induced injury and neuroprotection have attracted extensive attention. However, the findings are controversial. A pharmacological study indicated that the GLT-1 blocker reduces the ischemia-

Correspondence to: Professor Jianqiang Feng, Department of Physiology, Zhongshan School of Medicine, Sun Yat-Sen University, 74 Zhongshan 2nd Road, Guangzhou, Guangdong 510080, P.R. China
E-mail: fengjq-sums@163.com

Professor Changran Zhang, Department of Internal Medicine, Region of Huangpu, The First Affiliated Hospital, Sun Yat-Sen University, Guangzhou, Guangdong 510700, P.R. China
E-mail: zhcr2303@sina.com

*Contributed equally

Key words: chemical hypoxia, hydrogen sulfide, glutamate transporter-1, apoptosis, oxidative stress

induced glutamate release in rat cortical superfusates (12), revealing that GLT-1 releases glutamate during ischemia. By contrast, Rao *et al* (13) reported that the antisense knockdown of GLT-1 exacerbates ischemia-triggered neuronal damage in the rat brain, suggesting that GLT-1 takes up glutamate to protect neurons during ischemia. In addition, ischemic preconditioning upregulates the GLT-1 protein which may play a role in the neuroprotective mechanism of preconditioning (14). In neonatal rats, it was shown that the neuroprotection of ceftriaxone preconditioning against hypoxia/ischemia-induced neuronal injury is associated with upregulation of GLT-1 expression (15). On the other hand, an association between change in GLT-1 expression and hypoxia/ischemia has been reported by several studies (16,17). Raghavendra *et al* (16) observed that the expression of GLT-1 is reduced following transient global ischemia. Conversely, chronic hypoxia upregulates the expression of EAAC1 and GLT-1, but not GLAST (17). These findings support the theory that GLT-1 has a complicated function (cytoprotective vs. cytotoxic effects) after hypoxic/ischemic episodes. Thus, it is necessary to explore the roles of GLT-1 in neuronal injury or the neuroprotective effects in different hypoxic/ischemic models.

Hydrogen sulfide (H₂S), recently considered a novel neuromodulator in the CNS, has been shown to protect astrocytes against H₂O₂-induced neural damage by enhancing glutamate uptake (18), suggesting an impact of H₂S on glutamate transporters. We have also demonstrated that H₂S protects PC12 cells against chemical hypoxia-induced injury by inhibiting reactive oxygen species (ROS) overproduction, extracellular signal-regulated kinase 1/2 (ERK1/2) and the p38 mitogen-activated protein kinase (MAPK) signaling pathways (19,20). Since it is reported that ROS and the activation of the ERK1/2 pathway are involved in the downregulation of GLT-1 protein expression induced by H₂O₂ or amyloid- β (A β) in astrocytes (19,21), we hypothesized that ROS and ERK1/2-mediated downregulation of GLT-1 might be implicated in chemical hypoxia-induced neuronal injury and that H₂S might confer neuroprotection by enhancing GLT-1 expression. To test this hypothesis, PC12 cells, which are derived from chromaffin cells of the adrenal medulla, were exposed to cobalt chloride (CoCl₂), a well-known hypoxia mimetic agent, to establish a model of chemical hypoxia injury. The effects of CoCl₂ and pretreatment with NaHS (a donor of H₂S) on GLT-1 expression were observed. We found that: i) CoCl₂ significantly inhibits the expression of GLT-1, ROS and the ERK1/2 pathway contribute to this inhibitory effect; ii) NaHS pretreatment clearly attenuates the inhibitory effect of CoCl₂ on GLT-1 expression; iii) DHK, a selective inhibitor of GLT-1, blocks the neuroprotection of H₂S against CoCl₂-induced injury in PC12 cells.

Materials and methods

Materials. NaHS, CoCl₂, N-acetyl-L-cysteine (NAC), Hoechst 33258, propidium iodide (PI), RNase and Rhodamine 123 (Rh123) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The cell counter kit-8 (CCK-8) was purchased from Dojindo Laboratories (Kumamoto, Japan). The DMEM medium and fetal bovine serum (FBS) were supplied by Gibco-BRL (Grand Island, NY, USA). Anti-GLT-1 antibody was purchased from Abcam (Cambridge, UK). DHK was

purchased from Merck Co. Anti- β -actin antibody, horseradish peroxidase (HRP)-conjugated secondary antibody and the BCA protein assay kit were purchased from KangChen Bio-tech, Inc. (Shanghai, China). Enhanced chemiluminescence (ECL) solution was purchased from Nanjing KeyGen Biotech Co., Inc. (Nanjing, China).

Cell culture and treatments. The rat pheochromocytoma cell line PC12 cells were purchased from the Sun Yat-Sen University Experimental Animal Center, and were grown in DMEM medium supplemented with 10% FBS at 37°C under an atmosphere of 5% CO₂ and 95% air. According to our previous study (20), chemical hypoxia was achieved by adding CoCl₂ at 600 μ M into the medium and cells were incubated in the presence of CoCl₂ for the indicated times. The cytoprotective effects of H₂S were observed by administering 400 μ M NaHS (a donor of H₂S) for 30 min prior to exposure to CoCl₂ for 24 h. NAC (a scavenger of ROS) or U0126 (a MEK1/2 inhibitor) was administered 60 or 120 min prior to exposure of the PC12 cells to 600 μ M CoCl₂ for 24 h.

Cell viability assay. The CCK-8 assay was employed to investigate the cell viability of PC12 cells cultured in 96-well plates. After the indicated treatments, 10 μ l CCK-8 solution was added to each well of the plate and the cells in the plate were incubated for 4 h in the incubator. The absorbance at 450 nm was measured with a microplate reader (Molecular Devices, Sunnyvale, CA, USA). Means of 4 well optical density (OD) in the indicated groups were used to calculate the percentage of cell viability according to the formula below: Percentage of cell viability (%) = (OD_{treatment group}/OD_{control group}) x 100%. The experiment was repeated 3 times.

Nuclear staining for assessment of apoptosis with Hoechst 33258. Morphological changes, such as chromosomal condensation and fragmentation in the nuclei of PC12 cells, were observed by Hoechst 33258 staining followed by photofluorography. Cells were plated at a density of 1x10⁶ cells/well in 35 mm dishes. Cells were preconditioned with 400 μ M NaHS for 30 min, and subsequently exposed to 600 μ M CoCl₂ for 48 h. To test the role of GLT-1 in H₂S-induced cytoprotection against chemical hypoxia-induced apoptosis, cells were treated with the GLT-1 inhibitor DHK for 30 min prior to preconditioning with NaHS. At the end of the indicated treatments, cells were harvested and fixed with 4% paraformaldehyde in 0.1 mol/l phosphate-buffered saline (PBS, pH 7.4) for 10 min. After rinsing with PBS, the nuclear DNA was stained with 5 mg/ml Hoechst 33258 solution for 10 min before being rinsed briefly with PBS and then visualized under a fluorescence microscope (Bx50-FLA; Olympus, Tokyo, Japan). Viable cells displayed a uniform blue fluorescence throughout the nucleus, whereas apoptotic cells showed condensed and fragmented nuclei.

Flow cytometric analysis of apoptosis. Treated PC12 cells were digested with trypsin (2.5 mg/ml), centrifuged at 350 x g for 10 min and the supernatant was removed. Cells were washed twice with PBS and fixed with 70% ice-cold ethanol. Cells were then centrifuged at 350 x g for 10 min, washed twice with PBS and adjusted to a concentration of

1×10^6 cells/ml. Subsequently, 0.5 ml RNase (1 mg/ml in PBS) was added to a 0.5 ml cell sample. After gentle mixing with PI (at a terminal concentration of 50 mg/l), mixed cells were filtered and incubated in the dark at 4°C for 30 min before flow cytometric analysis (FCM). The PI fluorescence of individual nuclei was measured by a flow cytometer (Beckman-Coulter, Los Angeles, CA, USA). Excitation, 488 nm; emission, 615 nm. The research software matched with FCM was used to analyze all the data of DNA labeling. In the DNA histogram, the amplitude of the sub-G1 DNA peak, which is lower than the G1 DNA peak, represents the number of apoptotic cells. The experiment was repeated 3 times.

Measurement of MMP. Mitochondrial membrane potential (MMP) was monitored using the fluorescent dye Rh123, a cell-permeable cationic dye that preferentially enters into the mitochondria based on the highly negative MMP. Depolarization of MMP results in loss of Rh123 from the mitochondria and a decrease in intracellular fluorescence. In the present study, PC12 cells were cultured in 24-well plates and treated with 400 μ M NaHS for 30 min prior to the administration of 600 μ M CoCl_2 for 24 h. DHK was administered 30 min prior to NaHS preconditioning. To evaluate MMP, Rh123 (100 μ g/l) was added to cell cultures for 45 min at 37°C and fluorescence was measured over the entire field of vision using a fluorescent microscope connected to an imaging system (BX50-FLA; Olympus). The mean fluorescence intensity (MFI) of Rh123 from 5 random fields was analyzed using ImageJ 1.410 software (National Institutes of Health, Bethesda, MD, USA), and the MFI was taken as an index of the MMP. The experiment was repeated 3 times.

Western blot assay for protein expression. After the cells were subjected to the indicated treatments, they were harvested and lysed with cell lysis solution. Total protein in the cell lysate was quantified using the BCA protein assay kit. Sample buffer was added to cytosolic extracts, and after boiling for 5 min, equal amounts of supernatant from each sample were fractionated by 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE). Total protein in the gel was transferred into polyvinylidene difluoride (PVDF) membranes. Membranes were blocked for 1.5 h at room temperature in fresh blocking buffer [0.1% Tween-20 in Tris-buffered saline (TBS-T) containing 5% fat-free milk] and then incubated with either anti-GLT-1 (1:2,500 dilution), or anti- β -actin antibodies (1:5,000 dilution) in freshly prepared TBS-T with 3% free-fat milk overnight with gentle agitation at 4°C. After 3 washes with TBS-T, membranes were incubated with HRP-conjugated goat anti-rabbit secondary antibodies (1:3,000 dilution; KangChen Bio-tech, Inc.) in TBS-T with 3% fat-free milk for 1.5 h at room temperature. Membranes were washed 3 times with TBS-T, developed in ECL solution and visualized with X-ray film. Each experiment was repeated at least 3 times. For quantification, the film were scanned and analyzed using ImageJ 1.410 software. The density of specific bands was measured and normalized with the bands of Actin. The experiment was repeated 3 times.

Statistical analysis. Data are representative of experiments performed in triplicate and are expressed as the mean \pm SE.




Figure 1. CoCl_2 induces downregulation of GLT-1 expression in PC12 cells in a time-dependent manner. (A) Time course for the effects of CoCl_2 on the expression of GLT-1 detected by western blot analysis. (B) Densitometric analysis for the results in (A). # $P<0.05$, ## $P<0.01$ vs. the control group.

Differences between groups were analyzed by one-way analysis of variance (ANOVA) using SPSS 13.0 software, followed by the LSD post hoc comparison test. $P<0.05$ was considered to indicate statistically significant differences.

Results

CoCl_2 reduces the level of GLT-1 expression in PC12 cells. In order to explore the effect of CoCl_2 on the GLT-1 expression level in PC12 cells, PC12 cells were exposed to 600 μ M CoCl_2 for the indicated times (i.e., 3, 6, 12 and 24 h). Western blot analysis revealed that treatment with 600 μ M CoCl_2 caused downregulation of GLT-1 expression in a time-dependent manner (Fig. 1). These data indicate that chemical hypoxia may reduce GLT-1 protein levels in PC12 cells.

H_2S reverses CoCl_2 -induced downregulation of GLT-1 expression in PC12 cells. After PC12 cells were exposed to 600 μ M CoCl_2 for 24 h, the levels of GLT-1 protein expression were markedly decreased (Fig. 2). However, pretreatment of PC12 cells with 400 μ M NaHS for 30 min before exposure to CoCl_2 reversed this effect, suggesting that NaHS preconditioning may enhance GLT-1 protein expression level in CoCl_2 -treated PC12 cells.

GLT-1 is involved in the cytoprotection of H_2S against CoCl_2 -induced injury. To explore whether GLT-1 is involved in the cytoprotection of H_2S against CoCl_2 -induced injuries, PC12 cells were pretreated with DHK (a inhibitor of GLT-1) at




Figure 2. Effects of H₂S and U0126 on the downregulation of GLT-1 expression induced by CoCl₂ in PC12 cells. (A) PC12 cells were treated with 600 μ M CoCl₂ for 24 h in the presence or absence of pretreatment with 400 μ M NaHS for 30 min or 10 μ M U0126, an inhibitor of MEK1/2, for 120 min. (B) Densitometric analysis for the results in (A). ^{##}P<0.01 vs. the control group; ^{**}P<0.01 vs. the CoCl₂ group.




Figure 3. Role of GLT-1 in the cytoprotection of H₂S against the cytotoxicity induced by CoCl₂ in PC12 cells. PC12 cells were treated with 600 μ M CoCl₂ for 24 h in the presence or absence of pretreatment with 400 μ M NaHS for 30 min. To inhibit GLT-1, cells were pretreated with DHK for 30 min before pretreatment with 400 μ M NaHS followed by exposure to 600 μ M CoCl₂ for 24 h. The CCK-8 assay was performed to detect cell viability. ^{##}P<0.01 vs. the control group; ^{**}P<0.01 vs. the CoCl₂ group; ^{\$\$}P<0.01 vs. the NaHS + CoCl₂ group.

400 μ M for 30 min prior to NaHS preconditioning followed by exposure to 600 μ M CoCl₂ for 24 h. As shown in Fig. 3, DHK pretreatment significantly blocked the protection of NaHS preconditioning against CoCl₂-induced cytotoxicity, the cell viability was considerably decreased, from 62 \pm 2.3% to 50 \pm 2.1% (P<0.01) (Fig. 3). Moreover, pretreatment with 400 μ M DHK also markedly inhibited H₂S-induced anti-apoptotic effects, increasing the number of apoptotic cells with nuclear condensation and fragmentation (Fig. 4A) as




Figure 4. Role of GLT-1 in the cytoprotection of H₂S against apoptosis induced by CoCl₂ in PC12 cells. (A) Morphological changes in apoptotic PC12 cells assessed by Hoechst 33258 staining. (B) Apoptosis percentage of PC12 cells was determined by flow cytometry (FCM) as described in Materials and methods. (C) The statistical results of the FCM analysis for the apoptosis of the PC12 cells. Control group, normal PC12 cells; NaHS group, cells were treated with 400 μ M NaHS for 30 min alone; DHK group, cells were treated with 400 μ M DHK for 30 min alone; CoCl₂ group, cells were treated with 600 μ M CoCl₂ for 48 h; NaHS + CoCl₂ group, cells were preconditioned with 400 μ M NaHS for 30 min followed by exposure to 600 μ M CoCl₂ for 48 h. DHK + NaHS + CoCl₂ group, cells were pretreated with 400 μ M DHK for 30 min prior to NaHS preconditioning followed by the same protocols as in the NaHS + CoCl₂ group. ^{##}P<0.01 vs. the control group; ^{**}P<0.01 vs. the CoCl₂ group; ^{\$\$}P<0.01 vs. the NaHS + CoCl₂ group.

well as the apoptotic percentage of PC12 cells compared with the NaHS pretreatment + CoCl₂ group (P<0.01) (Fig. 4B). Additionally, pretreatment of PC12 cells with DHK for 30 min before 400 μ M NaHS preconditioning clearly inhibited H₂S-induced preservation of MMP (Fig. 5). These findings




Figure 5. Role of GLT-1 in the cytoprotection of H₂S against mitochondrial damage induced by CoCl₂ in PC12 cells. MMP was measured by Rhodamine 123 (Rh123) staining followed by photofluorography. (A) Random micrographs of Rh123-derived fluorescence in PC12 cells. Control group, normal PC12 cells; NaHS group, cells were treated with 400 μ M NaHS for 30 min alone; DHK group, cells were treated with 400 μ M DHK for 30 min alone; CoCl₂ group, cells were treated with 600 μ M CoCl₂ for 24 h; NaHS + CoCl₂, cells were pretreated with 400 μ M NaHS for 30 min followed by exposure to 600 μ M CoCl₂ for 24 h. DHK + NaHS + CoCl₂ group, cells were pretreated with 400 μ M DHK for 30 min prior to NaHS preconditioning followed by the same protocols as in the NaHS + CoCl₂ group. (B) Quantitative analysis of the mean fluorescence intensity of Rh123 in each group, performed using ImageJ 1.410 software. ##P<0.01 vs. the control group; **P<0.01 vs. the CoCl₂ group; \$\$P<0.01 vs. the NaHS + CoCl₂ group.

suggest that GLT-1 contributes to the cytoprotection of H₂S against CoCl₂-induced injuries.

ROS are involved in the CoCl₂-induced downregulation of GLT-1 expression in PC12 cells. Since ROS generation inhibits glutamate uptake function (22), we examined whether ROS is involved in the CoCl₂-induced downregulation of GLT-1 protein expression in PC12 cells. Pretreatment of cells with 500 μ M NAC (a ROS scavenger) for 60 min prior to exposure to 600 μ M CoCl₂ for 24 h significantly blocked CoCl₂-induced downregulation of GLT-1 expression (Fig. 6). These data indicate that the inhibitory effect of CoCl₂ on GLT-1 expression may be associated with oxidative stress.

Activation of ERK1/2 contributes to the downregulation of GLT-1 expression induced by CoCl₂ in PC12 cells. Stimulation of ERK1/2/MAPK also contributes to the inhibition of glutamate uptake (23). In order to investigate the effect of ERK1/2 activation on the downregulation of GLT-1 expression induced by CoCl₂, PC12 cells were pretreated with 10 μ M U0126 (a MEK1/2 inhibitor) for 120 min prior to treatment with 600 μ M




Figure 6. Effect of NAC on the downregulation of GLT-1 expression induced by CoCl₂ in PC12 cells. (A) PC12 cells were treated with 600 μ M CoCl₂ for 24 h in the presence or absence of pretreatment with 500 μ M NAC for 60 min. (B) Densitometric analysis for the results in (A). ##P<0.01 vs. the control group; **P<0.01 vs. the CoCl₂ group.

CoCl₂ for 24 h. U0126 significantly reversed the inhibitory effect of CoCl₂ on the expression of GLT-1 in PC12 cells, suggesting that activation of ERK1/2 contributes to the downregulation of GLT-1 expression induced by CoCl₂ in PC12 cells (Fig. 2).

Discussion

GLT-1 has been classified as an astroglial transporter due to its predominant and widespread expression in astrocytes. In the present study, we found that PC12 cells expressed GLT-1, suggesting that GLT-1 may be involved in maintaining a normal level of glutamate in PC12 cells, which is consistent with a previous study (17). It is well known that GLT-1 plays a major role in glutamate re-uptake from the synaptic cleft after neuronal transmission (6,24,25). Lack of GLT-1 has indeed been shown to promote extracellular glutamate accumulation, excitotoxicity and, ultimately, cell death (26,27). GLT-1 has been estimated to represent up to 1% of total brain protein (6). The expression of GLT-1 is reduced in several animal models of neurodegenerative diseases, including traumatic brain injury (28) and hypoxic/ischemic insults (16,29). The levels of the GLT-1 and/or GLAST protein are also lower in the brain tissue from the patients with Alzheimer's disease (AD) (30) and Huntington's disease (31). The results of the present study showed that CoCl₂, a well-known hypoxia mimetic agent, attenuates expression of GLT-1 in a time-dependent manner. Our findings are comparable with a study showing that transient global ischemia reduces GLT-1 expression (16). Similarly, it was reported that GLT-1 protein levels are reduced in the brain in various models of central hypoxia/ischemia (16,29,32). Under hypoxic conditions (2.5 and 1% O₂ exposure for 24 h), glutamate uptake and GLT-1 protein levels

are significantly decreased in astrocytes (33). These studies all support our results. By contrast, Kobayashi and Millhorn (17) indicated that exposure of PC12 cells to hypoxia (1% O₂) for 6 to 24 h increases GLT-1 protein levels. Therefore, it is likely that the effects of hypoxia/ischemia on the expression of GLT-1 may be affected by many factors, including tissue or cell types, the level of hypoxia, manner of hypoxia induction and also the period of hypoxia/ischemia.

To clarify the mechanisms underlying the inhibitory effect of chemical hypoxia on GLT-1 expression, we tested the possible involvement of ROS. Several previous studies have shown that oxidative stress is implicated in glutamate clearance impairment and reduction of GLT-1 expression (18,21,34). Our recent studies have demonstrated the promotive effects of CoCl₂ on ROS production (19,20). In this study, we found that NAC, a ROS scavenger, can significantly block the inhibition of GLT-1 expression induced by CoCl₂, revealing that ROS partly contribute to the inhibitory effect of chemical hypoxia on the expression of GLT-1 in PC12 cells. We provide novel evidence for the role of ROS in CoCl₂-induced neuronal injury. Additionally, there is currently a lot of data demonstrating that oxidative stress may trigger and modulate the MAPK signaling pathways (21,35,36). We recently demonstrated that ROS can activate the MAPK pathway (20), linking to the possibility of an altered ERK1/2 activation that ultimately affects the expression of GLT-1. To confirm this possibility, we observed the effects of pretreatment of PC12 cells with U0126 (an inhibitor of MEK1/2) on the inhibition of GLT-1 expression by CoCl₂. Our results showed that U0126 clearly suppressed the CoCl₂-induced decrease in the expression of GLT-1, suggesting that the ERK1/2 pathway is involved in the inhibitory effect of CoCl₂ on GLT-1 expression. This is also a novel finding showing that the ROS-activated ERK1/2 pathway plays a role in the inhibition of GLT-1 expression by CoCl₂. Our findings are supported by previous studies (18,35). Lu *et al* (18) reported that PD98059, a specific ERK1/2 inhibitor, significantly reverses the reduction of trafficking of GLT-1 from cytoplasm to plasma membrane.

Although research on the regulatory mechanisms for GLT-1 expression has intensified, scarce data are available regarding the regulatory effect of gasotransmitter on the expression of GLT-1. H₂S, recently recognized as the third gasotransmitter alongside nitric oxide (NO) and carbon monoxide (CO) (39), has attracted extensive attention due to its multiple physiological and pathophysiological roles in various body systems (18-20,37-41). Kimura and Kimura (38) demonstrated the neuroprotective effect of H₂S against oxidative stress-induced injury in primary rat cortical neurons. H₂S also protects astrocytes from H₂O₂-induced neural injury (18). We recently found that H₂S protects PC12 cells against CoCl₂-induced damage by enhancing heat shock protein 90 (HSP90) (19), inhibiting the ROS-activated ERK1/2 and p38MAPK signaling pathways (20) and scavenging ROS (19,20). In the present study, we provide evidence for the first time that NaHS (a donor of H₂S) pretreatment prevents the CoCl₂-induced downregulation of GLT-1 expression in PC12 cells. Our results are in line with a recent study that H₂S protects astrocytes against oxidative stress-induced neural damage by increasing glutamate uptake (18). Based on our recent results (19,20,39-41) and other studies (18,21,35,36,38,42), there are several possible mechanisms

responsible for the regulatory effect of H₂S on the expression of GLT-1: i) its antioxidation, by which H₂S can protect PC12 cells from CoCl₂-induced suppression of GLT-1 expression; ii) its inhibitory effect on the ERK1/2 pathway (20); and iii) H₂S functions as an ATP-sensitive potassium (K_{ATP}) channel opener (43). Hu *et al* (42) indicated that iptakalim, a K_{ATP} channel opener, can reverse the inhibition of glutamate uptake induced by N-methyl-4-4-phenylpyridinium (MPP⁺) [used to stimulate Parkinson's disease (PD)-like conditions], revealing a role of the K_{ATP} channel opener in the functional regulation of glutamate transporter. Further research is required to confirm these findings.

We further explored the role of GLT-1 in the neuroprotection of H₂S against chemical hypoxia-induced injury. We found that pretreatment with DHK, a selective inhibitor of GLT-1, significantly reversed the protective effect of H₂S against CoCl₂-induced injuries, evidenced by a decrease in cell viability and an increase in apoptotic PC12 cells as well as MMP loss, suggesting that upregulation of GLT-1 expression may play an important role in the neuroprotective effects of H₂S.

In summary, in the present study, we have demonstrated for the first time that: i) both ROS and the ERK1/2 pathway contribute to the downregulation of GLT-1 expression induced by CoCl₂; ii) H₂S, a novel gaseous neuromodulator, reverses CoCl₂-induced downregulation of GLT-1 expression; and iii) upregulation of GLT-1 expression may play a crucial role in the neuroprotective effects of H₂S against chemical hypoxia-induced neuronal injury in PC12 cells. The findings of the present study may provide a potential neuroprotective therapeutic approach for treatment of hypoxia/ischemia-related neuronal injury. In addition, based on the notable findings that both levels of endogenous H₂S and GLT-1 are reduced in neurodegenerative diseases, such as AD and PD, we speculate that endogenous H₂S may be an important modulator of GLT-1. These findings remain to be confirmed in future studies.

Acknowledgements

The present study was supported by the Guangdong Science and Technology Planning project (nos. 2010B080701105, 2009B080701014 and 2007B080701030).

References

1. Nilsson GE and Lutz PL: Release of inhibitory neurotransmitters in response to anoxia in turtle brain. *Am J Physiol* 261: R32-R37, 1991.
2. Richter DW, Lalley PM, Pierrefiche O, *et al*: Intracellular signal pathways controlling respiratory neurons. *Respir Physiol* 110: 113-123, 1997.
3. Nicholls D and Attwell D: The release and uptake of excitatory amino acids. *Trends Pharmacol Sci* 11: 462-468, 1990.
4. Rothman SM and Olney JW: Glutamate and the pathophysiology of hypoxic - ischemic brain damage. *Ann Neurol* 19: 105-111, 1986.
5. Sattler R, Xiong Z, Lu WY, *et al*: Distinct roles of synaptic and extrasynaptic NMDA receptors in excitotoxicity. *J Neurosci* 20: 22-33, 2000.
6. Lehre KP and Danbolt NC: The number of glutamate transporter subtype molecules at glutamatergic synapses: chemical and stereological quantification in young adult rat brain. *J Neurosci* 18: 8751-8757, 1998.
7. Tanaka K: Expression cloning of a rat glutamate transporter. *Neurosci Res* 16: 149-153, 1993.

8. Attwell D, Barbour B and Szatkowski M: Nonvesicular release of neurotransmitter. *Neuron* 11: 401-407, 1993.
9. Kanai Y, Smith CP and Hediger MA: A new family of neurotransmitter transporters: the high-affinity glutamate transporters. *FASEB J* 7: 1450-1459, 1993.
10. Anderson CM and Swanson RA: Astrocyte glutamate transport: review of properties, regulation, and physiological functions. *Glia* 32: 1-14, 2000.
11. Gegelashvili G and Schousboe A: High affinity glutamate transporters: regulation of expression and activity. *Mol Pharmacol* 52: 6-15, 1997.
12. Phillis JW, Ren J and O'Regan MH: Transporter reversal as a mechanism of glutamate release from the ischemic rat cerebral cortex: studies with DL-threo-beta-benzoyloxyaspartate. *Brain Res* 868: 105-112, 2000.
13. Rao VLR, Dogan A, Todd JG, *et al*: Antisense knockdown of the glial glutamate transporter GLT-1, but not the neuronal glutamate transporter EAAC1, exacerbates transient focal cerebral ischemia-induced neuronal damage in rat brain. *J Neurosci* 21: 1876-1883, 2001.
14. Zhang G, Raol YS, Hsu FC and Brooks-Kayal AR: Long-term alterations in glutamate receptor and transporter expression following early-life seizures are associated with increased seizure susceptibility. *J Neurochem* 88: 91-101, 2004.
15. Mimura K, Tomimatsu T, Minato K, *et al*: Ceftriaxone preconditioning confers neuroprotection in neonatal rats through glutamate transporter 1 upregulation. *Reprod Sci* 18: 1193-1201, 2011.
16. Raghavendra Rao VL, Rao AM, Dogan A, *et al*: Glial glutamate transporter GLT-1 downregulation precedes delayed neuronal death in gerbil hippocampus following transient global cerebral ischemia. *Neurochem Int* 36: 531-537, 2000.
17. Kobayashi S and Millhorn DE: Hypoxia regulates glutamate metabolism and membrane transport in rat PC12 cells. *J Neurochem* 76: 1935-1948, 2001.
18. Lu M, Hu LF, Hu G and Bian JS: Hydrogen sulfide protects astrocytes against H₂O₂-induced neural injury via enhancing glutamate uptake. *Free Radic Biol Med* 4: 1705-1713, 2008.
19. Meng JL, Mei WY, Dong YF, *et al*: Heat shock protein 90 mediates cytoprotection by H₂S against chemical hypoxia-induced injury in PC12 cells. *Clin Exp Pharmacol Physiol* 38: 42-49, 2011.
20. Lan A, Liao X, Mo L, *et al*: Hydrogen sulfide protects against chemical hypoxia-induced injury by inhibiting ROS-activated ERK1/2 and p38MAPK signaling pathways in PC12 cells. *PLoS One* 6: e25921, 2011.
21. Matos M, Augusto E, Oliveira CR and Agostinho P: Amyloid-beta peptide decreases glutamate uptake in cultured astrocytes: involvement of oxidative stress and mitogen-activated protein kinase cascades. *Neuroscience* 156: 898-910, 2008.
22. Sun XL, Zeng XN, Zhou F, *et al*: K(ATP) channel openers facilitate glutamate uptake by GluTs in rat primary cultured astrocytes. *Neuropsychopharmacology* 33: 1336-1342, 2008.
23. Figiel M, Maucher T, Rozyczka J, *et al*: Regulation of glial glutamate transporter expression by growth factors. *Exp Neurol* 183: 124-135, 2003.
24. Lehre KP, Levy LM, Ottersen OP, *et al*: Differential expression of two glial glutamate transporters in the rat brain: quantitative and immunocytochemical observations. *J Neurosci* 5: 1835-1853, 1995.
25. Kugler P and Schmitt A: Glutamate transporter EAAC1 is expressed in neurons and glial cells in the rat nervous system. *Glia* 27: 129-142, 1999.
26. Tanaka K, Watase K, Manabe T, *et al*: Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 276: 1699-1702, 1997.
27. Vorwerk CK, Naskar R, Schuettauf F, *et al*: Depression of retinal glutamate transporter function leads to elevated intravitreal glutamate levels and ganglion cell death. *Invest Ophthalmol Vis Sci* 41: 3615-3621, 2000.
28. Rao VL, Başkaya MK, Doğan A, *et al*: Traumatic brain injury downregulates glial glutamate transporter (GLT-1 and GLAST) proteins in rat brain. *J Neurochem* 70: 2020-2027, 1998.
29. Torp R, Lekieffre D, Levy LM, *et al*: Reduced postischemic expression of a glial glutamate transporter, GLT1, in the rat hippocampus. *Exp Brain Res* 103: 51-58, 1995.
30. Li S, Mallory M, Alford M, Tanaka S and Masliah E: Glutamate transporter alterations in Alzheimer disease are possibly associated with abnormal APP expression. *J Neuropathol Exp Neurol* 56: 901-911, 1997.
31. Lipton SA and Rosenberg PA: Excitatory amino acids as a final common pathway for neurologic disorders. *N Engl J Med* 330: 613-622, 1994.
32. Martin LJ, Brambrink AM, Lehmann C, *et al*: Hypoxia-ischemia causes abnormalities in glutamate transporters and death of astroglia and neurons in newborn striatum. *Ann Neurol* 42: 335-348, 1997.
33. Dallas M, Boycott HE, Atkinson L, *et al*: Hypoxia suppresses glutamate transport in astrocytes. *J Neurosci* 27: 3946-3955, 2007.
34. Brera B, Serrano A and de Ceballos ML: beta-amyloid peptides are cytotoxic to astrocytes in culture: a role for oxidative stress. *Neurobiol Dis* 7: 395-405, 2000.
35. McCubrey JA, Lahair MM and Franklin RA: Reactive oxygen species-induced activation of the MAP kinase signaling pathways. *Antioxid Redox Signal* 8: 1775-1789, 2006.
36. Zhu X, Lee HG, Raina AK, *et al*: The role of mitogen-activated protein kinase pathways in Alzheimer's disease. *Neurosignals* 11: 270-281, 2002.
37. Wang R: The gasotransmitter role of hydrogen sulfide. *Antioxid Redox Signal* 5: 493-501, 2003.
38. Kimura Y and Kimura H: Hydrogen sulfide protects neurons from oxidative stress. *FASEB J* 18: 1165-1167, 2004.
39. Yang Z, Yang C, Xiao L, *et al*: Novel insights into the role of HSP90 in cytoprotection of H₂S against chemical hypoxia-induced injury in H9c2 cardiac myocytes. *Int J Mol Med* 28: 397-403, 2011.
40. Chen SL, Yang CT, Yang ZL, *et al*: Hydrogen sulphide protects H9c2 cells against chemical hypoxia-induced injury. *Clin Exp Pharmacol Physiol* 37: 316-321, 2010.
41. Yang CT, Yang ZL, Zhang MF, *et al*: Hydrogen sulfide protects against chemical hypoxia-induced cytotoxicity and inflammation in HaCaT cells through inhibition of ROS/NFκB/COX-2 pathway. *PLOS One* 6: e21971, 2011.
42. Hu LF, Wang S, Shi XR, *et al*: ATP-sensitive potassium channel opener iptakalim protected against the cytotoxicity of MPP⁺ on SH-SY5Y cells by decreasing extracellular glutamate level. *J Neurochem* 94: 1570-1579, 2005.
43. Johansen D, Ytrehus K and Baxter GF: Exogenous hydrogen sulfide (H₂S) protects against regional myocardial ischemia-reperfusion injury - evidence for a role of K_{ATP} channels. *Basic Res Cardiol* 101: 53-60, 2006.