

Protective effect of 7-difluoromethoxy-5,4'-Di-hydroxyl isoflavone against the damage induced by glutamate in PC12 cells

XING FENG, QIN ZHOU, CHONG LIU, MEI-LING TAO, JIAN-GUO CAO and ZHI-HONG ZHONG

College of Medicine, Hunan Normal University, Changsha, Hunan 410006, P.R. China

Received May 24, 2012; Accepted July 13, 2012

DOI: 10.3892/ijmm.2012.1109

Abstract. 7-difluoromethoxy-5,4'-Di-hydroxyl isoflavone (dFGEN), prepared by the difluoromethylation of genistein, is an active chemical entity. In this study, our main purpose was to investigate whether dFGEN had an effect on glutamate-induced apoptosis in cultured PC12 cells. The PC12 cells were treated with different glutamate concentrations for 24 h *in vitro*. The PC12 cells impaired by glutamate were used as the cell model of excitability. Cells were incubated for 30 min with genistein, dFGEN, vitamin E, and exposed to 10 mM glutamate for 24 h. Cell morphology was observed by light microscopy. The growth and proliferation of PC12 cells were detected by 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) assay. Cell apoptosis was determined by flow cytometry (FCM) with propidium iodide (PI) staining. The activity of lactate dehydrogenase (LDH), superoxide dismutase (SOD) and the content of malondialdehyde (MDA) were measured by kits, respectively. Acridine orange (AO) staining was used to detect characteristics of cell apoptosis. When PC12 cells were incubated with glutamate for 24 h, cells appeared to have significant changes in shape. The cellular viability was reduced and the apoptotic rate was increased. The levels of LDH and the content of MDA were increased. The activity of SOD was decreased. After PC12 cells were pretreated with dFGEN, dFGEN significantly improved cell morphology, cell growth and proliferation, suppressed apoptosis of cells, reduced the release of LDH, improving SOD activity and decreased MDA content in a concentration-dependent manner. AO staining displayed that apoptosis was decreased. These results suggested that dFGEN has a protective effect against glutamate-induced damage in PC12 cells. dFGEN seemed to have a better protective effect than the lead compound genistein in a concentration-dependent manner. The mechanism of protective effect of dFGEN may be mainly related to its antioxidative activity.

Introduction

With the improvement of people's living standard, social aging trend will be more obvious, so senile disease rates are on the rise. Neurodegenerative diseases is a progressive disease associated with age in nervous system, including Alzheimer's disease (AD), Parkinson's disease (PD), amyotrophic lateral sclerosis (ALS) and multiple sclerosis (MS). Statistics indicate that thirty-four million people will have AD in the world by the year 2025 (1). It is estimated that the central nervous system (CNS), such as AD, PD, brain cancer, and stroke, will cost trillions of dollars for their treatment (2). Alzheimer's disease is the most common type of dementia. The United States statistics show that AD affects 5.4 million people and ~200,000 people under the age 65 have younger-onset AD (3). The clinical presentation is mainly memory loss and cognitive decline. Currently available drugs for the treatment of AD are purely for symptoms (4) and among these drugs are most the cholinesterase inhibitors (5). Another type of drug available for AD patients is an N-methyl-d-aspartate (NMDA) receptor antagonist named memantine (6). Many drugs are available to treat AD currently, but the effect is not significant or serious side effect may be occurs. We still lack specially effective drug treatment for the AD patients. Therefore, it is a very important responsibility and obligation for the pharmaceutical industry to study a new central nervous system drugs with high quality, high efficiency and low side effects.

In recent years, a large number of research data show that many acute and chronic neurodegenerative diseases are association with extracellular abnormally gathered glutamic acid in the brain (7). Glutamate, an excitatory amino acid, is one of the major neurotransmitters in the central nervous system (CNS). Glutamate participate in multiple physiological pathology processes. The right amount of glutamate is required to maintain normal cell physiological activities. However, high concentrations of glutamate in the brain can lead to neuronal damage. Glutamate are thought to be involved in the etiology of a number of neurodegenerative disorders including AD, PD, ALS, MS. A large amount of glutamate cause excessive activation of NMDA receptors, leading to calcium overload which can trigger a cascade of events eventually leading to apoptosis or necrosis (8).

Genistein (4',5,7-trihydroxyisoflavone) (Fig. 1), plentiful in soybeans, has estrogenic activity. Physiological levels of estrogen have many significant biological activities which are beneficial for treatment of osteoporosis and menopausal symptoms (9), AD in women after menopause (10), inflammatory response (11,12),

Correspondence to: Dr Xing Feng, College of Medicine, Hunan Normal University, Changsha, Hunan 410006, P.R. China
E-mail: fengxing01@hotmail.com

Key words: 7-difluoromethoxy-5,4'-Di-hydroxylisoflavone, genistein, glutamate, PC12 cells

atherosclerosis (13), liver cancer (14), colon cancer (15), and it is neuro-protective (16). Genistein has a strong protective effect on the damage induced by oxidative stress (17). However, the absorption of genistein in the gastrointestinal tract is poor, resulting in low biological activity (18). To overcome this problem, the CHF2 group was introduced into the lead compound genistein, which can change its chemical and physical properties (19). After the introduction of CF₂ into genistein, Fu *et al.* (20) design and synthesized a series of difluoromethyl-derivatives of genistein to screen an effective drug with oxidative stress injury model. The results confirmed that 7-difluoromethyl-genistein is a protective new chemical entity for injury model induced by oxidative stress (21).

PC12 is a cell line derived from a rat adrenal medulla pheochromocytoma. PC12 cells have typical characteristics of nerve cells, which can be widely used as a nerve cell model. There are many mechanisms of nerve cell injury, one of which is the glutamate-induced damage. In this study, we used MTT assay to measure cell growth and proliferation activity, flow cytometry (FCM) with propidium iodide (PI) staining and acridine orange (AO) staining to detect cell apoptosis, assay kits to detect LDH activity, SOD activity and MDA content. Then we investigated the effect of 7-difluoromethoxy-5,4'-Di-hydroxyl isoflavone (dFGEN) on PC12 cells induced by glutamate.

7-difluoromethoxy-5,4'-dihydroxy isoflavone (dFGEN) is a genistein derivative (332 kDa) that had 7-OH group substituted by -OCHF₂. Its formula is shown in Fig. 2.

Materials and methods

Reagents. dFGEN (98% pure) was synthesized as reported in our laboratory (20). Genistein and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) were purchased from Sigma (USA). Glutamate, vitamin E and Dimethyl sulfoxide (DMSO) were purchased from Genview (USA). Dulbecco's minimum essential medium (DMEM) were obtained from Hyclone (USA). Fetal calf serum (FBS) was purchased from Hangzhou Sijiqing Biological Engineering Materials Co. (Hangzhou, China). Acridine orange was purchased from Sinopharm Chemical Reagent Co., Ltd. (China). Cell lysis buffer, trypan blue staining, penicillin and streptomycin were purchased from Beijing Dingguo Changsheng Biotech Co., Ltd. (China). Lactate dehydrogenase (LDH) assay kit, superoxide dismutase (SOD) assay kit, lipid peroxidation (MDA) assay kit were purchased from Nanjing Jiancheng Bioengineering Institute (Nanjing, China). BCA protein assay kit and trypsin were purchased from Beyotime Institute of Biotechnology (Shanghai, China). Cell culture plates and cell culture dishes were purchased from Corning Inc. (USA).

Cell culture and treatment. PC12 cells (rat adrenal pheochromocytoma cells) were purchased from Cell Bank, Chinese Academy of Sciences (Shanghai, China). Cells were cultured in DMEM medium supplemented with 10% FBS, 100 U/ml penicillin and 100 µg/ml streptomycin at 37°C in a humidified atmosphere of 5% CO₂ incubator. When cells were ~80% confluent, new media with 1% newborn calf serum were added before the drug treatment. Glutamate was added in final concentrations ranging from 1 to 20 mM in

the pilot study, and the concentration (10 mM) was selected by determining dose-response curves. When needed, cells were incubated for 30 min with genistein (0.1, 1.0 and 10 µM), dFGEN (0.1, 1.0 and 10 µM), vitamin E (10 µM), and then exposed to 10 mM glutamate for 24 h.

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. The PC12 cells were seeded in 96-well plates at a density of 1x10⁴ cells/well. After 24 h, the cells were treated with various concentrations of genistein, dFGEN, vitamin E for 30 min prior to glutamate (10 mM) treatment for 24 h. Briefly, 20 µl MTT was added to each well at a final concentration of 0.5 mg/ml, and afterwards the cells were cultured for 4 h at 37°C. The medium was then carefully removed and 150 µl of DMSO was added to each well. The absorbance at 490 nm wavelength (A₄₉₀) was measured with enzyme-linked immunosorbent instrument (ELX800, Bio-Tek, USA). Experiment was divided into zero setting group, control group, and experimental group. The cell viability was expressed as a percentage of the viability of the control culture. Relative cell proliferation inhibition rate (IR) = (1 - average A₄₉₀ of the experimental group/average A₄₉₀ of the control group) x 100%.

Flow cytometry (FCM) with propidium iodide (PI) staining. The PC12 cells were seeded in 6-well plates at a density of 2x10⁵/ml (2 ml/well) and incubated for 24 h. Cells were treated by drugs in the same way as described above. Cells were treated for 24 h, then collected and harvested with 0.25% trypsin and made into a single cell suspension, washed with cold PBS twice. Cells were resuspended as a single cell suspension with 50 µl PBS, fixed with 1 ml of cold 70% ethanol, stained with propidium iodide (PI) and cell apoptosis was detected using flow cytometry (FC 500 American Beckman Coulter Co.).

Morphological observations and acridine orange staining. The PC12 cells were seeded in 6-well plate at a density of 2x10⁵/ml (2 ml/well) and incubated for 24 h. Cells were treated by drugs in the same way as described above. After cells were treated for 24 h, cell morphological changes were observed under an optical microscope. Then the cells were washed two times with cold PBS and incubated with AO (5 µg/ml) at room temperature for 10 min in the dark. The stained cells were observed using fluorescence microscope (Olympus BX41, Japan Olympus Co.) and images were taken.

Detection of LDH activity. The PC12 cells were seeded in 24-well plates at a density of 1.2x10⁵/ml (1 ml/well) and incubated for 24 h. Cells were treated by drugs in the same way as described above. Cells were treated for 24 h, the culture supernatants were collected to a 1 ml Eppendorf tube, and LDH activity was detected at 450 nm by the assay kit.

Detection of SOD activity and MDA content. The PC12 cells were seeded in 66-mm culture dish at a density of 1x10⁶/ml (4 ml/dish) and incubated for 24 h. Cells were treated by drugs in the same way as described above. Cells were treated for 24 h, then collected, washed two times with cold PBS, lysed at 4°C for 30 min in lysate and then centrifuged at 12,000 x g for 10 min at 4°C. SOD activity and MDA contents were measured according to the direction of the assay kit.




Figure 1. The structure of genistein.




Figure 2. The structure of 7-difluoromethoxy-5,4'-Di-hydroxyl isoflavone (dFGEN).




Figure 3. The effect of dFGEN on the proliferation of PC12 cells by MTT assay. The PC12 cells were treated with various concentrations of dFGEN (0.1, 1.0 and 10 μ M), genistein (0.1, 1.0 and 10 μ M) or vitamin E (10 μ M) for 30 min followed by the addition of glutamate to a final concentration of 10 mM, and incubated for 24 h. The cell inhibitory rate was detected using MTT assay. * $p < 0.05$ vs glutamate 10 mM; ** $p < 0.01$ vs glutamate 10 mM; # $p < 0.05$ vs the same concentration of genistein + glutamate 10 mM; ▲ $p < 0.05$ vs the same concentration of VE + glutamate 10 mM.

Statistical analysis. Data are presented as the mean \pm SD. The database were set up with the SPSS 16.0 software package for

analysis. The means of multiple groups were compared with one-way ANOVA, the two-two comparisons among the means were performed with LSD t-test. $P < 0.05$ was considered as statistically significant.

Results


Effects of dFGEN on glutamate-induced PC12 cell proliferation. As shown in Fig. 3, the MTT assay demonstrates that glutamate obviously inhibited proliferation of PC12 cells compared with the control group ($p < 0.01$). Results showed that dFGEN could effectively increased proliferation of PC12 cells in a concentration-dependent manner. Compared with the same concentration of genistein (1.0 and 10 μ M), dFGEN (1.0 and 10 μ M) could distinctly increase proliferation of PC12 cells ($p < 0.05$). The cell viability of 10 μ M dFGEN was $> 10 \mu$ M vitamin E ($p < 0.05$).

Effects of dFGEN on glutamate-induced PC12 cell apoptosis. As shown in Fig. 4, the percentage of apoptotic cells was increased from 0.29 ± 0.06 to $18.82 \pm 1.47\%$ after the cells were exposed to 10 mM glutamate for 24 h ($p < 0.05$). dFGEN effectively decreased apoptosis of PC12 cells in a concentration-dependent manner. dFGEN (10 μ M) distinctly decreased glutamate-induced PC12 cell apoptosis compared to the same concentration of genistein and vitamin E ($p < 0.05$).

Effects of dFGEN on glutamate-induced PC12 cell morphology and apoptotic morphology. As shown in Fig. 5, glutamate-induced PC12 cells exhibited significantly morphological alterations under microscopy and they became smaller and irregularly shaped. dFGEN obviously improved cell morphology. DNA-binding dye acridine orange (AO) was used to observe the morphological characteristic of apoptotic cells. As shown in Fig. 6, AO staining displayed karyopyknosis, chromatin condensation and apoptotic bodies by fluorescence microscopy in glutamate treated group. Dense staining yellow-green fluorescence and granules could be seen under the fluorescence microscope. dFGEN evidently decreased the glutamate-induced apoptosis of PC12 cells.

Effects of dFGEN on glutamate-induced PC12 cell LDH activity. As shown in Fig. 7, while glutamate (10 mM) increased the release of LDH to 492.58 ± 8.48 U/l ($p < 0.01$), dFGEN reduced their release in a concentration-dependent manner. Compared with the same concentration of genistein (1.0 and 10 μ M), dFGEN (1.0 and 10 μ M) distinctly decreased LDH activity of PC12 cells ($p < 0.01$). The LDH activity of 10 μ M dFGEN group was lower than in the 10 μ M vitamin E group ($p < 0.01$).

Effects of dFGEN on glutamate-induced PC12 cell SOD activity and MDA content. As shown in Fig. 8, glutamate treatment for 24 h induced an increase in MDA formation but decreased SOD enzyme activity in PC12 cells ($p < 0.01$). While dFGEN significantly attenuated lipid peroxidation in a concentration-dependent manner. Compared with the same concentration of genistein (1.0 and 10 μ M), dFGEN (1.0 and 10 μ M) distinctly improved SOD activity and decreased MDA content ($p < 0.01$). dFGEN attenuated lipid peroxidation (10 μ M) compared with 10 μ M vitamin E ($p < 0.01$).



Discussion

Aging is accompanied by many neurodegenerative diseases. A consistent number of neurons undergo apoptosis in the aged brain, which may lead to neurodegeneration in the long-term (22). Glutamate is the major excitatory neurotransmitter in the CNS. In many neurodegenerative diseases, glutamate causes over-excited glutamate receptors, causing glutamate excitotoxicity, which can trigger a cascade of events eventually leading to apoptosis or necrosis (23). Data suggest that glutamate excitotoxicity is an apoptotic process (24), while some data suggest that the excitotoxicity death is only due to the mechanism of necrosis (25). Although mechanism of glutamate-induced excitotoxicity is not clear, a large number of data suggest that the mechanism is related to glutamate-induced oxidative stress (26,27). In this study, we used glutamate (10 mM) to establish PC12 cell injury




Figure 4. The effect of dFGEN on glutamate-induced PC12 cell apoptosis. The PC12 cells were treated with various concentrations of dFGEN (0.1, 1.0 and 10 μ M), genistein (0.1, 1.0 and 10 μ M) or vitamin E (10 μ M) for 30 min followed by the addition of glutamate to a final concentration of 10 mM, and incubated for 24 h. Cell apoptosis was detected by FCM with PI staining. * p <0.05 vs glutamate 10 mM; ** p <0.01 vs glutamate 10 mM; # p <0.05 vs the same concentration of genistein + glutamate 10 mM; ▲ p <0.05 vs the same concentration of VE + glutamate 10 mM. (b) A, Control; B, Vehicle; C, Glu 10 mM; D, genistein 0.1 μ M + Glu 10 mM; E, genistein 1 μ M + Glu 10 mM; F, genistein 10 μ M + Glu 10 mM; G, dFGEN 0.1 μ M + Glu 10 mM; H, dFGEN 1 μ M + Glu 10 mM; I, dFGEN 10 μ M + Glu 10 mM; J, VE 10 μ M + Glu 10 mM.




Figure 5. The effect of dFGEN on glutamate-induced PC12 cell morphology (10x10). The PC12 cells were treated with various concentrations of dFGEN (0.1, 1.0 and 10 μ M), genistein (0.1, 1.0 and 10 μ M) or vitamin E (10 μ M) for 30 min followed by the addition of glutamate to a final concentration of 10 mM, and incubated for 24 h. (A) Control; (B) 10 mM Glu; (C) 10 μ M genistein; (D) 10 μ M dFGEN; (E) 10 μ M VE.




Figure 6. Morphological assessment of apoptosis by AO staining. a fluorescence microscope (10x10) images. The PC12 cells were treated with various concentrations of dFGEN (0.1, 1.0 and 10 μ M), genistein (0.1, 1.0 and 10 μ M) or vitamin E (10 μ M) for 30 min followed by the addition of glutamate to a final concentration of 10 mM, and incubated for 24 h. (A) Control; (B) 10 mM Glu; (C) 10 μ M genistein; (D) 10 μ M dFGEN; (E) 10 μ M VE.

model, and then to explore whether dFGEN have protective effect on damaged nerve cells compared with genistein and vitamin E.

Mitochondria is considered to be very important during cell apoptosis. Mitochondrial activity can be measured with MTT assay. In this study, pretreatment of dFGEN significantly improved cell survival and reduced the rate of proliferation inhibition in a concentration-dependent manner, suggesting that dFGEN could effectively maintain mitochondrial structure and function. dFGEN antagonized the effect of glutamate on cell proliferation inhibition rate, which is more effective than genistein and vitamin E. This study provided the experimental basis for inhibition of apoptosis and antioxidant function of dFGEN.

Propidium iodide (PI) is a nuclear fluorescent dye used for staining DNA. PI can not pass through viable cell membrane, but it can pass through the damaged membrane, used in cell apoptosis detection, which can be used for detection of cell apoptosis. Our study showed that dFGEN decreased the cell apoptosis rate in a concentration-dependent manner. Acridine orange (AO) staining showed that dFGEN evidently decreased the glutamate-induced apoptosis of PC12 cells. We proposed that dFGEN may play a positive role in the process of glutamate-induced PC12 cell apoptosis.




Figure 7. The effect of dFGEN on glutamate-induced LDH activity in PC12 cells. The PC 12 cells were treated with various concentrations of dFGEN (0.1, 1.0 and 10 μ M), genistein (0.1, 1.0 and 10 μ M) or vitamin E (10 μ M) for 30 min followed by the addition of glutamate to a final concentration of 10 mM, and incubated for 24 h. The cell culture supernatant LDH activity was detected by LDH assay kit. * p <0.01 vs glutamate 10 mM; # p <0.01 vs the same concentration of genistein + glutamate 10 mM; Δ p <0.01 vs the same concentration of VE + glutamate 10 mM.




Figure 8. The effect of dFGEN on glutamate-induced SOD activity and MDA content in PC12 cells. The PC12 cells were treated with various concentrations of dFGEN (0.1, 1.0 and 10 μ M), genistein (0.1, 1.0 and 10 μ M) or vitamin E (10 μ M) for 30 min followed by the addition of glutamate to a final concentration of 10 mM, and incubated for 24 h. The activity of SOD and the content of MDA were detected by SOD assay kit and MDA assay kit. * $p < 0.01$ vs glutamate 10 mM; # $p < 0.01$ vs the same concentration of genistein + glutamate 10 mM; ## $p < 0.01$ vs the same concentration of VE + glutamate 10 mM.

LDH activity is a classic indicator that reflects cell function. The more LDH release into extracellular space the greater damage to cells. Schreihöfer and Redmond (28) reported that phytoestrogen genistein can reduce LDH release of cortical cells induced by glutamate. The results in this study showed that genistein, dFGEN and vitamin E could reduce LDH release of PC12 cells to different degrees. Compared with the same concentration of genistein and vitamin E, dFGEN could distinctly decrease LDH activity of PC12 cells and reduce the degree of cell damage. Our results indicated that dFGEN has a protective effect better than the lead compound genistein and vitamin E in a dose-dependent manner.

Oxidative stress is an important factor in a neurodegenerative disease. The loss of redox system balance will lead to oxidative damage of the body. SOD activity and MDA content are the indicators reflecting free radical damage. Our study showed that dFGEN increase SOD activity and decrease MDA content more than the same concentrations of genistein and vitamin E. The results suggested that dFGEN could protect PC12 cells against oxidative stress by increasing capacity of scavenging the free radicals, reducing the damage of free radicals on the cells, inhibiting changes in cell membrane permeability. This might represent one of the mechanisms for dFGEN against glutamate induced damage.

The process of glutamate-induced PC12 cell injury involves multiple signal transduction pathways, and the mechanism is very complex. Numerous studies show that Bcl-2 gene family (29) and caspase proteinase family (30) play important roles in the apoptotic process. Some reports have shown that apoptosis of PC12 cells may be related to the ERK1/2 and P38 signaling pathway (31). Our present preliminary study shows that dFGEN protects PC12 cells against glutamate toxicity. Its anti-apoptotic mechanism and *in vitro* experiments still need further research, so that ultimately our goal for dFGEN applied to neurodegenerative disease treatment can be achieved.

In conclusion, dFGEN can improve cell morphology, improve cell viability, suppress the apoptosis of cells, reduce the release of LDH, improve SOD activity and decrease MDA content in a concentration-dependent manner. These data demonstrated that dFGEN provide better protection against PC12 cell injury caused by glutamate than the lead compound genistein in a concentration-dependent manner. The mechanism of protective effect of dFGEN may be mainly related to its antioxidative activities.

Acknowledgements

This project was supported by Provincial Science and Technology Plan of Hunan, China (No2010FJ3017), Research Foundation of Education Bureau of Hunan Province, China (No11B081), Chinese Traditional Medicine Administration of Hunan Province, China (No 2009102), Program Excellent Talent Hunan Normal University (2011).

References

- Hynd MR, Scott HL and Dodd PR: Glutamate-mediated excitotoxicity and neurodegeneration in Alzheimer's disease. *Int J Neurochem* 45: 583-595, 2004.
- Ghose AK, Herbert T, Hudkins RL, *et al*: Knowledge-based, central nervous system (CNS) lead selection and lead optimization for CNS drug discovery. *ACS Chem Neurosci* 3: 50-68, 2012.
- Hebert LE, Scherr PA, Bienias JL, *et al*: Alzheimer disease in the US population: prevalence estimates using the 2000 census. *Arch Neurol* 60: 1119-1122, 2003.
- Hampel H and Broich K: Enrichment of MCI and early Alzheimer's disease treatment trials using neurochemical and imaging candidate biomarkers. *J Nutr Health Aging* 13: 373-375, 2009.
- Kadir A, Andreassen N, Almkvist O, *et al*: Effect of phenserine treatment on brain functional activity and amyloid in Alzheimer's disease. *Ann Neurol* 63: 621-631, 2008.
- Roberson ED and Mucke L: 100 years and counting: prospects for defeating Alzheimer's disease. *Science* 314: 781-784, 2006.

7. Bobich JA, Zheng Q and Campbell A: Incubation of nerve endings with a physiological concentration of Abetal-42 activates CaV2.2 (N-Type)-voltage operated calcium channels and acutely increases glutamate and noradrenalin release. *J Alzheimers Dis* 6: 243-255, 2004.
8. Leist M and Nicotera P: Apoptosis, excitotoxicity and neuropathology. *Exp Cell Res* 239: 183-201, 1998.
9. Dixon RA and Ferreira D: Molecules of interest: genistein phytochemistry. *Phytochemistry* 60: 205-211, 2002.
10. Kuiper GG, Lemmen JG, Carlsson B, *et al*: Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* 139: 4252-4263, 1998.
11. Polkowski K and Mazurek AP: Biological properties of genistein. A review of in vitro and in vivo data. *Acta Pol Pharm* 57: 135-155, 2002.
12. Wang X, Chen S, Ma G, *et al*: Genistein protects dopaminergic neurons by inhibiting microglial activation. *Neuroreport* 16: 267-270, 2005.
13. Si H and Liu D: Phytochemical genistein in the regulation of vascular function: new insights. *Curr Med Chem* 14: 2581-2589, 2007.
14. Chodon D, Ramamurty N and Sakthisekaran D: Preliminary studies on induction of apoptosis by genistein on HepG2 cell line. *Toxicol In Vitro* 21: 887-891, 2007.
15. Samuels Y and Ericson K: Oncogenic P13K and its role in cancer. *Curr Opin Oncol* 18: 77-82, 2006.
16. Malinowska M, Wilkinson FL, Langford-Smith KJ, *et al*: Genistein improves neuropathology and corrects behaviour in a mouse model of neurodegenerative metabolic disease. *PLoS One* 5: e14192, 2010.
17. Liang HW, Qiu SF, Shen J, *et al*: Genistein attenuates oxidative stress and neuronal damage following transient global cerebral ischemia in rat hippocampus. *Neurosci Lett* 438: 116-120, 2008.
18. Kroom PA, Clifford MN, Crozier A, *et al*: How should we assess the effects of exposure to dietary polyphenols in vitro. *Am J Clin Nutr* 80: 15-21, 2004.
19. O'Hagan D, Schaffrath C, Cobb SL, *et al*: Biochemistry: biosynthesis of an organofluorine molecule. *Nature* 416: 279, 2002.
20. Fu XH, Wang L, Zhao H, *et al*: Synthesis of genistein derivatives and determination of their protective effects against vascular endothelial cell damages caused by hydrogen peroxide. *Bioorg Med Chem Lett* 18: 513-517, 2008.
21. Wang CY, Xia H, Tu YL, *et al*: The protective effect of 7-difluoromethyl-genistein on oxidative stress injury induced by H₂O₂ in PC12 cells. *J Hunan Normal Univ (Med Sci)* 5: 4-6, 2008.
22. Floyd RA and Hensley K: Oxidative stress in brain aging. Implications for therapeutics of neurodegenerative diseases. *Neurobiol Aging* 23: 795-807, 2002.
23. Seyfried J, Evert BO, Rundfeldt C, *et al*: Flupirtine and retigabine prevent L-glutamate toxicity in rat pheochromocytoma PC12 cells. *Eur J Pharmacol* 400: 155-166, 2000.
24. Higuchi Ya and Marsukawa S: Active oxygen-mediated chromosomal 1-2 Mbp gaint DNA fragmentation into internucleosomal DNA fragmentation in apoptosis glioma cells induced by glutamate. *Free Radic Biol Med* 24: 418-426, 1998.
25. Tyurin VA, Tyutina YY, Quinn PJ, *et al*: Glutamate-induced cytotoxicity in PC12 pheochromocytoma cells: role of oxidation of phospholipids, glutathione and protein sulfhydryls revealed by bcl-2 transfection. *Brain Res Mol Brain Res* 60: 270-281, 1998.
26. Pi R, Li W, Lee NT, *et al*: Minocycline prevents glutamate-induced apoptosis of cerebellar granule neurons by differential regulation of p38 and Akt pathways. *J Neurochem* 91: 1219-1230, 2004.
27. Parathath SR, Gravanis I and Tsirka SE: Nitric oxide synthase isoforms undertake unique roles during excitotoxicity. *Stroke* 38: 1938-1945, 2007.
28. Schreihöfer DA and Redmond L: Soy phytoestrogens are neuroprotective against stroke-like injury in vitro. *Neuroscience* 158: 602-609, 2009.
29. Jang JH and Surh YL: Potentiation of cellular antioxidant capacity by Bcl-2: implications for its antiapoptotic function. *Biochem Pharmacol* 66: 1371-1379, 2003.
30. Pike BR, Zhao X, Newcomb JK, *et al*: Stretch injury causes calpain and caspase-3 activation and necrotic and apoptotic cell death in septo-hippocampal cell cultures. *J Neurotrauma* 17: 283-298, 2000.
31. Su QJ, Chen XW, Chen ZB and Sun SG: Involvement of ERK1/2 and p38 MAPK in up-regulation of 14-3-3 protein induced by hydrogen peroxide preconditioning in PC12 cells. *Neurosci Bull* 24: 244-250, 2008.